



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



THE GIFT OF
Mrs. Benj. Dickson.

10-11-12

QK

281

647

81







CAROLI LINNÆI
SYSTEMA
PLANTARUM EUROPÆ

TOMUS III.

2011-2012

2011-2012

2011-2012

2011-2012

CAROLI LINNÆI
BOTANICORUM PRINCIPIS
SYSTEMA
PLANTARUM EUROPÆ,
EXHIBENS

*CHARACTERES naturales generum, characteres
essentiales generum & specierum, synonyma anti-
quorum, Phrases specificas recentiorum HALLERI,
SCOPOLI, &c. Descriptiones rariorum, nec-non
Floras tres novas, Lugdunæam, Delphinalem,
Lithuanicam, non omissis plantis exoticis in hortis
Europæ vulgo obviis.*

^{Jean} DURANTE (JOAN) ^{ur} EMMAN. GILIBERT, MED.
DOCT. BOTAN. PROFESS. PROVINCIAE LUGD.
PRO EPIDEMIIS PROTO-MEDICO. NOSO-
COMII GENERALIS LUGD. MEDICO. ACAD.
SCIENT. LUGD. &c.

TOMUS III.

COLONIÆ-ALLOBROGUM,
Sumptibus PIESTRE & DELAMOLLIÈRE.

1785.

L I N N Æ I
S P E C I E S P L A N T A R U M
E U R O P Æ.
P A R S. I I.

Linnæi spec. plant. Europ. Pars II.

CLASSIS XIV.

DIDYNAMIA.
GYMNOSPERMIA.

* *Calyces subquinquefidi.*

- | | |
|-----------------|--|
| 780. LEONURUS. | <i>Antheræ</i> punctis ossis ad-
spersæ. |
| 773. GLECOMA. | <i>Antherarum</i> paria cruciata. |
| 767. HYSSOPUS. | <i>Filam.</i> distantia, recta. <i>Cor.</i>
ringens. |
| 771. MENTHA. | <i>Filam.</i> distantia, recta. <i>Cor.</i>
subæqualis. |
| 770. SIDERITIS. | <i>Stigma</i> alterum vaginans alte-
rum. |
| 769. LAVENDULA. | <i>Corolla</i> resupinata. |
| 764. TEUCRIUM. | <i>Cor.</i> labium superius nullum,
sed bipartitum. |
| 763. AJUGA. | <i>Cor.</i> lab. superius staminibus
brevius. |
| 781. PHLOMIS. | <i>Cor.</i> lab. superius hirtum. |
| 776. BETONICA. | <i>Cor.</i> lab. superius planum,
adscendens, tubo cylin-
drico. <i>Stam.</i> longitudine
faucis. |
| 774. LAMIUM. | <i>Cor.</i> lab. inferius utrinque
dente setaceo. |
| 775. GALEOPSIS. | <i>Cor.</i> lab. inferius supra biden-
tatum. |
| 777. STACHYS. | <i>Cor.</i> lab. inferius lateribus
reflexum. <i>Stam.</i> deflorata
ad latera deflexa. |
| 768. NEPETA. | <i>Cor.</i> lab. inferius crenatum.
<i>Faux</i> margine reflexo. |
| 765. SATUREIA. | <i>Corolla</i> laciniis subæqualibus.
<i>Stam.</i> remota. |

Linnaei spec. plants. Europ. Pars II.

A

2 DIDYNA M I A:

- 778 **BALLOTA.** *Cal.* 10. striatus. *Cor.* labium
superius fornicatum.
779. **MARRUBIUM.** *Cal.* 10. striatus. *Cor.* lab. su-
perius rectum.
782. **MOLUCELLA.** *Cal.* campanulatus, corolla
amplior, dentibus spinosis.

* *Verbenæ species aliquot. Monarda didyma.*

* * *Calyces bilabiati.*

792. **SCUTELLARIA.** *Cal.* fructiferus opercula-
tus.
785. **THYMUS.** *Cal.* fauce villis clausus.
790. **OCYIMUM.** *Cal.* resupinata. *Filamenta*
bina basi processu.
793. **PRUNELLA.** *Filamenta* omnia apice bi-
furca.
787. **DRACOCEPHALUM.** *Corollæ* faux inflato - dila-
tata.
784. **ORIGANUM.** *Strobilus* calyces colligens.
783. **CLINOPODIUM.** *Involucrum* calyces colli-
gens.
766. **THYMBRA.** *Calyx* utrinque linea ciliata
carinatus. *Stylus* semibifidus. *Cor.* labia plana.
789. **MELITTIS.** *Cal.* tubo corolla amplior.
Corollæ lab. superius planum, integrum. *Antheræ*
cruciatae.
786. **MELISSA.** *Cal.* angulatus, scariosus,
labio superiore adscenden-
te.
788. **HORMINUM.** *Cor.* quatuor laciniae subæ-
quales: quinta major
emarginata.
795. **PRASIUM.** *Semina* baccata.

ANGIOSPERMIA.

* *Calyces bifidi.*

841. OROBANCHE. *Caps.* 1. *locularis.* *Cor.* subæqualis, 4. *fida.* *Glandula* sub basi germinis.
857. ACANTHUS. *Caps.* 2. *locularis.* *Cor.* 1. *labiata,* 3 *fida.* *Antheræ* villosæ.

*** *Calyces quadrifidi.*

801. LATHRÆA. *Caps.* 1. *locularis.* *Cor.* personata. *Glandula* sub germine.
797. BARTSIA. *Caps.* 2. *locularis.* *Cor.* personata. *Cal.* coloratior.
799. EUPHASIA. *Caps.* 2. *locularis.* *Cor.* personata. *Antheræ* inferiores hinc spinosæ.
798. RHINANTHUS. *Caps.* 2. *locularis.* *Cor.* personata. *Caps.* compressa.
800. MELAMPYRUM. *Caps.* 2. *locularis.* *Cor.* personata. *Sem.* bina, gibbosa.
824. LANTANA. *Drupa* nucleo. 2. *locul.* *Cor.* hypocraterif. *Stigma* uncinatum.

**** *Calyces quinquefidi.*

803. TOZZIA. *Caps.* 1. *locul.* *Cor.* hypocraterif. *Sem.* unicum.
837. LIMOSELLA. *Caps.* 1. *locul.* *Cor.* campanul. regular. *Sem.* plurima.
834. BROVALLIA. *Caps.* 1. *locul.* *Cor.* hypocrat. *Sem.* numerosa.
828. LINDERNIA. *Caps.* 1. *locular.* *Cor.* ringens. *Stam.* inferiora dente terminali.

814. SCROPHULARIA. *Caps.* 2. locul. *Cor.* resupinata. *Lab.* segmento intermedio interno.
815. CELSIA. *Caps.* 2. locul. *Cor.* rotata. *Filamenta* lanata.
827. CAPRARIA. *Caps.* 2. locul. *Cor.* campanulata. *Stigm.* cordatum bivalve.
816. DIGITALIS. *Caps.* 2. locularis. *Cor.* campanulata subtus ventricosa. *Stam.* declinata.
817. BIGNONIA. *Caps.* 2. locul. *Cor.* campanulata. *Sem.* alata, imbricata.
832. ERINUS. *Caps.* 2. locul. *Cor.* bilabiata: labio superiore brevissimo reflexo.
808. ANTIRRHINUM. *Caps.* 2. locul. *Cor.* personata, subtus Nectario prominente.
804. PEDICULARIS. *Caps.* 2. locul. *Cor.* personata. *Sem.* tunicata.
811. MARTYNIA. *Caps.* 5. locul. *Cor.* campan. *Rudim.* filamentis 5.
835. LINNÆA. *Bacca* 3. locularis, sicca. *Cor.* campan. *Cal.* superus.
853. VITEX. *Bacca* 4. sperma. *Cor.* ringens: lab. super. 3. fido.

DIDYNAMIA.

GYMNOSPERMIA.

763. AJUGA.

Corollæ labium superius minimum. *Stamina* labio superiore longiora.

1. *AJUGA orientalis*. A. floribus resupinatis. *Fabric. helmst.* 79. *Kniph. cent.* 3. n. 4. *Sabbat. hort.* 3. t. 100.

A. foliis ovatis, corollis obliquis. *Schreb. unilab.* p. 22.

Bugula. *Mill. dict.* n. 5.

Teucrium staminibus tubo corollæ breviorib. *Roy. lugdb.* 306.

Bugula orientalis villosa, flore inverso candido cum oris purpureis. *Tournef. cor.* 14.

β. *Bugula orientalis villosa*, flore inverso cæruleo : alba macula notato. *Tournef. cor.* 14. *Dill. elsh.* 60. t. 63. f. 61.

Habitat in Oriente. 24

2. *AJUGA pyramidalis*. A. tetragono-pyramidalis villosa, foliis radicalibus maximis. *suec. * paris. Oed. dan.* 185. *sil. Ludw. ed.* t. 8. *lithu. burg. lugd.*

Bugula foliis obtuse dentatis, caule simplici. *Mill. dict.*

Consolida media pratensis cærulea. *Bauh. pin.* 260. *non synon.*

Phyllochnois. *Renal. spec.* 125. *

Habitat in Suecia, Helvetia Germania. ♂

3. *AJUGA alpina*. A. caule simplici, foliis caulinis radicalia æquantibus. *Gunn. noru. Gmel. tub.*

Bugula foliis oblongo-ovatis, caulibus decumbentibus, verticillis distantibus. *Mill. dict.*

Bugula folio maximo, flore pallide cæruleo. *Bærh. lugdb.*

Bugula alpina maxima. *Tournef. inst.* 209.

Habitat in Alpibus helveticis, austriacis. *Jacq. Tubingæ.* 24

Differt a præcedente Foliis radicalibus vix reliquis majoribus; Floribus utrinque 10, nec 3. iisque saturate cæruleis, nec albis; Bracteis viridibus nec coloratis.

Distinctissima ab A pyramidali.

4. *AJUGA genevensis*. A. foliis tomentosis, lineatis, calycibus hirsutis. *herborn. carn. lugd. lithu. burg.*

A 3.

6 DIDYNAMIA. GYMNOSPERMIA.

A. pyramidalis. Pollich. pal.

Bugula foliis angulosis hirsutis calycibus villosis. Hall. helu. n. 283.

Teucrium (*pyramidale*) foliis angulose dentatis, caule simplici Crantz. austr. *A. genevensis* & *pyramidalem* conjungit. R.

Bugula montana. Riu. monop. t. 140. f. 2. R.

Consolida media genevensis. Bauh. hist. 3. p. 432. R.

Habitat in Europæ pratis editioribus, collibus. 24

Creditur varietas A. pyramidalis Gerardo, A. reptantis Hallero.

Ajugam pyramidalem, alpinam & genevensis, tanquam varietates jungit Schreberus sub nomine:

A. pyramidalis caulibus simplicibus erectis, foliis oblongis dentatis. Schreb. pl. vert. unil. Differunt tamen in eo inter se ut *A. pyramidalis* sint *Folia infima maxima*. *A. genevensis*. *Folia infima angustiora, floralia plerumque triloba*. *A. alpinæ*. *Folia infima angustiora, floralia integra*. Schreb. l. c. R.

3. *AJUGA reptans*. *A. stolonibus reptantibus*. Crantz. austr. p. 251. sub *Teucrio gallob. carn. pal. sil. tub. nass. Blackw.* t. 64. f. 1. Kniph. cent. 3. n. 6. Knorr. delic. 2. t. A. 8. Sabbat. hort. rom. 3. t. 99. Regn. bot. lithu. lugd. burg.

A. foliis obovatis crenatis, caule simplicissimo, stolonibus reptatricibus. Gort. gelr.

A. caule simplici erecto, stolonibus repentibus, foliis ovatis crenatis. Schreb. unilab.

Bugula flagellifera, foliis ovatis dentatis. Hall. helu. n. 282.

Teucrium foliis obverse ovatis crenatis, caule simplicissimo, stolonibus reptatricibus. Hort. cliff. Dalib. paris.

Consolida media pratensis cærulea. Bauh. pin. 260.

Bugula. Dod. pempt. 135. Riu. t. 75.

Habitat in Europa australiori. 24

764. TEUCRIUM.

Corollæ labium superius (nullum) ultra basin bipartitum, divaricatum ubi *Stamina*.

1. *TEUCRIUM campanulatum*. T. foliis multifidis, floribus lateralibus solitariis. Schreb. unilab.

T. foliis cuneiformibus trifidis laciniis divisis, floribus solitariis. Roy. lugdb.

T. calyce campanulato, laciniatum, flore parvo subcæruleo. Bærh. lugdb.

Iva moschata flore albo. Riu. monopet. t. 24.

DIDYNAMIA. GYMNOSPERMIA. 7

6. *T. supinum* perenne palustre apulum glabrum, foliis laciniatis, flore albo. *Till. pis.* 193. t. 49. f. 1. *Allion. taur.* 53.

Habitat in Orientis, Apuliæ humentibus. 24

Simillimum T. botryi, sed planta glabra. Radix perennis: Flores solitarii, Pedunculis flore brevioribus; Calyces campanulati, nec basi gibbi; Corollæ albo-margaritaceæ, folia incumbentes. Caules seniores prostrati, repentes. Folia bis trifida. Semina scabra.

2. **TEUCRIUM orientale.** *T. foliis multifidis, floribus racemosis.*

T. foliis multifidis linearibus, racemo composito, pedicellis brevibus. Schreb. unilab.

T. foliis pinnatifidis: laciniis linearibus, infimis divisis, floribus racemosis. Hort. cliff.

T. orientale angustifolium laciniatum, flore magno sub-cæruleo. Tournef. cor. 14. *Comm. rar.* 25. t. 25. *Riv. monop.* t. 19.

Habitat in Oriente.

An satis distincta a præcedenti?

3. **TEUCRIUM Botrys.** *T. foliis multifidis, floribus axillaribus ternis pedunculatis. pal. carn. Kniph. cent.* 9. n. 88. *Sabbat. hort.* 3. t. 91. *burg. lugd.*

Chamædrys foliis pinnatis, pinnis dentatis, trilobatis, verticillis paucifloris. Hall. helv. n. 289.

Teucrium foliis aliquoties trifidis, floribus verticillatis. Hort. cliff. Dalib. paris. Sauv. monsp.

Teucrium foliis multifidis, verticillis dimidiatis. Schreb. unil.

Chamædrys laciniatis foliis. Lob. hist. 209. ic. 385. *Tournef. inst.* 205. *Vaill. paris.* 33. *Mill. ic.* 264.

Iva moschata, folio multifido. Riv. t. 14.

Botrys chamædryoides. Bauh. pin. 138.

Chamæpithys altera. Dod. pempt. 46.

Habitat in Germaniæ, Galliæ, Helvetiæ, Italiæ apricis cultis. ☉

4. **TEUCRIUM Chamæpithys.** *T. foliis trifidis linearibus integerrimis, floribus sessilibus, lateralibus solitariis, caule diffuso. austr. gallob. pal. Kniph. cent.* 8. n. 92. *Blackw. t.* 528. *Ludw. ed.* t. 130. *Regn. bot. Flor. dan. t.* 733. *burg. lugd.*

T. foliis simpliciter trifidis. Hort. cliff. stamp. paris. monsp.

Ajuga (chamæpithys) foliis trifidis, floribus axillaribus solitariis folio brevioribus, caule diffuso. Schreb. unil.

8. DIDYNAMIA. GYMNOSPERMIA.

Bugula foliis imis linearibus, caulinis tripartitis. *Hall. helu.* n. 284.

Bugula Chamæpithys. *Scop. carn.*

Chamæpithys vulgaris odorata, flore luteo. *Bauh. hist.* 3. p. 295.

Chama pithys lutea vulgaris f. folio trifido. *Bauh. pin.* 249.

Chamæpithys prima. *Dod. pempt.* 46. *Riu. t.* 24.

β. *Chamæpithys vulgaris* folio trifido, flore roseo, lusitanica. *Tournef. inst.* 208. *excluso synonymo*. *Schreb. R.*

Habitat in Italiae, Galliae, Angliae, Hungariae, Helvetiae, Germaniae arvis. ☉

Calycis dentes inermes. Pedunculi non raro floribus marcescentibus elongatur. *Gouan. illustr.* 35. *R.*

Teucrium chamæpithys *Gmel. it.* 1. t. 26. f. 1. *floribus magnis longitudine fere foliorum a nostrate diversum; an Ajuga chia?* *Schreb. l. c. R.*

5. *TEUCRIUM nissolianum*. *T.* foliis trifidis quinquefidisque filiformibus, floribus pedunculatis solitariis oppositis, caule decumbente.

T. supinum annuum lusitanicum, foliis laciniatis. *Tournef. inst.* 208.

T. foliis quinquefidis trifidisque linearibus muticis, pedunculis solitariis brevissimis, caule decumbente. *Schreb. unil.*

Chamaedrys annua multiflora tenuifolia hispanica *Tournefortii*. *Morif. hist.* 3. p. 423. f. 11. t. 22. f. 19.

Habitat in Hispania, Lusitania. ☉

A Morfeno & Millero cum T. multifloro cumfunditur. *Schreb. l. c. R.*

6. *TEUCRIUM pseudo-chamæpithys*. *T.* foliis tripartitis trifidis linearibus, floribus racemosis, caule hirsuto. *Ger. prou. Gou. hort.*

T. foliis tripartitis trifidis linearibus integerrimis, floribus pedunculatis solitariis oppositis, caule patulo. *Spec. pl. Sabb. hort.* 3. t. 95.

T. foliis trifidis integrisque linearibus mucronatis, pedunculis solitariis elongatis, caule diffuso. *Schreb. unilab.*

T. supinum perenne, foliis laciniatis. *Tournef. inst.* 208.

Chamæpithys spuria, multifido folio, lamii flore. *Bauh. pin.* 250.

Pseudochamæpithys. *Clus. hist.* 2. p. 185.

Chamæpithys alia. *Cam. epit.* 680.

Habitat in Hispania, Lusitania, Marsilia. 24

Calyces aristati.

DIDYNAMIA. GYMNOSPERMIA. 9

7. TEUCRIUM *Iva*. T. foliis subtricuspidatis linearibus, floribus sessilibus lateralibus solitariis.

T. foliis lanceolatis obtusis dentatis, floribus solitariis sessilibus. *monsp.*

Ajuga (Iva) foliis linearibus acute dentatis, floribus axillaribus solitariis. *Schreb. unil.*

Anthyllis Chamæpityides minor. *Lob. hist. 208, ic. 384. bon.*

Chamæpithys moschata foliis serratis. *Bauh. pin. 429.*

Chamæpithys spuria prior. *Dod. pempt. 47.*

Anthyllis altera. *Clus. hist. 2. p. 186.*

Habitat in Lusitania, G. Narbonensi, Monspeli. ☉

9. TEUCRIUM *fruticans*. T. foliis integerrimis ellipticis subtus tomentosis, floribus lateralibus solitariis pedunculatis. *Kniph. cent. 1. n. 90. Sabbat. hort. 3. t. 96.*

T. foliis lanceolatis integerrimis subtus niveis, floribus solitariis. *Schreb. unilab.*

T. *fruticans boeticum.* *Dill. elth. t. 284. f. 366. 368. Clus. hist. 1. p. 348. Barrel. rar. 348. t. 512.*

T. foliis ovato-oblongis integerrimis petiolatis, floribus solitariis pedunculatis. *Hort. cliff.*

T. *peregrinum*, folio sinuoso. *Bauh. pin. 247.*

Habitat in Bæticæ, Siciliæ maritimis. ☿

10. TEUCRIUM *latifolium*. T. foliis integerrimis rhombeis acutis villosis: subtus tomentosis.

T. *fruticans bæticum*, ampliore folio. *Dill. elth. 379. t. 284. f. 368.*

T. *bæticum humilius & ramosus*, folio subrotundo. *Hort. angl. t. 3.*

T. *bæticum flore cœruleo.* *Riu. mon. t. 18.*

T. *hispanicum latiore folio.* *Tournef. inst. 208.*

Habitat in regione Bætica. ☿

Varietas est præcedentis. R.

12. TEUCRIUM *Marum*. T. foliis integerrimis ovatis acutis petiolatis: subtus tomentosis, floribus racemosis secundis. *Blackw. t. 47. Kniph. cent. 3. n. 93. Ludw. ed. t. 14. Sabbat. hort. 3. t. 90. Regn. bot. sub nomine Nepetæ Catarinæ.*

T. foliis ovatis utrinque acutis integerrimis, floribus solitariis utrinque digestis. *Hort. cliff.*

T. foliis omnibus parvis integerrimis, margine revolutis, spicis fere secundis, floribus solitariis pedunculatis. *Medic. Aët. Palat. Vol. 3. phys. p. 207. **

10 DIDYNAMIA. GYMNOSPERMIA.

T. foliis ovatis integerrimis subtus incanis, floribus racemosis oppositis. *Schreb. unilab.*

Chamædrys maritima incana frutescens, foliis lanceolatis. Tournef. inst. p. 205.

Marum cortusi. Bauh. hist. 3. p. 242. Riv. monop. t. 23.

β. *Chamædrys cretica saxatilis tenuifolia, foliis subtus incanis. Tournef. inst. 206.*

Habitat in regno Valentino. h

Caules erecti, fruticulosi, rigiduli. Folia petiolata, acuta. Flores secundi. Totum fragantissimum, ut natura vix simile produxerit. Bærh.

13. TEUCRIUM *multiflorum*. T. foliis ovatis: superne glabris serrato-dentatis, floribus racemosis: verticillis sexfloris.

T. foliis ovatis aut integris, aut superius bi-vel tridentatis, glabris, bracteis sessilibus magnis cochleatis, ovato-lanceolatis, integerrimis. *Medic. Aḡ. Palat.*

B. foliis ovalibus antice dentatis, floralibus integerrimis petiolatis, verticillis racemosis, caule ramosissimo. *Schreb. vert. unil.*

T. foliis ovatis utrinque acutis: superne serrato-dentatis, floribus racemosis. *Roy. lugdb.*

T. foliis ovatis utrinque acutis utrinque bidentatis, verticillis laxis. *Hort. cliff.*

T. floribus verticillatis pedunculatis senis, calycibus inferne ventricosus, foliis ovatis glabris apice serratis. *Gouan. monsp.*

T. foliis ovatis acutis denticulatis; floribus solitariis. *Zinn. gatt.*

Marum hispanicum nigrum, flore purpureo. Hall. gatt. 314.

Chamædrys multiflora tenuifolia hispanica. Tournef. inst. 205. Bærh. lugdb. I. p. 182.

Marum hispanicum nigrum. Bocc. mus. t. 117.

γ. *Chamædrys hispanica tenuifolia, latiori folio, multiflora. H. R. P. Bærh. lugdb.*

Habitat in Hispania. h

22. TEUCRIUM *hircanicum*. T. foliis cordato-oblongis obtusis, caule brachiato dichotomo, spicis longissimis terminalibus sessilibus spiralibus. *Schreb. vert. unilab.*

T. foliis cordatis crenatis rugosis, spicis florum longissimis, cylindricis bracteis linearibus. *Medic. Aḡ. Palat. Vol. 3. phys. p. 206.*

T. foliis cordatis crenatis petiolatis, spicis oblongis den-

DYDINAMIA. GYMNOSPERMIA. 11

fissimis. Hall. comm. gætt. tom. 2. p. 343. t. 13. Arduin. spec. 13. t. 4.

Habitat in Hircania. 24

Folia Botanicae officinalis. Spicae longissima, floribus undique spiraliter dense congestis, ex dichotomis ultimis caulibus.

23. TEUCRIUM Scorodonia. T. foliis cordatis serratis petiolatis, racemis lateralibus secundis, caule erecto. *carn. pal. Schreb. vert. unil. Fl. dan. t. 485. Kniph. cent. 11. n. 96. gallob. Regn. bot. burg. lugd.*

T. foliis cordatis crenatis petiolatis, spicis laxis secundis. *Hort. cliff. paris.*

T. foliis oblongis ovatis, ad basin profunde emarginatis. rugosis, parum crenatis, racemis secundis. *Medic. Aë. Pal. Vol. 3. phys. p. 206.*

Chamædrys foliis cordatis productis, spicis longissimis nudis, heteromalis. *Hall. helv. n. 287.*

Scorodonia. *Riu. mon. t. 12. Gess. ic. 85. Blackw. t. 9.*

Scordium alterum f. *Salvia sylvestris. Bauh. pin. 247.*

Habitat in arenosis editis Germaniae, Helvetiae, Galliae, Angliae, Belgii, in duris praedii Hartecampi, beati quondam mihi Paradisi. 24

Corollae albae. Filamenta purpurascens.

24. TEUCRIUM massiliense. T. foliis ovatis rugosis incisocrenatis incanis, caulibus erectis, racemis rectis secundis. *Jacq. hort. t. 94. Schreb. unilab.*

T. foliis parvis cordato-ovatis, rugosis crenatis, crenis reflexis, floribus racemosis, racemis, omnibus secundis. *Medic. aë. Pal. Vol. 3. t. 11. R.*

T. foliis ovatis crenatis, floribus racemosis staminibus corolla brevioribus. *Ger. prou. t. 11.*

T. creticum. *Clas. hist. 1. p. 348.*

Chamædrys fruticosa, insularum Stœchadum, melissae folio minore pomum redolens. *Tournef. inst. p. 205. huc pertinet secundum Schreb. Gerard. Medic. R.*

T. fruticosum incanum creticum, flore purpureo. *Barr. ic. 806.*

Chamædrys fruticosa cretica, purpureo flore. *Tournef. inst. 205.*

Habitat in Creta, Gallia australi: villa Franca. 25

Caules suffruticosi, vix pedales. Folia ovata f. cordato-ovata, rugosa inaequaliter serrata, obtusiuscula, parva, margine reflexa, subtus venosa. Racemi terminales, secundi.

14 DIDYNAMIA. GYMNOSPERMIA:

Flowers opposite pedunculati. Calyces fere Scutellariæ ut in *T. hircanico*: Labio superiore subcordato, patente, acuto. Corolla purpurascens Labio inferiore valde concavo. Odor *Nepetæ*.

Neque figura Barrelieri, neque synonymon Tournefortii a Linnaeo citata huc pertinent. Medic. Schreb. l. c. R.

25. *TEUCRIUM Scordium*. T. foliis oblongis sessilibus dentato-serratis, floribus geminis axillaribus pedunculatis, caule diffuso. *ged. carn. gallob. Schreb. unil. pal. sil. Flor. dan. t. 593. Regn. bot. lugd. burg.*

T. foliis ovato-lanceolatis serrulatis sessilibus floribus sæpius binatis. *succ. paris.*

Chamædrys foliis mollibus, hirsutis, ellipticis, crenatis, verticillis paucifloris. *Hall. helu. n. 288.*

Scordium. *Bauh. pin. 247. Cam. epit. 588. Dod. pempt. 126. Riu. t. 11. Blackw. t. 475.*

Habitat in Europæ paludosis. 24

Conf. Scordium lanuginosum C. C. 247. Varietas hirsutior, ramosior, foliis minoribus.

Scordium lanuginosum C. B. est distincta species. Schreber. l. c. R.

26. *TEUCRIUM Chamædrys*. T. foliis cuneiformi-ovatis incisfis crenatis petiolatis, floribus ternis, caulibus procumbentibus subpilosis. *ged. Regn. bot. austr. carn. pal. Kniph. cent. 11. n. 95. Sabb. hort. 2. t. 88. lugd. burg.*

Chamædrys foliis firmis, ovatis crenatis verticillis paucifloris. *Hall. helu. n. 286.*

T. foliis cuneiformi-ovatis incisfis crenatis petiolatis, floribus verticillatis ternis petiolatis. *Hort. cliff. paris. monsp.*

T. foliis oblongis basi integerrimis profunde serratis, verticillis dimidiatis, caulibus procumbentibus pilosis. *Schreb. unil.*

T. caule procumbente, per totam longitudinem florente, foliis ovato-ellipticis, pilosis inciso-crenatis, vix petiolatis. *Medic. in Ad. Pal.*

Chamædrys major repens. Bauh. pin. 248.

Chamædrys. Riu. mon. 10. Lob. hist. 260. ic. 491.

β. *Chamædrys minor repens. Bauh. pin. 248. Tournef. inst. 205. t. 98.*

Chamædrys. Fuchs. 335. ic. 498. Clus. hist. 1. p. 351. t. 17. Blackw. t. 80.

γ. *Chamædrys alpina hirsuta repens. Moris. hist. 3. p. 422. Rai. hist. 3. p. 282. Schreb. R.*

DIDYNAMIA. GYMNOSPERMIA. 13
Habitat in Germania, Helvetia, Gallia. 24

Bractea serrata. Corolla purpurea labio plano obsolete trifido.

27. TEUCRIUM lucidum. T. foliis ovatis acute inciso-ferratis glabris, floribus axillaribus ternis, caule erecto lævi.

T. foliis ovatis acute inciso-ferratis glabris, floribus axillaribus geminis, caule erecto. *Spec. pl.*

T. foliis ellipticis, obtuse crenatis sessilibus splendentibus fragilibus, foliis floralibus dentatis, verticillis sex aut quatuor floribus. *Medic. aed. pal. vol. 3. t. 12. f. 7. 8. 9.*

T. foliis ovatis inciso-ferratis glabris, verticillis dimidiatis, caulibus erectis lævibus. *Schreb. unil. 22. Ger. prov.*

Chamædrys alpina major & elatior, foliis profundius dissectis. Moris. hist. 423. Rai. hist. 3. p. 282.

Teucrium regium flore rubro. Riu. monop. t. 80. Schreb. l. c. R.

Chamædrys alpina frutescens, folio splendente. Magn. hort. 32. t. 52.

Habitat in alpinis vallis Barfilionensis Sabaudia; in monte Lachin Galloprovinciæ. Gerard. 24

Radix perennis, stolones exserens, e quibus Caules recti, glabri, nigricantes. Folia petiolata, oblonga, acute crenata, nervosa, glabra, internodiis breviora. Flores axillares, utrinque gemini, purpurei: Calyces fusi.

Differt a T. Chamædry: Caule pedali, erecto, glabro, nec palmari, diffuso, piloso. Syst. Veg. 440.

28. TEUCRIUM flavum. T. foliis cordatis obtuse serratis: bracteis integerrimis concavis, caule fruticoso, floribus racemosis ternis. *carn. Kniph. cent. 9. n. 89. Sabb. hort. 3. t. 89. burg. lagd.*

T. foliis subovatis crenatis, floribus laxè spicatis pedunculatis, caule frutescente. *Hort. upf. monsp.*

T. foliis cordatis obtuse crenatis crassis, foliis floralibus magnis integerrimis, verticillis sexfloris. *Medic. aed. Pal. vol. 3.*

T. foliis ovatis crenatis: floralibus integerrimis, verticillis dimidiatis in racemos digestis, caule fruticoso frutescente. *Schreb. unil.*

Teucrium. Bauh. pin. 247. hist. 2. p. 290. Riu. mon. t. 10.

Chamædrys fruticosior, flore ochroleuco. Moris. hist. 3. p. 421. f. 11. t. 22. f. 1.

Teucrium vulgare fruticans. I. Clus. hist. 1. p. 348.

β. Teucrium lucidum, parvo folio flore venuste purpureo.

Pluk. alm. 367. t. 65. f. 2.

24 DIDYNAMIA. GYMNOSPERMIA:

Habitat in Italia, Sicilia, Melita, Hispania, Monspeli. *H.*

Flores racemosi, utrinque terni, pedunculati. Corolla alba labio integro concavo.

Varietas *β.* pro distincta specie habetur a B. Jussieu & Schrebero.

Morisoni nomen huc non pertinere monet cl. Medicus. l.c. R.

Tanquam varietas ab ill. Schrebero huc refertur:

Chamædrys fruticosa cretica, purpureo flore. *Tournef. inst.* 205. *Barr.* 318. ic. 896.

Teucrium atticum molli lanuginé incanum. *Pluk. phyt. t.* 65. f. 2. R.

29. *TEUCRIUM montanum.* T. corymbo terminali, foliis lanceolatis integerrimis subtus tomentosis. *car. burg. lugd.*

T. foliis lanceolatis integerrimis petiolatis, spicis laxis subrotundis. *Hort. cliff. paris. monsp.*

T. corymbis terminalibus glabris, foliis petiolatis ellipticis margine revolutis subtus tomentosis, caule procumbente. *Schreb. unil. cum Teucr. supino conjungit.*

Polium (montanum) foliis lanceolatis integerrimis, caulibus procumbentibus, floribus corymbosis terminalibus. *Mill. dict.*

Chamædrys foliis ellipticis integerrimis, contractis, subtus incanis, floribus capitatis. *Hall. hesl. n.* 285.

Polium VIII. Clus. hist. 363. R.

Polium lavendulæ folio. *Bauh. pin.* 220.

Polium majus. *Cord. hist.* 124. b.

Ajuga folio integro. *Riu. mon.* 15.

Habitat in siccis Germaniæ, Monspeli, Genevæ, Helvetiæ, Parisiis. 24

Flores albi. Folia lanceolata, remota.

30. *TEUCRIUM supinum.* T. corymbo terminali, foliis linearibus margine revolutis.

Polium montanum repens. *Bauh. pin.* 220.

Polium montanum supinum minimum, *Lob. ic.* 488.

Habitat in agro Viennensi.

Flores albi. Folia linearia, integerrima, acuta, conferta.

Præcedenti affinis.

31. *TEUCRIUM pyrenaicum.* T. corymbo terminali, foliis cuneiformi-orbiculatis crenatis.

T. foliis cuneiformibus crenatis, corymbo terminali, foliolis floralibus linearibus. *Roy. lugdb.*

T. foliis suborbiculatis crenatis, spica laxa orbiculata depressa. *Hort. cliff.*

DIDYNAMIA. GYMNOSPERMIA. 15

T. corymbis terminalibus, foliis cuneiformi-orbiculatis crenatis pilosis. *Schreb. unil.*

Polium (*pyrenaicum*) caulibus procumbentibus hirsutissimis, foliis cuneiformi-orbiculatis crenatis. *Mill. dict.*

Polium pyrenaicum supinum, hederæ terrestris folio. *Tournef. inst.* 206.

Polium saxatile purpureum chamædryoides, ampla coma. *Bocc. mus.* 2. t. 61. *Barr. rar.* 315. t. 1086. *folia male.*

Habitat in Pyrenæis. 24

32. TEUCRIUM Polium. Teucrium capitulis subrotundis, foliis oblongis obtusis crenatis tomentosis sessilibus, caule prostrato. *Hort. cliff. monsp. carn. delph.*

α. Polium montanum. luteum *Bauh. pin.* 220. *Tour. inst.* 206. *Rai. hist.* 1. p. 525. *Barrel. obs.* 320. ic. 1082. *Moris. hist.* 3. p. 455. f. 11. t. 2. f. 1. *Blackw. t.* 456.

Polium (luteum) spicis longis, foliis obtusis crenatis tomentosis. *Mill. dict.*

Teucrium (aureum) corymbis terminalibus pilosis, foliis ovatis serratis, caulibusque dense lanatis, summitatibus luteis nitidis. *Schreb. unil. distincta species.* R.

β. Polium montanum album. *Bauh. pin.* 221.

Polium (album) caule ramoso procumbente foliis lineari-lanceolatis dentatis, floribus corymbosis terminalibus. *Mill. dict.*

Teucrium (Teuthrion) capitulis subrotundis breviter pedunculatis, foliis lanceolatis crenatis tomentosis incanis, caule decumbente. *Schreb. unil.*

Polium montanum 1. *Clus. hist.* 1. p. 361. fig. mala. *Lob. hist.* 257. ic. 486. fig. ead. *Schreb. R.*

Polium montanum album ferratifolium latifolium majus. *Barrel. obs.* 326. ic. 1074. & minus ejusd. *obs.* 329. f. 1078. *Schreb. R.*

γ. Polium montanum supinum alterum. *Bauh. pin.* 221.

δ. Polium maritimum supinum venetum. *Bauh. pin.* 221. *hist.* 3. p. 300. ic. mala. *Moris. hist.* 3. f. 11. t. 2. f. 12. *Tournef. inst.* 206.

T. (Achaemenis) corymbis densis, foliis inverse lanceolatis tomentosis incanis floralibus longe longioribus. *Schreb. unil. distincta species.* R.

Polium montanum 5. *Clus. hist.* 1. p. 362.

ε. **Hyssopium** dioscoridis. *Col. ecphr.* 1. p. 59. t. 67.

T. (Pseudhyssopus) capitulis rotundatis laxis, foliis oblongis antice crenatis tomentosis incanis, caule lanato corymbifero. *Schreb. unil. distincta species.*

26 DIDYNAMIA. GYMNOSPERMIA:

Polium montanum incanum &c. *Mich. hort. flor. p. 75.*
n. 2. *Schr. R.*

Habitat in Italia, Hispania, Lusitania, Narbona, Libano. 24

Polii species, a varietatibus distinctas, hodie confusas, inquirant solidi Botanici in solo natali.

33. *TEUCRIUM capitatum*. T. capitulis pedunculatis, foliis lanceolatis crenatis tomentosis, caule erecto.

T. (*Belion*) capitulis subrotundis breviter pedunculatis, foliis lanceolatis crenatis tomentosis niveis, caule erecto.
Schreb. unil.

Polium dasphyllum gnaphalodes flore albo. *Barr. rar.*
324. ic. 1047. *Schreb. R.*

Polium montanum album serratum latifolium. *Barr. rar.*
325. t. 1079. *sec. Linné, sed ad T. Teuthrion refert.*
Schreber. R.

β. *Polium maritimum erectum monspeliacum.* *Bauh. pin. 221.*

Polium (capitatum) caule erecto diffuso, foliis lineari-lanceolatis crenatis, corymbis terminalibus lateralibusque.
Milk. dict.

Polium monspeliacum. *Bauh. hist. 3. p. 699.*

Habitat in Hispania, Gallia, Sibiria. 24

Habitus Origani Majoranae, sed tomentosum & angustifolium.

Varietas β. ad T. Polium refert Gerard.

34. *TEUCRIUM pumilum*. T. capitulis terminalibus, sessilibus, foliis linearibus planis quadrifariam confertis, caule procumbente tomentoso. *Amæn. acad. 4.* *
Schreb. unil.

Polium hispanicum montanum pumilum, rosmarini folio, flore rubro. *Tournef. inst. 207.*

Polium montanum pumilum rubrum, viridi stœchadis folio, caule tomentoso. *Barr. ic. 192.*

γ. *Polium montanum pumilum, angusto viridique folio, caule incano.* *Barr. rar. 357. ic. 1093. Schreb. R.*

Habitat in Hispania.

Planta Tournefortii nova species est & ill. Schrebero audit:

Teucrium (Libanotis) spicis subrotundis, foliis confertis linearibus revolutis integerrimis cauleque tomentosis.
Schreb. unil 48. n. 52. R.

35. *TEUCRIUM spinosum*. T. spinosum, calycibus labio superiore ovato, corollis relupinatis, pedunculis geminis. *Lafl. it.*
T.

DIDYNAMIA. GYMNOSPERMIA. 17

T. ramulis spinulentibus, corollis resupinatis, pedunculis axillaribus geminis. *Schreb. unil.*

Chamædrys multifida spinosa odorata. Tournef. inst. 208.

Chamædrys spinosa. Bauh. pin. 248. prodr. 117. f. 117. Moris. hist. 3. p. 423. f. 11. t. 22. f. 17.

β. *Teucrium mucronatum spec. pl.*

Scordium spinosum odoratum. Corn. can. 123. Barr. rar. 319. t. 20.

Habitat in Lusitaniæ collibus, arvis. ☉

Corollæ resupinatae, albæ. Stamina brevissima.

765. SATUREIA.

Corollæ laciniae subæquales. Stamina distantia.

1. *SATUREIA juliana. S. verticillis fastigiatis, foliis linearilanceolatis.*

Satureia floribus verticillatis, foliis lanceolatis glabris Roy. lugdb.

Satureia spicata. Bauh. pin. 218.

Satureia perennis, verticillis spicatis & densius dispositis. Moris. hist. 3. p. 412. f. 11. t. 17. f. 4.

Habitat in Hetruria, in Thyrrheni maris asperis, Florentiæ. 24

2. *SATUREIA Thymbra. S. verticillis subrotundis hispidis, foliis oblongis acutis. Blackw. t. 318. Sabb. hort. 3. t. 71. S. cretica. Bauh. pin. 218.*

Thymus frutescens, verticillis fere nudis globosis, foliis ovato-lanceolatis. Hort. cli. f. 306.

Thymbra legitima. Clus. hist. 3. p. 358.

Thymum creticum pone verticillatum, Barr. rar. 279. t. 898.

Habitat in Creta, Tripoli.

Verticilli in hac 4. f. 5. in præcedenti 9. f. 20.

4. *SATUREIA montana. S. pedunculis lateralibus solitariis, floribus fasciculatis fastigiatis, foliis mucronatis linearilanceolatis. Sabb. hort. 3. t. 64. delph.*

S. racemis lateralibus solitariis: foliis linearilanceolatis punctatis duris. Scop. carn. t. 30. R.

S. pedunculis lateralibus solitariis trifloris, foliis carinatis punctatis ciliatis mucronatis. Zinn. gætt.

S. pedunculis dichotomis, verticillis concatenatis. Hort. upf. monsp.

S. montana. Bauh. pin. 218.

S. perennis. Riv. mon. 41.

Melissa foliis linearibus integerrimis. Hort. cliff. 308.

Linnaei spec. plant. Europ. Pars II.

B

19 DIDYNAMIA. GYMNOSPERMIA!

Saxifraga secunda. *Cam. epit.* 717. R.

Habitat in Hetruria, Narbona. 24

5. *SATUREIA hortensis*. S. pedunculis bifloris. *monsp.*
Ludv. ed. t. 199. *Kniph. cent.* 3. n. 80. *Sabb. hort.* 3. t.
70. *Regn. bot.*

S. *hortensis*. *Bauh. pin.* 218. *Blackw. t.* 419.

S. *altera*. *Cam. epit.* 487.

Thymus erectus annuus, foliis lanceolato-linearibus. *Hort.*
cliff. 306.

Habitat in G. Narbonensi & Italia, circa Herborn *quasi*
indigena. *Leers flor. herb. n.* 463. ☉

6. *SATUREIA capitata*. S. floribus spicatis, foliis carinatis
punctatis ciliatis.

Thymus capitatus qui *dioscoridis*. *Bauh. pin.* 219.

Thymum legitimum. *Clus. hist.* 1. p. 357.

Thymum creticum incanum capitatum. *Barr. ic.* 897.

Habitat in Creta, Bætica, Hispali, Græcia, Palestina. 24

766. THYMBRA.

Cal. subcylindricus, bilabiatus, utrinque linea
villosa exaratus. *Stylus* semibifidus

1. *THYMBRA verticillata*. T. floribus verticillatis.

Hyssopus angustifolia montana aspera. *Bauh. pin.* 218.

Hyssopus montana. *Dalech. hist.* 394.

Habitat in Europa australi. ☿

Caules *vix pedales*. Folia *opposita*, *lineari-lanceolata*,
integra, *punctata*: *Floralia paulo latiora*, *acutiora*, *sed*
ciliata. *Hæc omnino verticillata*, *floribus sessilibus*; *Præ-*
cedens obsolete spicata, *spica oblonga bracteis s. foliis ciliatis*
imbricata. *Forte prioris sola varietas*.

767. HYSOPOS.

Cor. labium inferius lacinula intermedia crenata.

Stam. recta, distantia.

1. *HYSSOPUS officinalis*. H. spicis secundis, foliis lanceo-
latis. *Gouan. hort.* *Jacq. austr. t.* 254. *Blackw. t.* 296.
Kniph. cent. 6. n. 47. *Ludw. ed. t.* 40. *Sabb. hort.* 3.
3. t. 79. *Regn. bot.*

H. foliis linearibus punctatis, verticillis in spicam conti-
nuatis. *Hall. helv. n.* 249. *Riu. t.* 68.

H. *officinarum cærulea s. spicata*. *Bauh. pin.* 217.

H. *vulgaris*. *Dod. pempt.* 287.

DIDYNAMIA GYMNOSPERMIA. 19

Hyssopus rubro flore. Bauh. pin. 217. Mill. dict. n. 2.
Habitat in Valle angusta, Austria, Sibiria. 24

768. NEPETA.

Corollæ labium inferius lacinula intermedia crenata, faux margine reflexo. Stamina approximata.

1. *NEPETA Cataria. N. floribus spicatis, verticillis subpedicellatis, foliis petiolatis cordatis dentato-serratis. ged. carn. gallob. pal. sil. Kniph. cent. 4. n. 53. Flor. dan. t. 580. Blackw. t. 455. Sabb. hort. 3. t. 80. Regn. bot. sub titulo Teucris Mari. lithu. lugd. burg.*

N. floribus interrupte spicatis pedunculatis. Fl. suec. paris.

Cataria foliis cordatis, verticillis spicatis. Hall. helu. n. 246.

Nepeta foliis cordatis petiolatis, verticillis spicatis subpedicellatis: corollarum labio superiore papillonaceo. Crantz. austr.

Mentha cataria vulgaris & major. Bauh. pin. 228.

Cataria herba. Dod. pempt. 99.

β. Mentha Cataria minor. Bauh. pin. 228.

Nepeta (minor) floribus spicatis, spicis interruptis, verticillis pedicellatis, foliis subcordatis serratis petiolatis. Mill. dict.

Habitat in Europa. 24

Varietas β. dimidio minor est, non ramosa.

2. *NEPETA pannonica. N. floribus paniculatis, foliis cordatis petiolatis obsolete crenatis. Jacq. vind. t. 24. austr. t. 129.*

N. caule paniculato, pedunculis partialibus ramosis multifloris. Roy. lugdb.

N. (paniculata) foliis cordatis petiolatis. Crantz. austr.

N. pannonica major & elatior. Moris. hist. 3. p. 415. f. II. t. 6. f. 6.

Mentha montana verticillata. Bauh. pin. 227.

Menthastrum montanum 1. Clus. hisp. 2. p. 32.

Habitat in Austria, Pannonia, Sibiria. 24

3. *NEPETA violacea. N. verticillis pedunculatis corymbosis, foliis petiolatis cordato-obl. dentatis. Scop. carn. delph. N. foliis ovatis petiolatis glabris verticillis diffuse spicatis glabris. Sauv. monsp.*

N. montana purpurea major, sparsa spica. Barr. ic. 601. Bocc. mus. 2. p. 46. t. 36.

Cataria hispanica, betonica folio angustiore, flore cæruleo. Tournes. inst. 202.

Habitat in Hispania, Sibiria, Carniolia. 24

10 DIDYNAMIA. GYMNOSPERMIA.

5. **NEPETA** *Nepetella* N. floribus racemosis : racemulis quinquefloris , foliis cordato-lanceolatis dentatis tomentosis. *delph.*

Habitat in Europa australi.

Planta triplo minor *Cataria*. Folia *angustissima* , acute sinuato-dentata. Flores *laxe ramosi* : racemulis *sub-quinquefloris* , nec ut in *N. violacea* floribus 30. Calyces nudi Corollæ *recentes rubræ* , nec *albide aut cæruleæ*. Antheræ *cæruleæ*. Bractææ *angustissimæ* , *patentes*.

6. **NEPETA** *nuda*. N. florum racemis verticillatis nudis , foliis cordato-oblongis sessilibus serratis. *Jacq. austr. t. 24. delph.*

Cataria foliis imis cordatis obtusis , superioribus ovatis , verticillis sessilibus , densissimis , spicatis. *Hall. helv. n. 248.*

Nepeta foliis sessilibus nudis , calycibus fructus ovatis æqualibus. *Hort. upf. **

Cataria hispanica *Betonica* folio angustiori. *Tourn. inst. R.*

Mentha Cataria hispanica , olim *Sideritis altissima*. *Bauh. hist. 3. p. 226. quoad descriptionem.*

Mentha cataria hispanica , angustioribus foliis. *Bauh. hist. 3. p. 236. quoad figuram.*

Mentha montana purpurea major , spica sparsa. *Barr. ic. 601. sed foliæ paulo breviora.*

Habitat in Helvetia , Hispania. 2x

Caules *bipedales* , *glabri* , *stricti* , *quadriusculi* : *seniores atropurpureascentes*. Folia (*stachydis palustris*) *sessilia* , *cordata* , *oblonga* , *serrata* , *obtusa* , *venosa* , *nuda* , *utrinque scabra*. Racemi *brachiati* , Bracteis *linearibus* , Floribus *distinctis* Corolla *albido-rufescens* , *palati barba albida* , *fauce punctis purpureis*.

Flores *cærulei & albi*. *Hall. R.*

7. **NEPETA** *hirsuta*. N. floribus sessilibus verticillato - spicatis , verticillis tomento obvolutis.

Hormium spicatum , *lavendulæ flore & odore*. *Bocc. fic. 48. t. 25. f. 2.*

Habitat in Sicilia.

8. **NEPETA** *italica*. N. floribus sessilibus verticillato-spicatis , bracteis lanceolatis longitudine calycis , foliis petiolatis. *Jacq. hort. t. 112.*

Mentha Cataria minor alpina. *Bauh. pin. 228. prodr. 110.*

Habitat in Italia. D. Rathgeb. 2x

Radix *perennis* Caulis *pedalis*. Folia *cordata* , *obtusa* , *crenata* , *petiolata* , Spicæ *terminates* , *verticillatæ* : *verti-*

DIDYNAMIA. GYMNOSPERMIA. 21

ovilli remoti, sessiles, obvallati Bracteis lanceolatis, viridibus, lateribus albidioribus, longitudine calycum, acuminatis. Calyces erecti, aequales. Corollæ albæ.

9. *NEPETA tuberosa*. N. spicis sessilibus terminalibus, bracteis ovatis coloratis, foliis summis sessilibus. *Kniph. cent. 9. t. 71.*

Mentha tuberosa radice. Bauh. pin. 217.

β. *Nepeta tuberosa spicata hispanica. Bocc. mus. 2. p. 44. t. 36. Barr. ic. 1131. Sabb. hort. 3. t. 81.*

Habitat in Hispania, Lusitania. 24

Caules simplices. Folia cordato-oblonga, rugosa, obtusiuscula, grosse serrata, pubescentia. Spica terminalis, longa, subinterrupta. Bracteis numerosis, erectis, ovatis, acutis, nervosis, rubro, coloratis, margine pubescentibus, calyces obtegentibus, Flores violacei β. procumbens est.

769. LAVANDULA.

Cal. ovatus, subdentatus, bractea suffultus. Corolla resupinnata. Stamina intra tubum.

1. *LAVANDULA Spica*. L. foliis lanceolatis integerrimis, spicis nudis. *monsp. Sabb. hort. 3. t. 72. Kniph. cent. 4. n. 39. Regn. bot. delph.*

L. foliis linearibus, spicis nudis. *Hall. helv. n. 232.*

α. L. angustifolia. *Bauh. pin. 216.*

L. altera. *Dod. pempt. 273. Blackw. t. 294.*

β. L. latifolia. *Bauh. pin. 116. Blackw. t. 296.*

Spik oder gemeiner Lavendel. Linn. Pflanzen syst. 4. p. 41.

Habitat in Europa australi. 5

Varietas α. Fruticans foliis linearibus. β. Annua foliis lanceolatis.

2. *LAVANDULA multifida*. L. foliis duplicato-pinnatifidis. *Burm. ind. t. 38. f. 1. Kniph. cent. 5. n. 38.*

L. folio dissecto. *Bauh. pin. 216.*

L. multifido folio. *Clus. hist. 1. p. 345.*

β. *Lavandula folio longiore tenuius & elegantius dissecto T. Comm. rar. 27. t. 27.*

L. (*canariensis*) foliis duplicato-pinnatifidis hirsutis, spicis fasciculatis. *Mill. dict.*

γ. *Lavandula canariensis maritima, spica multiplici cærulea Pluk. alm. 209. t. 303. f. 5.*

Vielspaltiger Lavendel. Linn. Pflanzen syst. 4. p. 43.

Habitat in regione Bætica. ♂

22. DIDYNAMIA. GYMNOSPERMIA.

3. *LAVANDULA dentata*. L. foliis pinnato dentatis : spicis ramosis. *Kniph. cent.* 2. n. 38

Stœchas folio serrato. *Bauh. pin.* 216.

Stœchas crispo folio. *Clus. hist.* 1. p. 346.

Gezæhnter Lavendel. *Linn. Pflanzensyst.* 1. p. 43.

Habitat in Hispaniæ Calape ; Oriente. ☿

4. *LAVANDULA Stœchas*. L. foliis lanceolato-linearibus , integerrimis , spicis comosis. *monsp. Blackw.* 1. 241.

Kniph. cent. 3. n. 52. *Regn. bot.*

Stœchas (officinarum) foliis lanceolato-linearibus , pedunculis brevioribus. *Mill. dict.*

Stœchas purpurea. *Bauh. pin.* 216.

Stœchas brevioribus ligulis. *Clus. hist.* 1. p. 344.

ß. *Stœchas cauliculis non foliatis*. *Bauh. pin.* 216.

Stœchas (pedunculata) foliis lanceolato-linearibus , pedunculis longissimis. *Mill. dict.* n. 2. R.

Stœchas longioribus ligulis. *Clus. hist.* 1. p. 344.

Stœchaskraut. *Lin. Pflanzensyst.* 4. p. 44.

Habitat in Europa australi. ☿

770. SIDERITIS.

Stamina intra tubum corollæ. Stigma brevius involvens alterum.

* *Ebracteata.*

5. *SIDERITIS montana*. S. herbacea ebracteata , calycibus corolla majoribus spinosis : labio superiore trifido. *Kniph. cent.* 7. n. 87.

S. herbacea ex decumbenti erecta , calycibus corolla fere majoribus spinosis : labio superiore trifido. *Crantz. austr.*

Cunila calycum labio spinoso : superiore trifido , inferiore bipartito. *Hort. ups.*

Sideritis montana parvo varioque flore. *Bauh. pin.* 233.

Sideritis montana : parvo flore nigro purpureo , capite medio croceo. *Col. ecphr.* 1. p. 198. t. 196.

Habitat in Italia. ☉

Corollæ flavæ margine ferrugineo , calyce minores. Verticillis sexfloris absque bracteis.

In saxosis erecta , in pratis procumbens.

6. *SIDERITIS romana*. S. herbacea decumbens ebracteata , calycibus spinosis : labio superiore ovato.

Cunila calycum lacinia superiore latiori ovata trinervi. *Hort. cliff.*

Sideritis genus verticillis spinosis. *Bauh. hist.* 3. p. 428.

DIDYNAMIA. GYMNOSPERMIA. 23

Sideritis, verticillis spinosis, minor procumbens. *Morif. hist.*

3. p. 388. f. 11. t. 12. f. 5.

Habitat in agris Europæ australis. ♂

Calycis lobus 5tus. f. superior ovatus reliquis duplo latior. Corollæ albæ *Marubii*, Labio superiore tantum emarginato. Verticilli sexflori absque bracteis. Caules depressi, rigiduli.

* *Bracteata*: bracteis dentatis.

7. *SIDERITIS incana*. S. suffruticosa tomentosa, foliis lanceolato-linearibus integerrimis, floribus bracteisque dentatis.

S. hispanica erecta, folio angustiore. *Tournef. inst.* 191.

Hyssopus montana verticillata major. *Barr. ic.* 239. *Bocc. mus.* 2. p. 77. t. 67. f. 2.

Habitat in Hispania, Læfling. Alstræmer. ♀

Habitus *Lavandulæ*, sed tomentoso-incanus. Truncus suffruticosus, basi plerumque decumbens, e quo Caules s. rami multi, tenuissime tomentosi albi, erecti, elongati, pedales, tetragoni. Folia minus explicata, integerrima, lineata, tomentosa, alba: inferiora lineari-lanceolata, supra nuda; superiora linearia, remotiora; summa versus verticillos interdum dentata. Bractea cordatæ, dentatæ, subpinosæ. Verticilli sexflori. Calyces quinque-dentati, incani. Corollæ flavæ: labio superiore erecto, lineari, sæpe bifido; inferiore trifido: intermedia quadruplo latiore.

8. *SIDERITIS hyssopifolia*. S. foliis lanceolatis glabris integerrimis, bracteis cordatis dentato-spinosis, calycibus æqualibus. *delph.*

Betonica foliis ellipticis, obiter crenatis, verticillis spicatis, bracteatum dentibus aristatis. *Hall. helu. n.* 260.

Sideritis foliis lanceolato-linearibus sessilibus glabris. *Hort. cliff.*

S. alpina hyssopifolia. *Bauh. pin.* 233.

Sideritis VII. Clus. hist. 2. p. 41.

Habitat in Hetruria, Pyrenæis, Thuri. 24

Similis sequenti, caule magis tomentoso. *Hall.*

9. *SIDERITIS scordioides*. S. foliis lanceolatis subdentatis supra glabris, bracteis ovatis dentato-spinosis, calycibus æqualibus.

Betonica foliis ellipticis, acute dentatis, bracteis serratis aristatis, verticillis spicatis. *Hall. helu. n.* 261.

Sideritis montana scordioides tomentosa. *Barrel. ic.* 343. *Hall. R.*

S. foliis hirsutis profunde crenatis. *Bauh. pin.* 233.

14 DIDYNAMYA. GYMNOSPERMIA:

Sideritis f. *ferruminatrix heraclea*. *Lob. is.* 528.

Habitat Monspelii. *D. Sauvages*; in Helvetia, &c.

Caules supra terram parum perennantes, adscendentes, simplices. Differt a S. hirsuta Caulibus erectioribus: Folia supra sæpe glabris, Corollis etiam labio superiore luteis; Spicis ovatis, non interruptis; Bracteis ovatis, dentato-ciliatis.

10. *SIDERITIS hirsuta*. *S. foliis lanceolatis obtusis dentatis pilosis, bracteis dentato-spinosis, caulibus hirsutis decumbentibus. burg. lugd.*

S. foliis lanceolatis obtusis dentatis pilosis, bracteis dentato-spinosis, caule hirsuto, spicis interruptis elongatis. Spec. pl. Kniph. cent. 4. n. 81. Sabb. hort. 3. t. 58.

Tetrahit caulibus procumbentibus, foliis sessilibus. Ger. prou.

Sideritis hirsuta procumbens. Bauh. pin. 432.

Sideritis III. Clus. hist. 2. p. 40.

Habitat in G. Narbonensi, Hispania, Italia.

Caulis rigidus, pilosus. Folia lanceolata, obtusiuscula, rugoso-plicata, ferraturis 3. f. 4. subacutis, Bractæ cordatæ ferrato-spinosæ. Verticilli remotissimi, sessiles, sexflori, caule inter verticillos hirsutior. Corollæ flavae, tubo incurvo; Labio superiore lineari, semibifido, reflexo, erecto, niveo; L. inferiore minimo. Calyces hirti, spinis 3. aequalibus. Stamina intra tubum.

771. M E N T H A.

Cor. subzqualis, 4. fida: lacinia latiore emarginata.

Stamina erecta distantia.

* *Spicatae.*

1. *MENTHA sylvestris*. *M. spicis oblongis, foliis oblongis tomentosis ferratis sessilibus, staminibus corolla longioribus. paris. gelr. angl. Oed. dan. t. 484. pal. sil. Kniph. cent. 9. n. 67. lugd. delph. lithu. burg.*

M. foliis elliptico-lanceolatis, ferratis, subtus tomentosis, spicis cylindricis, staminibus flore duplo longioribus. Hall. helv. n. 227.

M. candicans. Crantz. austr.

N. sylvestris, folio longiore. Bauh. pin. 227.

Menthastrum spicatum, folio longiore candicante. Bauh. hist. 3. p. 321. Blackw. t. 292. Riv. t. 31.

Habitat in Dania, Germania, Anglia, Gallia, Sibiria, &c. *Folia albida.*

3. *MENTHA viridis*. *M. spicis oblongis, foliis lanceolatis*

DIDYNAMIA. GYMNOSPERMIA. 25

¶ nudis ferratis sessilibus, staminibus corolla longioribus.

Scholl. barb. Lærh. nass. Regn. bot. burg. lugd.

M. spicis solitariis interruptis, foliis lanceolatis ferratis sessilibus. Hort. cliff. Hudf. angl.

M. foliis elliptico-lanceolatis glabris, spicis cylindricis. Hall. helu. n. 229.

M. spicata. Crantz. austr.

M. spicata, folio longo acuto glabro nigriori. Bauh. hist. 3. p. 220. Hall. R.

M. angustifolia spicata. Bauh. pin. 227.

Mentha. III, IV. Dod. pempt. 95.

Mentha aquatica spicata, foliis oblongis viridibus ferratis acuminatis. Gron. virg. 167.

Habitat in Germania, Anglia, Gallia, Helvetia. 24

Affinis minimum M. sylvestri, sed minor & glabra.

4. *MENTHA rotundifolia. M. spicis oblongis, foliis subrotundis rugosis crenatis sessilibus. Hudf. angl. ged. pal. burg.*

M. foliis ovatis, rugosis, spicis cylindricis, Hall. helu. n. 226.

M. spicis confertis, foliis ovatis rugosis sessilibus. Mill. dict.

M. sylvestris, rotundiore folio. Bauh. pin. 227.

Menthastrum folio rugoso rotundiore, spontaneum, flore spicato odore gravi. Bauh. hist. 3. p. 219.

Habitat in Angliæ aquosis. 24

* * *Capitata.*

5. *MENTHA crispa. M. spicis capitatis, foliis cordatis dentatis undulatis sessilibus, staminibus corollam æquantibus. Kniph. cent. 11. n. 75. delph.*

M. foliis rugosis, brevissime petiolatis, verticillis spicatis. Hall. helu. n. 230.

M. crispa rotundifolia spicata. Bauh. hist. p. 217. R.

M. crispa danica s. germanica speciosa. Moris. hist. 3. p. 367. f. 11. t. 6. f. 5.

M. crispa. Riv. mon. t. 50.

Habitat in Sibiria, Helvetia, Hircynia. Weber. spicil. p. 17. 24

6. *MENTHA hirsuta. M. floribus capitatis, foliis ovatis ferratis subsessilibus pubescentibus, staminibus corolla longioribus. Hudf. angl. fl. herborn.*

M. aquatica hirsuta s. Sisymbrium hirsutius. Bauh. hist. 3. p. 224. Moris. hist. 3. p. 370.

Sisymbrium hirsutum. Rai. angl. 3. p. 233.

¶ *M. Sisymbrium hirsuta, glomeratis & foliis rotundioribus. Rai. angl. 3. p. 233. t. 10. f. 1.*

66 DIDYNAMIA. GYMNOSPERMIA:

Habitat in aquis Angliæ; in Hollandia. Osbekh. Germania.

Differt a M. aquatica; Foliis magis sessilibus hirsutis. convenit staminibus corolla longioribus, & tota facie. Pistillum corolla longius. Corolla profunde 4. fida, jere regularis. Calyces ciliati.

M. aquaticæ varietatem facit Haller. R.

7. *MENTHA aquatica. M. floribus capitatis, foliis ovatis serratis petiolatis, staminibus corolla longioribus. Fl. suec. parif. austr. pal. gallob. Fl. dan. t. 673. fil. Blackw. t. 32. Kniph. cent. 11. n. 74. lithu. burg. lugd.*

M. foliis ovato-lanceolatis, serratis, verticillis paucissimis, terminante maximo. Hall. helv. n. 225.

M. floralibus lateralibus, glomeratis, foliis ovatis, serratis; staminibus corolla longioribus. Scop. carn.

M. palustris. Mill. dict. n. 11.

M. palustris spicata, Riv. mon. t. 49. R.

M. rotundifolia palustris s. aquatica major. Bauh. pin. 227.

Symbrium sylvestre. Dalech. hist. 677.

Habitat in Europa ad aquas. 2.

Planta non hirta. Verticilli terminales in capitulum s. spicam abusam conferti; Stamina corolla longiora.

Datur varietas, ubi stamina corolla breviora sunt. Pollich. l. c. R.

8. *MENTHA piperita. M. spicis capitatis, foliis ovatis serratis petiolatis, staminibus corolla brevioribus. Regn. bot.*

M. spicis brevioribus & habitioribus, foliis menthæ fuscæ; sapore fervido piperis. Rai. angl. 3. p. 234. t. 10. f. 2.

Habitat in Anglia. 2.

*** *Verticillata.*

9. *MENTHA sativa. M. floribus verticillatis, foliis ovatis acutiusculis serratis, staminibus corolla longioribus.*

M. crispa verticillata. Bauh. pin. 217.

M. crispa verticillata, folio rotundiore. Bauh. hist. 3. p. 415.

Habitat in Europa australi. 2.

10. *MENTHA gentilis. M. floribus verticillatis, foliis ovatis acutis serratis, staminibus corolla brevioribus. austr.*

Scholl. barb. Sabb. hort. 3. t. 50. Flor. dan. t. 736.

M. hortensis verticillata, ocyimi odore. Bauh. pin. 217.

Mentha verticillata minor acuta non crispa, odore ocyimi. Bauh. hist. 3. p. 216.

Mentha quarta. Fuchs. hist. 291.

Habitat in Europa australiore. 2.

Simillima M. arvensi, at minus hirta, Caulibus rubris;

DIDYNAMIA. GYMNOSPERMIA. 27

Stamina Tubo corollæ, non longiora: Verticilli laterales. Corollæ infima lacinia acuta. Calyces adpersi punctis resinosis.

II. MENTHA arvensis. *Mentha floribus verticillatis, foliis ovatis acutis serratis, staminibus corollam æquantibus. suec. parif. carn. pal. sil. Flor. dan. t. 512. lugd. lithu. burg. M. floribus verticillatis, foliis hirsutis; ovato-lanceolatis, verticillis tomentosis. Hall. helu. n. 223. R.*

Mentha arvensis verticillata procumbens. Moris. hist. 3. p. 369. f. 11. t. 9. f. 5.

Calamintha arvensis verticillata. Bauh. pin. 229.

Habitat in Europæ agris, frequens post messem. 24

Caules diffusi; Verticilli laterales; Planta tota hirta; Stamina corollæ limbo longitudine æqualia. Corollæ lacinia infima obtusa. Calyces canescentes.

Stamina persæpe quoque intra tubum latent, ut introspectanti appareant, quam M. gentilem L. esse putaverim, nisi Barbæ media lacinia obtusa fuisset. Pollich. l. c. R.

12. MENTHA exigua. *M. floribus verticillatis, foliis lanceolato-ovatis glabris acutis integerrimis. gallob. burg. lugd.*

M. floribus verticillatis, foliis glabris, ovato-lanceolatis, verticillis subhirsutis. Hall. helu. n. 224. R.

M. aquatica exigua. Rai. angl. 3. p. 232.

M. hortensis. 4. Fuchsf. hist. 291.

Balamintha aquatica belgarum & matthioli. Lob. ic. 505.

Habitat in Anglia. Miller. 24

14. MENTHA Pulegium. *M. floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus, staminibus corolla longioribus. parif. gelr. Cramz. carn. sil. Ludw. ed. t. 195. Sabb. hort. 3. t. 49. Regn. bot. burg. lugd.*

M. caule prostrato, foliis subrotundis, obiter dentatis; staminibus exsertis. Hall. helu. n. 221.

Pulegium (vulgare) foliis ovatis obtusis, staminibus corollam æquantibus. Mill. dict. n. 1.

Pulegium latifolium. Bauh. pin. 222.

Pulegium. Riv. mon. t. 23. f. 1. Blackw. t. 302. Fuchsf. hist. 199.

Habitat in Germaniæ, Galliæ, Angliæ, Helvetiæ, inundatis. 24

15. MENTHA cervina. *M. floribus verticillatis, bracteis palmatis, foliis linearibus staminibus corolla longioribus. Sauv. monsp.*

28 DIDYNAMIA. GYMNOSPERMIA:
M. foliis lanceolatis punctatis, sub verticillo palmatis. *Hall.*
helu. n. 222.

Pulegium (cervinum) foliis linearibus, floribus verticillatis.
Mill. dict.

Pulegium angustifolium. *Bauh. pin. 222. Moris. hist. 3.*
p. 371. f. 11. t. 7. f. Riv. mon. t. 23. f. 2. Blackw.
t. 304.

Habitat Monspeli & ad Rhodanum. 24

773. GLECOMA.

Antherarum singulum par in formam crucis connivens.
Cal. 5. fidus.

1. GLECOMA *hederacea*. G. foliis reniformibus crenatis.
suec. pal. ingr. gallob. sil. nass. ludw. ed. t. 62. Knorr. del.
2. t. G. Regn. bot. delph. lugd. burg. lithu.

Chamæclema caule procumbente, radicato, foliis reniformibus, rotunde crenatis. *Hall. helu. n. 245.*

Calamintha hederacea. *Scop. carn.*

Glecoma caulibus procumbenti-erectis radicatis, foliis renum aut cordis forma, crenatis. *Crantz. austr.*

Hedera terrestris vulgaris. *Bauh. pin. 306. Blackw. t. 225.*
Riv. t. 67.

Chamæ Cissus. *Fuchsf. hist. 876.*

Habitat in Europæ septentrionalioris sepibus. 24

Variat flore albo; alia datur varietas flore majore. *Vaill.*
par. t. 6. f. 4. Hall. R.

774. LAMIUM.

Corollæ labium sup. integrum, fornicatum; lab. inf.
2. lobum; faux utrinque margine dentata.

1. LAMIUM *Orvala*. L. foliis cordatis inæqualiter arguteque ferratis, corollis fauce inflata, calyce colorato.

L. (*pannonicum*) foliis cordatis petiolatis, corollæ galea ferrata, tubo longiore. *Scop. carn. t. 27. R.*

L. maximum sylvaticum alterum. *Bauh. pin. 231.*

L. montanum saxatile ferme glabrum, flore amplo purpureo & labio inferiore crenulato. *Till. pisf. 92. t. 34. f. 1.*

Galeopsis maxima pannonica. *Clus. hist. 2. p. 35.*

β. *Orvala garganica* *Sp. pl.*

Papia garganica, foliis urticæ alternis, & eleganter incisæ; flore purpureo. *Michel. gen. 20. t. 17.*

Habitat in Pannonia, Italia, Istria. 24

DIDYNAMIA. GYMNOSPERMIA. 29.

Orvala garganica mera varietas est Lamii Orvalæ, ex loco umbroso rorido lecta, confirmante Horto Bononiensi litteris D. Blassii.

2. **LAMIUM lævigatum.** L. foliis cordatis rugosis, caule lævi, calycibus glabris longitudine tubi corollæ. *Sabb. hort. 3. t. 34. burg.*

L. foliis cordato-lanceolatis, serratis, verticillis paucifloris. *Hall. helu. n. 270. cum L. maculato conjungit. R.*

L. purpureum foetidum, folio parvo acuminato, flore majore. *Pluk. phyt. 198. f. 1.*

L. purpureum non foetens, folio oblongo. *Bauh. pin. 231.*

L. subrotundo rugoso folio, flore rubro. *Bocc. mus. 2. t. 23.*

Galeopsis flore purpurascente majore, folio non maculato. *Bauh. hist. 3. p. 321. Rai. hist. 559.*

Habitat in Italia, Sibiria, Helvetia. 24

Caules purpurascens. Folia serrata, vix pubescentia. Verticilli utrinque subquinqueflori. Calycum dentes, ventre suo duplo longiores, absque maculis fuscis ad basin. Corollæ dentes laterales latiores, lanceolati nec setacei, antrorsum porrecti; Labio inferiori patente, nec lobis reflexis parallelis. Calyx similis L. orvalæ.

3. **LAMIUM garganicum.** L. foliis cordatis pubescentibus, corollis fauce inflata, tubo recto, dente utrinque gemino. *Mill. dict. n. 3.*

L. garganicum subincanum, flore purpurascente: labio superiore crenulato. *Till. pis. 93. t. 34. f. 2.*

L. catarinæ folio, flore purpureo. *Ad. parif. 1717. p. 351. Habitat in Gargano monte ad castrum angeli. 24*

Caules pubescentes. Folia obtusiuscula, sublobato-serrata, pubescentia, non vero scabra. Verticilli multiflori. Corollæ tubi recti, nec ad basin ut in reliquis curvati. Labium superius 4. dentatum, patens, nec compressum.

2. **LAMIUM maculatum.** L. foliis cordatis acuminatis, verticillis decemfloris. *herb. sil. Kniph. cent. 3. n. 50. delph.*

L. alba linea notatum. *Bauh. pin. 231.*

L. plinii montanum campoclarense. *Col. ecphr. 1. p. 190. t. 192.*

L. maculatum. *Bauh. pin. 231.*

Habitat in Italia, Germania, Silesia. 24

Affine minimum L. albo, diversum Corollis purpureis; Foliis area longitudinali alba; Petiolis non dilatatis; Floribus utrinque 5. nec 10. Corollæ faux utrinque dentibus 2, superiore setaceo. Macula foliorum alba æstate disparet.

32 DIDYNAMIA. GYMNOSPERMIA:

Habitat inter Europæ segetes & olera.

Caulis geniculis tumidis. Calycis dentes spinosi s. pungentes. Corollæ galea crenata.

Gal. *h. canabina* differt. Corolla duplo majore flava (non purpurea); Labii intermedio purpureo. *Hallerus* separat, ego conjugo.

3. GALEOPSIS *Galeobdolon*. G. verticillis sexfloris involucro tetraphyllo. *ged. herborn. pal. sil. Kniph. cent. 3. n. 40. Knorr. del. hort. 1. t. N. 9. n. 1.*

Leconurus foliis ovatis serratis, acutis. *Fl. suec. paris. lithu. lugd. delph.*

Cardiaca foliis petiolatis, cordatis, verticillis foliosis. *Hall. helu. n. 274.*

Lamium (*Galeobdolon*) verticillis dimidiatis subsexfloris: ore edentulo. *Crantz. austr.*

Leovurus (*Galeobdolon*) foliis cordatis. *Scop. carn.*

Lamium flore luteo. *Riu. t. 20.*

Lamium folio oblongo luteum. *Bauh. pin. 231.*

Urtica iners tertia s. *Lamium* flore luteo. *Dod. pempt. 153.*

Habitat in Europæ nemoribus. 24

Genus obscurum, cum corollæ labium æquale, dentibus destitutum. Folia interdum albo-maculata. Flagella longa e radice.

776. B E T O N I C A.

Cal. aristatus. Corollæ lab. sup. adscendens, planiusculum; tubus cylindricus.

1. BETONICA *officinalis*. B. spica interrupta, corollarum labii lacinia intermedia emarginata. *suec. ingr. fl. ged. pal. gallob. sil. Ludw. eccl. t. 2. Blackw. t. 46. Kniph. cent. 5. n. 18. & cent. 11. n. 16. Knorr. del. hort. 2. t. B. 4. Sabb. hort. t. 82. Regn. bot. Flor. dan. t. 726. lithu. burg. lugd.*

B. foliis petiolatis, imis cordatis, superioribus crenatis, spica brevi, infidente foliis. *Hall. helu. n. 264.*

B. verticillorum calycibus bracteisque spinulosis, tubo longissimo. *Crantz. austr.*

B. alis septemfloris; foliis imis petiolatis, rugosis, ovatis, crenatis. *Scop. carn.*

Betonica. Dod. pempt. 40. Riu. t. 28.

Betonica purpurea. *Bauh. pin. 234.*

β. *Betonica* alba. *Bauh. pin. 235.*

Habitat in Europa. 24

DYDINAMIA. GYMNOSPERMIA. 33

3. **BETONICA alopecuros.** B. spica basi foliosa, corollis galea bifida. *Jacq. aust. t. 78. fil. delph.*

B. spicæ calycibus bracteisque mollioribus, tubo brevi, galea bifida. *Crantz. austr.*

B. alis multifloris; foliis petiolatis, cordatis, crenatis. *Scop. carn.*

Sideritis alopecuros. *Scop. carn.*

Horminum minus album, betonicae facie. *Bauh. pin. 239. prodr. 114.*

Betonica montana lutea. *Barr. icon. 339? Bocc. mus. 2. p. 82. t. 7.*

Betonicae folio, capitulo alopecuri. *Bauh. pin. 235.*

Betonicae folio, Alopecuros quorundam. *Bauh. hist. 3. p. 303.*

Alopecuros alpina, betonicae folio, capitulo alopecuri. *Pont. anth. 253. Segu. ver. 318.*

Horminum alpinum luteum, betonicae spica. *Rai. hist. 547.*

Habitat in montibus Sabaudicis, Silesiæ, Austriæ superioris, Lessanensibus, Italicis, Galloprovinciæ. 24

Folia caulina duabus oppositionibus. Bractææ integerrimæ. Corollæ ochroleucæ.

4. **BETONICA hirsuta.** B. spica basi foliosa, corollis galea integra. *delph.*

B. foliis ovatis, rotunde crenatis, spica ovata, obesa. *Hall. helv. n. 265. R.*

B. (Monieri) spica basi foliosa, corollis galea integra, foliis cordato-oblongis. *Gouan illustr.*

B. alopecuros minor purpureo rubicundo flore, montis aurei. *Le Monnier. obs. 146. Gou. R.*

B. alpina, incana, purpurea. *Barr. ic. 340. bene Gou. R.*

B. foliis hirsutis, floribus purpureis amplicissimis. *Mentz. pug. Zanon. t. 40. p. 46.*

Habitat in Apenninis. D. Basse hanc resuscitavit. In Pyrenæis. Gouan. 2.

Simillis B. alopecuroidi, sed torosior, hirsutior, robustior, spica brevior, crassior. Corollæ galea integra, nec bifida; colore purpurascens, nec flavescens. Spica ovata, nec ut in B. alopecuroidi interrupta. Bractææ inter flores oblongæ: majores serratæ, minores tantum subciliatæ, nec ovatæ. Folia in caule similiter uno alterove pari petiolato, præter par sessile sub spica.

34 DIDYNAMIA. GYMNOSPERMIA.

777. STACHYS.

Cor. lab. superius fornicatum, lab. infer. lateribus reflexum: intermedia majore emarginata. *Stam.* deflorata versus latera reflexa.

1. *STACHYS sylvatica*. *S.* verticillis sexfloris, foliis cordatis petiolatis. *suec. paris. ged. carn. gallob. noru. austr. pal. sil. Sabb. hort. rom. 3. t. 38. Regn. bot. burg. lithu. delph. lugd.*

St. foliis cordatis acuminatis petiolatis, floribus laxè spicatis. *Hort. cliff.*

Stachys sylvatica. *Riu. t. 26.*

Cordica foliis cordatis serratis, verticillis nudis, spicatis. *Hall. helu. n. 275.*

Lamium maximum sylvaticum foetidum. *Bauh. pin. 231.*

Galeopsis legitima. *Clus. hist. 2. p. 35. Blackw. t. 84.*

Habitat in Europæ nemoribus umbrosis. ☉

2. *STACHYS palustris*. *St.* verticillis subsexfloris, foliis lineari-lanceolatis semi-amplexicaulibus sessilibus. *suec. paris. ged. gallob. pal. sil. Læsel. pruss. ic. 41. Blackw. t. 273. delph. lithu. burg. lugd.*

St. foliis lineari-lanceolatis sessilibus basi emarginatis. *Hort. cliff.*

St. foliis subhirsutis, elliptico-lanceolatis, breviter petiolatis, verticillis spicatis. *Hall. helu. n. 257.*

St. foliis subhirsutis ellipticis ad basin emarginatis, verticillis ad 10. floris. *Crantz. austr.*

St. verticillis sexfloris, foliis lanceolatis. *Scop. carn.*

St. palustris foetida. *Bauh. pin. 236. Riu. t. 26.*

St. aquatica. *Tabern. ic. 577. bona.*

Habitat in Europa ad ripas, inque cultis humidiusculis. 2x
Verticilli 6. ad 10 floribus.

3. *STACHYS alpina*. *St.* verticillis multifloris, foliorum serraturis apice cartilagineis, corollis labio plano. *Fl. suec. herb. carn. Sabb. hort. 3. t. 41. burg. delph.*

St. verticillis duodecimfloris, caule villoso. *Hort. upf. 170. **

St. foliis lanceolato-ovatis basi emarginatis, verticillis tomentosis. *Hort. cliff. Dalib. paris.*

St. foliis hirsutis cordatis, verticillis subtomentosis. *Hall. helu. n. 256. R.*

St. latifolia major foliis obscure virentibus, flore galeato ferrugineo. *Pluk. alm. 356. t. 317. f. 4.*

Pseudo stachys alpina. *Bauh. pin. 236. prodr. 113.*

DIDYNAMIA. GYMNOSPERMIA. 35
Habitat in Germania, Helvetia, Carniola, &c. 24

Plantæ color obscurus. Folia flavo-virescentia serraturis apice cartilagineo fusco: Floralia superiora ovata, margine plano integerrimo. Verticilli involucrati, ut in 4. 5. Corollæ trisite purpureæ galea horizontali, nec erecta ut 4. 5. Labio non ad latera reflexo.

4. **STACHYS germanica.** St. verticillis multifloris, foliorum serraturis imbricatis, caule lenato. *ped. carn. pal. sil. Flor. dan. t. 684. Jacq. aust. t. 319. Kniph. cent. 10. n. 83. delph. burg. lugd.*

St. tomentosa, foliis imis cordatis, superioribus elliptico-lanceolatis, verticillis densissimis. *Hall. helu. n. 255.*

St. lanata. *Crantz. austr.*

St. verticillis quadragintafloris; caule lanato. *Hort. upf. 170.*

St. foliis oblongo-cordatis, floribus verticillatis. *Hort. cliff. paris.*

St. major germanica. *Bauh. pin. 236.*

Stachys. Fuchs. hist. 766. montana. Riu. t. 27. optima.

Habitat in Germania, Anglia, Gallia, Sibiria, &c. 24

Caulis lana alba vestitus, ut tota planta albus. Folia serraturis subimbricatis: floralia lanceolata, reflexa. Antherae albo-flavescentes.

Variat flore albo, foliisque angustioribus. *Bærh. R.*

10. **STACHYS maritima.** St. foliis cordatis obtusis tomentosis crenatis bracteis oblongis integerrimis. *Jacq. hort. t. 70.*

St. verticillis sexfloris, foliis radicalibus ovalibus crenatis: summis ovatis integerrimis, corollis calyce duplo longioribus. *Gouan. monsp.*

Betonica maritima, flore ex luteo pallescente. Dillen. elih. 50. t. 42. f. 50.

Betonica heraclea latifolia discoridis. Donat. venet. 84. Bocc. mus. 2. p. 164. t. 127.

Sideritis salviæ folio, nostras. Magnol. monsp. 203.

Habitat in littoribus maris, Venetiis, Monspeli, Orientis. D. Jacquin.

Folia radicalia, cordata, oblonga, obtusa, venosa, crenata, tomentosa. Caulis suffruticosus, spithameus, erectus, pubescens. Spica e verticillis interrupta. Bractea longitudine florum, ovals, oblonga, integerrima. Verticilli sexflori. Calyces lanati, quinquedentati, acuti, vix aristati. Corollæ luteæ.

Obs. Sideritis heraclea Col. ecphr. 1. p. 128. t. 131. hanc refert, sed flores purpurei. H. V.

36 DIDYNAMIA. GYMNOSPERMIA.

12. **STACHYS** *hirta*. St. verticillis sexfloris, caulibus prostratis, corollis labio superiore bifido divaricato reflexo.

Galeopsis caule piloso, calycibus labio corollæ superiore longioribus. *Sp. pl. Allion. nicæns.*

Marubium nigrum rotundifolium. Bauh. pin. 230.

Ocymum valentinum. Clus. hist. 2. 1. p. 42.

Habitat in Orientis, Hispaniæ, Italiæ collibus. 24

Caules prostrati, tetragoni, fistulosi, fragiles, pilis rarioribus adspersi. Folia petiolata, cordata, utrinque pilosa, venosa: inferiora obtuse serrata, obtusa, longius petiolata; Superiora acuta, acute serrata, Flores verticillati, utrinque 3. absque involucro. Calyx aristatus. Corollæ tubus calyce brevior, incurvatus. Labium superius fornicatum, semibifidum: lobis divaricatis obtusis; Lob. inferius obtusissimum, planum, obsolete trilobum, ad basin utrinque lacinula acuta vix manifesta. Puncta purpurascencia ad margines baseos labii superioris & in disco baseos labii inferioris.

13. **STACHYS** *recta*. St. verticillis subspicatis, foliis cordato-ellipticis crenatis scabris, caulibus adscendentibus. *pal. rub. aust. 1. 359. lugd. burg. lithu. delph.*

Betonica foliis hirsutis, ovatis, rotunde crenatis, bracteis ovato-lanceolatis, aristatis, integerrimis. Hall. helu. n. 262.

St. Betonica. Crantz. austr. non descriptis. Jacq. R.

Sideritis flore luteolo. Riu. mon. 1. 70.

Sideritis vulgaris hirsuta. Bauh. hist. 3. p. 425. Magn. bot. 240. R.

Stachys vulgaris hirsuta erecta. Bauh. pin. 233.

Habitat in Europa australi ad vias, in sterilibus. 24

Radix perennis. Caules adscendentes, rigidiusculi, pilis raris adspersi. Folia subpetiolata, cordata, elliptica, vel saepe lanceolata, profunde crenata, utrinque pilis raris scabra, subrugosa. Spicæ terminales, in verticillos distitos extrescentes, everticillis decemfloris. Calyces subspinosi. Flores lutei.

14. **STACHYS** *annua*. St. verticillis sexfloris, foliis ovato-lanceolatis trinerviis lævibus petiolatis, caule erecto.

Scholl. barb. n. 1012. pal. sil. Jacq. austr. 1. 360. burg. lugd. lithu.

St. foliis lanceolatis sessilibus basi attenuatis. Hort. cliff. paris.

Betonica foliis ovatis crenatis, bracteis ovato-lanceolatis. Hall. helu. n. 263.

Sideritis flore albo, barba luteola. Riu. 1. 69. Hall. R.

Betonica verticillata, calycibus spinosis. Sp. pl.

Sideritis arvensis latifolia glabra. Bauh. pin. 233.

DIDYNAMIA. GYMNOSPERMIA. 37

Alyssum majus. Tabern. hist. 929.

Habitat in Germania, Helvetia, Gallia, ad agrorum margines. ☉

Stamina ad latera reflexa genus indicant. Folia inferiora ovata, rugosa, obtusissima: Media oblonga, petiolata, Suprema sessilia, lanceolata, trinervia. Corollæ flavæ: labium superius marginibus lateralibus ferrugineis, recurvatum, emarginatum. Calyces mucronati, non tamen spinosi.

Caulis in montosis erectus, in pratis procumbens. Mygind. Sub stachide Betonica hanc describit Crantz. austr. p. 265. Jacq. R.

15. **STACHYS arvensis.** S. verticillis sexfloris, foliis obtusis nudiusculis, corollis longitudine calycis, caule debili. *Oed. dan. t. 887. pal. Sabb. hort. rom. 3. t. 51. delph.*

Trixago foliis hirsutis, ovato-cordatis, rotunde dentatis. Hall. helu. n. 231. R.

Stachys verticillis sexfloris, foliis obtusis nudiusculis: corollarum labio superiore integerrimo. De Neck. gallob.

Glecoma foliis cordato-oblongis crenatis. Hort. cliff.

Sideritis calycibus subinermibus bispidis. Fl. suec.

Sideritis albes trixagynis folio. Bauh. pin. 233. prodr. III.

Sideritis humilis, lato obtuso folio. Rai. hist. 565.

Stachys arvensis minima. Riv. t. 27. Petiv. t. 33.

Marubiastrum vulgare. Tournef. inst. 190.

β. *Lamium paludosum belgicum. Herm. lugdb. 531. t. 353. Habitat in Europæ arvis.*

Caulis tetragonus, obtusus, brachiatus, hirtus uti Petioli & calyces. Folia cordata, obtusa, obtuse serrata minus hirta. Calyces sessiles, semiquinquesidi, æquales, mucronati. Corolla albida, calyce fere minor: galea integerrima. Labio 3. fido: intermedio latiore purpurasente punctato, non emarginata. (Flos carneus. Hall. R.)

778. BALLOTA.

Cal. hypocrateriformis, 5. dentatus, 10. striatus.

Corollæ lab. superius crenatum, concavum.

1. **BALLOTA nigra.** B. foliis cordatis indivisis serratis, calycibus acuminatis. *suec. ged. carn. gallob. pal. sil. Kniph. cent. 6. n. 15. Sabb. hort. rom. 3. t. 35. Regn. bot. lugd. burg. lithu.*

B. foliis cordatis serratis, verticillis nudis. Hall. helu. n. 259. Ballote. Fuchf. hist. 154. Cæsalp. plant. 445.

Marrubium nigrum. Crantz. austr.

38 DIDYNAMIA. GYMNOSPERMIA.

Marrubiastrum. Riv. t. 66.

Marrubium nigrum foetidum. Bauh. pin. 230. Blackw. t. 136.

Habitat in Europæ ruderatis. 2.

2. **BALLOTA alba.** B. foliis cordatis indivisis serratis, calycibus subtruncatis.

Ballote flore albo. Tournef. inst 185. Vaill. paris. 20.

Ballote. Cam. epit. 572.

Habitat in Europa. 2.

Affinis nimium B. nigrae, an varietas.

Est omnino præcedentis varietas quum calyx non truncatus. Hall. Hanc etiam pingit. Sabbati in hort. rom. tom.

3. t. 36. R.

779. MARRUBIUM.

Cal. hypocrateriformis, rigidus, 10. striatus.

Corollæ lab. sub. bifidum, lineare, rectum.

** Calycibus 5. dentatis.*

1. **MARRUBIUM Alyssum.** M. foliis cuneiformibus quinque-dentatis plicatis, verticillis involucro destitutis. *Savv. monsp. Kniph. cent. 9. n. 63.*

M. album, foliis profunde incisissimis, flore cæruleo. Moris. hist. 3. p. 377. f. 11. t. 10. f. 12.

Alysson verticillatum, foliis profunde incisissimis. Bauh. pin. 232.

Alysson galeni. Clus. hist. 2. p. 35.

Habitat in Hispania. 2.

Flores utrinque 3, purpurei, parvi. Calyces acuti densibus patentibus.

2. **MARRUBIUM peregrinum.** M. foliis ovato-lanceolatis serratis, calycum denticulis setaceis. *Jacq. aust. t. 160. Kniph. cent. 7. n. 54.*

M. hispanicum supinum, foliis sericeis argenteis. Tournef. inst. 192. Dill. elsh. 219. t. 174. f. 215.

M. album latifolium peregrinum. Bauh. pin. 230. Moris. hist. 3. p. 377. f. 11. t. 9. f. 8.

M. alterum pannonicum. Clus. hist. 2. p. 34.

β. M. album angustifolium peregrinum. Bauh. pin. 230.

M. (creticum) foliis lanceolatis dentatis, verticillis minoribus, dentibus calycinis setaceis erectis. Mill. dict. n. 3.

M. creticum. Dalech. hist. 962.

Habitat in Siciliae, Cretæ, Austriae siccis β. in Libano 2.

4. **MARRUBIUM supinum.** M. dentibus calcynis setaceis rectis villosis. *monsp. carn.*

DIDYNAMIA. GYMNOSPERMIA. 39

M. album, sericeo parvo & rotundo folio. *Barr. ic.* 685.

Bocc. mus. 2. p. 78. t. 69.

M. album hispanicum majus. *Barr. ic.* 686.

Habitat in Hispania, G. Narbonensi, Carniolia. 24

* * *Calycibus 10. dentatis.*

5. **MARRUBIUM vulgare.** *M. dentibus calycinis setaceis uncinatis. suec. paris. ged. gallob. carn. pal. sil. Ludw. t. 145. Kniph. cent. 5. n. 56. Sabb. hort. 3. t. 59. Regn. bot. lithua. burg. lugd.*

M. dentibus calycinis setaceis uncinatis, medio corollarum segmento orchideo. *Crantz. austr.*

M. album vulgare. *Bauh. pin.* 230. *Riu. t.* 67. *Blackw. t.* 479.

M. vulgare. *Clus. hist.* 2. p. 34.

6. *M. album villosum.* *Bauh. pin.* 230. *prodr.* 110. *Burser*

Habitat in Europæ borealioris ruderatis. 24

7. **MARRUBIUM crispum.** *M. foliis cordatis subrotundis crenato-subdentalis, calycibus decemdentatis muticis.*

M. dictamni spurii foliis & facie. *Herm. par.* 200. t. 200.

Pseudo-dictamnium nigrum, rotundo crispo folio. *Boc. mus.* 2. p. 152. t. 129. & in titulo.

Habitat in Italia, Hispania. *Arduini.* 24

Caulis suffruticosus, erectus, hirtus. Folia petiolata, rugosissima, crenata, subtomentosa, subtus pilosa. Calyx infundibuliformis, 10. dentatus. Corolla purpurascens, non pilosa: Labio superiore subrotundo, obtuso, integro; L. inferiore trifido: intermedia bifida: lateralibus porrectis. Antheræ non ultra faucem.

Diversa species à Marrubio dictamini spurii foliis & facie. Herm. parad. t. 200. cui Lab. superius lanceolatum barbatum & stamina longitudine labii.

8. **MARRUBIUM hispanicum.** *M. calycum limbis patentibus, denticulis acutis.*

M. album rotundifolium hispanicum. *Herm. par.* 201. t. 201.

M. subrot. folio. *Barr. ic.* 797. *Bocc. mus.* 2. p. 167. t. 122.

Habitat in Hispania. 24

Folia subrotunda, plana, crenata. Corollæ lab. sup. 3. f. 4. fidum. Calycis dentes alterni minores.

9. **MARRUBIUM Pseudo-Dictamnus.** *M. calycum limbis planis villosis, foliis cordatis concavis, caule fruticoso.*

Kniph. cent. 8. n. 65. Sabb. hort. rom. 3. t. 47.

Pseudo-Dictamnus verticillatus inodorus. *Bauh. pin.* 222.

Pseudo-Dictamnium. Dpt. pempt. 281.

40 DIDYNAMIA. GYMNOSPERMIA.

Habitat in Creta. 7

Planta tota tomento denso compacto vestita.

780. LEONURUS.

Antheræ punctis nitidis adpersæ.

1. **LEONURUS Cardiacæ.** L. foliis caulinis lanceolatis trilobis. *suec. parif. pal. aust. ged. gallob. sil. Ludw. ed. t. 5. Kniph. cent. 4. n. 40. Reg. bot. Flor. dan. t. 727. lugd. burg. litua.*

L. foliis trilobis, lobis acutis. *Scop. carn.*

Cardiacæ foliis petiolatis, cordatis, verticillis foliosis. *Hall. helu. n. 274.*

Marrubium cardiacæ dictum. *Bauh. pin. 230.*

Cardiacæ. *Fusch. hist. 395. Blackw. t. 171.*

β. Cardiacæ crispa. *Rai. hist. 572. Sabb. hort. 3. t. 43.*

Habitat in Europæ ruderatis. 24

Var. β *distincta species est & cl. Mauray audit: Leonurus (crispus) foliis omnibus acute serratis, margine irregulariter reflexis, rugosissimis, caulinis quinquelobis. Vid. nov. Comm. Gætt. t. 8. p. 44. t. 4. R.*

2. **LEONURUS Marrubiastrum.** L. foliis ovatis lanceolatisque serratis, calycibus sessilibus spinosis. *carn. Kniph. cent. 8. n. 57.*

Sideritis foliis ovatis lanceolatis inciso-serratis. *Hort. cliff. mons.*

Marrubiastrum foliis cardiacæ. *Bocc. mus. 2. p. 120. t. 98.*

Habitat in Bohemia, Verania, Germania, etiam Java.

Calyx quinquesfidus, aristatus; aristæ etiam sub verticillis. Corollæ vix calyce longiores, parvæ, facie Nepetæ: galea integra; barbæ intermedia non crenata.

781. PHLOMIS.

Cal. angulatus, Corollæ labium superius incumbens, compressum, villosum.

1. **PHLOMIS fruticosa.** P. foliis subrotundis tomentosis crenatis, involucris lanceolatis, caule fruticoso. *Kniph. cent. 1. n. 61. Sabbat. hort. rom. 3. t. 15.*

P. involucris tomentosis lanceolatis, foliis cordatis villosis, caule fruticoso. *Hort. ups.*

P. involucri radiis lanceolatis villosis. *Hort. cliff.*

Verbascum latis salviæ foliis. *Bauh. pin. 240.*

Verbascum sylvestre alterum. *Dod. pempt. 146.*

β. **Phlomis cretica** fruticosa, folio subrotundo, flore luteo. *Tournef. cor. 10. Kniph. cent. 3. n. 71.*

DIDYNAMIA. GYMNOSPERMIA. 41

P. (latifolia) foliis oblongo ovatis petiolatis tomentosis, floribus capitatis, caule fruticoso. *Mill. dict. n. 3.*

P. latifolia capitata lutea grandiflora. *Dill. elth. 316. t. 237. f. 306.*

Habitat in Sicilia, Hispania. 2.

2. PHLOMIS purpurea. P. involucris linearibus obtusis calyce brevioribus, foliis cordatis oblongis tomentosis, caule suffruticoso.

Verbascum subrotundo salviæ folio. Bauh. pin. 240.

Salvia fruticosa, cisti folio haud incano, floribus purpureis.

Pluk. alm. 329. t. 57. f. 6.

Phlomis fruticosa lusitanica, flore purpurascente, foliis acutioribus. Tournef. inst. 178.

Habitat in Lusitania, Italia.

Caulis fere suffruticosus, tomentosus; plerumque debilis. Folia cordata, oblonga, utrinque tomentosa, albida. Petioli inferiores folio longiores. Verticilli floribus sex. Involucra calyce dimidio breviora, linearia, obtusa, tomentosa. Corolla labio inferiore supra purpurascente.

4. PHLOMIS lychnitis. P. foliis lanceolatis tomentosis, floralibus ovatis, involucris setaceis lanatis. *Mill. dict. n. 6. et. ic. t. 203.*

P. foliis ligulatis utrinque tomentosis, radiis involucris setaceis villosis. Sauv. monsp.

Verbascum angustis salviæ foliis. Bauh. pin. 240.

Verbascum sylvestre, foliis calviæ tentifoliæ. Lob. ic. 558.

Habitat in Europa australi D. Sauvages.

Habitus speciei 1. sed Folia angustiora. Cotollæ vix calyce majores. Involucra linearia, pilis longis crinita.

7. PHLOMIS herba venti. P. involucris setaceis hispidis, foliis ovato-oblongis scabris, caule herbaceo. *Sauv. monsp. Sabb. hort. 3. t. 17. delph.*

P. involucris radiis setaceis hispidis. Hort. cliff. 315.

Marrubium nigrum longifolium. Bauh. pin. 230.

Habitat in Persia, Tartaria, Narbona ad aggeres. 2.

8. PHLOMIS tuberosa. P. involucris hispidis subulatis, foliis cordatis scabris, caule herbaceo. *Mill. dict. n. 5. Gmel. it. 2. præf. p. VI. Pall. it. 1. p. 319. Kmph. cent. 4. n. 61.*

P. involucris radiis setaceis, foliis ex cordato lanceolatis.

Gmel. sib. 3. p. 241. n. 73.

P. foliis cordiformibus, galea lacera. Ad. gatt. 211. t. 12.

P. urticæ foliis, glabra. Amm. ruth. 49. et. n. 48.

42 DIDYNAMIA. GYMNOSPERMIA.

Galeopsis maxima, foliis hormini. *Buxb. cent. 1. p. 4. t. 6.*
Habitat in Sibiriae campestribus. 2.

Involucra subulata & Calycis dentes patentes, hispidi.
Corollæ galea dentata. Radix tuberosa.

12. *PHLOMIS Leonurus*. P. foliis lanceolatis serratis, caly-
cibus decagonis decedentatis muticis, caule fruticoso.

Knorr. del. hort. 2. t. L. Sabb. hort. 3. t. 44. Kniph. cent.
2. n. 54.

Leonurus foliis lanceolatis obruse serratis. Hort. Cliff.

Sideritis africana, flore aureo oblongo. *Barrth. act. 2. p.*
57. fig.

Leonurus capitis bonæ spei. Breyn. cent. 171. t. 86. Seb.
mus. 1. p. 22. t. 14. f. 1.

Stachys africana frutescens angustifolia, flore longissimo
phoeniceo; *Leonurus dicta. Moris. hist. 3. p. 383. g. 11.*
t. 19. f. 17.

Habitat ad Cap. b. Spei. 7

Planta spectatissima, dum floret. Calyx muticus. Involucrum
calyce longius, polyphyllum, subulatum, reflexum. Corolla
fulva, holosericea: Labio sup. longo, tomentoso, ciliato,
integerrimo; Lab. infer. brevi, nudo, membranaceo; Late-
ralibus reflexis, aridis: intermedia trifida: media emarginata.
Filamenta nivea. Antheræ bilobæ, flavæ, basi tantum adpersæ
farina globulari. Stigma superius brevius.

782. MOLUCELLA.

Calyx campanulatus, ampliatus, corolla latior,
spinosus.

1. *MOLUCCELLA lævis*. M. calycibus campanulatis sub-
quinquedentatis: denticulis æqualibus. *Kniph. cent. 11.*
n. 77. Sabb. hort. 3. t. 45. Regn. bot.

Molucca lævis. Dod. pempt. 91.

Melissa moluccana odorata. Bauh. pin. 229.

Habitat in Syria. ☉

Calyx corollis longior.

2. *MOLUCCELLA spinosa*. M. calycibus ringentibus octo-
dentatis. *Sabb. hort. 3. t. 46.*

Molucca spinosa. Dod. pempt. 92.

Melissa moluccana foetida. Bauh. pin. 229.

Habitat in Moluccis. ☉

Verticilli sexflori, sessiles. involucrum Spinis 4. ternatis;
linearibus, patentibus, spinosis. calycis Lab. superius majus,

DIDYNAMIA. GYMNOSPERMIA. 43

acuminato-spinosum ; *L. inferius rotundatum, integrum spinis septem. Corolla calyce longior.*

3. **MOLUCCELLA** *frutscens*. *M. calycibus infundibuliformibus quinquefidis, corollis calyce longioribus. Allion. nicæens. 51. **

M. persica. Burm. ind. 128. t. 38. f. 2. ex sicco.

Scordium spinosum, floris labio superiore s. galea lanuginosa s. villosa. Rai suppl. 311.

Habitat in Persia, ex qua in Italiam migravit. ¶

Fruticulus Ramulis dichotomis. Folia ovata, pollicaria, pilis raris pubescentia, utrinque 2. s. 3. profundis serraturis notata, petiolata. Spinæ axillares, utrinque solitariae, ternatae ut in Berberide, sed tantum ad florum verticillos. Calyces infundibuliformes, parvi, decagoni, regulares limbo quinquefido, spinis terminalibus 10. Corolla Cardiacæ, galea lanata. Media inter Ballotam Leonurum & Moluccellam.

Specimen Persicum Foliis magis pubescentibus, serraturis acutioribus, Calycis laciniis obovatis, membranaceis, acutioribus, subtridentatis; certe eadem. Cl. Burmannus misit.

783. CLINOPODIUM.

Involucrum multifetum, verticillo subiectum.

1. **CLINOPODIUM** *vulgare*. *C. capitulis subrotundis hispidis, bracteis setaceis. ged. aust. pal. sil. Scop. carn. Kniph. cent. 11. n. 31. Sabb. hort. 3. t. 65. flore albo. t. 66. burg. lugd. lithu.*

C. foliis ovatis, capitulis verticillatis. Fl. suec.

C. foliis ovatis lanceolatis obiter dentatis, stipulis setaceis. Hall. helu. n. 239.

C. origano simile. Bauh. pin. 224.

Clinopodium. Cam. epit. 563. Riu. t. 43.

β. *Clinopodium foliis ovatis rugosis verticillis omnibus distantibus. Mill. ic. 63. t. 95.*

Habitat in rupestribus Europæ, Canadae, Ægypti. ¶

Canadense ex seminibus duplo minores gerit corollas, cæterum idem.

Ægyptiacum β. corollis incarnatis, nec purpureis est.

784. ORIGANUM.

Strobilus tetragonus, spicatus, calyces colligens.

2. **ORIGANUM** *Dictamnus*. *O. foliis inferioribus tomentosis, spicis nutantibus. Kniph. cens. 4. n. 58. Luchw. ed. t. 134. Blachw. t. 462. Sabb. hort. 3. t. 76. Regn. bot.*

Origanum hybridum & Dictamnus. Mill. dict. n. 10. & 12. R.

44 DIDYNAMIA GYMNOSPERMIA.

Dictamnus creticus. Bauh. pin. 222.

Dictamnus cretense. Cam. epit. 472.

Dictamnus. Lob. ic. 502.

Habitat in Cretæ monte Ida. H

4. **ORIGANUM creticum.** O. spicis aggregatis longis prismaticis rectis, bracteis membranaceis calyce duplo longioribus. *Hall. helu. n. 234.*

O. foliis ovatis acutis glabris: venis scabris, spicis tetragonis. *Gron. orient.*

O. foliis ovatis obtusis scabris integerrimis, spicis confertis compactis glabris. *Amœn. acad. 1.*

O. creticum. *Bauh. pin. 223.*

β. O. folio subrotundo. *Bauh. pin. 223.*

O. monspeliense pulchrum. *Cam. epit. 468.*

Habitat in Europa australi, Palæstina, Basileæ. 24

6. **ORIGANUM heracleoticum.** Origanum spicis longis pedunculatis aggregatis, bracteis longitudine calycum.

O. heracleoticum, Cunila gallinacea Plinii. *Bauh. pin. 223.*

Orega, Origanum heracleoticum, Cunila. *Lob. ic. 492.*

Habitat in Europa australi, Græcia. 24

Habitus O. sylvestris, sed spicæ digitales, laxè imbricatæ; at differt ab O. cretico, quod bractææ vix lycibus longiores & demum quod spicæ pedunculatæ.

7. **ORIGANUM vulgare.** O. spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. *ged. gallob. austr. carn. pal. sil. Flor. dan. t. 638. Ludw. eccl. t. 90. Kniph. cent. 4. n. 59. Sabb. hort. 3. t. 75. Regn. bot. lithu. burg. lugd.*

O. foliis ovatis, spicis laxis erectis confertis paniculatis. *Fl. suec.*

O. foliis ovatis, umbellis coloratis staminibus exsertis. *Hall. helu. n. 233.*

O. sylvestre. *Bauh. pin. 243. Blackw. t. 280.*

O. sylvestre s. vulgare. *Fusch. hist. 522. Riv. t. 60.*

Habitat in Europæ, Canadæ rupestribus.

Americanum bracteis coloratissimis gaudet.

Datur varietas umbellis decoloribus, floribus albis. Crantz austr. l. c.

9. **ORIGANUM syriacum.** O. spicis longis ternatis pedunculatis villosis, foliis ovatis villosis.

Majorana syriaca vel cretica. *Bauh. pin. 224.*

Marum syriacum. *Lob. ic. 499. **

Habitat. . . .

DIDYNAMIA. GYMNOSPERMIA. 43

Habitus *Majoranae*. Folia subrotundo ovata, acuta, pilis vagis adspersa. Racemi ex axillis. Corymbus terminales, ex ramis longioribus brachiatus. Spicae saepius ubique 3, terminales, longae, erectae, tetragonae, villosae.

11. *ORIGANUM Majorana*. O. foliis ovatis obtusis, spicis subrotundis compactis pubescentibus. *Blackw. t. 319. Regn. bot.*

Majorana vulgaris. Bauh. pin. 224.

Amaracus vulgarior. Lob. ic. 498.

β. *Majorana tenuifolia. Bauh. pin. 224. Sabb. hort 3. t. 78.*

Majorana hortensis odorata perennis. Moris. hist. 3. p. 359.

Habitat in Lusitania, Palæstina. ☉

785. THYMUS.

Calycis bilabiati faux villis clausa.

1. *THYMUS Serpyllum*. T. floribus capitatis, caulis repentibus, foliis planis obtusis basi ciliatis. *Snec. gallob. aust. carn. pal. sil. Ludw. ed. t. 121. Kniph. cens. 6. n. 91. Regn. bot. lithu. burg. lugd.*

T. foliis ovatis, basi ciliatis. *Hall. heru. n. 235.*

T. repens, foliis planis, floribus verticillato-spicatis. *Hort. cliff.*

Serpyllum vulgare minus. Bauh. pin. 220.

Serpyllum vulgare. Dod. pempt. 277. Blackw. t. 418.

β. *Serpyllum vulgare majus. Bauh. pin. 220. Sabb. hort. 3. t. 69.*

γ. *Serpyllum vulgare minus, capitulis lanuginosis. Tournef. inst. 197.*

δ. *Serpyllum angustifolium hirsutum. Bauh. pin. 220. Kniph. cens. 6. n. 92.*

ε. *Serpyllum foliis citri odore. Bauh. pin. 220.*

ζ. *Serpyllum angustifolium glabrum. Bauh. pin. 220.*

Habitat in Europæ aridis apricis. ☿

Calycis dentes ciliati. Stamina longitudine corollæ.

Variat staminibus corollæ minoribus, flore minore & flaminibus corollæ longioribus, flore majore. Potlich. l. c.

2. *THYMUS vulgaris*. T. erectus, foliis revolutis ovatis, floribus verticillato spicatis. *monsp. Kniph. cens. 5. n. 89.*

Blackw. t. 211. Ludw. ed. t. 123. Reg. bot. delph.

T. vulgaris, folio tenuiore. *Bauh. pin. 219. Mill. dict. n. 2.*

β. T. vulgaris, folio latiore. *Bauh. pin. 219. Mill. dict. n. 1. Sabb. hort. 3. t. 68.*

Thymum durius. Dod. pempt. 276.

46 DIDYNAMIA. GYMNOSPERMIA.

γ. *Thymus supinus candidans odoratus. Tournef. inst. 196.*

δ. *Thymus capitulis minoribus, massiliensis. Tournef. ibid.*

Habitat in Hispaniæ montosis saxosis, in Gallia Narbonensi. nulla alia planta frequentior; in Sibiria. H. 24

3. **THYMUS** *Zygis*. T. floribus verticillato-spicatis, caule suffruticoso erecto, foliis linearibus basi ciliatis. *Læst. Kniph. cent. 8. n. 94.*

Thymo vulgatori rigidiori simile. Bauh. hist. 2. p. 371.

Thymum angusto longioreque folio. Barr. ic. 777.

Serpillum sylvestre Zygis dioscoridis. Clus. hist. 1. p. 358.

Serpillum folio thymi. Bauh. pin. 220?

Habitat in Hispania. H

Facie T. vulgaris. Caulis suffruticosus, erectus, palmaris; Rami multi, recurvi ante florescentiam. Folia linearia, longitudinaliter revoluta. supra pubescentia, subtus rubra. Petioli ciliati. Calyces bilabiati. Corollæ albæ cum staminibus. Mant. 413.

5. **THYMUS** *Acinos*. T. floribus verticillatis, pedunculis unifloris, caulibus erectis subramosis, foliis acutis serratis. *suec. pal. ged, austr. gallob. carn. sil. Kniph. cent. 8. n. 93 lugd. lithu. burg.*

T. caulibus vix ramosis, foliis ovatis acutis, pedunculis plurimis unifloris. Hort. cliff.

Clinopodium foliis ovatis dentatis, flore folio brevior. Hall. helu. n. 237.

Clinopodium arvense, ocymi facie. Bauh. pin. 225.

Clinopodium vulgare. Lob. ic. 506.

Acinos. Riu. t. 43.

Habitat in Europæ glareosis, cretaceis, siccis. ☉

5. **THYMUS** *alpinus*. T. verticillis sexfloris, foliis obtusiusculis concavis subserratis. *Jacq. austr. t. 97. cara. delph.*

T. montanus. Crantz. austr.

Clinopodium foliis ovatis acutis serratis, flore folio majori. Hall. helu. n. 238.

Clinopodium austriacum. Riu. pann. p. 622. 623. Hall. R.

Clinopodium montanum. Bauh. pin. 225. Bocc. mus. 2. p. 50, t. 45.

Acini pulchra species. Bauh. hist. 3. p. 620.

Habitat in Alpibus Helveticis, Austriacis, Monspeli. D. Rathgeb. ☉

Affinis nimium antecedenti, sed major & Flores tripla majores, pedunculati, utrinque quaterni.

DIDYNAMIA. GYMNOSPERMIA. 47

6. **THYMUS** *Piperella*. T. pedunculis multifloris lateralibus, foliis ovatis obtusis glabris nervosis integerrimis. Marum hispanicum nigrum. *Bocc. mus.* 2. p. 166. t. 117. *Barr. rar.* 354. t. 694.

Habitat in Hispania.

Similis *Melissæ Nepetæ*, sed glabra, Foliis minutis. *Dau. Royen.*

7. **THYMUS** *cephalotus*. T. capitulus imbricatis magnis, bracteis ovatis, foliis lanceolatis.

T. lusitanicus cephalotos, squamis capitulorum amplioribus. *Tournef. inst.* 196.

Tragoriganum, dictamni capite, hispanicum. *Barr. ic.* 787. *Bocc. mus.* 2. p. 50. t. 43.

β. Thymus lusitanicus cephalotos, capitulo majore. T.

γ. Thymus lusitanicus cephalotos, capitulo minori. V.

Habitat in Hispania, Lusitania. ♀

8. **THYMUS** *villosus*. T. capitulis imbricatis magnis, bracteis dentatis, foliis fetaceis pilosis.

T. lusitanicus, folio capillaceo villoso capite magno, purpurascens oblongo. *Tournef. inst.* 196.

β. T. lusitanicus, folio capillaceo villoso, capite magno purpurascens rotundo. *Tournef. inst.* 190.

Habitat in Lusitania. ♀

9. **THYMUS** *Mastichina*. T. floribus verticillatis, calycibus lanuginosis dentibus: calycinis fetaceis villosis.

Sampucus s. Marum mastichen redolens. *Bauh. pin.* 224.

Marum vulgare s. Clinopodium. *Dod. pempt.* 271. *Blackw.* t. 134.

Habitat in Hispaniæ petrosis. ♀

Ambiget media inter *Satureiam* & *Thymum*, sed cum *Stamina* delitescant in fundo corollæ, & *stilus* corollæ longior ad *thymum* refero. Verticilli plures diffusi. Corollæ albæ, longitudine calycis.

786. MELISSA.

Calyx aridus, supra planiusculus: labio superiore subfastigiato. *Corollæ* lab. superiore subfornicatum 2. fidum, lab. inf. lobo medio cordato.

1. **MELISSA** *officinalis*. M. racemis axillaribus verticillatis: pedicellis simplicibus. carn. *Kniph. cent.* 4 n. 48. *Sabb. hort.* 3. t. 61. *Blackw.* t. 27. *Regn. bot.*

M. floribus ex alis inferioribus subsessilibus. *Hort. ups. monsp.*

48 DIDYNAMIA. GYMNOSPERMIA.

M. foliis cordatis ferratis, petiolis foliis brevioribus. Hall. helv. n. 242.

M. hortensis. Bauh. pin. 229. Riu. t. 45.

Apiastrum f. Melissophyllum. Lob. ic. 227.

Habitat in montibus Genevensibus, Allobrogicis, Italicis. 24

2. *MELISSA grandiflora. M. pedunculis axillaribus dichotomis longitudine florum. Kniph. cent. 7. n. 56. delph. lugd.*

Thymus racemis lateralibus, sparsis: corollis calyce triplo longioribus. Scop. carn.

Calamintha magno flore. Bauh. pin. 229. Riu. mon. 43.

Calamintha montana præstantior.. Best. cyst. est. 7. t. 7. f. 1.

Habitat in Hetruriæ montosis. 24

3. *MELISSA Calamintha. M. pedunculis axillaribus dichotomis longitudine foliorum. austr. sil. Kniph. cent. 4. n. 47 Ludw. cat. t. 33. Regn. bot. delph. lugd. burg.*

M. floribus ex alis superioribus, pedunculo communi confertis. Hort. cliff. monsp.

M. floribus subhirsutis, ovato lanceolatis, acute ferratis, petiolis florigeris ramosis, folia superantibus. Hall. helv. n. 241.

Thymus racemis lateralibus, secundis, foliis ovatis, ferratis Scop. carn. non synonyma.

Thymus Calamintha. Ei. carn. ed. 2.

Calamintha vulgaris & officinarum germaniæ. Bauh. pin. 228.

Calamintha montana. Dod. pempt. 98. Riu. t. 46. Blackw. t. 166.

Habitat in Italiæ, Hispaniæ, Galliæ, Helvetiæ, Austriæ clivis, saxosis. 24

Caulis erectus.

4. *MELISSA Nepeta. M. pedunculis axillaribus dichotomis folio longioribus, caule adscendente hirsuto. Gisecke. icon. fasc. I. n. 14. Blackw. t. 167. delph.*

M. foliis hirsutis, ovato-lanceolatis, serratis, petiolis florigeris ramosis, folia superantibus. Hall. helv. n. 240.

M. floribus ex alis superioribus, pedunculo dichotomo, caule procumbente. Hort. cliff. Sauv. monsp. Gort. gelr.

Calamintha pulegii odore f. Nepeta. Bauh. pin. 228.

5. *Calamintha montana præalta, pulegii odore. Bocc. mus. 2. p. 45. t. 40. & 38. Riu. t. 47. Kniph. cent. 7. n. 57.*

Habitat in Italiæ, Galliæ, Angliæ, Helvetiæ aggeribus glareosis.

Caules duriusculi, uti tota planta hirsuti, Folia subcordata,

DIDYNAMIA. GYMNOSPERMIA. 49
data, supra glabra. Bractææ subulatae. Corolla subcærulea, palato albo cæruleoque marginato.

5. **MELISSA cretica.** *M. racemis terminalibus, pedunculis solitariis brevissimis.*

Calamintha incana, ocymi foliis. Bauh. pin. 228.

Calamintha, pulegii odore minor. Barr. ic. 1166.

Calamintha clinopodii austriaci foliis, odore pulegii. Pluk. alm. 75. t. 163. f. 4?

Habitat Monspelii, in Hispania.

Similis M. Nepetæ. Caulis palmaris, humilis, rectus, atropurpureascens. Folia ovata, parce obtuseque serrata. Verticilli plurimi, e pedunculis unifloris, aphillis, instar racemi ramos terminantes. Bracteis lanceolatis. Calyces fusci. Corollæ albo-purpureascentes.

6. **MELISSA fruticosa.** *M. ramis attenuatis virgatis, foliis subtus tomentosis, caule fruticoso.*

M. caule fruticoso, racemis verticillatis. Roy. lugdb.

Calamintha hispanica frutescens, mari folio. Tournef. inst. 194.

Calamintha montana incana minor. Moris. hist. 3. p. 413. n. 6.

Habuat in Hispania. ♂

787. DRACOCEPHALUM.

Corollæ faux inflata, lab. super. concavum.

** Spicata.*

1. **DRACOCEPHALUM virginianum.** *D. floribus spicatis, foliis lanceolatis serratis. Kniph. cent. 2. n. 22. Sabb. hort. 3. t. 26.*

D. foliis simplicibus, floribus spicatis. Hort. cliff.

Dracocephalum. Breyn. ic. 33. t. 27. Hir. act. 1712. p. 276. t. 11. Riu. monop.

Dracocephalus angustifolius. folio glabro serrato. Moris. hist. 2. p. 407. f. 11. t. 4. f. 1.

Pseudodigitalis persicæ foliis. Bocc. sic. 12. t. 6 f. 3.

Digitalis americana purpurea, foliis serratis. Dod. mem. 272.

Lysimachia galericulata spicata purpurea canadensis. Barr. ic. 1152.

Habitat in America septentrionali. 2

2. **DRACOCEPHALUM canariense.** *D. floribus spicatis, foliis compositis. Hort. cliff. 308. Mat. med. 153. Mill. dict. n. 2. Kniph. cent. 6. n. 38.*

Dracocephalo affinis americana trifoliata, terebenthinæ odore. Volk. norib. 145. t. 145.

Camphorosma. Moris. hist. 3. p. 366. f. 11. t. 11. f. ult.

Linnaei spec. plant. Europ. Pars II.

D

50 DIDYNAMIA. GYMNOSPERMIA.

Melissa forte canariana triphyllos, odorem camphoræ spirans penetrantissimum. *Pluk. alm.* 401. t. 315. f. 5. *Sabb. hort.* 3. t. 27.

Cedronella canariensis viscosa, foliis plerumque ex eodem pedicello ternis. *Comm. hort.* 2. p. 81. t. 41.

Habitat in Canariis? Americana. *H*

Morison descripsit. 1690.

5. DRACOCEPHALUM austriacum. D. floribus spicatis, foliis bracteisque linearibus partitis spinosis. *delph.*

Ruschiana laciniata. *Mill. dict.*

Hyssopus spicis interruptis. *Hort. cliff.*

Hyssopus, austriacus magno flore, folio chamæpithyos, *Herm. lugdb.* 330.

Ruyschiana hirsuta, foliis laciniatis. *Amm. ruth.* 50.

Chamæpithys cærulea austriaca. *Bauh. pin.* 250.

Chamæpithys austriaca. *Clus. hist.* 2. p. 185.

Habitat in Austria. *2*

Non vidit Crantzius in Austria. R.

6. DRACOCEPHALUM Ruyschiana. D. floribus spicatis, foliis bracteisque lanceolatis indivisis muticis. *Fl. suec. Flor. dan.* t. 121. *Kniph. cent.* 9. n. 33. *delph. lith.*

D. foliis linearibus indivisis integerrimis, floribus spicatis. *Hort. ups.* *

D. floribus verticillato-spicatis, foliis linearibus confertis. *Gmel. sib.* 3.

Dracocephalus foliis ellipticis, fasciculatis, verticillis spicatis, bracteis ovato-lanceolatis. *Haller. helv.* n. 254.

Ruyschiana (*spicata*) floribus spicatis, foliis bracteisque linearibus glabris indivisis. *Mill. dict.* n. 1.

Ruyschiana glabra, foliis integris. *Amm. ruth.* 50.

Ruyschiana flore cæruleo magno. *Bærh. lugdb.* 1. p. 172. *Zanon. nov.* t. 146.

Pseudo Chamæpithys austriaca. *Riu. mon.* 146.

Prunella hyssopi folio viridi, amplo flore cæruleo. *Morif. hist.* 3. p. 364. f. 11. t. 6. f. 9.

Habitat in Sibiria, Suecia, Helvetia, Dania. *2*

* * *Verticillata.*

9. DRACOCEPHALUM Moldavica. D. floribus verticillatis, bracteis lanceolatis: serraturis capillaceis. *Kniph. cent.* 4. n. 28. *Ludw. ed.* t. 15. *Blackw. t.* 551. *lith.*

D. floribus verticillatis, foliis ex ovato lanceolatis, bracteis longe ciliatis. *Gmel. sib.* 3 p. 230. R.

D. floribus verticillatis, foliis ovato-lanceolatis. *Hort. cliff.*

DIDYNAMIA. GYMNOSPERMIA. 52

Melissa peregrina, folio oblongo. *Bauh. pin.* 229.

Melissa moldavica. *Gam. epit.* 576.

Habitat in Moldavia, Sibiria. ☉

10. *DRACOCEPHALUM canescens*. D. floribus verticillatis, bracteis oblongis: ferraturis spinosis, foliis subtomentosis. *Hort. ups.* *Mill. dict. n.* 5. & *ic. t.* 129. *Kniph. cent.* 9. n. 31.

D. floribus verticillatis, foliis lanceolatis, floribus oblongis. *Hort. cliff.*

Moldavica orientalis, betonicae folio, flore magno violaceo.

Tournef. cor. 11. *Comm. rar.* 28. t. 18.

Sideritis incana, oleae folio, flosculis ex incarnato candidantibus, montis libani. *Volk. norib.* 353. t. 353.

Sideritis annua, flore luteo, utriculis & foliis longioribus. *Morif. hist.* 3. p. 389. f. 11. t. 8. f. 18.

Habitat in Oriente. ☉

11. *DRACOCEPHALUM peltatum*. D. floribus verticillatis, bracteis orbiculatis ferrato-ciliatis. *Hort. ups.* 166. *Mill. dict. n.* 8. *Kniph. cent.* 8. n. 38.

D. floribus verticillatis, foliis floralibus orbiculatis. *Hort. cliff.* 309. * *Roy. lugdb.* 312.

Moldavica orientalis, salicis folio, parvo flore caeruleo.

Tournef. cor. 11.

Habitat in Oriente. ☉

14. *DRACOCEPHALUM thymiflorum*. D. floribus verticillatis, bracteis oblongis integerrimis, corollis vix calyce majoribus. *Hort. ups.* 167. * *Mill. dict. n.* 7. *Kniph. cent.* 9. n. 34.

D. floribus verticillatis, bracteis lanceolatis integerrimis, foliis ovatis, superioribus integris, floribus vix e calyce eminentibus. *Gmel. sib.* 3. p. 233. t. 50.

Cedronella tatarica perennis, urticae foliis, flosculis minoribus, ex caeruleo rubentibus. *Roy. lugdb.* 537.

Moldavica orientalis minima, ocymi folio. *Comm. rar.* 29. t. 29 ?

Moldavica betonicae folio, floribus minimis pallide caeruleis.

Amm. ruth. 46. n. 58.

Habitat in Sibiria. ☉

52 DIDYNAMIA. GYMNOSPERMIA.

785. H O R M I N U M.

Calyx campanulatus : laciniis 4. subæqualibus ; quinta majore emarginata. *Corollæ* labium superius concavum.

1. *HORMINUM pyreanicum*. H. foliis ovatis obtusis, caule nudo. Hall. helu. n. 243.

Melissa pyrenaica. Jacq. hort. t. 183. R.

Callitrichum folio rotundo, flore magno violaceo. Bauh. hist. 3. p. 312. Moris. hist. 3. p. 395. Rai. hist. 546.

Melissa pyrenaica, caule brevi, plantaginis folio T. magn. hort. 133. t. 133.

Habitat in altis montium Tyrolensium, Pyræneorum. 24

Bractææ ovatæ, acuminatæ, integerrimæ. Melissa pyrenaica Scop. ann. 2. p. 55. Calyx 10-striatus : striis alternis crassioribus : Labio superiore 3-dentato ; lab. inferiore bifido : segmentis cruciformiter incumben- tibus. Corolla grandis, rubro-cærulea : Lab. super. bifido ; Lab. infer. trifido : segmentis subrotundis. Syst. Veg. p. 455.

789. M E L I T I S.

Calyx tubo corollæ amplior. *Corollæ* labium superius planum ; lab. inf. crenatum. *Antheræ* cruciatæ.

1. *MELITTIS Melissophyllum*. Melittis. monsp. ged. fil. Jacq. aust. t. 26. Kniph. cent. 6. n. 60. Reyn. bot. lithu. delph. burg. lugd.

Melissophyllum. Hall. helu. n. 244. Riv. mon. t. 21.

Melissa Fuchsi. Cam. ept. 99. 30.

Lamium montanum, melissæ folio. Bauh. pin. 231.

Habitat in dumetis, subalpinis Germaniæ, Helvetiæ, Austriæ, Angliæ, Monspelii. 24

Calyx campanulatus, erectus, corolla latior, trifidus : lacinia suprema saepe utrinque dente parvo notata. Corolla alba ; Tubo duplo longiore calyce ; Limbo quadripartito, patente, ex Labio superiore subrotundo, recto, integro ; Lab. inferiore trifido : intermedia majore, plana, integra, purpurea. Antheræ flavæ, corolla breviores.

Variat floribus totis niveis. Non est Lamium montanum melissæ folio Miller. t. 158. Hall. R.

790. OCYMUM.

Calyx labio superiore orbiculato; inferiore quadrifido. *Corollæ* resupinatae alterum labium 4-fidum; alterum indivisum. *Filamenta* exteriora basi processum emittentia.

2. OCYMUM *monachorum*. O. staminibus edentulis: alternis basi barbatis.

O. caryophyllatum monachorum. *Bauh. hist. 3. p. 260.*

Habitas -- ☉

Caulis erectus, pedalis, teretiusculus, subpilosus: Ramis fere supra-axillaribus. Folia petiolata, ovata, obtusiuscula, nuda, serrata, vix ciliata. Racemi verticillis sexifloris. Calycis labium superius cordatum striis tribus. Bractææ cordatae, caducae. Corollæ minores, albidæ labio inferiore purpurascens. Filamenta purpurea, longitudine corollæ, absque dentibus ad basin, at horum loco alterna ad basin villosa. H. V.

5. OCYMUM *basilicum*. O. foliis ovatis glabris, calycibus ciliatis. *Kniph. cent. 4. n. 56. Regn. bot.*

O. caryophyllatum majus. *Bauh. pin. 226.*

Ocymum. *Camm. epit. 308. Sabb. hort. 3. t. 84. Blackw. t. 104.*

β. Ocymum caryophyllatum maximum. *Bauh. pin. 226.*

λ. Ocymum latifolium maculatum s. crispum. *Bauh. pin. 226.*

δ. Ocymum viride, foliis bullatis. *Bauh. pin. 225.*

Habitas in India, Persia. ☉

6. OCYMUM *minimum*. O. foliis ovatis integerrimis.

O. foliis ovatis incanis. *Hort. cliff.*

O. minimum. *Bauh. pin. 226. Moris. hist. 3. p. 407. t.*

11. t. 10. f. 17. Sabb. hort. 3. t. 85.

Habitas in Zeylona. ☉

792. SCUTELLARIA.

Calyx ore integro: post florescentiam clauso, operculato.

3. SCUTELLARIA *alpina*. S. foliis cordatis inciso-serratis crenatis, spicis imbricatis rotundato-tetragonis. *Kniph. cent. 9. n. 87. delph.*

S. foliis ovatis, spicis imbricatis. *Hort. cliff.*

Cassida procumbens, foliis ovatis crenatis, spicis raris, foliosis. *Hall. helu. n. 281.*

54 DIDYNAMIA. GYMNOSPERMIA.

Teucrium alpinum inodorum, magno flore. *Bauh. pin.* 472.
prodr. 116. *Bursf. XV.* 35.

Habitat in Alpibus Helvetiæ, Vallis angustæ. H

Simillima S. lupulinæ, sed Corollæ violaceæ labio albida nec flavæ; Bractææ ovatae, acutæ, nec acuminatæ, subcorollis breviores, nec æquales; Folia minus oblonga.

6. *SCUTELLARIA galericulata. S. foliis cordato-lanceolatis crenatis, floribus axillaribus. Fl. suec. gallob. austr. pol. sil. Flor. dan. t. 637. Blackw. t. 516. Kniph. cent. 8. n. 8; Sabb. hort. 3. t. 30. delph.*

Cassida foliis oblonge cordatis, crenatis, verticillis nudis, bifloris. Hall. helv. n. 280.

Cassida galericulata. Scop. carn.

Scutellaria. Riu. t. 77. R.

Lysimachia cærulea galericulata f. Gratiola cærulea. Bauh. pin. 246.

Lysimachia galericulata. Dalech. hist. 1060.

Habitat in Europæ littoribus. 2f

7. *SCUTELLARIA hastifolia. S. foliis integerrimis: inferioribus hastatis, superioribus sagittatis. Fl. suec. * austr. carn. sub. Cassida. lithu.*

Scutellaria folio non serrato. Riu. mon. 77.

Habitat ad littora Sueciæ rarius; in Austria, Germania. 2f
An varietas præcedentis? certe maxime affinis.

8. *SCUTELLARIA minor. S. foliis cordato ovatis subintegerrimis, floribus axillaribus. Hudsf. angl. lugd. burg. delph. lithu.*

Cassida palustris minima, flore purpurascente. T. Rai. angl. 3. p. 244.

Habitat in Angliæ paludibus. 2f

Affinis nimium præcedentibus duabus.

12. *SCUTELLARIA peregrina. S. foliis subcordatis serratis, spicis elongatis secundis. 1. Kniph. cent. 5. n. 83.*

S. foliis cordatis obtusis, obtuse serratis, spica foliosa in summo caule. Gmel. sib. 3.

Cassida latifolia, menthæ facie, semine flavicante. Amm. ruth. Gmel. R.

Cassida. Colum. ecphr. 1. p. 187. t. 189.

Lamium peregrinum f. Scutellaria. Bauh. pin. 231.

Habitat in nemoribus Florentiæ, Liburni, in Sibiria.

Alia huic simillima Calycibus lanatis, pilis griseis: Corollæ galea apice pilosa. Bracteis ovatis brevibus.

Varietas floribus albis crassior gaudet foliis subpilosis.

793. PRUNELLA.

Filamenta bifurca : altero apice antherifera. *Stigma* bifidum.

1. PRUNELLA *vulgaris*. P. foliis omnibus ovato-oblongis serratis petiolatis. *gallob. carn. pal. fil. Ludw. ed. t. 19. Kniph. cent. 5. n. 72. Sabb. hort. 3. t. 32. lugd. lithu. burg.*

P. bracteis cordatis. *Fl. suec.*

Prunella foliis ovatis oblongis, calycibus superne truncatis. *Hall. helu. n. 277.*

Prunella officinalis. *Crantz. austr.*

Brunella major, folio non dissecto. *Bauh. pin. 260.*

Brunella. *Dod. pempt. 136. Blackw. t. 24. Riu. t. 29.*

β. Prunella. *vulgaris grandiflora. Jacq. austr. 377. Gmel. sub. Kniph. cent. 6. n. 74. lithu. burg. lugd.*

P. (*grandiflora*) foliis subintegris, ex subrotundo ovatis, calycibus superne profundius tridentatis. *Pollich. pal.*

Brunella foliis ovatis oblongis calycibus superne tridentatis *Hall. helu. n. 278.*

Brunella flore majore. *Riu. t. 29.*

Brunella cœrulea, magno flore. *Bauh. pin. 261. It. gotl. 219. 228.*

Brunella I, non vulgaris. *Clus. hist. 2. p. 42.*

Habitat in Europæ pascuis. 24

Plantæ. in apricis procumbens, pollicaris; in nemoribus erecta pedalis; β. grandiflora in rupium fissuris plerumque nascitur.

2. PRUNELLA *laciniata*. P. foliis ovato-oblongis petiolatis; supremis quatuor lanceolatis dentatis. *austr. herb. Jacq. austr. t. 378. Sabb. hort. 3. t. 31. delph.*

Brunella foliis imis ovatis oblongis, caulinis dentatis & semipinnatis. *Hall. helu. n. 279.*

Brunella folio laciniato. *Bauh. pin. 261. Riu. t. 30.*

Brunella II, non vulgaris. *Clus. hist. 2. p. 43.*

γ. Brunella minor alba laciniata. *Bauh. pin. 261. Kniph. cent. 10. n. 72.*

γ. Brunella verbenulæ folio, flore cæruleo. *Vaill. paris. 22. t. 5. f. 1.*

Habitat in Europæ pascuis rarius. 24

Fructificatio omnino P. vulgaris, a qua olim orta; fructura hodie persistens; adeoque tantillum distincta.

3. PRUNELLA *hyssopifolia*. P. folis lanceolatis integerrimis sessilibus, caule erecto. *burg.*

56 DIDYNAMIA. GYMNOSPERMIA.

P. foliis lanceolato-linearibus ciliatis subsessilibus. *Spec. pl. monsp.*

Brunella hyssopifolia. *Bauh. pin.* 261.

Brunella angustifolia integra hirsutior. *Moris. hist.* 3. p. 364. f. 11. t. 5. f. 7.

Habitat Monspeli. ☿

Differt a *P. vulgari*: *Caule erecto, nec adscendente; Foliis lanceolatis sessilibus integerrimis scabris, nec ovatis petiolatis serratis lævibus; Corollis quadruplo majoribus labii superioris carina pilosa.*

794. CLEONIA.

Filamenta bifurca; apice altero antherifera. Stigma quadrifidum.

1. *CLEONIA lusitanica.* *Prunella bracteis pinnato-dentatis ciliatis.* *Læfl. it. Mill. ic.* 47. t. 70.

Prunella odorata lusitanica, flore violaceo. *Barr. ic.* 561.

Clinopodium lusitanicum spicatum, & verticillatum. *Tournef. inst.* 195.

Bugula odorata lusitanica. *Corn. canad.* 46. *Moris. hist.* 3. p. 391. f. 11. t. 5. f. 4.

Prunella lusitanica, capite reticulato, folio pedicularis Tournesortii. *Moris. hist.* 3. p. 363.

Habitat in Lusitania, Hispania. ☉

Caulis erectus, brachiatus, inferne pilosus. Corollæ labium superius erectum: laciniis apice reflexis; Lab. inferius obcordatum, complicato-patens, basi superiori gibba. Filamenta omnia spinosa. Antheræ a tergo cristatæ.

795. PRASIUM.

Baccæ quatuor monospermæ.

1. *PRASIUM majus.* *P. foliis ovato-oblongis serratis.* *Kniph. cent.* 4. n. 66. *Sabb. hort.* 3. t. 37.

Lamium arboreum perenne creticum. *Zann. hist.* 112. t. 46. *Rai. hist.* 561.

Melissa fruticosa cretica sempervirens, teucris facie, flore albo. *Moris. hist.* 3. p. 408. f. 11. t. 21. f. 3.

Teucrium fruticans, amplo & albo flore, italicum. *Barr. ic.* 895.

Habitat in Sicilia, Romæ & in agro Tingitano. ☿

2. *PRASIUM minus.* *P. foliis ovatis crena utrinque duplici. Lamium fruticans, teucris folio lucido, calyce & flore magno candido: tantilla purpura varie notato.* *Cup. cathol.* 106.

Habitat in Sicilia. ☿

ANGYOSPERMIA.

797. BARTSIA.

Cal. 2. lobus, emarginatus, coloratus. *Corolla*
minus ipso calyce colorata: labio superiore
longiore. *Capsula* 2. locularis.

3. *BARTSIA viscosa*. B. foliis superioribus alternis serratis,
floribus distantibus lateralibus. *Huds. angl. Ger. prov.*

Euphrasia latifolia viscata ferrata. Moris. blas. 364.

*Euphrasia major lutea latifolia palustris. Rai. hist. 772. angl.
3. p. 285.*

Euphrasia lutea palustris. Pluk. alm. 142. t. 27. f. 5.

Euphrasia lutea latifolia palustris. Moris. hist. 3. p. 432.

Alectorolophos italica luteo-pallida. Barr. rar. 209. t. 665.

Habitat in Angliæ. Galliæ, Italiæ paludibus ad rivulos. ☉

4. *BARTSIA alpina*. B. foliis oppositis cordatis obtuse ser-
ratis. *succ. carn. delph.*

*Stæhelinia foliis cordatis, amplexicaulibus, serratis, flora-
libus coloratis. Hall. helu. n. 312.*

Stæhelinia alpina. Crantz. austr.

*Euphrasia caule simplici, foliis cordatis sessilibus obtuse
serratis. Fl. lapp.*

*Euphrasia rubra Westmorlandica, foliis brevibus obtusis.
Rai. angl. 4. p. 312*

Pedicularis genus montanum. Oed. dan. t. 43.

*Chamædrys vulgari falsæ aliquatenus adfinis alpina. Bauh.
hist. 3. p. 289. Hall. R.*

*Clinopodium alpinum hirsutum. Bauh. pin. 225. Pluk. alm.
110. t. 163. f. 5.*

Clinopodium alpinum. Pon. bald. 343.

Teucrium alpinum, coma purpureo-cærulea. Bauh. pin. 247.

*Habitat in Alpibus Lapponicis, Helveticis, Allobrogicis,
Baldo, Vallicis. 24*

798. RHINANTHUS.

Cal. 4. fidus ventricosus. *Caps.* bilocularis, obtusa,
compressa.

2. *RHINANTHUS Elephas*. R. corollis labio superiore
subulato recto,

58 DIDYNAMIA. ANGIOSPERMIA:

Elephas italica; flore magno proboscide surrecta. *Tournef. cor.* 48.

Elephas campoclarensum. Col. ecphr. 1. p. 186, t. 188.

Euphrasia lutea, floribus elephantis caput cum proboscide referentibus. *Morif. hist.* 3. p. 432. f. 11. t. 24. f. 14

Scordio Affinis Elephas ob florem. *Bauh. pin.* 248.

β. *Elephas orientalis*, flore parvo, proboscide surrecta. *Tournef. cor.* 48. itin. 3. p. 54. *

Habitat in umbrosis Italiae, Sibiriae. ☉

Habitus *præcedentis*, at Flos *diversus Calyce trifido: laciniis duabus reflexis: tertia majore erecta, plicata, acuta.*

3. **RHINANTHUS** *Crista galli*. R. corollarum labio superiore compresso brevior. *suec. gallob. sub. sil. pal. carn. Var. α. Crantz. austr. lugd. lithu. burg.*

Alectorolophus calycibus glabris. Hall. helu. n. 313.

Mimulus Crista galli. Scop. carn.

Crista galli. Dod. cer. 219. 220. *Riu.* 1. 29.

Crista galli femina. Bauh. hist. 3. p. 476. *Dill. giss. app. p.* 44.

Pedicularis pratensis lutea f. Crista galli. Bauh. pin. 163.

β. *Crista galli angustifolia montana. Bauh. pin.* 163. *prodr.* 86.

γ. *Crista galli mas. Bauh. hist.* 3. p. 436. *Dill. giss. p.* 80.

Rhinanthus (alectorolophus) corollarum labio superiore compresso brevior, calycibus villosis. *Pollich. pal. delph.*

Alectorolophus calycibus hirsutis. Hall. helu. n. 314.

Mimulus Alectorolophus. Scop. carn.

Habitat in Europæ pratis. ☉

4. **RHINANTHUS** *Trixago*. R. calycibus hirsuto-tomentosis, foliis oppositis obtuse serratis, caule simplicissimo.

Bartisia foliis oppositis lanceolatis obtuse serratis. Sp. pl. 1.

Trixago apula unicaulis. Colum. ecphr. 1. p. 199. t. 197.

Morif. hist. 3. p. 527. f. 11. t. 24. f. 8.

Crista galli spicata, flore luteo magno, messanensis. Rai. hist. 769.

Chamædrys unicaulis spicata. Bauh. pin. 248.

Anthirrinum folio dissecto. Bauh. pin. 211.

Habitat in Palestinæ, Italiae, Monspeliæ maritimis humen-sibus inter juncos. ☉

799. EUPHRASIA.

Cal 4. fidus, cylindricus. *Caps.* 2. locularis, ovato-oblonga. *Antheræ* inferiores altero lobo basi spinosæ.

1. **EUPHRASIA** *latifolia*. E. foliis dentato palmatis, floribus subcapitatis. *Sabb. horr.* 3. t. 7. *delph. burg.*

DIDYNAMIA. ANGIOSPERMIA. 59

E. foliis ovatis, floribus spicatis. *Sauv. monsp.*

E. purpurea minor. Bauh. pin. 111. Magn. monsp. 95. t. 94.

E. purpurea minor. Bauh. prodr. 111. Rai. hist. 774.

E. pratensis latifolia italica. Bauh. pin. 124. Moris. hist. 3. p. 430. f. 11. t. 24. f. 8. Rai. hist. 773.

E. tertia latifolia pratensis. Col. ecphr. 200. t. 302. f. 2.

Odontites foliis circa radicem ovatis ferratis, ceteris lanceolatis extremitate trifidis. Segu. ver. 1. p. 270.

Habitat in Apulia, Italia, Monspeli. ☉

Bractæ palmatæ, subhirsutæ.

2. *EUPHRASIA officinalis. E. foliis ovatis lineatis argute dentatis. suec. Hall. helu. n. 303. ged. austr. gallob. carn. pal. sil. Blackw. t. 427. Ludw. ed. t. 145. Kniph. cens. 8. n. 40. Sabb. hort. 3. t. 6. Regn. bot. lithu. burg. lugd.*

E. caule ramoso, foliis ovatis acute dentatis, Hort. cliff.

E. officinarum. Bauh. pin. 233.

E. minor. Dill. app. 53.

Euphrasia. Cam. epit. 767. Tabern. 362. Riu. t. 90.

Habitat in Europæ pascuis aridis. ☉

Variat flore albo; flore flavo. R.

3. *EUPHRASIA tricuspidata. T. foliis linearibus tricuspidatis.*

E. angustis & tricuspidatis foliis, floribus ex albo, purpureis. pluk. alm. 142. t. 177. f. 1. Zann. hist. 110. t. 76.

Habitat in Italia. ☉

Folia linearia dente utrinque unico. Corolla similis E. officinali.

4. *EUPHRASIA Odontites. E. foliis linearibus, omnibus ferratis. suec. ged. austr. gallob. pal. herb. sil. Flor. dan. t. 625. Kniph. cens. 12. n. 44. lithu. lugd. burg.*

E. foliis lineari-lanceolatis, ferratis. Scop. carn.

Odontites bracteis ferratis hirsutis. Hall. helu. n. 304.

Euphrasia pratensis rubra. Bauh. pin. 234.

E. altera. Dod. pempt. 55. Riu. t. 89.

β. *Euphrasia sylvestris major purpurea latifolia. Col. ecphr. 1. p. 201. t. 202. f. 1.*

Habitat in Europæ arvis pascuisque sterilibus. ☉

Variat floribus albis. R.

5. *EUPHRASIA lutea. E. foliis linearibus ferratis: superioribus integerrimis. Leyf. hal. carn. pal. Jacq. austr. t. 498. lugd. burg. delph.*

E. (Cort.) foliis inter flores ferratis. Crantz. austr.

Odontites bracteis glabris integerrimis. Hall. helu. n. 305.

60 DIDYNAMIA. ANGIOSPERMIA.

Odontites flore luteo. *Riu.* t. 91.

Euphrasia pratensis lutea. *Bauh.* pin. 244. *Moris. hist.* 3. p. 432. f. 11. t. 24. f. 16.

Euphrasia lutea montana angustifolia major altera. *Col. ecphr.* 1. p. 204. t. 203.

Habitat in Europæ australis montosis aridis. ☉

6. *EUPHRASIA linifolia* E. foliis linearibus : omnibus integerrimis calycibus glabris. *Ger. prov.*

E. foliis linearibus *Sauv. monsp.*

E. foliis lini angustioribus. *Bauh.* pin. 234.

E. linifolia. *Col. ecphr.* 2. p. 68. t. 69.

Habitat in Italia, Gallia. ☉

7. *EUPHRASIA viscosa.* E. foliis linearibus, calycibus glutinoso-hispidis. *Ger. prov. delph.*

Odontides foliis viscidis, rariter serratis, *Hall. helu.* n. 306.

Pedicularis annua lutea tenuifolia viscosa, pomum redolens. *Garid. aix.* 351. t. 78.

Habitat in Galloprovinciæ glareosis, sterilibus, Helvetiæ. ☉

Differt. ab. E. lutea foliis lineari-lanceolatis, nec striatis linearibus; Calycibus villosis viscidis, nec glabris: Corollis clausis, nec stamine brevioribus. *Distinguenda ab* E. linifolia.

800. MELAMPYRUM.

Cal. 4. fidus. *Cor.* lab. sup. compressum, margine replicato. *Caps.* 2. locularis, obliqua, hinc dehiscens. *Sem.* 2. gibba.

1. *MELAMPYRUM cristatum.* M. spicis quadrangularibus: bracteis cordatis compactis denticulatis imbricatis. *Suec. austr. carn. pal. Rivin.* t. 80. *Kniph. cent.* 11, n. 71. *delph. burg. lithu.*

M. foliis integerrimis, floribus spicatis, bracteis duplicatis, cristatis. *Hall. helu.* n. 311.

M. luteum angustifolium. *Bauh.* pin. 234.

M. luteum, linariæ folio. *Bauh. prodr.* 112.

M. angustifolium cristatum. *Pluh. alm.* 249. t. 99. f. 2.

Habitat in Europæ borealis pratis asperis. ☉

Variat floribus albis.

2. *MELAMPYRUM arvense.* M. spicis conicis laxis: bracteis dentato-setaceis, coloratis. *suec. ged. austr. gallob. pal. sil. Kniph. cent.* 1. n. 58. *Riu.* t. 79. *Sabb. kors.* 3. t. 8. *burg. lithu. lugd.*

M. foliis integerrimis, bracteis semipinnatis. *Hal. helu.* n. 310.

DIDYNAMIA. ANGIOSPERMIA. 61

M. foliis laciniatis laxè spicatis. Roy. lugdb.

M. purpurascènte coma. Bauh. pin. 234.

Triticum vaccinum. Dod. pempt. 541.

Melampyrum lanuginosum bæticum. Bauh. pin. 234.

Habitat in Europæ agris. ☉

3. *MELAMPYRUM nemorosum. M. floribus secundis lateralibus : bracteis dentatis cordato - lanceolatis : summis coloratis sterilibus , calycibus lanatis. succ. ingr. gei. austr. sil. Oed. flor. dan. t. 305. Kniph. cent. II. n. 71. lithu. delph.*

M. foliis ovato-lanceolatis , floralibus hamatis , acutatis , patulis , calycibus hirsutis. Hall. helu. n. 309.

M. bracteis unifloris coloratis , cordatis ; calycibus hirsutis. Scop. carn.

M. sylvaticum. Riv. t. 81. Hall. R.

M. coma cærulea. Bauh. pin. 234.

Parietaria sylvestris. 1. Clus. hist. 234.

Habitat in Europæ borealis nemoribus. ☉

4. *MELAMPYRUM pratense. M. floribus secundis lateralibus , conjugationibus remotis , corollis clausis. succ. austr. gallob. carn. pal. sil. Kniph. cent. II. n. 73. burg. lithu.*

M. foliis imis integerrimis , mediis dentatis , floralibus hastatis. Hall. helu. n. 308.

M. foliis lanceolatis , florum paribus remotis. Fl. lapp.

M. luteum latifolium Bauh. 243.

Habitat in Europæ borealis siccis. ☉

5. *MELAMPYRUM sylvaticum. M. floribus secundis lateralibus conjugationibus remotis , corollis hiantibus. Fl. succ. austr. carn. tub. Flor. dan. t. 145. Kniph. cent. 9. n. 65. delph. lithu.*

M. foliis integerrimis , floribus hiantibus. Hall. helu. n. 307.

M. latifolium , floribus parvis luteis. Fl. lapp. Hall. it. herc. n. 9.

Habitat in Europæ borealis sylvis. ☉

Differt a præcedente corollis dimidio brevioribus , limba hiantibus : iisque totis flavis , nec tubo albis.

801 LATHRÆA.

Cal. 4. fidus. Glandula depressa , ad basin futuræ germinis. Caps 1. locularis.

1. *LATHRÆA clandestina. L. caule ramoso subterrestri , floribus erectis solitariis.*

62 DIDYNAMIA. ANGIOSPERMIA.

L. foliis cordatis, floribus fasciculatis. *Guett. stamp. Dalib. paris. delph.*

Clandestina flore subcæruleo. *Tournef. inst. 952.*

Dentaria aphyllus purpurea, cespite denso. *Rai. extr. 116.*

Dentaria aphyllus, flore purpureo. *Rai. hist. 1230.*

Orobanche montana, *Rudb. elys. 2. p. 233. f. 14.*

Orobanche flore majore ex cæruleo purpurascente. *Bauh. pin. 87. prodr. 31. Rai. hist. 1227. Moris. hist. 3. p. 502.*

Orobanche s. Dentaria aphyllus purpurea, cespite denso. *Moris. hist. 3. p. 503. f. 12. s. 16. f. 13.*

Planta clandestina, s. madrona. *Dalech. hist. 960.*

β. Clandestina, Madrona, Occulta. *Borel. obs. 38.*

Habitat in umbrosis Galliae, Pyrenæorum, Italiae. 24

Caulis infra terram corymbosus, ut fructificationes solæ supra terram emineant solitariae in singulo ramo.

2. LATHRÆA *Phelypæa*. L. corollis campanulato - paten-
tibus.

Phelypea lusitanica, flore luteo. *Tournef. cor. 47.*

Orobanche elegantissima verna, flore luteo. *Grisl. lusit. Moris. hist. 3. p. 502. **

β. *Phelypea orientalis*, flore coccineo. *Tournef. cor. 47.*

Habitat in Lusitaniae umbrosis. 24

4. LATHRÆA *Squamaria*. L. caule simplicissimo, corollis pendulis labio inferiore trifido. *succ. paris. herb. sil. Flor. dan. t. 136. Blackw. t. 430. burg. lithu.*

L. caule subterrestri simplicissimo, floribus unilateralibus in series digestis. *Crantz. austr.*

Squamaria caule simplicissimo, floribus nutantibus, labello trifido. *Hall. helv. n. 297.*

Squamaria Orobanche. Scop. carn.

Squamaria. Riv. t. 89.

Dentaria. Mathiol. 964.

Orobanche, radice dentata, major. *Bauh. pin. 88.*

Orobanche, radice squamata, foliis rotundis, flore pendente & suave rubente, fundboënsis. *Rudb. elys. 2. p. 234. f. 17.*

Orobanche radice dentata altius radicata, foliis & floribus albo-purpureis. *Mentz. pug. t. 3. f. 3. Moris. hist. 3. p. 503. f. 12. s. 16. f. 14.*

Habitat in Europæ frigidioris umbrosissimis. 24

DIDYNAMIA. ANGIOSPERMIA. 63

803. TOZZIA.

Calyx 5. dentatus. *Capsula* unilocularis, globosa, monosperma.

1. TOZZIA *alpina*. Tozzia. Hall. helu. n. 228. Jacq. austr. t. 165. delph.

T. *alpina* lutea, alfine folio, radice squamata. Mich. gen. 20. t. 16.

Euphrasia lutea alfinefolia, radice squamata. Bauh. pin. 234. prodr. 111.

Dentaria buguloides, radice globosa: squamulis myontoideis, alpina. Mentz. pug. t. 9. f. 3. 4.

Orobanche buglossoides, radice rotunda, squamulis myontoideis, alpina. Moris. hist. 3. f. 12. t. 16. f. antepenult.

Anonyma f. gregorii, radice dentariæ. Col. ecphr. 2. p. 49. t. 50.

Habitat in Alpibus Helveticis, Austriacis, Italicis, Pyrenæis, locis asperis, humidis. 24

804. PEDICULARIS.

Calyx 5. fidus. *Capsula* bilocularis, mucronata, obliqua. *Semina* tunicata.

* *Caule ramoso.*

1. PEDICULARIS *palustris*. P. caule ramoso, calycibus oblongis cristatis calloso-punctatis corollis labio obliquis. Oed. dan. 255. pal. noru. carn. gallob. sil. Rivin t. 92. lugd. burg. lithu. delph.

P. caule ramoso, foliis pinnatis, pinnis pinnatis, dentatis, calyce bifido. Hall. helu. n. 320.

P. caule ramoso erecto; calycibus bifidis, 4. fidisque, laciniis fimbriatis. Crantz. austr.

P. caule ramoso erecto; calycibus bifidis crenatis. Fl. suec.

P. pratensis rubra elatior. Rai. hist. 700.

P. danica maxima. Lob. illust. 147.

Habitat in Europæ septentrionalis paludibus. ☉

Corollæ purpureæ.

2. PEDICULARIS *sylvatica*. P. caule ramoso, calycibus oblongis angulatis lævibus, corollis labio cordato. ged. gallob. pal. sil. nass. Fl. dan. t. 225. burg. lithu. delph.

P. caule ramoso, procumbente, foliis pinnatis, pinnis dentatis. Hall. helu. n. 321.

P. caule ramoso, floribus solitariis, calycibus quinquefidis. Fl. suec.

64 DIDYNAMIA. ANGIOSPERMIA.

P. caule ramoso, floribus solitariis remotis. *Hort. cliff.*

P. pratensis purpurea. *Bauh. pin.* 163.

P. minor. *Riu.* 1. 92. *Dill. giff. app.* p. 40.

Fistularia, *Dod. pempt.* 556.

Habitat in Europæ sylvis paludosis.

Corollæ purpureæ. (Variat & floribus albis. R.)

3. PEDICULARIS rostrata. *P.* caule declinato subramoso, corollis galea rostrato-acuminatis, calycibus cristatis subhirsutis. *Jacq. austr.* 1. 205. *carn. delph.*

P. caule procumbente, ramoso, floribus rostratis sparsis. *Hall. helu.* n. 322. 1. 8. f. 1.

P. (rostrato-capitata) foliis in utramque paginam versis crispatis; spica capitata foliosa: floribus rostrato-acuminatis. *Crantz. austr. sec. Jacq.* R.

P. caulibus reflexis, spica laxa purpurea. *Segu. ver.* 3. p. 125.

P. alpina, filicis folio, minor. *Bauh. pin.* 163.

Alectorolophus alpina 3. minor. *Clus. hist.* 2. p. 210.

Habitat in Alpibus, helveticis, austriacis.

Caulis simplex s. rarius ramosus, sæpe procumbens. Calyx hirsutus. Corolla rostrata.

* * * *Caule simplicissimo*

4. PEDICULARIS Sceptum Carolinum. *P.* caule simplici, floribus terno-verticillatis, corollis clausis, calycibus cristatis, capsulis regularibus. *Fl. succ. ingr. noru. Oed. dan.* 1. 26. *Kniph. cent.* 9. n. 75. *lithu.*

P. capsulis subrotundis regularibus. *Roy. lugdb.*

Sceptum carolinum. *Fl. lapp.* 1. 4. f. 4.

Habitat in Sueciæ, Borussæ, Rutheni spongiosis; sylvaticis, riguis. 24

Caulis tripedalis speciosissimus. Corollæ clausæ luteæ ore sanguineo.

5. PEDICULARIS verticillata. *P.* caule simplici, foliis quaternis. *carn. Jacq. austr.* 1. 206. *delph.*

P. foliis pinnatis quaternis, floribus spicatis. *Hall. helu. ed.* 2. n. 318. 1. 9. f. 1.

P. alpina, floribus purpureis spicatis, foliis pinnatis verticillatis. *Hall. helu. ed.* 1. 623. 1. 17. f. 1.

Alectorolophus alpina 2, floribus purpureis. *Clus. hist.* 2. p. 210.

Filipendula montana altera. *Bauh. pin.* 163.

Habitat in Sibiria, Helvetia, Austria. 24

Corollæ obtusæ purpureæ. Variat albo flore. *Gmel.*

7. PEDICULARIS recutita. *P.* caule simplici, foliis pinnatifidis

DIDYNAMIA ANGIOSPERMIA. 65

rifidis ferratis, spica foliosa, calycibus corolatis corilos obtusis. *Jacq. austr. t. 258.*

P. caule simplici, staminibus floris galea longioribus. *Sp. pl.*

P. caule simplici, foliis pinnatis, floribus spicatis, staminibus eminentibus. *Hall. helu. n. 316. t. 8. f. 2.*

P. (*obsoleta*) radice tuberosa; staminibus floribus minoribus obsoletis majoribus. *Crantz. austr. sec. Jacq. R.*

Habitat in summis Helvetiæ alpibus; inque Austria.

Caulis simplex. Folia pinnatifida, serrata. Spica sub foliosa.

Calyces colorati. Cor. purpurea: galea obtusa, nec rostrata.

9. PEDICULARIS flammea. P. caule simplici, foliis pinnatis retro-imbricatis. *Oed. t. 30. lugd.*

P. caule simplici, foliis semipinnatis obtusis: laciniis, imbricatis crenatis. *Fl. lapp. 244. t. 11. f. 2.*

P. caule simplici, foliis pinnatis, pinnis dentatis, denticulis retroversis. *Hall. helu. n. 315. t. 8. f. 3.*

Pedicularis alpina, folio ceterach. *Bauh. pin. 163.*

Habitat in Alpibus Lapponiæ, Helvetiæ. 24

Corollæ atropurpureæ præsertim apice.

10. PEDICULARIS hirsuta. P. caule simplici, foliis dentatopinnatis linearibus, calycibus hirsutis. *delph.*

P. caule simplici, calycibus villosis, foliis linearibus dentatis crenatis. *Fl. lapp. 245. t. 4. f. 3. Fl. suec. 508. 555.*

Oed. dan. t. 30.

P. pinnis foliorum acute pinnatis, floribus obtusis purpureis spicatis. *Allion. pedem. 52. t. 11. f. 1.*

Habitat in Lapponiæ Alpibus. 24

Corollæ incarnatæ.

11. PEDICULARIS incarnata. P. caule simplici, foliis pinnatis ferratis, calycibus rotundatis glabris, corollis galea uncinata acuta. *Jacq. austr. 140. delph.*

P. foliis pinnatis, pinnis dentatis, floribus rostratis spicatis, calycibus tomentosis. *Hall. helu. n. 319. sec. Jacq. R.*

P. (*rostrato-spicata*) caule strigoso recto spica oblonga canescente, flor. rostrato acuminatis. *Crantz. austr. Jacq. R.*

P. foliis alternis, pinnis semipinnatis, floribus laxè & longissime spicatis. *Allion pedem. 54. t. 12. f. 2. sed. calyces villosi acutiores.*

Habitat in Sibiria, Austria, Helvetia.

Folia Radicalia pinnata, pedalia, pinnis pinnatifidis, ferratis; Caulina pinnatifida fere ad basin, serrata. Caulis lævis, plusquam pedalis, Foliis alternis. Spica nuda, Calycibus semiglobosis, quinquedentatis, glabris. *Co-*
Linnaei, Spec. plant. Europ. Pars II.

E

86 DIDYNAMIA. ANGIOSPERMIA:

rolla incarnata: *Galea falcata & uncinata, acuta, valde adunca.*

Calycem tomentosum faciunt Jacquin, Haller, Crantz; hinc etiam nomen alionii etiam. R.

12. *PEDICULARIS lapponica*. P. caule simplici, foliis pinnatifidis ferratis, calycibus bifidis obtusis. *Oed. dan. 1. 2.*

P. caule simplici, foliis lanceolatis semipinnatis ferratis acutis. *Fl. lapp. 242. t. 4. f. 1.*

Habitat in Alpibus Lapponicis frequens. 24

Corollæ flavæ.

13. *PEDICULARIS comosa*. P. caule simplici, spica foliosa, corollis galea acuta emarginata, calycibus quinquedentatis. *delph.*

P. foliis bispinnatis, calyce non cristato, floribus ochroleucis in spicam nudam congestis. *Allion pedem 50. t. 12. f. 1.*

β. *Pedicularis alpina*, foliis alternis, pinnis semipinnatis. floribus rostratis ochroleucis, cet. *Allion pedem. 50. t. 12. f. 2. ex sententia Gouani. R.*

Habitat in Alpibus, italicis, Pyrenæis. 24

Calyx 5. dentatus, tubulosus integerrimus, parum pilosus. Corolla lutea labio superiore acuto, sed emarginato: dentibus duobus acutis, nutantibus.

14. *PEDICULARIS foliosa*. P. caule simplici, spica foliosa, corollis galea obtusissima integra calycibus quinquedentatis. *Jacq. austr. 2. t. 139. delph.*

P. spica foliosa. *Hall. helu. n. 31. t. 9. f. 2.*

P. comosa. *Scop. carn.*

P. alpina, filicis folio, major. *Bauh. pin. 163.*

P. alpina flore luteo, radice nigra. *Bauh. hist. 3. p. 43. 438. Hall. R.*

Alectorolophus alpina prima. Clus. hist. 2. p. 210.

Habitat in Alpibus Helvetiæ, Austriæ. 24

Folia bipinnatifida. Calyces angulis oreque pilosis. Corolla ochroleuca: Galea non rostrata, obtusissima, adunca. Caulis sæpe pedalis.

16. *PEDICULARIS tuberosa*. P. caule simplici, calycibus cristatis, corollis galea rostrato-adunca. *Scop. carn. delph.*

P. caule simplici, floribus spicatis, rostratis, calycibus glabris. *Hall. helu. n. 323. tab. 10.*

P. alpina lutea. C. B. *Seguier. ver. 3. p. 126.*

Filipendula montana, flore pediculariæ. Bauh. pin. 163.

Alectorolophus montana, flore luteo. Barr. ic. 466. Bacc. musc. 315. t. 8.

DIDYNAMIA. ANGYOSPERMIA. 67
Habitat in Alpibus Helveticis, Italicis, Sibiricis. 24.

Folia Millefolii. Lanugo ad folia & calyces. Corollæ luteæ.

808. ANTIRRHINUM

Cal. 5. phyllus. Corollæ basis deorsum prominens; nectarifera. Caps. bilocularis.

** Foliis angulatis.*

1. ANTIRRHINUM Cymbalaria. A foliis cordatis quinquelobis alternis, caulibus procumbentibus. *stamp. ingr. gallob. pal. Regn. bot. delph.*

A. caule repeate, foliis reniformibus quinquelobatis. Hall. helu. n. 339.

A. floribus solitariis, foliis quinquelobis, alternis, cordatis. Scop. carn.

Cymbalaria. Bauh. pin. 306. Cam. epit. 860. Dalech. hist. 1322. Riv. t. 86.

Habitat in rupibus & muris antiquis Germaniæ, Helvetiæ, Galliæ, Harlemi, &c. ☉

Semina rugosa, ut nuciei in glandis. Flore albo variat.

2. ANTIRRHINUM pilosum. A. foliis reniformibus pilosissimis alternis, caulibus procumbentibus. *Jacq. obs. 2. p. 29. t. 48. †*

Linaria hederæ folio villoso f. Cymbalaria alpina. Tourn. inst. 169.

Habitat in Alpibus.

Statura & filia A. Cymbalariae, sed tota pilosissima. Folia reniformi-subrotunda, margine in 9 f. 10. lobos obtusos crenulatos divisa. Jacq.

3. ANTIRRHINUM Elatine. A. foliis hastatis alternis, caulibus procumbentibus. *gallob. carn. pal. Oed. Fl. dan. 4. t. 6. delph. burg. lugd.*

A. caule procumbente, foliis hastatis, imis conjugatis, superioribus alternis. Hall. helu. n. 340.

Elatine folio acuminato in basi auriculato, flore luteo. Bauh. pin. 253.

Elatine. Cam. epit. 754. Blackw. t. 170.

6. Elatine folio acuminato; flore cæruleo. Bauh. pin. 253.

Habitat in Germaniæ, Angliæ, Galliæ, Italiæ arvis. ☉ Floribus cæruleis variat. R.

4. ANTIRRHINUM spurium. A. foliis ovatis alternis,
E 2

68 DIDYNAMIA. ANGIOSPERMIA.

caulibus procumbentibus. *austr. gallob. carn. lich. Regn. bot. delph. burg. lugd.*

A. caule procumbente, foliis villosis ovatis, imis conjugatis, superioribus alternis. *Hall. helu. n. 341.*

Elatine. Riu. 1. 86.

Elatine folio subrotundo. Bauh. pin. 253.

Veronica femina. Dod. pempt. 42.

Habitat in Germaniæ, Angliæ, Galliæ, Italiæ arvis. ☉

Peloriam & in hac videre Ramspeck. (comm. gætt. 1. p. 351. t. 15 f. 4.) & Stehelin. (Act. helu. 2. p. 25. t. 4.) R.

* * *Foliis oppositis.*

7. ANTIRRHINUM *triphyllum*. A. foliis ternis ovatis. *Sauv. Monsp.*

Linaria triphylla minor lutea. Bauh. pin. 212.

Linaria hispanica. Clus. hist. 1. p. 320.

Linaria ficula latifolia triphylla. Bocc. sic. 44. t. 22.

β. *Linaria triphylla cæulea. Bauh. pin. 312.*

Habitat in umbrosis montibus Valentinis & Hyblæis circa Syracusas. ☉

8. ANTIRRHINUM *triornithophorum*. A. foliis quaternis lanceolatis, caule erecto ramoso, floribus pedunculatis.

Linaria americana maxima, purpureo flore. Herm. lugd. 376. t. 377. Rai. hist. 1884.

Linaria, flore purpureo, americana. Riu. mon. 81.

Habitat in Lusitania, America.

Corollæ labium superius fornicatum est. Planta humanæ altitudinis.

9. ANTIRRHINUM *purpureum*. A. foliis quaternis linearibus, caule florifero erecto spicato.

A. foliis linearibus sparsis, nectariis subulatis recurvis, floribus laxè spicatis. *Hort. cliff.*

L. flore purpureo minore. *Riu. mon. 82.*

Linaria purpurea magna. Bauh. hist. 3. p. 460.

Linaria purpurea major odorata Bauh. pin. 213.

Linaria altera purpurea. Dod. pempt. 183.

Habitat ad radices Vesuvii.

Caulis (radice perenni) bipedalis, erectus, teres, levis. Rami seriores ex alis superioribus minores, florentes. Folia lanceolato-linearibus, lævia, subtus trinervia, patentia, alterna: inferiora verticillata. Racemi terminales, simplices, erecti, longi: pedicellis flore longioribus. Calyx

DIDYNAMIA. ANGIOSPERMIA. 69
*minutus. Corolla violacea tota, extus pallidior, palato
margine pubescente. Calcar longitudine corollæ, extrorsum
curvatum. Antheræ luteæ.*

10. ANTIRRHINUM repens. A. foliis linearibus confertis:
inferne quaternis calycibus capsulæ æqualibus, *Neck.
gallob. delph.*

A. foliis linearibus alternis sæpius oppositis acuminatis, flo-
ribus laxè spicatis. *Roy. lugdb.*

A. foliis lanceolato-linearibus sparsis, calycinis laciniis
capsulæ æqualibus. *Guett. stamp. Dalib. paris.*

*Linaria angustifolia, flore cinereo striato. Dill. elik. 198
t. 163. f. 197.*

Habitat in Anglia, Gallia, Italia. ☉

Flores inodori.

11. ANTIRRHINUM monspessulanum. A. foliis lineari-
bus confertis, caule nitido paniculato pedunculis spicatis
nudis. *Sauv. monsp. delph.*

*Linaria, capillaceo folio, odora. Bauh. pin. 213. prodr.
106. n. 4. Dill. elth. 199.*

Linaria odorata monspessulana. Bauh. hist. 3. p. 459.

Habitat in Gallia. 24

*Calcar corollæ rectum brevissimum, ut ipso calyce bre-
vius. Flores odorati. Simillimum A. sparteo, cui calcar
longitudine floris. Maxime etiam affine A. repenti.*

12. ANTIRRHINUM sparteum. A. foliis subulatis cana-
liculatis carnosis: inferioribus ternis, caule paniculato,
corollisque glaberrimis.

Habitat in Hispania. Læfl. ♂

*Radix altero anno florida. Caulis pedalis, glaberrimus,
paniculatus, erectus, minus strictus: ramis virgatis. Folia
primordialia, antequam caulem erigit, terna, oblonga,
reliqua alterna, subulata, canaliculata, glabra, carnosae,
rectae. Flores racemosi. Calyces glabri. Corollæ flavæ,
palato glabro cum tinctura rubicunda in retuso gibbo palati.
Differt ab A. bipunctato floribus pedunculatis, triplo
majoribus, ut tota planta multoties majore.*

13. ANTHIRRHINUM bipunctatum. A. foliis linearibus
glabris: inferioribus quaternis, caule erecto paniculato,
floribus spicato-capitatis.

A. foliis lanceolato-linearibus, floribus in summitate caulis
confertis parvis. *Guett. stamp.*

A. erectum, foliis linearibus: imis verticillatis, pauciflorum.
Sauv. monsp.

70 DIDYNAMIA. ANGIOSPERMIA:

Linaria lutea parva annua. Bauh. hist. 3. p. 357.

Habitat in Hispania, Italiæ arvis. ☉

Caulis spithameus, Ramis strictis, ad angulum rectum exeuntibus. Folia lanceolata, sessilia, plana, patenula, vix carnosæ, fere lævia: inferiora quaternæ. Flores subspicati, pauci, sessiles. Corollæ flavæ, punctis duobus fulvis in palato glabro versus labium superius; Labium superius bipartitum, replicatum, lanceolatum, erectum.

14. ANTIRRHINUM triste. *A. foliis linearibus sparsis: inferioribus oppositis, nectariis subulatis, floribus, sub-sessilibus. Fol. lanceolatis.*

Linaria hispanica procumbens. Mart. dec. 35. f. 2.

Linaria tristis hispanica. Dill. elth. 201. t. 264. f. 199. Mill. ic. t. 166.

Habitat Gibraltariæ.

Caules radicales plures, glabri, sæpius decumbentes. Folia linearia, subcarnosa, convexa, rore glauca. Capitulum ante anthesin simile A. arvensi, calyce & bracteis solis pubescentibus. Corolla colore Loti jacobæi, calcare striato, sæpe variat colore cinereo s. luteo, s. purpureo. Capsula craniolaris.

15. ANTIRRHINUM supinum. *A. foliis subquaternis linearibus, caule diffuso, floribus racemosis, calcar recto. delph. burg. lugd.*

A. foliis subquaternis linearibus, caule diffuso, floribus subspicatis nectario recurvato. Roy. lugdb. paris. monsp.

Linaria pumila supina lutea. Bauh. pin. 213.

Linaria hispanica. 5. Clus. hist. 1. p. 321.

Habitat in Galliæ, Hispaniæ arenosis.

Calcar reflexum est. Folia fere filiformia ut in A. sparteo, at racemi stricti, Corollæ calcar recurvatum & Calycis forium superius reliquis longius.

Simile præcedenti, sed Caulis non glaber. Syst. Veg. l. c.

16. ANTIRRHINUM arvense. *A. foliis sublinearibus: inferioribus quaternis, calycibus piloso-viscidis, floribus spicatis, caule erecto. gallob. pal. herb. Sabb. hort. 3. t. 5. delph. lugd. burg.*

A. foliis linearibus alternis recurvis, ramis caule brevioribus. Roy. lugdb.

Linaria arvensis cærulea. Bauh. pin. 213. prodr. 107. Dill. elth. 199. t. 163. f. 198.

β. Linaria pumila, foliolis carnosis, flosculis minimis flavis. Bauh. pin. 213. Dill. elth. 200.

DYDINAMIA. ANGYOSPERMIA. 71

7. *Linaria quadrifolia lutea. Bauh. pin. 213.*

Linaria tetraphylla lutea. Col. ecphr. 1. p. 299. t. 300. f. 1.

Habitat in Angliæ, Galliæ, Italiæ, Germaniæ arvis. ☉

Flores spicati Bractea reflexa, minuti: Calcare recurvo.

Folia glabra.

17. *ANTIRRHINUM pelisserianum. A. foliis caulinis linearibus alternis: radicalibus lanceolatis ternis, floribus corymbosis. stamp. paris. Sabb. hort. rom. 3. t. 4. burg. lugd.*

A. foliis caulem linearibus, ad radicem ovatis sæpe ternis Sauv. monsp.

Linaria foliis caulinis linearibus sparsis, radicalibus rotundis. Mill. dict.

Linaria annua purpureo-violacea, calcaribus longis foliis imis rotundioribus. Vaill. paris. 118. Magn. monsp. 159. Barrel. ic. 1162.

Linaria cærulea minor. Lob. illustr. 103.

Linaria cærulea calcaribus longis. Bauh. hist. 3. p. 461.

Habitat in Gallia, Italia. ☉

Caulis spithameus, erectus, teres, glaberrimus, subramosus. Folia alterna, remotissima, sublinearia, subcarnosa, glabra: radicalia s. infima terna s. quaterna. Flores parvi in capitulo subcorymboso. Calyx erectus, non appressus, sed laciniis distantibus. Corolla violacea palato albo, venis violaceis obsoletis: labio superiore longiore. Calcar rectum, corolla fere longius.

18. *ANTIRRHINUM saxatile. A. foliis lanceolato-linearibus sparsis villosis: inferioribus quaternis, caule decumbente, floribus spicatis.*

Linaria minor viscosa, flore luteo. Moris. bles. 280.

Linaria maritima minima viscosa, foliis hirsutis, floribus luteis. Moris. hist. 2. p. 499.

Habitat in Hispania. 24

Caules adscendentes. Folia sparsa linearia, acuta, pubescenti-viscida, apice patula. Flores spicati. Calyces lanati. Corolla flavissima: punctis 2 fulvis in palato & 2 in faucibus.

19. *ANTIRRHINUM viscosum. A. foliis caulinis linearibus alternis, radicalibus lanceolatis quaternis, calycibus villosis cauli approximatis.*

Habitat in Hispania. ☉

Planta erecta, statura A. pelisseriani, at flores magnitudine Linariæ.

72 DIDYNAMIA. ANGYOSPERMIA.

20. ANTIRRHINUM *multicaule*. A. foliis quinis linearibus carnosiss, floribus capitatis.

Linaria sicula multicaulis, molliginis folio. *Bocc. succ.* 38. t. 19. f. 1.

Habitat in Sicilia, Oriente. ☉

Caules *junci*, *laxi*. Folia *linearia*, *obtus*.

21 ANTIRRHINUM *glaucum*. A. foliis quaternis subulatis carnosiss, caulibus erectis floribus spicatis.

Linaria foliis confertis linearibus carnosiss. *Hall. Cat.* 306.

Linaria maritima foliis succulentis. *Buxb. cent* 4. p. 23 t. 37.

Linaria foliis carnosiss cinereis. *Bauh. pin.* 213?

Habitat in Europa australi, Oriente. ☉

Calcar cœruleis striis. Nimis affine priori.

22. ANTIRRHINUM *alpinum*. A. foliis quaternis linearilanceolatis glaucis, caule diffuso, floribus racemosis calcari recto. *monsp. Jacq. aust.* t. 58. *delph. burg.*

A. caule procumbente, rariter spicato, foliis verticillatis. *Hall. helv.* n. 338.

A. racemis terminalibus; foliis verticillatis; caule diffuso. *Scop. carn.*

Linaria quadrifolia supina. *Bauh. pin.* 213.

Linaria 3 stiriaca. *Clus. hist.* 1. p. 322.

β. Linaria cœrulea repens. *Bauh. pin.* 213.

γ. Linaria foliis carnosiss cinereis. *Bauh. pin.* 213. *ex Hall.*

Habitat in Helvetia, Austria, Baldo, Styria, Pyræneæis.

Folia lanceolata verticillis remotissimis; calycum una lacinia longior.

Varietas flore purpureo, ridu non aureo est Linaria parva purpurea, foliis sine ordine dispositis. *Bauh. hist.* 3. p. 460. *Hall. R.*

24. ANTIRRHINUM *villosum*. A. foliis omnibus oppositis ovatis villosis, caulibus simplicibus, floribus oppositis lateralibus.

A. saxatile minus, origani folio viscoso & villoso, flore albo amplo. *Barr. ic.* 597.

Linaria hispanica, nummulariæ folio villoso. *Tournef. inst.* 169.

Habitat in Hispania. Cl. Alstræmer. 24

Folia minuta, pedunculis breviora. Tota planta densissime pilosa.

25. ANTIRRHINUM *organifolium*. A. foliis plerisque oppositis oblongis, floribus alternis. *delph.*

A. foliis oppositis lanceolatis, floribus laxe racemoso *R.. ugd.*

DIDYNAMIA. ANGIOSPERMA. 73

A. saxatile, foliis serpilli. *Bauh. pin.* 212. *prodr.* 106. *Burs.* XII. 8, 9.

A. saxatile villosum, origani folio glutinoso. *Barr. ic.* 598.

A. saxatile, alines folio unctuosus & villosus. *Barr. ic.* 1102, 1103.

Orontium saxatile thymi folio, flore rubello. *Barr. ic.* 1313.

Habitat in Pyrenæis, Monte aureo, Massiliæ. ☉

Planta polymorpha, varians foliis ovatis & lanceolatis, forte filia A. minoris, à quo distinguitur ramis lignosis; Foliis caulinis omnibus oppositis, floriferis tamen alternis; Floribus duplo majoribus.

26. *ANTIRRHINUM minus*. *A.* foliis plerisque alternis lanceolatis obtusis, caule ramosissimo, diffuso. *suec. austr. gallob. pal. sil. Flor dant. t.* 502. *burg. lithu.*

A. viscidum, foliis inferioribus conjugatis, ellipticis, obtusis, calcare dimidii floris longitudine. *Hall. helv. n.* 335.

A. floribus solitariis; foliis sublinearibus alternis obtusis, Scop. carn.

A. arvense minus. *Bauh. pin.* 212.

A. tertium. *Cam. epit.* 922.

Linaria arvensis minima. *Riu. mon. t.* 85.

Habitat in Europæ cultis, ruderatis, glareosis. ☉

Folia infima, per tria paria, tantum opposita.

* * * *Foliis alternis.*

27. *ANTIRRHINUM dalmaticum*. *A.* foliis alternis cordatis amplexicaulibus. *Pall. it. 3. p.* 590.

Linaria foliis lanceolatis alternis, caule suffruticoso. *Mill. dict. n.* 13.

Linaria latifolia dalmatica, magno flore. *Bauh. pin.* 212, *prodr.* 106.

Linaria maxima, folio lauri. *Bauh. hist. 3. p.* 458. *Buxb. cent. 1. p.* 95. *t.* 24.

Habitat in Creta, Armenia unde innotuit 1594. *C. B.* ☉.

Flores ex alis foliorum, solitarii, pedunculati.

28. *ANTIRRHINUM hirtum*. *A.* foliis lanceolatis hirtis alternis, floribus spicatis: foliolo calycino supremo maximo.

A. foliis hirtis, radicalibus quaternis ovatis, caulinis lanceolatis alternis, floribus spicatis. *Gouan. illustr. 38. R.*

Habitat in Hispania. ☉

Caules aliquot simplices, spithamæi, non ramosi. Folia lanceolata (Lavandulæ,) alterna. Flores dense spicati, colore & magnitudine A. Linariæ. Calycis foliolum superius duplo majus quam reliqua. Capsulæ subrotundæ.

74 DIDYNAMIA. ANGYOSPERMIA.

29. ANTIRRHINUM *genistifolium*. A. foliis lanceolatis acuminatis, panicula virgata flexuosa. *Jacq. austr. t. 244. delph.*

A. foliis linearibus adscendentibus, floribus spicatis, confertis, calcare flore longiori. *Hall. helu. n. 337. sec. De la Chenal in act. helu. 8. p. 142. R.*

A. corollarum labio superiore maxilla destituto, panicula virgata flexuosa. *Crantz. austr.*

Linaria genistæ folio glauco, flore luteo. Herm. par. app. 9. Dill. elth. 202. Rai. suppl. 391.

Linaria flore pallido: rictu aureo. Bauh. pin. 213.

Linaria pannonica 1. Clus. hist. 1. p. 321.

Habitat in Sibiria, D. Gmelin. Austria inferiore, Helvetia, circa Dresdam. 24

Caules paniculati, subdecumbentes vel erectiusculi. Folia alterna, lanceolata, glabra, acuminata, omnino Genistæ tinctoriæ. Racemi longi, in paniculam dispositi, a calyce ad calycem flexuosi: Floribus alternis, distantibus, pedicellis longitudine calycis. Calyx quinquepartitus, glaber, acutus, longitudine capsulæ. Corollæ luteæ, magnitudine A. *Linaria.*

30. ANTIRRHINUM *junceum* A. foliis linearibus alternis, caule paniculato virgato, floribus racemosis.

Habitat in Hispania. Læsling. ☉

31. ANTIRRHINUM *Linaria*. A. foliis lanceolato-linearibus confertis, caule erecto, spicis terminalibus sessilibus, floribus imbricatis. *suec. austr. gallob. pal. sil. Ludw. ed. t. 12. Kniph. cent. 6. n. 9. Regn. bot. lithu. burg. lugd.*

A. foliis linearibus adscendentibus congestis, caule erecto spicato. *Hall. helu. n. 336.*

A. racemis terminalibus; floribus imbricatis, foliis linearibus confertis. *Scop. carn.*

A. foliis linearibus sparsis. *Hort. cliff.*

A. foliis lanceolato-linearibus sparsis, calycis laciniis capsula dimidio brevioribus. *Guett. stamp.*

Linaria vulgaris lutea, flore majore. Bauh. pin. 212. Riv. t. 83. Black. t. 115. Knorr. del. hort. 2. t. L. 6.

Osyris. Fuchf. hist. 543. Cam. epit. 390.

Habitat in Europæ ruderatis. 24

Palatum villosum fulvum.

Conf. Linaria lutea montana, genistæ tinctoriæ folio. Bauh. pin. 213. a Bursero lecta in monte S. Bernardi, que parte augustam vallem spectat, videtur hujus varietas Floribus tantum dimidio minoribus.

DIDYNAMIA. ANGIOSPERMIA. 75

Peloria. *Amen. acad.* 1. p. 55. t. 3. *Linariæ proles hybrida ejusdemque qualitatis & constans, radicibus infinite se multiplicans, charactere fructificationis diversissima, Corolla regulari, quinquesida, quinquecorniculata, pentandra, ut genus proprium constitueret & distinctissimum, nisi fructus semper abortiret; naturæ prodigium.*

Peloriam hanc in Germania etiam legerunt *Leers fl. herb.* n. 492. & *Scholl. barb.* n. 507. Iconem dedit *Kniph. cent.* 9. n. 9. R.

32. **ANTIRRHINUM linifolium.** A. foliis lanceolatis alternis trinerviis, floribus racemosis, pedunculis distantibus bractea brevioribus.

Linaria americana, parvo flore luteo. Bauh. pin. 212. *prodr.* 106.

Linaria constantinopolitana, lini sativi folio, flore luteo.

Tournef. cor. 9. *Buxb. cent.* 1. p. 16. t. 25. f. 2. *mala.*

Habitat in Italiæ maritimis. 24

Caules teretes, erecti, laves, Ramis alternis. Folia lanceolata s. angusto-lanceolata, alterna, remota, lævia, obsolete trinervia, exquisite acuminata. Racemus rectus, longus. Bracteæ lanceolatæ, sæpe flore longiores. Pedunculi longitudine calcaris. Calyces acuti, breves; Corollæ luteæ palato interius villoso; calcar recto, longitudine floris.

33. **ANTIRRHINUM chalepense.** A. foliis lineari-lanceolatis alternis, floribus racemosis, calycibus corolla longioribus, caule erecto.

A. foliis inferioribus quaternis, cæteris alternis, floribus racemosis, calycibus corolla duplo longioribus *Gouan. illustr.*

Linaria flore albo. Riv. mon. t. 80. f. 2.

Linaria chalepensis minor erecta, flore albo lineis violaceis. Moris. hist. 3. p. 502. f. 5. t. 35. f. 9.

Linaria annua angustifolia, flosculis albis longis caudatis.

Triumf. obs. 87. t. 87. f. 2. *Rai. hist.* 1884.

Habitat in Italia, Monspelii. ☉

34. **ANTIRRHINUM reflexum.** A. foliis ovatis alternis glabris, pedunculis axillaribus: fructiferis recurvatis elongatis, caule procumbente.

A. procumbens ramosum, foliis alternis ovatis acuminatis integerrimis, flor. cordatis axillaribus. *Allion. cors.* 205. f. 1.

Linaria pusilla procumbens latifolia, flore pallido: rictu luteo. Rai. hist. 755.

76 DIDYNAMIA. ANGIOSPERMIA.

Habitat in Barbaria. Brander. ☉

Caules filiformes, spithamæi. Folia alterna, ovata, subpetiolata, remota. Pedunculi axillares, solitarii, uniflori, longitudine foliorum; absoluta florescentia duplo longiores reflexi versus terram. Calyx acuminatus. Corollæ calcar longitudine floris. Capsula globosa.

35. ANTIRRHINUM *pedunculatum*. A. foliis alternis linearibus remotis, floribus paniculatis, pedunculis folio longioribus strictis.

Linaria lusitanica maritima, polygalæ folio. Tournef. inf. 169?

Habitat in Hispania. C. Alstræmer.

Caules spithamæi, lignosi, ramosissimi. Folia lanceolata, lævia, facie polygoni avicularis. Pedunculi stricti, foliis triplo longiores. Calyx lævis. Corollæ luteæ extus cæruleæ striatæ.

* * * * Corollis hiantibus aut ecaudatis.

36. ANTIRRHINUM *majus*. A. corollis ecaudatis, floribus spicatis, calycibus rotundatis. *Regn. ged. carn. gallob. Regn. bot. Kniph. cent. 1. t. 7. & cent. 3. n. 12. Rix. s. 82. Knorr. del. 1. t. L. 12. Ludw. ed. s. 51. delph. burg. lugd.*

A. foliis lanceolatis petiolatis, calycibus brevissimis, racemo terminali. *Hort. Cliff.*

A. foliis ellipticis obtusis, floralibus spicatis, calycibus obtusis, calcare brevissimo. *Hall. helu. n. 333.*

A. majus rotundiore folio. *Bauh. pin. 211.*

A. luteo flore. *Bauh. pin. 211.*

A. latifolium, pallido amplo flore. *Bocc. mus. 2. p. 49. t. 41.*

A. (*latifolium*) foliis lanceolatis glabris, calycibus hirsutis, racemo longissimo. *Mill. dict.*

β. A. majus alterum, folio longiore. *Bauh. pin. 211.*

A. foliis lanceolatis petiolatis, calycibus brevissimis, racemo terminali. *Mill. dict.*

Antirrhinum. Dod. pempt. 182. Lob. ic. 404.

Habitat in Europæ australis maceris, sepibus. ♂

Variat a Caule recto. Foliis ovatis.

β. Caule diffuso virgato; Foliis lanceolatis.

36. ANTIRRHINUM *Orontium*. A. corollis ecaudatis, floribus subspicatis, calycibus digitatis corolla longioribus. *austr. carn. gallob. pal. sil. Kniph. cent. 12. n. 13. delph. burg. lithu.*

DIDYNAMIA. ANGIOSPERMIA. 77

A. foliis lanceolatis petiolatis, calycibus flores superantibus.
Fl. suec.

A. foliis ellipticis obtusis, floribus sparsis, calycibus subulatis longissimis, calcare brevissimo. *Hall. helu. n. 334.*

A. arvense majus. *Riu. t. 82. Bauh. pin. 212.*

A. sylvestre, *Phyteuma. Dod. pempt. 212.*

Habitat in Europæ agris & arvis. ☉

Capsula dehiscens refert cranium Simiæ, venenata planta.

Datur varietas albo flore.

39. ANTIRRHINUM *Asarina.* **A.** corollis ecaudatis, foliis oppositis cordatis crenatis, caulibus procumbentibus. *Kniph. cent. I. t. 6.*

Asarina (procumbens) caule decumbente, foliis oppositis reniformibus crenatis. *Mill. dict. n. 1.*

Asarina S. Hederula saxatilis. Lob. ic. 601.

Hedera saxatilis, magno flore. Bauh. pin. 306.

Habitat in rupibus Végano conterminis, Genevæ. 24

Caules prostrati, dependentes. Folia obtuse lobata s. profunde crenata, pilosa. Corollæ absque calcari.

40. ANTIRRHINUM *molle.* **A.** corollis ecaudatis, foliis oppositis ovatis tomentosis, caulibus procumbentibus. *Kniph. cent. 10. n. 9.*

A. hispanicum villosum, origani folio glutinoso. *Tournef. inst. 168. Barr. ic. 598. (cui perperam calcar).*

Habitat in Hispania. 24

Caules prostrati, fragiles. Folia caulina opposita, mollissima. Pedunculi calyce longiores. Corolla Asarine villosa, alba palato flavo, Labio superiore rubro-striato. Calcar nullum.

** * * * * Corollis hiantibus.*

41. ANTIRRHINUM *bellidifolium.* **A.** foliis radicalibus lingulatis dentatis lineatis; caulinis partitis integerrimis. *delph. burg. lithu.*

A. foliis imis elliptico-lanceolatis, acute dentatis, caulinis capillaribus trifidis. *Hall. helu. n. 342.*

A. foliis radicalibus spathulatis, duplicato-serratis, caulinis digitatis, laciniis lanceolato-linearibus integerrimis, intermedia longiore. *Gouan. illustr.*

A. foliis radicalibus oblongis serratis, corollis calcaratis, patulis. *Spec. pl. 2.*

A. foliis imis spathulatis, summis digitatis. *Sauv. monsp.*

Linaria bellidis folio. Bauh. pin. 212. prodr. 106. t. 106.

Linaria odorata. Dod. pempt. 184.

Dodartia foliis radicalibus oblongo-ovatis serratis, caulinis

78 DIDYNAMIA. ANGIOSPERMIA.

linearibus integerrimis, floribus spicatis terminalibus
Mill. dict.

β. *Linaria lusitanica*, bellidis folio ampliore & villosa
Tournef. 169. ex Gouan. R.

Habitat Monspeli, in agris inter Lugdunum & Viennam, interque Lugdunum & Gratianopolin. ♂

Caulis erectus, rigens: Rami e summitate plures, simplices. Caules plures, breviores e radice. Folia Radicalia lingulata, obtusa, dentata, lineata, glabra; Caulina 3, 5, 7. partita, linearia, integerrima. Spicæ terminales, rectæ. Bractæ unifloræ, lineares, longiores. Flores caeruleo-violaceæ, nutantes, minuti, 5-fidi: laciniæ 2. erectæ; 3, nutantes. Palatum obsoletum, faucem non claudens. Calcar reflexum, minutum. Antheræ reflexæ, atrocaeruleæ.

43. *ANTIRRHINUM. linarioides. A. foliis linearilanceolatis ternis, caule erecto racemoso.*

Habitat in Europa australi.

Simillimum A. Linariæ, sed Caules angustiores; Folia ternaria. Flores racemosi, simillimi, verum labium corollæ superius albidum, nec flavum. Reliqua conveniunt.

809. CYMBARIA.

Calyx decemdentatus. Capsula cordata, bilocularis.

1. *CYMBARIA daurica. C. daurica pumila incana, linaria folio, magno flore luteo guttato. Amm. ruth. 47. t. 1. f. 2.*

Habitat in Dauriæ montanis, apricis, saxosis.

Planta tota incana, proxima Antirrhino. Rami oppositi, steriles. Flores laterales, paucissimi, subsessiles, magni. Calyx decemdentatus, quo a ringentibus omnibus differt. Corolla ringens. Lab. superiori fornicato, apice bifido; Lab. inf. æquali, trifido; palato gibbo.

811. MARTYNIA.

Cal. 5. fidus. Cor. ringens. Caps. lignosa corticata, rostro hamato, 5. locularis, bivalvis.

2. *MARTYNIA annua. M. caule ramoso, foliis integerrimis angulatis. Kniph. cent. 8. n. 66. Sabb. hort. 2. 2. 91.*

M. caule ramoso, foliis cordato-ovatis pilosis. Mill. dict. n. 1. & ic. t. 286.

M. caule periolisque fistulosis, floribus bibracteatis. Gouan hort.

DDYNAMIA. ANGIOSPERMIA: 79

M. foliis dentatis. Hort cliff.

M. annua villosa & viscosa, folio subrotundo, flore magno rubro. Mart. cent. 42. t. 42. Ehret. pict. t. 1. f. 1.

Craniolaria. Læfl. it. 225.

Proboscidea. Schmid. ic. t. 12. R.

Habitat in Americæ Vera Cruce. ☉

Calyx exterior diphyllus: lanceolatus. Perianthium monophyllum, ovatum, subquinelobum, latere inferiore usque ad basin dehiscens. Fructus rostro incurvo, longo, uncinato. Stigma tactu clauditur. Turra.

814. SCROPHULARIA.

Cal. 5. fidus. Cor. subglobosa, resupinata. Caps. 2. locularis.

1. *SCROPHULARIA marilandica. S. foliis cordatis serratis acutis basi rotundatis, caule obtusangulo. Kniph. cent. 2. n. 80.*

S. foliis cordatis oppositis, racemo terminali. Gron. virg. 71.

S. marilandica, longo profunde serrato urticæ folio. Rai. supp. 396.

Habitat in Virginia. 24

Simillima sequenti; duplo altior. Folia basi cordata, sed non trinervata seu ad nervos usque excisa. Petioli utrinque connexi barba minima.

2. *SCROPHULARIA nodosa. S. foliis cordatis trinervatis, caule obtusangulo. norv. t. 4. f. 1-3. carn. gallob. pal. sil. Lugdw. cæ. t. 72. Kniph. cent. 2. n. 81. lugd. burg. lithu.*

S. foliis cordatis, racemo terminali. Fl. suec. paris.

S. caule quadrangulo, paniculato, foliis cordatis oblongis. Hall. helv. n. 326.

. foliis cordatis basi transversis; caule & ramis spica nuda paniculata terminatis. Crantz. austr.

S. nudosa foetida. Bauh. pin. 235.

Scrophularia. Cam. epit. 866. Riv. t. 106. Blackw. t. 87.

Habitat in Europæ succulentis. 24

Folia trinervata ut in Lappa, Petasitide.

Variat foliis ternis; floribus viridibus. Hall. l. c. R.

3. *SCROPHULARIA aquatica. S. foliis cordatis petiolatis decurrentibus obtusis, caule membranis angulato, racemis terminalibus austr. carn. gallob. pat. sil. Flor.*

80 DIDYNAMIA. ANGIOSPERMIA.

dan. t. 507 Blackw. t. 86. Kniph. cent. II. n. 92. Sabb. hort. 2. t. 89. Regn. bot. lugd. burg. lithu.

S. caule quadrangulo, alato, paniculato, foliis ovato-lanceolatis. Hall. helu. n. 325.

S. foliis cordatis, petiolorum alis in caulem decurrentibus. Guett. stamp.

S. aquatica major. Bauh. pin. 235. caule fimbriato. Læf. prujf. 248. t. 75.

Scrophularia femina. Cam. epit. 867.

Habitat in Angliæ, Helvetiæ, Galliæ humidis. ♂

4. *SCROPHULARIA auriculata. S. foliis cordatis subtus tomentosis basi appendiculatis, racemis terminalibus. Læfl.*

Betonica aquatica septentrionalium. Lob. ic. 533. fig.

Habitat in Hispania.

Statura S. nodosæ. Caulis tetragonus, minime pilosus, brachiatus. Folia cordata, duplicato-serrata, obsa ubique, opaca, nuda, ad basin foliolis 2. ovatis accessoriis, fere ternata; in minoribus ramis folia simplicia. Corollæ purpureæ tristiores: labio revoluta livido-virescente.

5. *SCROPHULARIA. Scorodonia. S. foliis cordatis duplicato-serratis, racemo composito foliis interstincto. Hudf. angl.*

S. Melissæ folio. Tournef. inst. 166.

S. scorodonix folio. Pluk. alm. 338. t. 59. f. 5. Rai. hist. 766 Moris. hist. 2. p. 482. f. 5. t. 35 f. 6.

Habitat in Lusitania & Jersea insula Angliæ; in Sibiria.

Similis S. nodosæ. Caulis valde pilosus. Folia obtuse serrata. Racemus foliis interstinctus; quod non in prioribus. Syst. Veget. ed. Murray. p. 469.

Obs. In Mantissa p. 418. ita planta breviter describitur: Folia cordata sinu hiantia, duplicato-serrata, angulis seu serraturis acutis (nec obtusis S. auriculatæ). Caule valde piloso & defectu auricularum foliorum differt a simili S. auriculata.

7. *SCROPHULARIA betonicifolia. S. foliis cordatis oblongis dentatis: dentibus integerrimis; baseos profundioribus.*

S. betonicæ folio. Tournef. inst. 166.

S. aquatica montana mollior. Barrel. ic. 274. mala.

Habitat in Lusitania. 24

Caulis bipedalis, erectus, tetragonus, subpubescens, quasi purpureascens. Folia opposita; petiolata, subcordata, oblonga s. triplo longiora, quam lata; rugosa, subpubescentia.

DIDYNAMIA. ANGIOSPERMIA. 81

Subpubescentia, argute dentata dentibus acuminatis, integerrimis (adeoque Ebuli simillimis) ad basin profundioribus. Petioli foliis breviores. Panícula terminalis, subfoliosa. Corollæ triste purpurascens labello virescente. Antheræ flavissimæ. H. V.

7. *SCROPHULARIA orientalis*. S. foliis lanceolatis serratis petiolatis : caulinis ternis, rameis oppositis. *Kniph. cent. 5. n. 80.*

S. foliis lanceolato-linearibus acute serratis : inferne incisís, racemo composito. *Roy. lugdb.*

S. orientalis, foliis cannabinis. *Tournef. cor. 9.*

Habitat in Oriente. 24

Racemus compositus e pedunculis 3: f. 4. dichotomis.

Folia caulina variant terna aut quaterna.

8. *SCROPHULARIA frutescens*. S. foliis subcarnosis sessilibus lævibus apice recurvis.

S. foliis lanceolatis obtusis serrato-dentatis, pedunculis bifidis. *Roy. lugdb.*

S. lusitanica frutescens, verbenacæ foliis. *Tournef. inst. 1673 Rai. suppl. 396.*

S. peregrina frutescens, foliis Teucrii crassiusculis. *Herm. lugdb. 545. t. 547.*

Habitat in Lusitania. 7

Caulis perennans, tetragonus, acutangulus, inferne brachiatus. Folia sessilia, ovata, dentata, subcarnosa, lucida, glabra, parva, opposita: inferiora obovata; superiora oblonga, apice recurva. Racemus terminalis, foliatus: bracteis integerrimis: Pedunculis oppositis, trifidis, longitudine bractearum. Corollæ parvæ, atropurpureæ: laciniis lateralibus albis; lacinia infima minuta.

9. *SCROPHULARIA vernalis*. S. foliis cordatis, caulinis ternis, pedunculis axillaribus solitariis bifidis. *Oed. dan: 411. delph.*

S. foliis longe petiolatis, cordatis, subhirsutis, petiolis in alis ramosis. *Hall. helv. n. 327.*

S. foliis cordatis, pedunculis axillaribus solitariis dichotomis. *Hort. ups.*

S. flore luteo. *Bauh. pln. 236. prodr. 112. t. 112. Riv. t. 107.*

S. montana maxima latifolia. *Barrel. ic. 273 bene.*

Lamium pannonicum 2. exoticum. Clus. hist. 1. p. 38.

Habitat in Italia, Helvetiâ, Austria. ♂

Folia hirta ut caulis. Corollæ ovatae, flavae,

Linnaei spec. Plant. Europ. Pars II.

F

82. DIDYNAMIA. ANGIOSPERMIA.

10. SCROPHULARIA *trifoliata*. S. foliis glabris, inferioribus ternato-pinnatis obtusis; superioribus simplicibus, pedunculis subtrifloris axillaribus.

S. sambucifolia, flore rubro luteo vario pulchro. *Pluk. alm.* 338. t. 313. f. 6.

S. subrotundo crasso nigricante folio, flore luteo pallido, capsula turgida. *Bocc. mus.* 2. p. 65. t. 60.

Habitat in Africa, Corsica. ♂

11. SCROPHULARIA *sambucifolia*. S. foliis interrupte pinnatis cordatis inæqualibus, racemo terminali: pedunculis axillaribus geminis dichotomis. *Kniph. cent.* 7. n. 84.

S. foliis difformibus, pedunculis axillaribus aggregatis. *Roy. lugdb.*

S. sambucifolia. *Alp. exot.* 203. t. 202. *Mill. dict.* t. 231.

S. foliis laciniatis. *Bauh. pin.* 236. *prodr.* 112.

S. sambuci foliis, capsulis maximis. *Moris. hist.* 2. p. 483. f. 5. t. 8. f. 6.

♂ S. lusitana, maximo flore, foliis dissectis. *Rei. suppl.* 396.

Habitat in Hispania, Lusitania, Oriente. ♀

Caulis erectus, tetragonus e petiolis in angulos decurrentibus. Folia pinnata: foliolis 5. f. 7. præter alternas oppositiones minutas, cordatis, rugosis, supra glabris, serratis, impari majore. Racemus terminalis ex Pedunculis axillaribus, geminis, brevissimis, subdichotomis: Floribus magnis purpurascens labio inferiore virescente.

12. SCROPHULARIA *canina*. S. foliis pinnatis, racemo terminali nudo, pedunculis bifidis. *monsp. Kniph. cent.* 4. n. 75. *Sabb. hort.* 2. t. 90. *delph. lugd. burg.*

S. foliis inferioribus pinnatis, summis integris duplicato-ferratis, racemis axillaribus. *Mill. dict.*

S. foliis pinnatis, pinnis lobatis, circumferratis, caule paniculato. *Hall. helv.* n. 328.

S. foliis pinnatis, caule paniculato. *Scop. carn.*

Scrophularia Ruta canina dicta. *Bauh. pin.* 236.

Ruta canina. *Clus. hist.* 2. p. 209. *Lob. ic.* 2. p. 55.

Habitat in Helvetia, Narbona, Italia. ☉

Pedunculi bifidi: Flores sessiles: primo axillari; reliqui alternis.

13. SCROPHULARIA *lucida*. S. foliis inferioribus bipinnatis subcarnosis glaberrimis racemis bipartitis.

S. foliis bipinnatis glaberrimis lucidis, racemis bipartitis

DIDYNAMIA. ANGIOSPERMIA. 83

terminalibus. *Mill. dict.*

S. foliis filicis modo laciniatis f. *Ruta canina latifolia*. *Bauh. pin.* 236.

S. saxatilis lucida, *laferpitii massiliensis folio*. *Bocc. mus.* 2. p. 166. t. 117.

S. glauco folio in amplas lacinias diviso. *Tournef. cor.* 9. it. 1. p. 84. t. 85.

Habitat in Oriente, Creta, Neapoli, in ipsis muris Hydrunti. 4

Caulis teres rectus viridis. Folia similia S. caninae, sed subcarnosa. Racemus rigidus, erectus. Corollae pallidae labio superiore rubro.

Scrophulariam foliis filicis modo laciniatis. Bauh. distinc- tam speciem esse monet Millerus eamque dicit: Scroph. (filicifoliam) foliis pinnatis, foliolis acute dentatis, racemis terminalibus. Mill. dict. n. 10. R.

16. *SCROPHULARIA peregrina. S. foliis cordatis lineatis lucidis, pedunculis axillaribus bifloris, caule sexan- gulari.*

S. foliis cordatis: superioribus alternis, pedunculis axilla- ribus bifloris. Spec. pl. 2. Gouan. monsp.

S. urticae folio. Bauh. pin. 236.

S. peregrina. Cam. hort. 157. t. 43.

Habitat in Italia. ☉

Caulis atropurpureus, acutangulus: angulis 4. 5. s. 6. Pedunculi 2-5-flori. Corolla purpurea.

815. CELSIA.

Cal. 5. partitus. Cor. rotata. Filamenta barbata.

Caps. 2. locularis.

1. *CELSIA orientalis. C. foliis bipinnatis.*

Verbascum orientale, sophiae folio. Tournef. cor. 8. *Buxb. cent.* 5. p. 17.

Blattaria orientalis, agrimoniae folio. Buxb. cent. 1. p. 14. t. 20.

Habitat in Cappadocia, Armenia. ☉

2. *CELSIA Arcturus. C. foliis radicalibus lyrato-pinnatis, pedunculis flore longioribus.*

Verbascum (Arcturus,) foliis lyrato-pinnatis. Sp. pl.

Verbascum humile creticum laciniatum. Bauh. pin. 240.

Verbascum brassicae folio. Colum. ecphr. 2. p. 81. t. 82.

84 DIDYNAMIA. ANGIOSPERMIA.
Verbascum sylvestre creticum. Alp. exot. 133. t. 122.
Habitat in Creta.

Folia opposita in mea (Auctorum vero alterna) ; infima pinnata s. sublyrata , foliolis lateralibus minutis ; superiores simplicia. Floralia sessilia , cordata , serrato-dentata. Racemus terminalis , longus. Pedunculi solitarii , uniflori , foliis floralibus longiores. Corollæ flavæ. Filamenta 4 barba purpurea.

816. DIGITALIS.

Cal. 5. partitus. Cor. campanulata , 5. fida , ventricosa. Caps. ovata , 2. locularis.

1. DIGITALIS purpurea. D. calycinis foliolis ovatis acutis , corollis obtusis : labio superiore integro. *pal. Flor. dan. t. 74. Kniph cent. 3. n. 37. Blackw. t. 16. Reg. bot. lugd. delph. burg.*

D. foliolis calycinis ovatis. *Dalib. paris. 192. acuminati. gelr.*

D. foliis calycinis ovatis , galea integra. *Hall. helv. n. 330.*

D. purpurea , folio aspero. *Bauh. pin. 243. Knorr. del. 1. t. F. 10.*

D. purpurea. *Ded. pempt. 168. Riv. t. 104.*

2. Digitalis alba , folio aspero. *Bauh. pin. 229. Vaill. p. 80. Knorr. del. 1. t. F. 9.*

Habitat in Europa australiore. ♂

Corollæ venter intus adpersus maculis ocellaribus. Folia rugosa.

2. DIGITALIS minor. D. corollis obtusis ; labio superiore subbilobo , foliis lævibus.

D. hispanica purpurea minor. *Tournef. inst. 165.*

Habitat in Hispania. 2℥

Caulis lævis , præcedente dimidio brevior , Pedunculi villosi. Folia sessilia , lanceolata , non rugosa , lævia , subintegerrima. Flores simillimi D. purpureæ , figura , magnitudine , colore , sed puncta copiosiora palati absque inde pallido. Labium superius profundius bilobum ; Lab. inferius obtusissimum : lateralia reflexa. Antheræ punctis ferrugineis.

3. DIGITALIS Thapsi. D. foliis decurrentibus.

D. hispanica purpurea minor, *Tournef. inst. 165.*

DATE: 3-22-2013 TIME: 11:00 AM

86 DIDYNAMIA. ANGIOSPERMIA.

obtusis patentibus, corollæ labio inferiore barbato. *Sabb. hort. 2. t. 86.*

D. foliolis calycinis ovatis obtusis, corollis villosis: labio longissimo Buttn. cunon.

D. foliolis calycinis ovatis obtusis. Roy. lugdb.

D. angustifolia, flore ferrugineo. Bauh. pin. 244. Rin. mon. 98.

D. latifolia, flore ferrugineo. Moris. hist. 2. p. 478. f. 5. t. 8. f. 2. 3.

D. lutea non ramosa, scorzonera folio. Buxb. cent. 5. p. 25. t. 49.

Habitat in Italia, Constantinopoli. 24

Caulis strictus, lævis, sexpedalis. Folia sessilia, lanceolata, lævia, lineata, integerrima. Racemus e singulis alis supremis, erectus. Bractea lineari-lanceolata, reflexa. Pedicelli brevissimi, solitarii. Calycis folia superiora 3 approximata. Corolla ventricosa, inferne gibba, calyce paulo longior, pubescens, intus lutescens: laciniis 2 superioribus obsoletis; lateralibus acutis; infima longiore, barbata. Stamina flexuosa, absque rudimento quinsi. Stylus longitudine floris.

7. *DIGITALIS obscura. D. foliis lineari-lanceolatis integerrimis glabris basi adnatis. Jacq. hort. t. 91.*

D. hispanica angustifolia, flore nigricante. Tournef. inf. 166.

D. angustifolia hispanica. Bocc. mus. 2. p. 136. t. 98.

Habitat in Hispania. 7

Caulis suffruticosus lignosus; Rami oppositi, pauci. Folia angusto-lanceolata, basi quasi adnata, glabra, ubi non flores conferta, alias alterna. Racemi terminales. Bractea lanceolata. Calyx foliolis ovatis, acutiusculis, parvis. Corollæ cernuæ, obtusæ, intus flavescens fundo reticulato-rufa; extus pallide rufæ, præsertim dorso. Labium superius semibifidum recurvum; inferius tripartitum medio paulo longiore.

817. BIGNONIA.

Cal. 5. fidus, cyathiformis. Cor. fauce campanulata, 5. fida, subtus ventricosa. Siliqua 2. locularis. Sem. membranaceo alata.

1. *BIGNONIA Catalpa. B. foliis simplicibus cordatis ternis, caule erecto, floribus diandris. Jacq. amer. 25.*

DYDINAMIA. ANGYOSPERMIA. 87

- B.* foliis simplicibus cordatis. *Hort. cliff.*
B. arborea, foliis ovatis verticillato-ternatis, siliqua gracili-longissima. *Bromw. jam.* 264.
B. arbor, folio singulari undulato., siliquis longissimis & angustissimis. *Pluk. spec.* 5. ic. 57.
B. americana, arbor syringæ cæruleæ folio, flore purpureo. *Duham. arb.* 1. p. 104 t. 41.
B. vruco foliis flore sordide albo intus maculis purpureis & luteis adperso, siliqua longissima & angustissima. *Catesb. car.* 1. p. 49. t. 29.
Habitas in Japona, Carolina. *H*
 Arbor erecta. Folia ternata. Cirrhi nulli. Flores diandri: staminibus duobus antheriferis; reliquis duobus castratis. Variat foliis cordatis & ovatis.
13. BIGNONIA radicans. *B.* foliis pinnatis: foliolis incisis, caule geniculis radicatis. *Mill. ic.* 43. t. 65. *Duham. arb.* 1. *Sabb. hort.* 2. t. 84. *Du Roi harpk.* 1. p. 116.
 Pseudo-Gelseminum siliquosum. *Riv. mon.* 101.
 Pseudo-Apocinum hederaceum americanum, tubuloso flore phœniceo, fraxini folio. *Morif. hist.* 3. p. 612. f. 15. t. 3. f. 1.
 Gelseminum hederaceum indicum. *Corn. can.* 102. t. 103.
B. Bignonia fraxini foliis, coccineo flore minore. *Catesb. car.* 1. p. 65. t. 65.
Habitas in America. *H*

824. LANTANA.

Cal. obsolete 4. dentatus. *Stigma* uncinato. refractum. *Drupa* nucleo biloculari.

- 7. LANTANA aculeata.** *L.* foliis oppositis, caule aculeato ramoso, spicis hemisphæricis.
L. foliis oppositis petiolatis, caule aculeato. *Hort. upf.*
L. caule quadrato & aculeato, floribus e colore citrino in croceum sese mutantibus. *Medic. Aët. Palat. Vol.* 3. *phys.* p. 231.
Myrobatindum viburnifolium spinosum, floribus coccineis. *Vaill. aët.* 1722. p. 276.
Viburnum americanum odoratum, urticæ foliis latioribus, spinosum, floribus miniatis. *Pluk. atm.* 385. t. 233. f. 5.
Habitas in America calidiora. *H*

38 DIDYNA MIA. ANGYOSPERMIA.

Calis frutex altus, hispidus: angulis aculeis brevissimis, reflexis, vagis. Folia ovata, basi acuta, petiolata, acuminata, scabra. Spicæ binæ, hemisphericæ, oppositæ, intersundæ bracteis parvis deciduis. Flores flavi, adultioribus coccinei.

827. CAPRARIA.

Cal. 5. partitus. Cor. campanulata. 5. fida, acuta. Caps. 2. valvis, 2. locularis, polysperma.

1. **CAPRARIA biflora.** C. foliis alternis, floribus geminis. *Jacq. amer. 182. t. 115.*

C. foliis alternis, corollis quinquefidis *Hort. cliff.*

C. curassavica. *Herm. parad. 110. t. 110.*

Gratiolæ affinis frutescens americana, foliis agerati s. veronicae erectæ majoris. *Comm. hort. 1. p. 79. t. 40.*

Lysimachiae peruvianæ affinis americana procumbens, ononidis vernæ frutescentis folio singulari glabro. *Pluk. alm. 237. t. 98. f. 4.*

Habitat in Curassao; in Græcia Wheler. 5

Conferatur *Capraria peruviana, agerati Foliis absque pediculis. Fevill. per. 1. p. 764. t. 84.*

828. LINDERNIA.

Cal. 5. partitus. Cor. ringens labio superiore brevissimo. Stam. 2. inferiora dente terminali antheraque sublaterali. Caps. 1. locularis.

1. **LINDERNIA Pyxidaria.** L. foliis sessilibus integerrimis, pedunculis solitariis. *burg.*

Lindernia. Allion. stirp. aliq. 178. t. 5.

Pyxidaria repens annua, floiculis monopetalis unilabiatis.

Lindern. alfat. 1. p. 152. t. 1. & 2. p. 267.

Capraria gratioloides. Spec. pl.

Gratiola floribus pedunculatis, foliis ovatis crenatis. Gron. virg.

Ruellia pedunculis solitariis unifloris longitudine foliorum.

*Gron. virg. **

Habitat in Virginiae, Alfatiae, Pedemontii paludibus spon-

gis inunclatis. ☉

Flores dilute purpurei.

832. ERINUS.

Cal. 5. phyllus. *Cor.* limbus. 5. fidus, æqualis : lobis emarginatis; Labio super. brevissimo reflexo. *Caps.* 2. locularis.

1. ERINUS *alpinus*. E. floribus racemosis, foliis spatulatis. *Hall. helv. n.* 30.. *delph.*

Erinus. Sauv. monsp.

Ageratum serratum alpinum. Bauh. pin. 221.

Ageratum purpureum. Dalech. hist. 1184.

β. *Ageratum minus saxatile*, flore albo. *Barr. rar.* 23. t. 1192.

Habitat in Alpibus Helveticis, Pyrenaicis, Monspeli. 2.

Folia radicalia in cæspitem congesta, omnia lineari-spatulata, pubescentia, apice utrinque serraturis aliquot. Caules plures, simplicissimi, palmares, teretes, pubescentes, erecti, foliis alternis remotis; laterales caules steriles, decumbentes. Racemus terminalis, simplex, erectus, longitudine caulis. Flores alterni, distincti foliis similibus minoribus. Calyx campanulatus, 5-partitus, persistens laciniis oblongis, erectis. Corolla infundibulif. Tubus reflexus. Limbus æqualis (æstivatione irregularis) laciniis obcordatis, oblongis, æqualibus. Stamina 4 intra tubum. Germen subrotundum. Stylus brevissimus. Stigma capitatum auriculis 2. subrotundis oppositis.

834.. BROWALLIA.

Cal. 5. dentatus. *Cor.* limbus. 5. fidus, æqualis patens: umbilico clauso. *Antheris* duabus majoribus. *Caps.* 1. locularis.

1. BROWALLIA *demissa*. B. pedunculis unifloris. *Hort. ups.* 179. * *Kniph. cent.* 8. n. 17. *Sabb. hort.* 2. t. 100.

Habitat in America australi, Panama. ☉

835. LINNÆA.

Cal. duplex; fructus 4. phyllus; floris 5. partitus; superus. *Corolla* campan. *Bacca* ficca, trilocularis.

1. LINNÆA *borealis*. L. floribus geminatis. *Fl. lapp.*

95 DIDYNAMIA. ANGIOSPERMIA.

250. l. 12. f. 4 Fl. succ. 1. 1. * Hall. helv. n. 199.
veron. ingr. Oed. dan. 1. 3. Blackw. 1. 597. Lathw. ed.
1. 142. monsp.

Valerianella nemorosa repens. Ann. ruth.

Obolaria. Siegesb. prim. 79.

Serpillifolia. Buxb. act. 2. p. 346. l. 21.

Campanulata serpillifolia. Buxb. pin. 93. prod. 35. ed.
upf. 1720. p. 96. l. 1.

Nannularia norvegica, flore purpureo. Kyt. act. dan.
2. p. 346. f. C.

Habitat in Suecia, Sibiria, Helvetia, Russia, Canada
sylvis antiquis, muscosis, acerosis, ferilibus & rudibus. N

Sempervirens. Bacca sicca, trilocularis. Semina gemina.

Æstivatio eadem cum *Lonicera*. Florescit mense *Flavescens*
cum *Pæonia* flore pleno.

837. LIMOSELLA.

Cal. 5. fidus. Cor. 5. fida, æqualis. Stam. per paria
approximata. Caps. 1. locularis. 2. valvis, po-
lyisperma.

1. LIMOSELLA *aquatica*. L. foliis lanceolatis.

Limosella. succ. Hall. helv. n. 300. paris. Fl. dan. 1. 69.
burg.

L. foliis longe petiolaris fine spatulatis. Crantz. austr.

L. annua. Lind. als. 156. 1. 5.

Plantaginella. Hall. jen. 23. 1. 6. f. 2.

Plantaginella palustris. Buxb. pin. 190. Moris. hist. 3. p.
605. f. 15. 1. 2. f. 1.

Spergula perpusilla, lanceolatis foliis. Lef. press. 261. 1.
81.

Alene palustris repens, foliis lanceolatis. Pink. stat.
2p. 1. 74. f. 4.

Alene palustris exigua, foliolis lanceolatis. Mentz. pag. 2.
1. 7. f. 6.

Habitat in Europæ septentrionalis inundatis. O. 24

841. OROBANCHE.

Cal. 2. fidus. Cor. ringens. Caps. 1. locularis;
2. valvis, polyisperma. Glandula sub basi ger-
minis.

1. OROBANCHE *levis*, O. caule simplicissimo levi,
ramulis exsertis: long. dupl.

DIDYNAMIA. ANGIOSPERMIA. 91

O. purpurea. *Jacq. vind. 252. austr. t. 276?* videtur ob synonymum *Halleri citatum. R.*

O. caule simplici, stipulis ternatis, calyce 5-fido. *Hall. helu. n. 294. ex Linn.*

O. caule simplici cæruleo, bracteis brevibus. *Sauv. monsp.*

O. majore flore. *Bauh. pin. 88. Moris. hist. 3. f. 12. t. 16. f. 2.*

O. magna purpurea monspessulana. *Bauh. hist. 2. p. 782.*

Habitat Monspeli, in Helvetia, Germania, Austria.

2. OROBANCHE major. O. caule simplicissimo pubescente, staminibus subexsertis. *Læfl. it. * carn. pal. sil. Kniph. cent. 1. n. 67. Sabb. hort. 3. t. 11. lithu. lugd. burg.*

O. caule simplicissimo. *Fl. suec. Dalib.*

O. caule simplicissimo, supra florifero. *Gmel. sib. 3. R.*

O. caule simplici, stipula unica, calyce quadrifido. *Hall.*

helu. n. 295.

O. caule simplici, bracteis longioribus. *Sauv. monsp. 5.*

O. major caryophyllum olens. *Bauh. pin. 87.*

Orobanche. 1. *Clus. hist. 1. p. 270. Riv. irreg. mon. t. 88.*

Rapum genistæ. *Lob. ic. 2. p. 89.*

Habitat in Europæ agris, pratis siccis, parasitica imprimis radicum *Diadelphicæ. Michel monograph. De Oro-*

banche.

4. OROBANCHE cernua. O. caule simplicissimo, corollis recurvatis, bracteis ovatis corolla brevioribus caule nudiusculo. *Læfl. it.*

Habitat in Hispania, Sibiria.

5. OROBANCHE ramosa. O. caule ramoso; corollis quinquefidis. *pal. delph. burg.*

O. caule ramoso. *Dalib. parif. Gelr.*

O. caule ramoso, flore quinquepartito. *Hall. helu. n. 296.*

O. ramosa. *Bauh. pin. 88.*

Orobanche. *Cam. epit. 311. Guett. mem. de 1746. t. 9.*

Habitat in Europæ siccis.

Flos purpurascens, vel cæruleus, vel albus. *R.*

847. RUELLIA.

Cal. 5. partitus. *Cor.* subcampanulata. *Stamina* per

paria approxim. *Caps.* dentibus elasticis diffiliens.

3. RUELLIA clandestina. *R.* foliis petiolatis, pedun-

92 DIDYNAMIA. ANGIOSPERMIA.

culis longis, subdivisis nudis. *Hort. ups.* 179. * *Gouan. illustr.* 39.

R. foliis sessilibus, pedunculis trifloris. *Hort. cliff.* 308. *Roy. lugdb.* 291.

R. capsulis teretibus. *Dill. elth.* 328. t. 248. f. 320.

Habitat in Barbados.

In Horto Monspeliensi primo jam anno floruit, corollis in floribus primordialibus calyce minoribus, in reliquis floribus eodem longioribus. Gou.

853. VITEX.

Cal. 5. dentatus. *Cor.* limbus 6. fidus. *Bacca* 4. *sperma.*

1. VITEX *Agnus castus.* V. foliis digitatis serratis, spicis verticillatis.

Vitex. Cam. epit. 105. *Scorp. carn. Ludw. ed.* t. 28. *Regn. bot.*

V. foliis angustioribus cannabis modo dispositis. *Bauh. pin.* 475. *Blackw.* t. 139.

2. Vitex latiore folio. *Bauh. pin.* 475. *Kniph. cent.* 2. n. 99.

V. (*lasifolia*) foliis digitatis serratis, spicis paniculatis. *Mill. dict.* n. 2.

V. folio serrato. *Bauh. hist.* 1. p. 205.

Keuschlamm, oder europäische Müllen. *Lin. Pfl. Syst.* 4. pag. 100.

Habitat in Sicilia & Neapolis paludosis. 7

857. ACANTHUS.

Cal. bifol. bifidus. *Cor.* 1. labiata, deflexa, 3-fida. *Caps.* 2. locularis.

1. ACANTHUS *mollis.* A foliis sinuatis inermibus. *Sauv. monsp. Lugdw. ed.* t. 126. *Kniph. cent.* 9. n. 1. *Sabb. hort.* 3. t. 13. *Regn. bot.*

A. *fativus* s. *mollis virgilii.* *Bauh. pin.* 383.

Carduus Acanthus s. *Branca urfi.* *Bauh. hist.* 3. p. 75. *Blackw.* t. 89.

Habitat in Italia, Sicilia humentibus, duris. 2

2. ACANTHUS *spinofus.* A. foliis pinnatis spinosis. *Hort.* 3. t. 14.

DIDYNAMIA. ANGIOSPERMIA. 93

A. aculeatus. Bauh. pin. 383.

A. sylvestris. Dod. pempt. 719.

Habitat in Italiæ humentibus. 24

859. MELIANTHUS.

*Cal. 5. phyllus : foliolo inferiore gibbo. Petala 4 ;
nectario infra infima. Caps. 4. locularis.*

1. MELIANTHUS major. *M. stipulis solitariis petiolo
adnatis. Kniph. cent. 12. n. 70.*

M. africanus Herm. lugdb. 414. t. 415.

Habitat in Æthiopiæ succulentis. 24

*Succussa stillat pluviam nectariferam florens. Venit in
Europam 1672. ad Thom. Bartholinum ab Hermanno
missa.*

2. MELIANTHUS minor. *M. stipulis geminis distinctis,
Kniph. cent. 8. n. 68.*

M. africanus minor foetidus. Comm. rar. 4. t. 4.

M. hyssiquanensis minor foetidus. Rai. dendr. 120.

Habitat in Æthiopia. 24

CLASSIS XV.

TETRADINAMIA.
SILICULOSÆ.* *Silicula integra, nec apice emarginata.*

- | | |
|-----------------|---|
| 864. DRABA. | <i>Sil.</i> valvulis planiusculis.
<i>Stylus</i> nullus. |
| 873. LUNARIA. | <i>Silic.</i> valvulis planis pedicel-
lata. <i>Stylus</i> exsertus. |
| 863. SUBULARIA. | <i>Silic.</i> valvulis semi-ovatis.
<i>Stylus</i> brevior filicula. |
| 860. MYAGRUM. | <i>Silic.</i> valvulis concavis. <i>Sty-
lus</i> persistens. |
| 861. VELLA. | <i>Silic.</i> valvulis dissepimentum
dimidio brevioribus. |

** *Silicula emarginata apice.*

- | | |
|------------------|---|
| 868. LIBERIS. | <i>Petal</i> a duo exteriora ma-
jora. |
| 869. ALYSSUM. | <i>Filamenta</i> quædam latere
interiore dente notata.
<i>Silicula</i> bilocularis. |
| 870. CLYPEOLA. | <i>Silic.</i> orbiculata: valvulis
planis, decidua. |
| 871. PELTARIA. | <i>Silic.</i> orbiculata, compresso-
plana, non dehiscens. |
| 867. COCHLEARIA. | <i>Silic.</i> cordata: valvulis ob-
tusis, gibbis. |
| 865. LEPIDIUM. | <i>Silic.</i> cordata: valvulis ac-
ute carinatis. |
| 866. THLASPI. | <i>Silic.</i> obcordata: valvulis
marginato-carinatis. |
| 872. BISCUTELLA. | <i>Silic.</i> biloba supra infraque,
margine carinato. |

862. ANASTATICA. *Silic.* retusa: valvulis diffimento mucronato longioribus.

SILICULOSÆ.

* *Calyx clausus foliolis longitudinaliter conniventibus.*

- | | |
|-------------------|---|
| 886. RAPHANUS. | <i>Silic.</i> articulata. |
| 878. ERYSIMUM. | <i>Silic.</i> tetragona. |
| 879. CHEIRANTHUS. | <i>Siliq.</i> germine utrinque glandula notato. |
| 881. HESPERIS. | <i>Glandula</i> intra stamina breviora: <i>Petala</i> obliqua. |
| 882. ARABIS. | <i>Glandulae</i> 4 intra foliola calycina. <i>Stigma</i> simplex. |
| 884. BRASSICA. | <i>Glandula</i> 2 intra stamina breviora; 2 extra stamina longiora. |
| 883. TURRITIS. | <i>Petala</i> erecta. |
| 875. DENTARIA. | <i>Siliq.</i> valvis revolutodehiscens. |
| 874. RICOTIA. | <i>Siliq.</i> unilocularis. |

** *Calyx hians foliolis superne distantibus.*

- | | |
|------------------|---|
| 889. CRAMBE. | <i>Siliq.</i> decidua, globosa, siccobaccata. <i>Filamenta</i> 4 apice bifurca. |
| 888. ISATIS. | <i>Siliq.</i> decidua, lanceolata, monosperma. |
| 887. BUNIAS. | <i>Siliq.</i> decidua, subrotunda, muricata. |
| 890. CLEOME. | <i>Siliq.</i> dehiscens, unilocularis. |
| 876. CARDAMINE. | <i>Siliq.</i> dehiscens: valvulis revolutis. |
| 885. SINAPIS. | <i>Siliq.</i> dehiscens. <i>Cal.</i> horizontaliter patens. |
| 877. SISYMBRIUM. | <i>Siliq.</i> dehiscens: valvis recurculis. <i>Cal.</i> patulus. |

TETRADINAMIA

SILICULOSÆ.

860. MYAGRUM.

Silicula stylo conico terminata, loculo submonospermo.

1. MYAGRUM *perenne*. M. siliculis biarticulatis monospermis, foliis extrorsum sinuatis denticulatis. *delph*
M. siliquæ articulo primo stricto, altero globoso striato
Hall. helu. n. 525.

M. siliculis articulatis binatis monospermis, foliis sinuatis subdentatis. *Ger. prou.*

M. siliculis biarticulatis dispermis, foliis extrorsum sinuatis denticulatis. *Spec. pl. carn.*

M. biarticulatum. *Crantz. austr.*

Crambe foliis lanceolatis dentato-sinuatis. *Hort. cliff.*

Rapistrum monospermum. *Bauh. pin. 95. prodr. 37. t. 37. Mapp. alsat. 266. t. 266.*

Habitat in Germania. ♀

Siliculæ articulus inferior strictus abortit; superior globosus, striatus, monospermus.

3. MYAGRUM *rugosum*. M. siliculis sulcatis pilosis rugosis, foliis oblongis obtusis dentatis. *Hort. ups.* *
Habitat in Europa australi. ☉

4. MYAGRUM *hispanicum*. M. siliculis lævibus subtorosis, foliis lyratis.

Sinapi hispanicum minus, raphani folio. Tournef. inst. 127?

Habitat in Hispania. ♂

Caulis pilis raris reflexis scaber. Racemi virgati, longi. Folia lyrata, dentata, petiolata, oblonga. Siliculæ cylindricæ, læves, nec sulcatæ rugosæ aut striatæ; constantes Corpore oblongo, pedicellato, cylindrico, sterili, cui Silicula ovata insidet collo angustiore: articulo inferiore biloculari inani, superiore monospermo. Corollæ luteæ.

5. MYAGRUM *perfoliatum*. M. siliculis obcordatis sessilibus, foliis amplexicaulibus, siliculis obcordatis

TETRADYNAMIA. SILICULOSA. 97

*Hort. ups. * monsp. Kniph. cent. 10. n. 64. burg. delph. lugd.*

M. loculo fecundo conico, sterili biloculari. *Hall. hel. n. 524.*

M. monospermum latifolium. *Bauh. pin. 109. prodr. 52. t. 51. Moris. hist. 2. p. 267. f. 3. t. 21. f. antepenult.*

Habitat inter Helvetiæ, Galliæ segetes. ☉

Siliculæ loculamenta lateralibus sunt inania.

6. MYAGRUM sativum. *M. siliculis obovatis pedunculatis polyspermis. Fl. suec. parif. ged. gallob. tub. pal. sil. Kniph. cent. 11. n. 78. lugd. burg.*

Alyssum foliis sagittatis, siliquis cordiformibus. *Hall. helu. n. 489.*

Alyssum sativum. *Scop. carn.*

Camelina sativa. *Crantz. austr.*

Myagrum sylvestre. *Bauh. pin. 109. Dill. giss. p. 134.*

β. Myagrum sativum. *Bauh. pin. 109. Pollich. l. c.*

Camelina s. Myagrion. *Dod. pempt. 532.*

γ. Alysson foliis auriculatis glabris profunde dentatis. *Hall. helu. ed. 1. n. 537. ed. 2. l. c. γ. Pollich. l. c.*

Myagrum fœtidum. *Bauh. pin. 109. Lind. alsat. 49. t. 1.*

Habitat in Europa inter linum. ☉

7. MYAGRUM paniculatum. *M. siliculis lentiformibus orbiculatis punctato-rugosis. Hort. cliff. 328. Fl. suec. parif. pal. sil. Flor. dan. t. 204. carn. burg. lithu. delph. lugd.*

Rapistrum foliis amplexicaulibus auriculatis acutis. *Hall. helu. n. 522.*

Nasturtium paniculatum. *Crantz. austr. p. 15.*

Myagrum monospermum hirsutum, siliquis rotundis. *Læf. pruss. 174. t. 56.*

Myagro similis siliqua rotunda. *Bauh. pin. 109. prodr. 52. t. 52.*

Habitat in Europa juxta agros. ☉

8. MYAGRUM saxatile. *M. siliculis lentiformibus obovatis glabris: foliis petiolatis oblongis serratis scabris, caule paniculato. austr. t. 128. delph. lugd.*

Alyssum foliis scabris, radicalibus petiolatis, ovatis dentatis, caule subnudo flexuoso. *Hall. helu. n. 490.*

Alyssum alpinum. *Scop. carn.*

Cochlearia siliculis sphericis, polyspermis, glabris; foliis radicalibus petiolatis, caulinis sessilibus. *Scop. carn.*

Nasturtium saxatile. *Crantz. austr. t. 1. f. 2. Hall.*

Linæi Spec. Plant. Europ. Pars II.

G

98 TETRADYNAMIA SILICULOSA.

Cochlearia foliis radicalibus lanceolatis ferratis, caulibus
linearibus. *Sauv. onop.*

Coccolia perennis laxanlis minima. *Magn. hort.*

Alysson ap. nam, foliis circa radicem subovatis ferratis in
ordem circumactis. *Segu. ver.*

Thlaspi petraeum myagroides. *Pon. bald.* 336. n. 338.

Thlaspi 3. laxanle *Carim. epis.* 338.

Thlaspi axiale rotundifolium. *Bauh. pin.* 106. *prodr.* 4.

Thlaspi alpinum minus, capitulo rotundo. *Bauh. pin.* 107.
prodr. 48. n. 6. *Act. ex.* 197.

Thlaspi alpinum majus. *Bauh. pin.* 106. n. 6. capitulo
rotundo. *Bauh. prodr.* 49. n. 49.

Habitat in Alpibus Helvetiae, Baldi, Carniole, Mont-
peki. 24

Filamenta longiora sunt approximata, ad latus alterum
alterius defixa, apice bifurco, cujus ramus interior vici-
catus, exterior extorsum arcuatus. *Jacq. unde affinis*
Alyss.

Plantae tres nobis allatae fuerunt, an specie diversa in-
dixerim.

a. Foliis Radicalibus spatulatis scabris, obtusiusculis,
integerrimis; Caulibus lingulatis, lacinisculis, Capi-
tulis simplicibus. *Allion.*

b. Foliis Radicalibus ut in a; caulibus lingulatis, et
basi sagittatis, caule aliquot racemis. *Sequier.*

γ. Foliis Radicalibus lanceolatis, subdentatis, scabris;
Caulibus lanceolatis, subdentatis, subpetiolatis. Capi-
tulis aliquot racemis. *Sequier.*

861. V E L L A.

Silicula dissepimento valvulis duplo majore, em-
ovato.

1. *VELLA annua.* V. foliis pinnatifidis, siliculis per-
dulis. *Knap. cens.* 10. n. 93. *Trew. ic.* 10. n. 8.

Nasturtium sylvestre, crucae facie. *Bauh. pin.* 105.

Nasturtium sylvestre valesatinum. *Clus. hist.* 2. p. 130.

Habitat in Hispania. ☉

2. *VELLA Pseudo-Cytisus.* V. foliis integris obtusis
ciliatis, siliculis erectis.

Pseudo-Cytisus flore luteo. *Bauh. pin.* 390.

Cytisif facie *Alysson* frut. 2. quorundam. *Lee. ic.* 2.

49. *Bauh. hist.* 2. p. 374.

TETRADYNAMIA. SILICULOSA.

Habitat in Hispania circa oppidum Aranivez. D. Minuart. f.
Planta frutescens Foliis integris, sessilibus, hirtis,
obverse ovatis. Filamentorum paria connata, castrata.

862. ANASTATICA.

Silicula retusa, margine coronata valvulis dissepimento duplo longioribus. Stylo intermedio mucronato, obliquo, loculis 1. spermis.

1. *ANASTATICA hierochuntica. A. foliis obtusis, spicis axillaribus brevissimis, siliculis unguulatis spinosis. Jacq. hort. t. 58.*

Thlaspi Rosa de hiericho dictum. Moris. hist. 2. p. 228. f. 5. t. 25. f. 2. 3.

Rosa hierochuntea vulgo dicta. Bauh. pin. 484.

Rosa hierochuntica. Comm. hort. t. 41. Best-eyst. ant. 4. t. 1. f. 3. 3.

Habitat in littoribus Maris rubri arenosis, Palæstinæ, Cairi. ☉

2. *ANASTATICA syriaca. A. foliis acutis, spicis folio longioribus, siliculis ovatis rostratis. Jacq. austr. tab. 6.*

A. hierochuntica. Crantz. austr.

Myagrūm rostratum. Scop. carn. t. 35.

Myagrūm e Sumatra & Syria, semine spinoso simili capiti aviculæ. Zan. hist. 142.

Thlaspi hierico sylvestri affine. Bocc. mus. 135. t. 98.

Thlaspi fruticosum moravicum. Bocc. ibid.

Anastatica quæ Myagrūm e Sumatra. Gron. orient. 78.

Rosa hieri contea sylvestris. Bauh. pin. 484.

Rosa hierochuntica alia. Cam. hort. t. 42.

Habitat in Austria, Syria, Sumatra. ☉

Folia lanceolata, serrata, petiolata, subscabra. Spicæ axillares, brevissimæ, terminatæ mucrone. Petala distantia, linearia, emarginata, calycis longitudine. Siliculæ sessiles, subovatae, scabriusculæ, mucronæ inarticulato.

863. SUBULARIA.

Silicula integra, ovata: valvis ovatis, concavis, dissepimento contrariis. Stylus filicula brevior.

1. *SUBULARIA aquatica. Subularia. Fl. suec. ingr. Fl. dan. t. 35.*

300 TETRADYNAMIA. SILICULOSA.

Alyssum palustre, folio junci. *Buxb. aca.* 2. p. 369 t. 2.
f. 1.

Graminifolia aquatica, thlaspeos capitulis rotundis: fermen-
mento siliculam dirimente. *Pluk. alm.* 180. t. 188.
5. *Ped. dan.* t. 35.

Gramen junceum hybernum, thlaspeos capitulis. *Morif.*
hist. 3. f. 8. t. 10. f. ult.

Habitat in Europæ borealis inundatis, lacustribus flu-
viis. ☉

864. D R A B A .

Silicula integra, ovali-oblonga, valvis planiusculi
dissejimento parallelis. *Stylus* nullus.

* *Caule nudo.*

1. *DRABA aizoides*. *D.* scapo nudo simplici, foliis
ensiformibus carinatis ciliatis. *Jacq. austr.* 2. t. 19
delph.

D. cauliculis nudis, foliis linearibus ciliatis. *Hall. bot.*
n. 498.

Draba alpina. *Scop. carn.*

Alyssum alpinum hirsutum luteum. *T. Mill. dict.* 1. t. 2
f. 2.

Sedum alpinum hirsutum luteum. *Bauh. pin.* 284.

Leucojum luteum aizoides montanum. *Col. eccl.* 1.
p. 62.

Bursa pastoris alpina rosea lutea. *Morif. hist.* 2. p. 306. f. 1.
t. 20. f. 6.

Habitat in Alpibus Europæ. 24

Simillima D. alpinae, a qua differt: Foliis levibus,
linearibus, carinatis, ciliatis; nec ovali-oblongis linearibus
absque carina. Differt a D. ciliari Foliis in cæspitem congestis
radicalibus, carina lævi; nec Foliis alternis in ramos
expansis, carinaque ciliata, Floribus ut in D. alpina
flavis, nec albis

2. *DRABA ciliaris*. *D.* caule subnudo, foliis linearibus
margine carinaque ciliatis, petalis integris. *Ger. bot.*
t. 13. f. 1.

Habitat in Alpibus Barcinonensibus. 7

Folia radicalia in cæspitem congesta, imbricata, conferta,
lævia, carinata, distincte ciliata. Petala oblonga,
integra, alba. Stylus staminibus longior.

TETRADYNAMIA. SILICULOSA. 101

Differt a precedente foliis angustioribus glabris, petalis minime emarginatis. Culta ramos exserit foliis racemosis.

3. *DRABA alpina*. D. scapo nudo simplici, foliis lanceolatis integerrimis. *Fl. suec. Oed. dan. t.*

Habitat in Alpibus Europæ. 24

Folia radicalia omnia, patentia, lanceolata, supra pilis adpersa, more *Androsaces* disposita; minime imbricata s. linearia s. ciliata, aut supra lævia. Caulis aphyllus, adpersus raris pilis: nec monophyllus s. lævis. Petala læviter emarginata, nec integra.

Folia persæpe apice dentata. *Gouan. l. c.*

4. *DRABA verna*. D. scapis nudis, foliis subserratis.

D. scapis nudis, foliis lanceolatis subincisis. *suec. ged. gallob. austr. norv. pal. sil. Kniph. cent. 1. n. 26. Scop. carn. lithu. lugd. burg.*

D. cauliculis nudis, foliis subhirsutis, tridentatis. *Hall. helu. n. 496.*

Cochlearia siliculis lanceolatis, polyspermiis, caule paniculato nudo. *Scop. carn.*

Bursa pastoris minor, loculo oblongo. *Bauh. pin. 108.*

Alysson vulgare, polygoni folio, caule nudo. *T. Segu. veron. 1. p. 575. t. 4. f. 3.*

Pilosella minor. *Thal. harc. 84. t. 7. g. E.*

Alfine minima. *Tabern. 108. R.*

Habitat in Europæ aridis & America septentrionali. ☉

3. *DRABA pyrenaica*. D. scapo nudo, foliis cuneiformibus palmatis trilobis. *prou. carn. Crantz. austr. p. 13. t. 1. f. 5. racq. austr. t. 228. delph.*

Alysson pyrenaicum perenne minimum, foliis trifidis. *Tournef. inst. 217. Allion. pedem. t. 6. f. 1.*

Habitat in Pyrenæis. 24

Folia inferiora 5-fida, superiora 3-fida. Flores purpurei.
** Caule folioso.

6. *DRABA muralis*. D. caule ramoso, foliis ovatis sessilibus dentatis. *lithu. delph. burg. lugd.*

D. caule ramoso, foliis cordatis dentatis amplexicaulibus. *suec. gallob. pal. Hall. helu. n. 499.*

Alysson folio Veronicæ, floribus luteis. *Buxb. æd. petrop. 3. p. 273. Gmel. R.*

Myagroides subrotundis serratisque foliis, flore albo. *Barrel. ic. 816. Hall. R.*

Bursa pastoris major loculo oblongo. *Bauh. pin. 108. prod. 50. t. 50.*

102 TETRADYNAMIA. SILICULOSA.

Bursa pastoris sublongo loculo affinis pulchra planta. *Bankh.* 2. p. 938.

β. *Draba nemorosa*. *Sp. plant.*

D. minima muralis discoides. *Col. ecphr.* 1. p. 274. t. 272.

Habitat in Europæ nemoribus. ☉

Pedunculi fructiferi horizontales siliculis longiores. Flores rosei aut albi.

7. *DRABA hirta*. *D.* scapo unifolio, foliis subhirsutis, siliculis, obliquis pedicellatis. *Flor. dan.* 1. 142. *delph.*

D. stellata. *Jacq. vind. f.* 3. *obs.* 1. p. 48.

D. cauliculis subnudis, foliis ovatis tomentosis. *Hall. herb.* n. 497.

D. (austriaca) rosulis hirsutis, incanis, raro dentatis, folio ad caulem ampliori, semper dentato. *Crantz. austr.* p. 12. t. 1. f. 4.

Bursa pastoris alpina hirsuta. *Bauh. pin.* 208. *prodr.* 1. 51.

β. *Draba norvegica*: *Flor. dan.* 1. 143. *Gunn. norv.* 346. R.

Habitat in Alpibus Helveticis, Lapponicis. *D. C. Scand.*

Simillima *D. incana*, sed Caulis nudus, in medio folio unico lanceolato. Petala alba, emarginata. Siliculæ nup̄ glabræ & ovatæ; pedunculo siliqua dimidio brevior; nup̄ sessiles.

8. *DRABA incana*. *D.* foliis caulinis numerosis incanis, siliculis oblongis obliquis, sessilibus. *Fl. succ. ang. norv.* *Flor. dan.* 1. 130.

D. caule ramoso folioso, foliis dentatis. *Fl. lapp.*

Leucoium s. *Lunaria* vasculo oblongo intorto. *Pluk. del.* 215. t. 42. f. 1.

Habitat in alpibus & alpinis Europæ. ♂

*865. LEPIDIUM.

Silicula emarginata, cordata, polysperma: valvæ carinatis, dissepimento contrariis.

1. *LEPIDIUM perfoliatum*. *L.* foliis caulinis pinnatis multifidis, ramiferis cordatis amplexicaulibus integris. *Jacq. austr.* 1. 346. *Kniph. cent.* 10. n. 54.

Nasturtium spicatum persicum perfoliatum maximum. *Muskh.* *hist.* p. 294. f. 3. t. 25. f. 17.

TETRADYNAMIA. SILICULOSA. 103

Thlaspi alexandrinum. Bauh. pin. 108. prodr. 50. Bauh. hist. 2. p. 933. Rai. hist. 834.

Thlaspi verum dioscoridis. 1 Zann. hist. 193. t. 129. Rai. hist. 930.

Habitat in Persia, Syria. ☉

3. LEPIDIUM nudicaule L. scapo nudo simplicissimo ; floribus tetrandris , foliis pinnatifidis. *Lafl. it. delph.*

L. foliis filiformibus apice , pinnatifidis , caule nudo.

Sauv. monsp.

Nasturtium minimum vernalis , foliis tantum circa radicem. *Magn. monsp. t. 186.*

Habitat Monspeli, & in Hispania. ☉

Folia radicalia multa , linearia , apice pinnatifida vel dentata. Siliculæ emarginatæ. Petala æqualia. Stamina 4.

Distinctæ planta ab Iberide nudicauli.

Variat foliis omnibus cordato-ovatis longe petiolatis. Stamina sæpe 6. Gouan. R.

4. LEPIDIUM procumbens. L. foliis sinuato-pinnatifidis ; impari majore , scapo nudo , caulibus prostratis racemiferis. *burg. delph.*

L. foliis pinnatis : impari majore , caule procumbente.

Sauv. monsp.

Nasturtium foliis omnibus alternatim pinnatis. Dalib. parif. 194.

Nasturtium pumilum vernalis. Magn. monsp. 185 t. 184.

Habitat Monspeli. ☉

Folia alia ovata integra ; alia triloba , quinqueloba , lyrata. Scapi radicales ; etiam pedunculi aliquot e caule , folio longiores. Calix foliolis concavis. Petala cuneata , alba , calyce non longiora. Stamina 6. Stylus vix ullus. Silicula Drabæ vernæ.

Ex collatis , ab Hallero , Jacquino communicatis specimenibus vidit Gouanus L. procumbens a L. petræo non differre. Gouan. illustr. p. 40. R.

5. LEPIDIUM alpinum. L. foliis pinnatis integerrimis , scapo subradicato , siliculis lanceolatis mucronatis. *Jacq. austr. t. 137. Flor. dan. t. 569 ? delph.*

L. foliis pinnatis , pinnis ovatis æqualibus. *Hall. helv. n. 316.*

L. (*Halleri*) petalis calyce longe majoribus , silicula tricuspidi mucronata. *Crantz. austr. p. 5. t. 1. f. 3.*

Draba Nasturtium. Scop. carn.

104 TETRADYNAMIA. SILICULOSA.

Nasturtium alpinum tenuissime divisum. *Bauh. pin.* 105.

Bauh. hist. 2. p. 918. *Rai. hist.* 826.

Thlaspi montanum minimum: *Dalech. hist.* 1180.

Cardamine alpina 3 minima. *Clus. hist.* 2. p. 128.

Habitat in Alpibus Schneeberg, Saltzburgensibus, Tyrolensibus, Helveticis, Baldo.

Petala calyce majora. Stamina 4. *Jacq.*

6. *LEPIDIUM petraeum*. L. foliis pinnatis integerrimis, petalis emarginatis calyce minoribus. *suec. Jacq. austr.* 1. 131. *delph.*

Nasturtium foliis pinnatis, radicalibus ovato-lanceolatis, caulinis linearibus. *Hall. helv. n.* 515.

Lepidium (*Linnaei*) petalis calyce vix majoribus; siliculo stylo carente. *Crantz. austr. p.* 7. 1. 2. f. 4. 5.

Cardamine pusilla saxatilis montana discoides. *Col. ecphr.* 1. p. 274. 1. 273. *bon.*

Nasturtium pumilum vernal. *Bauh. pin.* 105.

Habitat in lapidosis Oelandiae, Angliae, Austriae. ☉

Cum *Lep.* procumbente conjungit *Gouan. R.*

7. *LEPIDIUM cardamines*. L. foliis radicalibus pinnatis; caulinis lyratis. *ad. stockh.* 1755. p. 273. 1. 8; 9.

L. foliis inferioribus alterno-pinnatis carnosius glaucis, superioribus sessilibus dentatis apice ovatis. *Arduin. spec.* 1. p. 19. 1. 18.

Habitat in Hispaniae argillosis. ♂

9. *LEPIDIUM sativum*. L. floribus tetradynamis, foliis oblongis multifidis. *suec. Ludw. ed.* 1. 166. *Regn. bot.*

L. foliis varie divisis incisisque. *Hort. cliff.*

Nasturtium sativum. *Crantz austr.*

Nasturtium hortense vulgatum. *Bauh. pin.* 103.

Nasturtium hortense. *Dod. pempt.* 771. *Blackw.* 1. 23.

β. *Nasturtium hortense crispum.* *Bauh. pin.* 104. *prodr.* 44. 1. 43.

Habitat ☉

11. *LEPIDIUM latifolium*. L. foliis ovato-lanceolatis integris ferratis. *Fl. suec. paris. Hall. helv. n.* 505. sub *Nasturtio. austr. Flor. dan.* 1. 557. *Blackw.* 1. 448. *Kniph. cent.* 3. n. 53. *Regn. bot. burg. delph.*

L. latifolium. *Bauh. pin.* 97.

L. plinii. *Dod. pempt.* 716.

L. *Aeginetæ.* *Tabernæm.* 456. *Camer. epit.* 378. R.

Habitat in Galliae, Angliae, umbrosis, succulentis 24

12. *LEPIDIUM subulatum*. L. foliis subulatis indivisis sparsis, caule suffruticoso.

TETRADYNAMIA. SILICULOSA. 105

Nasturtium fruticosum, foliis glaucis, linearibus, subhirsutis. *Hall. Helu. n. 506.*

Lepidium, capillaceo folio, fruticosum hispanicum. *Tournef. inst. 216.*

Habitat in Hispania. Læfl. 24

Radix perennis, cespitem promens foliorum & ramulorum persistentium. Caules vix spissamæ, simplices, subtomentosi, plures. Folia sparsa, subulata, linearia, acuta, supra plana, subtus semiteretia. Racemi terminales & ex alis supremum foliorum simplices. Petala albæ Siliculæ omnino *L. luteræ*. Stamina sex.

13. *LEPIDIUM graminifolium*. L. foliis linearibus: superioribus integerrimis, caule paniculato virgato floribus, hexandris. *delph.*

Thlaspi lusitanicum umbellatum, gramineo folio, flore albo. *Tournef. inst. 213.*

Habitat in Europa australi. 24

Statura L. sativi: Caulibus paniculatis, virgatis, teretibus, laevibus. Folia Radicalia prima pinnato-lyrata; Caulina inferiora serrata; superiora linearia integerrima. Flores hexandri, parvi, albi. Calyces purpurascens.

Similis, diversa tamen ab Iberide graminifolia. Adulta dejectis folia.

14. *LEPIDIUM suffruticosum*. L. foliis lanceolato-linearibus tenuibus integerrimis, caulibus suffruticosis.

Habitat in Hispania. H

Caules suffruticosi, palmares, erecti, filiformes, virentes. Folia angusto-lanceolata, integerrima, lævia, tenuia, nec carnosæ, viridia, patentia, ad gemmas sæpe plura. Corymbi parvi, albi. Petala vix calyce longiora, distantia. Stamina 6 antheris flavis. Fructus *Nasturtii*.

Differt a L. graminifolio, quod est bipedale, foliis planis: inferioribus serratis, calycibus rubris, petalis latioribus, obovatis.

16. *LEPIDIUM ruderale*. L. floribus diandris apetalis, foliis radicalibus dentato-pinnatis; ramiferis linearibus integerrimis. *Fl. succ. ingr. ged. pal. gallob. sil. Flor. dan. t. 184. delph. lithu.*

Nasturtium apetalum, foliis pinnatis, radicalium pinnis semipinnatis, caulinis simplicibus. *Hall. helu. 508.*

Iberis (runderalis) floribus vix conspicuis, foliis imis multifidis, superioribus linearibus integerrimis. *Crantz. austr.*

306 TETRADYNAMIA. SILICULOSA.

Nasturtium ruderales. Scop. carn.

Nasturtium sylvestre, osyridis folio. Bauh. pin. 103.

Nasturtium angustifolium. Fuchs. hist. 307.

Habitas in Europæ ruderatis & ad vias. ☉

L. ruderales floribus tetrandris & petalis instructum non semel vidis. Gouan. illustr. p. 40. R.

Planta apud nos nec diandria, nec apetala. Scop. Petalis etiam vidis Hallerus. R.

18. **LEPIDIUM Iberis.** *L. floribus diandris tetrapetalis, foliis inferioribus lanceolatis serratis; superioribus linearibus integerrimis. lugd.*

L. foliis lanceolato-linearibus serratis. Hort. cliff.

Iberis latiore folio. Bauh. pin. 97.

Iberis. Dod. pempt. 714.

Iberis II. Tabern. 437.

Habitas in Germania, Gallia, Italia, Sicilia secus vias. ☉

Iberis latiore folio C. B. filiculis acuminatis ab Iber. nasturtii folio C. B. filiculis emarginatis compressis distinguit. Trew. Blackw. 1. 312.

Lepidium Iberis, quod in horto bot. Senkenbergiano colimus, annua est radice, floribus diandris, filiculis emarginatis. Quare vel nova planta est vel ad L. graminifolium pertineat: Lepidium Iberis! Pollich. pal. n. 607. Reich. fl. n. 441. Nasturtium foliis imis pinnatis incis, superioribus linearibus integerrimis. Hall. helv. n. 307. Hic enim radix perennis; Flores hexandri; calyx apice perparascens; Siliculæ integræ; Folia, qualia dicta Haller. Iconem non perfectam habet Blackw. 1. 312. R.

366. THLASPI.

'Silicula emarginata, obcordata, polysperma; valvulis navicularibus, marginato-carinatis.

1. **THLASPI peregrinum.** *T. filiculis suborbiculatis, foliis lanceolatis integerrimis. Scop. carn. ed. 2.*

T. capsula cordata, peregrinum. Bauh. hist. 2. p. 913. Moris. hist. 2. p. 297. f. 3. s. 18. f. 30.

Habitas in aridis collibus Carniolæ. Scop.

2. **THLASPI arvense.** *T. filiculis orbiculatis, foliis oblongis dentatis glabris. Flor. ged. gallob. p. 11. n. 1. Indw. ed. 1. 173. burg. lithu. lugd.*

Nasturtium filiquis orbiculatis, planis, foliis oblongis dentatis, glabris. Hall. helv. n. 311.

TETRADYNAMIA. SILICULOSA. 107
Thlaspi filiculis orbiculatis compressis emarginatis. Scop.
carn.

T. majus. Tabern. 452. Hall. R.

T. arvense, filiculis latis. Bauh. pin. 105.

T. latius. Dod. pempt. 712. Blackw. t. 68.

Habitas in Europæ agris. ☉

3. THLASPI alliaceum. D. filiculis subovatis ventricosis;
foliis oblongis obtusis dentatis glabris. austr. burg.

T. allium redolens. Moris. hist. 1. p. 297. f. 3. t. 18.
f. 28.

Scorodo-Thlaspi ulyssis aldrovandi. Bauh. hist. 2. p. 932.

Habitas in Europa australi.

Antecedenti similis, sed Siliculæ ovato-ventricosæ, mar-
gine minore.

4. THLASPI saxatile. T. filiculis subrotundis, foliis
lanceolato-linearibus obtusis carnosiss. monsp. Jacq. austr.
t. 236. Kniph. cent. 1. n. 92. delph.

Nasturtium foliis pulposis, ellipticis, filiculis alatis emar-
ginatis. Hall. helv. n. 512. gœtt. 245.

Thlaspi filiculis subrotundis, foliis integris glabris lan-
ceolato-linearibus, radicalibus ovatis. Ger. prou.

T. filiculis ellipticis, foliis lanceolato-linearibus integer-
rimis. Hebenstr. in Ad. Petrop. 5. p. 330. fig. 6.

T. montanum, pingui folio, carneo flore, planta & cordata
siliqua. Barr. ic. 245.

T. parvum saxatile, flore rubente. Bauh. pin. 107.

Lithonthlaspi parvum, carnosissimo rotundo folio. Col. ecphr.
1. p. 279. t. 277. f. 2.

Habitas in Italiæ, Narbonensis, Provinciæ, Austriæ
saxosis.

Similis Iberidi saxatili.

5. THLASPI hirtum. T. filiculis subrotundis pilosis, foliis
caulinis sagittatis villosis. monsp. austr. delph.

T. capsulis hirsutis. Bauh. hist. 2. p. 921.

T. villosum, capsulis hirsutis. Bauh. pin. 106. prodr. 47.
t. 47.

Habitas in Italia, Narbona, Austria, Monspeli. ♂

Folia Radicalia ovalia, nudiuscula, subciliata. Caulis
pubescens. Folia Caulina semiamplexicaulia, villosa mol-
lissima. Calyces albi. Petala alba, obovata, integra.

6. THLASPI campestre. T. filiculis subrotundis, foliis
sagittatis dentatis incanis. succ. parif. austr. carn. tub.
col. filiculis subrotundis, delph.

108 TETRADYNAMIA. SILICULOSA.

Nasturtium foliis imis petiolatis ovatis, reliquis amplexicaulibus cordatis. *Hall. helu. n. 509.*

Thlaspi arvense, vaccariæ incano folio, majus. *Bauh. pin. 106.*

T. latifolium. *Fuchsf. hist. 306.*

T. secundum. *Tabernæm. 458. Hall. R.*

β. *Thlaspi arvense* vaccariæ incano folio minus. *Bauh. pin. 106. prodr. 47. Gouan. R.*

γ. *T. arvense* acetosæ folio. *Bauh. pin. 105. Blackw. t. 407. Hæc varietas est folio glabro, vix serrato. R.*

T. primum. *Math. 566.*

Habitat in Europæ arvis, viis argillosis, apricis. ♂

Folia radicalia sinuato-lyrata.

7. *THLASPI montanum.* *T. filiculis obcordatis, foliis glabris: radicalibus subcarnosis obovatis integerrimis; caulinis amplexicaulibus, corollis calyce majoribus. pal. heborn. Jacq. austr. t. 237. burg. delph. lugd.*

Lepidium caule erecto, foliis radicalibus sagittatis. Hall. helu. n. 518.

Thlaspi filiculis emarginatis; foliis radicalibus ovatis; caulinis amplexicaulibus. Scop. carn.

T. filiculis obverse cordatis, foliis imis spathulatis; summis amplexicaulibus sagittatis. Sauv. monsp. 121. Gouan. monsp.

T. montanum, glasti folio, minus. Bauh. pin. 106.

T. montanum, bursæ pastoris fructu. Col. ecphr. 1. p. 275. t. 276.

T. montanum 2. Clus. hist. 2. p. 131.

β. *T. perfoliatum minus. C. B. 106. ? sec. Jacquini.*

Habitat in Helvetiæ, Austriæ, Italiæ, Monspeli petrosis, sylvaticis.

Corollæ subæquales. Fructus Faciesque thlaspeos ab Iberide hanc distinguunt.

8. *THLASPI perfoliatum.* *T. filiculis obcordatis, foliis caulinis cordatis glabris subdentatis, petalis longitudine calycis, caule ramoso. Jacq. austr. t. 337. pal. burg.*

Nasturtium foliis imis ovatis, reliquis amplexicaulibus cordatis. Hall. helu. n. 510.

Thlaspi foliis radicalibus ovatis, caulinis amplexicaulibus lanceolatis. stamp. Dalib. paris.

T. arvense perfoliatum majus. Bauh. pin. 106. Barr. ic. 815.

T. alterum minus, bursæ pastoris fructu. Col. ecphr. 278. t. 276. f. 2.

TETRADYNAMIA. SILICULOSA. 109

T. oleraceum. *Tabern. ic.* 462. *hist.* 851.

Pilosella filiqua. *Cam. hort.* t. 7. f. C.

Habitat in Helvetiæ, Germaniæ, Galliæ *agris apricis*. ♂

Caules *læves*, *oleracei*. Corollæ *albæ*, *vix calyce majores*. Stamina *flore longiora*, *albæ*, *Antheris lutescentibus*. Silicula *Bursæ pastoris*.

9. *THLASPI alpestre*. *T.* filiculis obcordatis, foliis subdentatis: caulinis amplexicaulibus, petalis longitudine calycis, caule simplici. *delph.*

T. filiculis obcordatis, foliis dentatis, radicalibus lanceolatis petiolatis, caulinis sagittatis, staminibus exsertis. *Gouan. monsp.*

Lepidium caule erecto, foliis radicalibus petiolatis caulinis ovatis amplexicaulibus. *Hall. helu. n.* 519.

Thlaspi minimum. *Arduin.* 2. p. 33. t. 15. f. 2. *Hall. R.*

T. vaccariæ folio, *bursæ pastoris* filiquis. *Bauh. pin.* 105. *prodr.* 47?

T. perfoliatum minus. *Bauh. pin.* 106.

T. pumilum 3. *Clus. hist.* 2. p. 31.

Habitat in Austria, Helvetia. 24

Cum T. perfoliato jungitur a Gerardo, & Jacquino; at separatur ab Hallero. Corolla perexigua, albæ. Stamina flore longiora.

Caulis simplicissimus, rectus. In horto culta multo minor evasit, sed perennis sempervirens, & caules ex eadem radice protulit plures, alios erectos, alios diffusos. Gouan. l. c. R.

10. *THLASPI Bursa pastoris*. *T.* filiculis obcordatis, foliis radicalibus pinnatifidis. *suec. parif. ged. carn. gallob. pal. sil. Blackw. t. 5. Ludw. cat. t. 186. Regn. bot. Flor. dan. t. 729. lithu. lugd. burg.*

Nasturtium filiquis triangularibus. *Hall. helu. n.* 514.

Iberis Bursa pastoris. *Crantz. austr.*

Bursa pastoris major, folio sinuato. *Bauh. pin.* 108.

Bursa pastoris. *Fuchs. hist.* 611.

β. *Bursa pastoris media*. *Bauh. pin.* 108.

γ. *Bursa pastoris major*, folio non sinuato. *Bauh. pin.* 108.

Habitat in Europæ *cultis, ruderatis*. ☉

867. COCHLEARIA.

Silicula emarginata, turgida, scabra: valvulis gibbis obtusis.

1. *COCHLEARIA officinalis*. *C.* foliis radicalibus cor-

110 TETRADYNAMIA. SILICULOSA.

dato-subrotundis, caulinis oblongis subsinuatis. *succ. anstr. Flor. dan. t. 133. Ludw. ed. t. 133. Kniph. cent. 3. n. 30. Hall. helu. n. 503. sub Nasturtio. Regn. bot.*

C. batava. Blackw. t. 227.

C. folio subrotundo. Bauh. pin. 110.

Cochlearia. Dod. pempt. 594.

Habitat in Europæ borealis littoribus marinis. ☉ ♂

Folia Radicalia cordata, obtusa, integra crenulis 7; callosis, obsolete. Caulina amplexicaulia, dense utrinque circa caulis latus. Caulis ramosus.

2. *COCHLEARIA danica. C. foliis hastato-angulatis: omnibus deltoideis. Fl. succ. Oed. dan. t. 100.*

C. aremorica. Tournef. inst. 215.

Thlaspi hederaceum. Lob. ic. 615.

B. Cochlearia danica repens. Bauh. pin. 110. prodr 53.

γ. *C. minor erecta. Bauh. pin. 110. prodr. 53.*

Habitat in Daniæ, Sueciæ littoribus marinis. ☉ ♂

Caulis non ramosus. Stolones procumbentes. Folia & radicalia & caulina hastata f. ovata angulo utrinque basico. Petiolus basi non dentatus f. amplexicaulis. Capsulæ ovata.

3. *COCHLEARIA anglica. C. foliis omnibus ovato-lanceolatis. Oed. dan. t. 329.*

C. foliis ovato-lanceolatis sinuatis. Hudf. angl.

C. foliis radicalibus lanceolatis integerrimis, caulinis sub-sinuatis. Mill. dict.

C. folio sinuato. Bauh. pin. 110. Rai. hist. 833.

C. britannica f. anglica. Lob. ic. 294. Blackw. t. 218.

Habitat in Angliæ littoribus marinis. ♂

Folia radicalia lanceolata; caulina sessilia.

4. *COCHLEARIA grœnlandica. C. foliis reniformibus carnosissimis integerrimis.*

C. minima repens insulæ alholmiæ. Barth. act. 3. p. 143. t. 144.

C. minima. Bærh. lugdb. 2. p. 10.

Habitat in Norvegia, Islandia, Grœnlandia. ☉

Folia radicalia minima, subtus admodum convexa & carnosissima, avenia, integerrima, longius petiolata.

5. *COCHLEARIA Coronopus. C. foliis pinnatifidis, caule depresso. succ. paris. monsp. carn. gallob. pal. Flor. dan. t. 202. lugd. burg.*

Coronopus. Hall. helu. n. 502. Blackw. t. 120. Ruell. 373.

Ambrosia campestris repens. Bauh. pin. 138.

TETRADYNAMIA. SILICULOSA. II.

Pseudo-Ambrosia. Cam. epit. 596.

Habitat in Europæ apricis, nudis. ☉

6. *COCHLEARIA Armoracia. cochlearia* foliis radicalibus lanceolatis crenatis, caulinis incis. *suec. austr. herb. Blackw. t. 415. Regn. bot. lithu. burg. delph.*

Nasturtium foliis radicalibus lanceolatis crenatis, caulinis incis. *Hall. helu. n. 504.*

Raphanus rusticanus. Bauh. pin. 96.

Raphanus magna. Dod. pempt. 678.

Habitat in Europæ fossis & ad rivulos. ♀

7. *COCHLEARIA glastifolia. C. foliis* caulinis obcordato-sagittatis amplexicaulibus. *Hall. gætt. 242.*

Lepidium glastifolium. Bauh. pin. 97. Moris. hist. 2. p. 312. f. 3. t. 21. f. 3.

Lepidium annuum. Dalech. hist. 1297.

Habitat in Ratisbonæ agris. ♂

Statura Turritidis.

8. *COCHLEARIA Draba. C. foliis* lanceolatis amplexicaulibus dentatis: *Jacq. austr. t. 315. delph.*

Lepidium foliis lanceolatis amplexicaulibus dentatis. *Hort. cliff.*

Draba umbellata f. major, capitulis donata. *Bauh. pin. 109. Moris. hist. 2. p. 313. f. 3. t. 21. f. 1.*

Draba l. vulgaris. Clus. hist. 2. p. 123.

Habitat in Austria, Gallia, Italia ad versuras. ♀

Caules prostrati, parum ramosi. Folia hastata, lato-lanceolata, incana.

Silicula apice integra, acuta. Crantz. l. c. R.

868. IBERIS

Cor. irregularis: petalis 2. exterioribus majoribus. Silicula polysperma, emarginata.

1. *IBERIS semperflorens. l. frutesces*, foliis cuneiformibus integerrimis obtusis. *Kniph. cent. 12. n. 59.*

Leucoium fruticosum umbellatum persicum, foliis leucii instar sempervirentibus. *Moris. hist. 2. p. 296.*

Thlaspi latifolium polycarpon, leucii foliis. Bocc. sic. 55. t. 22. f. a. 1.

Thlaspi persicum. Riu. tetrap. 224. f. 1.

Habitat in Sicilia, Persia. ♀

2. *IBERIS sempervirens. l. frutescens*, foliis linearibus acutis integerrimis.

112 TETRADYNAMIA. SILICULOSA:

Thlaspi montanum sempervirens. Bauh. pin. 106. Riv. tetr. 224. f. 1.

Thlaspi montanum candidum. Dalech. hist. 1180.

Thlaspi creticum perenne, flore albo. Barr. ic. 214. t. 734. Habitat in Cretæ rupestribus. H

3. *IBERIS gibraltarica. I. frutescens, foliis apice dentatis.*

Thlaspidium hispanicum, ampliore flore folio crasso dentato. Dill. elth. 382. t. 287 f. 37.

Habitat ad Gibraltariam Hispaniæ. H

4. *IBERIS saxatilis. I. suffruticosa, foliis lanceolato-linearibus carnosiss acutis integerrimis ciliatis. Ger. prodelph.*

Thlaspi fruticosum thymbræ folio hirsutum. Bauh. pin. 107. Gouan. R.

Thlaspi saxatile polygalæ foliis. Bauh. pin. 107. Gou. L.

Thlaspi saxatile, vermiculato folio. Bauh. pin. 107. Garbair. 466. t. 101.

Lithonthlaspi 3 fruticosius, vermiculato acuto folio. C. ecphr. 1. p. 278. t. 277. f. 1.

Habitat in Italia, Galliæ australis aridis sabulosis. H

5. *IBERIS rotundifolia. I. herbacea, foliis ovatis: caulinis amplexicaulibus lævibus succosis. Scop. carn. t. 3 prodelph.*

Thlaspi minimum, siliculis obcordatis, foliis inferioribus ovatis petiolatis, superioribus lanceolatis amplexicaulibus. Ard. spec. 2. p. 33. t. 15. f. 1.

Thlaspi montanum, serrato cepeæ folio, flore purpurascens umbellato. Barr. ic. 848.

Thlaspi alpinum, folio rotundiore carnosissimo, flore purpurascens. Tournef. inst. 212. Scheuch. alp. 50. 143.

Lepidium caule repente foliis ovatis amplexicaulibus. Helhelu. n. 517. Allion. pedem. 27. t. 4. f. 1.

Habitat in Helvetia. 2

Caulis repens. Folia glabra, mollia: Radicalia petiolatis ovata, integerrima; caulina amplexicaulia, ovata, glabra, integra. Flores fere regulares. Silicula subquadrangula. Similis Thlaspi saxatili.

6. *IBERIS umbellata. I. herbacea, foliis lanceolatis acuminatis: inferioribus serratis; superioribus integerrimis. Regn. bot.*

Thlaspi umbellarum. Crantz. austr. p. 25.

Iberis cretica. Riv. tetr. 225.

Thla

TETRADYNAMIA. SILICULOSA. 113

Thlaspi umbellatum creticum, iberidis folio. *Bauh. pin.* 106.

Draba s. *Arabis* s. *Thlaspi candidæ*. *Dod. pempt.* 713.

Habitat in Hetruria, Hispania, Creta. ☉

Fructus hujus corymbosus hemisphæricus persistit, I. amaræ vero racemosus evadit.

7. IBERIS amara. I. herbacea, foliis lanceolatis acutis subdentatis, floribus racemosis. *pal. Kniph. cent.* 9. n. 52. *Scop. carn. delph. lugd.*

I. foliis obverse lanceolatis, floribus umbellatis. *Guett. flam.*

I. foliis dilatatis, dentatis, floribus umbellatis. *Hall. helu.* n. 520.

Thlaspi siliculis orbiculatis, bidentatis; floribus racemosis; petalis exterioribus majoribus. *Scop. carn.*

Thlaspi amarum. *Crantz. austr.*

Thlaspi umbellatum arvense, iberidis folio. *Bauh. pin.* 106.

Thlaspidium folio iberidis. *Riv. hist.* 112.

Thlaspi umbellatum arvense amarum. *Bauh. hist.* 2. p. 925.

Thlaspi amarum. *Tabern. ic.* 462.

Habitat in Helvetia, Germania. ☉

8. IBERIS linifolia. I. herbacea, foliis linearibus integerrimis, caulinis serratis, caule paniculato, corymbis hemisphæricis. *delph. burg.*

I. umbellata var. β . *Gouan. hort. Flor. monsp.*

Thlaspi umbellatum arvense iberidis folio. *Magnol. bot.* 305. *Gou. R.*

Thlaspi IV. odorato flore. *Clus. hist.* 2. p. 132. *R.*

Thlaspi creticum foliis indivisis. s. lini amarum. *Bauh. hist.* 2. p. 212. *Gouan. R.*

Thlaspi lusitanicum umbellatum, gramineo folio, flore purpurascente. *Tournef. inst.* 213. *Garid. aix.* 459. t. 105.

Habitat in Hispania, Lusitania, Monspeli.

Simillima Lepidio graminifolio. Folia *Radicalia subdentata, glabra; Caulina linearia, integerrima.* Caulis *paniculatus.* Corymbi *hemisphærici.* Siliculæ *ovatae, apice truncato bidentatae, cum in Lepidio graminifolio sint ovatae acuminatae.* Planta *adultior sæpe omnia folia dejecit.*

9. IBERIS odorata. I. herbacea, foliis linearibus superne dilatatis serratis. *delph.*

Linnaei spec. Plant. Europ. Pars II.

H

114 TETRADYNAMIA, SILICULOSA.

Thlaspi umbellatum creticum, flore albo odore, minus.
Bauh. pin. 106.

Thlaspi parvum 4. odorato flore. *Clus. hist.* 2. p. 132.
Habitat in Alpibus. Allobrogicis. ☉

11. *IBERIS nudicaulis*. I. herbacea, foliis sinuatis, calice
nudo simplici. *suec. ged. pal. sil. Flor. dan.* 323.

I. foliis pinnatis, pinnis ovatis acutis. *Hall. helv. n.* 51.

Bursa pastoris minor, foliis incis. *Bauh. pin.* 108.

Bursa pastoris minor. *Dod. pempt.* 103.

Bursa pastoris minima. *Lob. ic.* 221. R.

Bursa pastoris parva, folio glabro spisso. *Bauh. hist.* 2.
p. 937. R.

Bursa pastoris minor. *Dod. pempt.* 103.

Nasturtium petræum. *Tabern. ic.* 451.

Habitat in Europæ arenosis, sylvaticis, nudis sterilissimis. ☉

12. *IBERIS pinnata*. I. herbacea, foliis pinnatifidis. *Herm. lugd.*

Thlaspi umbellatum, nasturtii folio, monspeliacum. *Bauh. pin.* 106.

Thlaspi alterum minus umbellatum, nasturtii hortensis folio
narbonense. *Lob. ic.* 218.

Habitat in Europæ australis maritimis. ☉

Corymbus non mutatur. Calyx rubicundus. *Pet. h.*

869. A LY S S U M.

Filamenta quædam introrsum denticulo notata.
Silicula emarginata.

* *Suffruticosa.*

1. *ALYSSUM spinosum*. A racemis fenilibus spinosis
mibus nudis. *Dalib. parif.*

Thlaspi fruticosum spinosum. *Bauh. pin.* 108.

Leucoium s. *Thlaspi spinosum.* *Bauh. hist.* 2. p. 931.

Thlaspi spinosum hispanicum. *Barr. ic.* 808.

Habitat in Hispaniæ, Galliæ castris, ☉

*Spinæ e ramis olim foliosis floridisque, etiamnum passim
sibus induratis, Petala alba, integra. Stamina educta.*

2. *ALYSSUM halimifolium*. A. caulibus procumbentibus
perequantibus, foliis lanceolato-linearibus acutis integris.

Thlaspi, halimi folio, sempervirens. *Herm. lugd.* 594.

Habitat in Europæ australis aridis. ☉

3. *ALYSSUM saxatile*. A. caulibus frutescentibus
culatis, foliis lanceolatis mollissimis repandis. *Herm.*

cent. 5. n. 11.

TETRADYNAMIA

Thlaspi. incanum, leucos folio. Jacq. num. 107.
A. folio leucos incanum, non incanum. Bern. alp.
 2. f. 3.

*Allyssum crenatum laxatum, folio lanceolato incanis. Tor-
 nes. cor. 15.*

Habitat in Creta, Austria. H

Petala lutea, integra.

4. *ALYSSUM alpestre. A. caulibus infruticulosum, foliis subrotundis incanis, calycibus coloratis. Ger. prom. 352. & 13. f. 2. delph.*

Habitat in Alpibus Galieprovincia versus Italianam. 2

Folia spatulata. ovulifera. Calyces persistentes. decidui. Petala lutea, integra. Stamina basi appendiculata membranula. Stylus deciduus. Silique ovata, incana. Mart. 9.

* * *Herbacea.*

6. *ALYSSUM incanum. Alyssum caule erecto, foliis lanceolatis incanis integerrimis, floribus compositis, petalis bifidis. succ. gel. austr. pal. sil. rure illuc.*

Thlaspi fruticosum incanum. Bauh. pin. 108.

Thlaspi incanum machliniense. Clus. hist. 2. p. 132.

Habitat in Europae septentrionalioris arenis, apricis. 2. ♂

Petala alba, Silicule integra.

7. *ALYSSUM minimum. A. caulibus herbaceis, diffusis foliis linearibus tomentosis, siliculis compressis.*

A. procumbens annum, foliis lineari-lanceolatis subtomentosis. Sauv. meth.

Lunaria annua minima hispanica, folio leucos maritimi. Bern. lugd. 1. p. 6.

Habitat in Hispania; Sibiria. Georgi. u. 1. p. 224, ☉

Petala lutea, submarginata.

8. *ALYSSUM calycinum. A. caulibus herbaceis, flaminibus omnibus dentatis, calycibus persistentibus. pal. gallob. Jacq. austr. c. 338. Kniph. cens. 11. n. 7. delph. burg. lugd.*

A. Raminibus quatuor dentatis: duobus squamula longitudinali, calycibus persistentibus. Jacq. vind.

A. foliis ellipticis scabris, calycibus persistentibus. Hall. helv. n. 494.

Adyseton calycinum. Scop. carn.

Clypeola alyssoides. Crantz austr.

Clypeola siliculis bilocularibus tetraspermis. Hort. cliff. 329.

Thlaspi Alysson dictum campestre majus. Bauh. pin. 107.

H 2

116 TETRADYNAMIA SILICULOSA.

Alysson minimum. *Clus. hist.* 2. p. 133.

Alysson. *Cam. epit.* 558. f. 1.

Habitat in Austria, Gallia, Germania. ☉

Stamina 4. in medio interius dentata ut 2. ex basi exserunt squamam lanceolatam longitudine germinis. Petala minima, vix manifeste emarginata, lutea, sed antiquata alba evadunt. Silicula emarginata. Calyx persistens.

Stamina non dentata Scholler? R.

9. *ALYSSUM montanum.* A. caulibus herbaceis diffusis, foliis sublanceolatis punctato-echinatis. *Jacq. austr.* 1. 37. *Kniph. cent.* 3. n. 10. *delph. burg. lugd.*

A. foliis lanceolatis obtusis incanis, caulibus procumbentibus, radice perenni. *Hort. cliff.*

A. caulibus herbaceis, foliis punctatis scabris; siliculis orbiculatis emarginatis nudis. *Ger. prou.*

A. ramulis suffruticosis diffusis, foliis obverse lanceolatis tomentosis. *Jacq. vind.* *

A. suffruticosum foliis ellipticis scabris. *Hall. helv.* 1. 491.

Clypeola montana. *Crantz. austr.*

Adyseton montanum. *Scop. carn.*

Thlaspi luteo flore incanum, discoides, umbellatum, montanum. *Colum. eschr.* 280. 281. *Hall. R.*

Thlaspi thymi folio, utriculo furcuto mugronio, hispanicum. *Barrel. ic.* 807. *Bocc. mus.* 71. 1. 67 *Hall. R.*

Thlaspi montanum luteum. *Bauh. hist.* 2. p. 928. 1. 99 *prodr.* 49.

Thlaspi alpinum repens. *Bauh. pin.* 107.

Habitat in Helvetia, Germaniae apricis siccis. ☿

Petala lutea. Stamina 4. apice bifida; 2. basi denticulo.

10. *ALYSSUM campestre.* A. caule herbaceo, staminibus stipatis pari setarum, calycibus deciduis. *pal. gallob. delph. lugd.*

A. caulibus herbaceis, foliis punctatis scabris, siliculis orbiculatis nudis. *Ger. prou.*

Clypeola annua, siliculis bilocularibus dispermis, calycibus caducis. *Sauv. monsp.*

Alysson incanum, serpilli folio, fructu nudo. *Tournef. inst.* 217. *Hall. helv.* n. 493.

Thlaspi Alysson dictum minus, capsulis majoribus rotundis non foliatis. *Magn. monsp.* 251.

TETRADYNAMIA. SILICULOSA. 117

Thlaspi montanum incanum luteum, serpylli folio, majus. Bauh. pin. 107. prodr. 49?

Habitat in Gallia, Germania, Helvetia. ☉

Simillimum A. calycino petalis caule foliis, sed magis decumbens foliis lanceolato-ovatis. Filamenta absque dentibus, sed 2. solitaria habent utrinque setam filamentum non adnatam, at receptaculo insertam. Siliculæ ovatae, vix compressæ, magis tomentosæ.

11. **ALYSSUM clypeatum.** A. caule erecto herbaceo, filiculis sessilibus ovalibus compresso-planis, petalis acuminatis linearibus.

Lunaria filiculis subsessilibus incanis lateralibus. Roy. lugdb.

Lunaria leucoii folio, siliqua oblonga majori. Tournef. inst. 218.

Leucoium alyssoides clypeatum majus. Bauh. pin. 201.

Alysson dioscoridis. Dod. pempt. 89.

Habitat in Europa australi, Libano. ☉

Denticulus in medio filamentorum.

**** Siliculis inflatis. s. calycibus oblongis clausis.*

12. **ALYSSUM sinuatum.** A. caule herbaceo, foliis lanceolato-deltoidibus, filiculis inflatis.

A. caule herbaceo, foliis lanceolatis dentatis, filiculis inflatis. Spec. pl.

A. foliis lanceolatis repando-dentatis. Hort. cliff.

Alyssoides incanum, foliis sinuatis. Tournef. inst. 218.

Leucoium incanum, siliquis rotundis. Bauh. pin. 201.

Erica peregrina. Clus. hist. 2. p. 134.

Habitat in Hispaniæ incultis, ad vias. ☉ ♂

Petala bifida. Filamenta lateralibus obsolete dentata.

13. **ALYSSUM creticum.** A. caule fruticoso, foliis lanceolatis subdentatis tomentosis, filiculis inflatis globosis.

Alyssoides fruticosum creticum, leucoii folio incano, Tournef. cor. 15.

Leucoium luteum, utriculato semine. Alp. exot. 117. t. 118.

Habitat in Hispania, Creta. ☿

14. **ALYSSUM gemovense.** A. caule herbaceo, ramis divaricatis, foliis radicalibus obovatis subtomentosis, filiculis inflatis. *Mant. 92. Ard. spec. 2. p. - - t. 10.*

Habitat - -

Differt ab A. cretico ramis divaricatis, floribus minoribus, saturate luteis.

III. TETRADYNAMIA SILICULOSE

870. CLYPEOLA

Silicula emarginata, orbiculata, compresso-plana, decidua.

1. **CLYPEOLA** *lonchitaspis*. C. siliculis orbiculatis, orbicularibus monospermis. *burg. lugd. delph.*
Thlaspi clypeatum, serpilli folio. Bauh. pin. 107.
Lonchitaspis minimum spicatum lunatum. Col. capm. 2. 1. 281. a. 284.

Habitat in Italia, Narbonae fabulosa. ☉
Flores minuti. Siliculae orbiculatae, emarginatae, orbiculares, dispermae. Mans. 426.
Stamina omnia habent dentem supra basin, ut si velis Alyssi speciem, non repugnabo.

3. **CLYPEOLA** *maritima*. C. perennis, siliculis orbicularibus ovatis dispermis. *Sauv. monsp.*
Alyssum caulibus diffusis, folijs linearibus. Gen. p. 350.

Thlaspi montanum perenne, flore albo. Barr. ic. 5.
Thlaspi Alysson dictum maritimum. Bauh. pin. 107.
Thlaspi narbonense, centunculi angusto folio. Tab. ic. 461.

Habitat in G. Narbonensis, Hispaniae maritimis. ☿
Caulis ramosissimus, diffusus, sempervirens. Folia ovata, lanceolata, albida, non rigida. Racemi striati. Calycis decidui. Petala alba, obovata, integra, unguibus tantum atropurpurea. Antherae luteae. Silicula in parvis loculis femine unico. Filamenta edentula, quae nota ab Alyssis quidem differt, sed posset etiam Alyssi associari.

871. PELTARIA

Silicula integra, suborbiculata, compresso-plana, non dehiscens.

1. **PELTARIA** *alliacea*. *Peltaria Jacq. vind. austr. 1. 123. Scop. carn.*
Bobadsechia foliis radicalibus cordatis, caulinis amplexicaulis lanceolatis. Crantz. austr. p. 3. 1. 1. f. 1.
Clypeola perennis, folijs inferioribus petiolatis, cernitibus.

TETRADYNAMIA SILICULOSA. 119

angulatis; superioribus amplexicaulibus lanceolatis, siliculis unilocularibus. *Ard. Spec.* 26 s. 6.

Thlaspi montanum, glasti folio, majus. *Bauh. pin.* 106.

Thlaspi montanum 1. *Clus. hist.* 2. p. 130.

Habitas in alpidibus Istriae inter & Croatiae, ad arcem Sichenstein versus Schneeberg Austria. Jacq. 24.

872. BISCUTELLA.

Silicula compresso - plana, rotundata, supra infraque biloba. Cal. foliola basi gibba.

1. **BISCUTELLA auriculata.** B. calycibus nectario utrinque gibbis, siliculis in stylum coeuntibus, *delph. Kniph. cent.* 12. n. 16.

Thlaspidium biseutatum villosum, flore calcari donato. *Bauh. pin.* 107. *prodr.* 49.

Leucoium montanum, flore pedato. *Col. esphr.* 2. p. 59. s. 61.

Ion Draba alyssoides lutea angustifolia. Barr. ic. 230.

Habitas in Italia, Galloprovincia. ☉.

Hae sola siliculæ lobis in stylum coit.

2. **BISCUTELLA apula.** B. siliculis scabris, foliis lanceolatis sessilibus serratis.

B. (didyma) siliculis orbiculato-didymis a stylo divergentibus. *Spec. pl. Savv. monsp.*

Thlaspidium apulum spicatum. Tournesf. inst. 214.

Thlaspi biseutatum asperum hieracifolium majus. Bauh. pin. 107.

Thlaspi clypeatum. Clus. hist. 2. p. 133.

Ion draba alyssoides apula spicata. Col. esphr. 1. p. 283. s. 284. f. 1.

Habitas in Italia. ☉ H. V.

Planta tota. Caulis. Foliisque hirta, pilis sparsis. Caulis erectus, pedalis, ramis duobus tribusve. Folia lanceolata s. oblonga, sessilia, remotissime serrata. Siliculæ orbiculato-didymæ, scabræ.

Conf. Bisc. (didymam) siliculis orbiculatis didymis, stylo interius longitudinaliter adhaerentibus. *Gouan illustr.*

41. Confundit R. apulam, lyratam & corono-pisoliā. R.

4. **BISCUTELLA lyrata.** B. siliculis scabris, foliis lyratis. *delph.*

110 TETRADYNAMIA. SILICULOSA.

Thlaspidium raphani folio. Tournef. inst. 214.

Thlaspi bicutatum, raphani f. irionis folio. Boerh. fr. 45. t. 23.

Habitas in Hispania, Sicilia, ☉

*Folia Radicalia, lyrata (Cardamines), parva, pilis
Caulis ramosus, subaphyllus, nudus, basi hispida. Peti-
celli divaricatissimi, capillares. Siliculæ parva (B. apulæ
paulo minores); scabriusculæ, orbiculatæ.*

4. *BISCUTELLA coronopifolia. B. siliculis glabris, foliis
dentatis hirtis.*

Habitas in Hispania, Italia, Germania. ☉

*Folia Radicalia lanceolata (Coronopi), ad medium
usque profunde dentata dentibus utrinque 3. f. 4. supra
infraque pilis distinctis hirta; Caulina amplexicaulia, cor-
data-lanceolata, minus dentata, pauca. Siliculæ immo
laeves.*

5. *BISCUTELLA lævigata. B. siliculis glabris, foliis
lanceolatis serratis. Jacq. austr. 4. t. 339. delph. bar.
B. didyma. Scop. carn.*

*B. silicula lævi, foliis dentatis asperis. Hall. helv. n. 301. B.
Clypeola didyma. Crantz. austr.*

*Leucium alyssoides umbellatum montanum. Col. cap.
1. p. 283. t. 284.*

Habitas in Italia, &c. ☉

*Folia radicalia, lanceolata, petiolata, acuminata,
remate serrata serraturæ parvis acutis serratis, nuda
subscabra, margine scabra. Caulis pedalis, fere totus
aphyllus, lævis, superne corymboso-ramosus. Siliculæ lævis
magnitudinis B. auriculatæ.*

*Columnæ planta differt caule foliisque magis pilosis.
Caule folioso.*

6. *BISCUTELLA sempervirens. B. siliculis scabriusculis,
foliis lanceolatis tomentosis.*

*Thlaspidium fruticosum, leucii folio, semperflorens. Tour-
nef. 214.*

7. *Thlaspi bicutellatum latum, anchusæ folio. Boerh. fr. 39
t. 841. Boerh. mus. 267. t. 122.*

*Sorores Biscutellæ didymæ, licet ab eadem matre esse
progenatæ, possunt discesci in 4; apulam, lyratam, cor-
onopifoliam, lævigatam, quamvis convenientes siliculis
& orbiculis confectis, uti B. sempervirens.*

TETRADINAMIA. SILICULOSA. 121

Habitat in Oriente, inque Hispania. Cl. Alstræmer. F.

Caules suffruticosi, palmares, prostrati, simplices, crassitie pennæ anserinæ, apice foliosi. Folia conferta, lanceolata, dense tomentosa (*Cheiranthi incani*), canescentia, dentato-serrata margine reflexo. Caulis floriferus inter folia enatus, simplex, subaphyllus. Racemus terminalis, compositus, minus elongatus. Siliculæ magnitudine & figura *B. Apulæ*.

873. LUNARIA.

Silicula integra, elliptica, compresso plana, pedicellata; valvis dissepimento æqualibus, parallelis, planis. Cal. foliolis faccatis.

1. *LUNARIA rediviva. L. foliis alternis. ingr. ged. aust. carn. Kniph. cent. 7. n. 46. Mathusch. sil. n. 482 lithu.*

L. foliis cordatis. Roy. lugdb. Best. cyst. ver. 7. f. 2.

L. perennis, siliquis ovato-lanceolatis. Hall. helu. n. 500.

Viola lunaria major, siliqua oblonga. Bauh. pin. 203.

Viola latifolia. Lunaria odorata. Clus. hist. 1. p. 297.

Habitat in Europa septentrionaliore, in Germania. Weber. spicil. 18. 24

Radix perennis. Flores odori. Folia alterna exceptis infimis & magis hirta.

2. *LUNARIA annua. L. foliis oppositis. Fl. succ. Best. cyst. vern. 7. f. 1.*

L. major, siliqua rotundiore. Bauh. hist. 2. p. 881.

Viola latifolia. Dod. pempt. 161. Dalech. hist. 805.

Habitat in Germania. Weber. spicil. p. 18.

Radix biennis. Flores inodori. Caulis hirsutus. Planta culta habuit folia opposita; an semper? præstantior nota inquirenda.

SILIQUOSA.

875. DENTARIA.

Siliqua elastice diffiliens valvulis revolutis. Stigma emarginatum, cal. longitudinaliter connivens.

1. *DENTARIA enneaphylla. D. foliis ternis ternatis. Jacq. austr. t. 316. sil. carn.*

122 PETRADYNAMIA. SILIQUOSA.

D. foliis omnibus ternatis. Roy. lugdb.

D. triphyllos. Bauh. pin. 322, Clus. hist. 2. p. 121. n. 4.

Turritis foliis verticillatis, ternis. Scop. carn. 1. p. 317.

n. 3.
Coralloides triphyllos. Gesn. fasc. 4. n. 2. f. 4.

Ceratia plinii. Col. ecphr. 1. p. 308. t. 307.

Habitat in Austria, Italia, montibus umbrosis floribus. 2f

2. DENTARIA bulbifera. D. foliis inferioribus pinnatis, summis simplicibus. fuer. Hall. helv. n. 472. Erant. carn. fil. Flor. dan. 1. 361.

D. heptaphyllos baccifera. Bauh. pin. 322.

Barbarea. Scop. carn. 1. p. 322.

Dentaria IV. baccifera. Clus. hist. 2. p. 122.

β. Dentaria baccifera, foliis ptermicæ. Bauh. pin. 322.

Habitat in Europa australi ad radices montium umbrosas. 2f

3. DENTARIA pentaphyllos. D. foliis summis digitatis. austr. Gouan. hort. fil. delph. burg. lugd.

α. D. foliis quinis digitatis. Hall. helv. n. 469.

D. foliis imis digitatis, caulinis ternis. Scop. carn.

D. foliis pinnatis & digitatis. Ger. prou.

D. foliis summis quinatis. Gesn. fasc. 1. n. 1. f. 1.

D. heptaphyllos. Bauh. pin. 322.

D. β. heptaphyllos. Clus. hist. 2. p. 123.

β. D. foliis omnibus quinatis-digitatis. Ger. prou. 396.

Saxifraga montana, radice squamis denticulatis. Gesn. fasc.

1. n. 1. f. 1.

D. pentaphyllos, foliis mollibus. Garid. prou. 152. n. 4.

D. pentaphyllos. Bauh. pin. 322.

γ. D. pentaphyllos, foliis asperis. Fournesi. inst. 125.

Dentaria 6. pentaphyllos 1. Clus. hist. 2. p. 122.

Habitat in Alpibus Helveticis, Allobrogis, aliisque.

Plantam β. pro distincta specie vocat Gerard.

876. CARDAMINE

Siliqua elastice diffiliens, valvulis revolutis. Siliqua integrum. cal. subhians.

* Foliis simplicibus.

1. CARDAMINE bellidifolia. C. foliis simplicibus ovali integerrimis: petiolis longis. Fl. lapp. 206. n. 2. f.

TETRADYNAMIA. SILIQUOSA. 123

tub. Fl. dan. t. 20. Jacq. miscell. austr. 1. t. 17. f. 2.

C. foliis radicalibus petiolatis ovatis, caulinis ovatis, obtusis. *Hall. helu. n. 476. a.*

Arabis bellidifolia. Scop. carn.

Nasturtium alpinum, bellidis folio, minus. Bauh. pin. 105.

Habitat in Alpibus Lapponiæ, Helvetiæ, Britanniæ. 24

2. CARDAMINE asarifolia. *C.* foliis simplicibus subcordatis.

Nasturtium montanum, asari folio, Bocc. sic. 5. t. 3. Herm. par. 203. t. 203. Rai. hist. 816.

Nasturtium alpinum, palustre rotundifolium, radice repente. Moris. hist. 2. p. 224.

Habitat in Alpibus Italicis.

4. CARDAMINE petræa. *C.* foliis simplicibus oblongis dentatis. *succ. angl. Flor. dan. t. 386.*

C. petræa cambrica, nasturtii facie. Dill. elth. 70. t. 61. f. 71.

Nasturtium petræum. Pluk. alm. 261. t. 101. f. 3. Pet. herb. 50. f. 3.

Habitat in Angliæ, Arvonæ, Merviniæ, Daniæ, Sueciæ rupibus excelsis. 24

Siliquæ valvulæ a basi dehiscunt, non vero revolvuntur. Mygind.

**** Foliis ternatis.**

5. CARDAMINE resedifolia. *C.* foliis inferioribus indivisis, superioribus trilobis pinnatisque. *tub. delph.*

Nasturtium alpinum minus; resedæ folio. Bauh. pin. 104. prodr. 45. t. 45. Bauh. mus. 2. p. 41. t. 46.

Habitat in Alpibus Helveticis, Pyrenæis, in Germania.

Folia inferiora ovata, indivisa; superiora trilobo-hastata, f. subternata lobis lateralibus minoribus. Statura C. bellidifolia.

Cum C. bellidifolia conjungit b. Hallerus.

6. CARDAMINE trifolia. *C.* foliis ternatis obtusis, caule subnudo. *Jacq. austr. t. 27. carn. Hall. helu. n. 475.*

C. foliis ternatis. Fl. succ.

C. foliis ternatis crenatis, caule simplici. Fl. lapp.

C. alpina trifolia. Clus. hist. 2. p. 127.

Nasturtium alpinum trifolium. Bauh. pin. 104.

Habitat in Alpibus Helveticis, Lapponicis, &c. locis umbrosis sylvaticis.

***** Foliis pinnatis.**

8. CARDAMINE chelidonia. *C.* foliis pinnatis: foliolis quinque incis.

124 TETRADYNAMIA. SILIQUOSA.

C. glabra, chelidonii folio. *T. Barr. ic.* 156.

Nasturtium pyrenæorum aquaticum latifolium, purpurascente flore. *Herm. par.* 203. t. 204.

Habitat in Sibiria, Italia.

9. *CARDAMINE impatiens*. *Cardamine* foliis pinnatis incisiss stipulatis, floribus apetalis. *aust. carn. herb. pal. Flor. dan.* t. 735. *delph. lithu. lugd.*

C. foliis pinnatis, imis obtuse lobatis, caulinis acute dentatis, floribus apetalis. *Hall. helu. n.* 471.

Cardamine foliis pinnatis: pinnis laciniatis. *Fl. succ. Sauv. monsp.*

Cardamine foliis pinnatis: foliolis laciniatis, petalis fugacissimis. *Huds. angl.*

Sisymbrium cardamines species quædam insipida. *Bauh. hist.* 2. p. 886. *Barr. ic.* 155.

Habitat in Europæ nemoribus ad radices montium. ♂

Florem unicum cum Corolla petalis albis, majoribus quam in *parviflora* reperi 1764; diversa tamen a *C. parviflora*.

Nulla petala vidit *Pollich. l. c. & Leers. Adesse petala*, sed fugacia alii mecum asserunt. R.

10. *CARDAMINE parviflora*. *C. foliis pinnatis exstipulatis foliolis lanceolatis*, obtusis, floribus corollatis. *gallob. herb. burg. lugd.*

Nasturtium pratense, parvo flore. *Bauh. prod.* 44.

Nasturtium pratense, flosculis minimis. *Morif. prat.* 290.

Habitat in Europa. ☉

Simillima *C. impatiens*, a qua differt: Caulis brevior, minus acutangulo. Foliis non stipulatis: foliolis lanceolatis obtusis vix dentatis: nec infimis palmatis, sed ovatis glabris. Stamina corolla breviora. Siliquæ dimidio minores, erectæ in pedicellis horizontalibus.

Hallerus pro varietate *C. hirsutæ* tradit. R.

11. *CARDAMINE græca*. *C. foliis pinnatis: foliolis palmatis æqualibus petiolatis.*

Nasturtium montanum nanum, rotundo thalictri folio, pyrenæum. *Bocc. mus.* 2. p. 171. t. 166.

Sic minimo prosperi alpini affinis, siliculis latis, *Bocc. sic.* 84. t. 44. f. 2.

Habitat in Sicilia, Corsica, insulis Græciæ, in qua Germania? ☉

Caulis brevis. Folia foliolis 9, ovatis, semipalmatis, obtusis. Calyx erectus. Corolla alba.

TETRADYNAMIA. SILIQUOSA. 125

12. CARDAMINE hirsuta. C. foliis pinnatis, floribus tetrandris. *succ. noru. herborn. pal. Flor. dan. t. 148. delph. litua. lugdb.*

C. caule foliisque pinnatis, hirsutis, pinnis subrotundis. *Hall. helu. n. 472.*

Nasturtium aquaticum minus. Bauh. pin. 104.

Sisymbrium aquaticum alterum. Cam. epit. 270.

Habitat in Europa arvis, hortis, arvis.

Flores tetrandri Hallero & Linneo; semper tetradynami Oedoro & Webero. R.

13. CARDAMINE pratensis C. foliis pinnatis: foliolis radicalibus subrotundis: caulinis lanceolatis. *ingr. austr. gallob. noru. pal. sil. Kniph. cent. 12. n. 20. Blackw. t. 223. carn. Regn. bot. litua. lugd. burg.*

C. foliis pinnatis: foliolis subrotundis integris. *Fl. succ. Dalib. parif.*

C. foliis pinnatis, caule erecto. *Fl. lap.*

C. foliis pinnatis, radicalibus subrotundis, caulinis linearibus. *Hall. helu. n. 473.*

Turritis foliis pinnatis; caule terete. Scop. carn.

Nasturtium pratense, magno flore. Bauh. pin. 104.

Flos cuculi. Dod. pempt. 592.

β. *Nasturtium pratense, folio rotundiore, flore majore. Bauh. pin. 104.*

Habitat in Europæ pascuis aquosis. 24

14. CARDAMINE amara. C. foliis pinnatis, axillis stoloniferis. *ingr. gallob. noru. pal. litua. delph. lugdb.*

C. foliis pinnatis, foliolis subrotundis angulosis. *Hall. helu. n. 474. Fl. succ. parif.*

Nasturtium pyrenaicum aquaticum latifolium, purpurascens flore. Hermann. parad. 203. t. 203.

Nasturtium aquaticum majus & amarum. Bauh. pin. 204.

Habitat in Europæ septentrionalioris nemoribus. 24

Stolones non vides Hallerus. l. c. R.

877. SISYMBRIUM.

Siliqua dehiscens valvulis rectiusculis. Calyx patens.

Corolla patens.

* *Siliquis declinatis brevibus.*

1. SISYMBRIUM Nasturtium. S. siliquis declinatis, se-

126 TETRADYNAMIA. SILIQUOSA.

liis pinnatis; foliolis subcordatis. *succ. parif. gallob. pal. carn. sil. Flor. dan. t. 660. delph. lugd. litua. burg.*

S. foliis pinnatis, pinnis subrotundis, spicis sessilibus. *Hall. helu. n. 482.*

Turritis foliis pinnatis, caule angulato. *Scop. carn.*

Sisymbria Cardamine. *Fuch. hist. 723.*

Nasturtium aquaticum supinum. *Bauh. pin. 104. Tabernam. 455. Blackw. t. 260.*

Habitat in Europa & America septentrionali ad fontes, inque Oriente.

2. SISYMBRIUM Sylvestre. S. siliquis declinatis oblongo-ovatis, foliolis lanceolatis ferratis. *Oed. dan. t. 409? pal. burg. litua. lugd.*

S. siliquis declinatis, foliis pinnatis: foliolis lanceolatis ferratis. *Fl. succ. parif. monsp. lips. geb. austr. carn. sil. Kniph. cens. 12. n. 93.*

S. foliis pinnatis, pinnis dentatis distitis. *Hall. helu. 485.*

Eruca palustris, nasturtii folio, siliqua oblonga. *Bauh. pin. 98.*

Eruca quibusdam sylvestris repens, flosculo luteo. *Bauh. hist. 2. p. 366.*

Eruca sylvestris minor, luteo parvoque flore. *Bauh. pin. 98.*

Eruca palustris minor. *Tabernam. 447. R.*

Eruca sylvestris. *Fuch. hist. 263.*

Habitat in Helvetiæ, Germaniæ, Galliæ ruderatis.

Sisymbrium islandicum *Oed. fl. dan. t. 403. a. S. sylvestri certe non differre putat Gunner. R.*

3: SISYMBRIUM amphibium. S. siliquis declinatis oblongo-ovatis: foliis pinnatifidis ferratis. *ged. gallob. sil. litua.*

α. Sisymbrium palustre.

S. foliis pinnatifidis ferratis. *Fl. succ.*

S. foliis infimis capillaceis, summis pinnatifidis. *Hort. cliff. parif.*

Radicula foliis integris & pinnatifidis, petalis calyce largioribus. *Hall. helu. n. 486.*

Sisymbrium Roripa. *Scop. carn.*

Raphanus aquaticus, foliis in profundas lacinias divisis. *Bauh. pin. 97. prodr. 38. t. 38. Fl. lapp. 262.*

β. Sisymbrium aquaticum.

TETRADYNAMIA. SILIQUOSA. 127

S. foliis simplicibus dentatis serratis. *Fl. suec.*

Raphanus aquaticus, rapistri folio. *Bauh. pin.* 97.

Raphanus aquaticus alter. *Bauh. prodr.* 38. t. 38.

γ. *Sisymbrium terrestre.*

S. aquaticum, foliis variis. *Vaill. paris.* 185.

Rapistrum aquaticum. *Tabernæm.* 408. R.

Habitat in Europæ septentrionalioris aquosis.

Dux plantæ specie distinctæ hic confunduntur a *S. Lin-*
næo, quarum prima est *Radicula*, *Hall. helv.* 486, de
qua diximus; altera vero est *Radicula* foliis pinnatis,
pinnis dentatis, petalis calyce brevioribus, *Hall. helv.* n.
487. Huc pertinet *Sisymbrium palustre* *Pollich. pal.* n.
625. *Raphanus aquaticus*, foliis in profundas lacimas
divisis. *Bauh. pin.* 97. *Radicula sylvestris* s. *palustris.*
Bauh. hist. 2. p. 868. *Sisymbrium palustre repens*, par-
vo flore. *Vaill. paris.* 186. Hæc a priori differt *Petalis*
calyce brevioribus. R.

4. **SISYMBRIUM pyrenaicum.** *S.* filiquis subovatis, foliis
inferioribus lyratis; superioribus bipinnatifidis amplexi-
caulibus, stylis filiformibus.

S. foliis pinnatis: imorum pinnis oblongis, superiorum
linearibus integris, filiquis ovalibus oligospermis. *De*
la Chenal. Act. helv. 4. p. 188. t. 15.

Alyssum foliis imis subrotundis, caulinis pinnatis, fili-
formibus, amplexicaulibus. *Hall. helv.* n. 488. R.

Raphanus minimus repens luteus, foliis tenuiter divisis.
Moris. hist. 2. p. 236. f. 3. t. 7. f. 1. *Rai. hist.* 809.

Alyssum foliis pinnatis multiformibus, floribus racemosis
luteis. *Alliogr. pedem.* 40. t. 7.

Habitat in Pyrenæis, Helveticis alpibus. 24

Strylus capillaris. Silicula ovalis. Petioli amplexicaules.

5. **SISYMBRIUM ranacetifolium.** *S.* foliis pinnatis: fo-
liolis lanceolatis inciso-serratis: extimis confluentibus.
Roy. lugdb. 659. delph.

Eruca foliis pinnatis, pinnis acutissime serratis, *Hall. helv.*
n. 460. R.

Eruca ranaceti foliis. *Moris. hist.* 2. p. 231. f. 2. t. 6.
f. 19.

Eruca fruticosa, foliis *tenaceti*, indica. *Zan. hist.* 86.
t. 33.

Habitat in Sabaudia, Helvetia.

6. **SISYMBRIUM tenuifolium.** *S.* foliis integerrimis,
infimis tripinnatifidis; supremis integris. *Stamp. par.*
pal. Regn. bot. burg. lugd. delph.

128 TETRADYNAMIA. SILIQUOSA.

Eruca foliis glabris pinnatis, pinnis linearibus difformibus.
Hall. helu. n. 461.

Eruca foliis sinuato-dentatis, caule mediusculo, radice perenni. *Ger. prov. 369.*

Sinapi erucz folio. *Bauh. pin. 99.*

Eruca tenuifolia perennis. *Bauh. hist. 2. p. 861. Vall. parif. 50.*

Eruca sativa. *Fuchs. 262.*

Sinapi sylvestre. *Dodon. 707.*

Eruca sylvestris. *Blackw. t. 166.*

Habitat in Germania, Gallia, Helvetia. 2.

* * *Siliquis sessilibus axillaribus.*

7. **SISYMBRIUM supinum.** *S.* siliquis axillaribus sessilibus solitariis, foliis dentato-sinuatis. *delph. hist.*

S. siliquis omnibus pedunculatis, intimis plurimis axillaribus foliis lyrato-pinnatifidis dentatis, caule hispido. *Gouan. illustr.*

Erysimum caule decumbente, siliqua laterali subsessili. *Rey. lugdb. Dalib. parif.*

Eruca supina alba, siliqua singulari e foliorum alis erumpente. *Isnard. act. 1724. p. 295. t. 12.*

Eruca palustris major flore albo. *Bauh. hist. 2. p. 876. Gouan. R.*

Eruca. *Hall. helu. n. 459. p. Gou. R.*

Eruca inodora. *Bauh. hist. 2. p. 862. Gou. R.*

Habitat Parisiis ad agrorum margines, inque Hispania, Gotlandia, Lugduni ad Rhodanum. ☉

8. **SISYMBRIUM polyceratum.** *S.* siliquis axillaribus sessilibus subulatis aggregatis, foliis repando-dentatis. *Sauv. monsp. Jacq. hort. t. 79.*

S. siliquis sessilibus extrorsum recurvis. *Hall. helu. n. 483. L.*

Erysimum siliquis in alis foliorum sessilibus. *Hort. cliff.*

Erysimum polyceratum f. corniculatum. *Bauh. pin. 101.*

Erysimum alterum Mathioli siliquis parvis, quibusdam *Dentaria.* *Bauh. hist. 2. p. 864. R.*

Iris altera. *Dalech. hist. 653.*

Habitat in Helvetia, Italiae ruderalis. ☉

9. **SISYMBRIUM bursifolium.** *S.* racemo flexuoso, foliis lyratis, caule erecto folioso. *Amern. acad. 4. * del.*

S. foliis radicalibus ovatis, dentatis, caulinis pinnatis, pinnis linearibus, extrema maxima. *Hall. helu. n. 466. Gouan. R.*

Hesperis flore albo minimo, siliqua longa, folio profunde decussato.

TETRADYNAMIA. SILIQUOSA. 129

dentato. *Bær. lugdb.* 2. p. 20. *Dill. elth.* 179. t. 148. f. 177.

Draba palustris siliquosa major alpina, bursæ pastoris folio. *Cup. sic.* 3.

Hesperis (*dentata*) foliis dentato-pinnatifidis, caule lævi. *Sp. pl.* 2. p. 928.

Hesperis foliis multifidis. *Roy. lugdb.* 338.

Habitat in Sicilia, Italia, Helveria, Pyrenæis. ☉

* * * *Caule nudo.*

10. **SISYMBRIUM murale.** S. subacaule, foliis lanceolatis sinuato-serratis læviusculis, scapis subscabris adscendentibus. *delph.*

S. foliis lanceolatis sinuato-serratis, caule subnudo erectiusculo. *Hort. cliff. Dalib. paris.*

Eruca viminea, iberidis folio, flore luteo. *Barr. rar.* 421. t. 131.

Habitat in Gallia, Italia. ☉

Caules subscabri, adpersi pilis raris rigidis, supra basin ramosi foliosique. Folia petiolata; lanceolata, oleracea, lævia, subtus pilis aliquot vagis adpersa, dentato-serrata. Scapi longi, primum decumbentes, dein adscendentes. Calyces semipatentes raris pilis, apice fornicati. Corollæ luteæ, obtusissimæ, magnæ diametro foliorum. Siliquæ crassiusculæ, compressiusculæ, valvulis obsolete carinatis.

Paucissimis differt a S. monensi. Folia obsolete s. obiter remotissimeque serrata.

11. **SISYMBRIUM monense.** S. acaule, foliis pinnatodentatis subpilosis, scapis lævibus. *delph.*

S. foliis sinuato-pinnatis, caulibus nudis simplicibus. *Ger. prou.*

Eruca monensis laciniata, flore luteo majore. *Dill. elth.* 135. t. 111. f. 135.

Eruca perennis & saxatilis, radice crassa e rupe victoriæ. *Fournes. schol.*

Habitat in insula Mona Angliæ, rupe Victoriæ Galloprovinciæ. 4

Simillimum præcedenti magnitudine plantæ, foliorum, florum, fructuum. Radix perennis. Folia fere pinnatifida, linearia, raris pilis adpersa: pinnis remotis, obtusiusculis, sæpius septenis. Scapi erecti, læves. Calyces clausi. Petala lutea, integra.

12. **SISYMBRIUM vimineum.** S. acaule, foliis lyratis lævibus, scapis adscendentibus, floribus minutis *delph.*
Linnaei spec. Plant. Europ. Pars II. 1

130 TETRADYNAMIA. SILIQUOSA.

Eruca pumila ficula, bursæ pastoris folio. *Bocc. sic.* 19. s. 10.

Eruca, minimo flore, monspeliensis. *Bauh. hist.* 2. p. 862. f. mala.

Eruca vinealis, parvis luteis floribus. *Marif. blas.* 263. *Habitat in Sicilia, Monspeli.*

Facies duorum præcedentium, sed Folia, licet parva t digiti longitudine, lyrata omnino *Barbarea*, obtusa, lævia, Scapi fissiformes, erectiusculi. Flores lutei, *Sophia* floribus non majores. Petala erecta, obtusa, vix calyce duplo longiora.

13. *SISYMBRIUM Barrelieri*. S. caule subnudo, ramoso, foliis radicalibus runcinatis dentatis hispidis.

Eruca sylvestris minor lutea, bursæ pastoris folio *italica*. *Barr. ic.* 1016. bona. *Bauh. pin.* 98. prodr. 39.

Habitat in Hispania, Italia. Læfling. ☉

Folia radicalia, plurima, lyrata, dentata ut in *Taraxaco*, licet vix pollicaria, hispida undique setis pilicidis. Caulis erectus, rarius hispidus, ramis paucis, foliis quibus folium subulatum. Calyces læves. Petala lata, oblonga, majuscula, integra. Siliquæ oblongæ, ad semina torulosæ.

14. *SISYMBRIUM arenosum*. S. caule subfolioso ramoso, foliis lyratis rectangulo-dentatis hispidis pilis ramosis. *Fl. suec. austr. ged. herb. sil. Kniph. cent* 8. n. 89. *edelph. lithu.*

Eruca foliis scabris, semipinnatis, pinnis rectangulis, minutis dentatis. *Hall. helu.* n. 462. R.

Arabis arenosa. *Scop. carn.* s. 40. R.

Eruca cærulea in arenosis proveniens. *Bauh. pin.* 98. prodr. 40. f. mala. *Barr. ic.* 196.

Eruca sylvestris major minorque, foliis subsessilibus orbem sparsis. *Læf. pruss.* 68. s. 13. bona.

Habitat in Germania, Helvetia. ☉

Folia Radicalia plurima, lanceolato lyrata, hispida utrinque pilis ramosis bifidis s. trifidis; Caulina lanceolata, dentata, similiter hispida. Caulis hirtus pilis simplicibus. Calyces glabri. Corollæ subviolaceæ.

15. *SISYMBRIUM valentinum*. S. caule simplici erecto, superne glabro, foliis lanceolatis hispidis antice dentatis.

Eruca hirsuta, floribus albis. *Barr. rar.* 423. s. 198. 1. *Bocc. mus.* 2. p. 84. s. 80.

TETRADYNAMIA. SILIQUOSA. 131

Habitat in Regno Valentino, Madridi. I. resling. ☉

Caulis erectus, serex, pedalis, tenuior, vir ramulosus, inferne raris pilis adpersus, superne lavis. Folia Radicalia angusto-lanceolata, dentibus quatuor validis antrosum exstantibus, hispida pilis albis simplicibus longioribus & brevioribus. Caulina rariora, lanceolata-linearia, non densata. Pedunculi filiformes, flore longiores.

* * * *Folius pinnatus.*

17. **SISYMBRIUM asperum.** S. filiquis scabris, foliis pinnatifidis: pinnis lineari-lanceolatis subdentatis, corollis calyce longioribus. *Sauv. monsp. † burg. delph.*

S. foliis pinnatifidis: foliolis inequaliter laciniatis, filiquis scabris. *Ger. pro.*

S. foliis pinnatifidis subdentatis angustissimis. *Sauv. monsp. 255.*

Sinapi parvum, filiqua aspera. *Bank. pin. 499. prod. 41.*

Sinapi monspellulanum, filiqua aspera hirsuta. *Bank. hist. 2. p. 857.*

Habitat in Monspelii palustribus. ☉

Planta palmaris.

18. **SISYMBRIUM Sophia.** S. petalis calyce minoribus, foliis decomposito-pinnatis. *saec. parif. austr. carn. gallol. sub. pal. fil. Flor. dan. t. 5:8. Ludw. ec. t. 73. Blackw. t. 440. Kniph. cent. 6. n. 86. burg. lishu. lugd.*

S. pinnis & pinnulis pinnatis, laciniis linearibus subhirsutis. *Hall. helv. n. 484.*

S. corolla calyce minore, foliis multifidis linearibus. *Fl. lapp. Descurea. Guett. stamp. 2. p. 164.*

Nasturtium sylvestre tenuissime divisum. *Bank. pin. 105.*

Sophia chirurgorum. *Lob. ic. 378. R.*

Seriphium Abinthium. *Fuchs. hist. 2.*

Habitat in Europæ maceris, muris, tectis. ☉

19. **SISYMBRIUM alissimum.** S. foliis runcinatis flaccidis: foliolis sublinearibus integerrimis, pedunculis laxis. *monsp. delph.*

S. *Waltheri*, foliis hastato-pinnatis, subasperis; filiquis longissimis, striato-subquadrangulis; caule glabro. *Cranz. austr.*

Erysimum filiquis laxis, foliis hastato-pinnatis. *Hort. cliff. Sauv. monsp.*

Erysimum foliis sinapi, filiquis longissimis & strictissimis. *Buxb. cent. 3. p. 26. t. 51.*

Erysimum monspellulanum, sinapios folio. *Walsh. hort. 55. t. 24.*

132 TETRADYNAMIA. SILIQUOSA.

Habitat in Armenia, Sibiria, Austria. ☉

Calyx patentissimus, unde Sinapios affinis.

20. **SISYMBRIUM Irio.** S. foliis runcinatis dentatis nudis, caule lævi, siliquis erectis. *Jacq. austr. t. 321. burg. lugd. delph.*

Erysimum foliis lyrato pinnatis: extimo hastato. Fl. sue. paris. monsp.

Erysimum latifolium majus glabrum. Bauh. pin. 101.

Erysimum II. Tabern. hist. 840.

rio lævis apulus, erucæ folio. Col. ecphr. I. p. 264. s. 265.

Habitat in Europæ cultis. ☉

Differt a S. altissimo Foliis summis non in pinna lineares longissimas sectis; Foliis omnibus magis dentatis; Caule brevior, minus glabro; Siliquis magis erectis, nec laxis subpendulis. confer. Rai. exir. 123.

Sisymbrium Irio. Crantz. austr. p. 49. non huc pertinet, monente Jacquin. R.

21. **SISYMBRIUM Læselii.** S. foliis runcinatis acutis hirtis, caule retrorsum hispido. *ged. I. pal. Jacq. austr. t. 324. Kniph. cent. 12. n. 92. delph. lithu.*

S. foliis hastato-pinnatis sæpe hirtis, caule ramisque p. albis splendentibus, hispidis. Crantz. austr.

Sinapi erysimo tragi cognatum. Bauh. hist. 2. p. 677.

Erysimum angustifolium majus. Bauh. pin. 107.

Erysimum hirsutum, filiqua crucæ. Læf. pruss. 69. t. 1.

Erysimum foliis pinnatis: extrema pinna triangula, pinnis longis hirsutis. Hall. Gatt. 248.

Rapistrum montanum, irionis folio, macroleptoceras. Colum. ecphr. I. p. 266. s. 268.

Habitat in Borussia, Gallia, Germania, &c. ☉

23. **SISYMBRIUM barbareæ.** S. foliis simplicibus hastato-ovatis dentatis amplexicaulibus nudis. *delph.*

Sisymbrium orientale, barbareæ facie, plantaginis simile. Tournef. cor. 16.

Habitat in Oriente. Burmannus.

Sinillimum Erysimi Barbareæ. Planta insipida. Caulis angulatus. Folia radicalia integerrima, cordato-ovata.

24. **SISYMBRIUM catholicum** S. siliquis filiformibus lævibus, foliis pinnatis denticulatis caulinis.

Habitat in Hispania, Lusitania.

Caules palmares s. pedales, læves, Folia radicalia caulina pinnata, lævia, fragilia: foliolis oblongis.

TETRADYNAMIA. SILIQUOSA. 133

ratis dentatis, ad apicem petioli sæpe confluentibus. Ramus oblongus. Flores longius pedunculati. Petala flava, obovata, integra, magna. Calyx lævis. Siliquæ remotæ.

* * * * * *Foliis lanceolatis integris.*

25. **SISYMBRIUM strictissimum.** S. foliis lanceolatis dentato-ferratis caulinis. *Jacq. austr. t. 194. delph.*

Eruca caule paniculato, foliis nervosis, ovato-lanceolatis, integerrimis. Hall. helu. n. 458. R.

Draba lutea, siliquis strictissimis. Bauh. pin. 110.

Arabis quibusdam dicta planta. Cam. epit. 342.

Habitat in Helvetiæ, Italiæ, montibus asperis apricis; in Germania. Reich. flor. n. 475. Austria. 24

Folia utrinque pubescentia valdeque venosa. Antheræ defloratæ violaceæ.

878. E R Y S I M U M.

Siliqua columnaris, exacte tetraëdra. Cal. clausus.

1. **ERISYIMUM officinale.** E. siliquis spicæ adpressis, foliis runcinatis. *succ. pal. fil. Flor. dan. t. 560. Ludw. ed. t. 187. Blackw. t. 28. Regn. bot. lithu. burg. lugd.*

Erysimum, foliis pinnatis, pinnis rectangulis acutis, extrema triangulari maxima, siliquis adpressis. Hall. helu. n. 478.

Sisymbrium (officinarum Erysimum) siliquis conicis, multangulis, spicæ adpressis. Crantz. austr.

Erysimum vulgare. Bauh. pin. 100.

Verbena mas. Fuchf. hist. 592.

Habitat in Europæ ruderatis. ☉

2. **ERYSIMUM Barbaræa.** E. foliis lyratis extimo subrotundo. *succ. ingr. austr. sub. Sisymbrio. gallob. carn. pak fil. bot. burg. lugd.*

E. foliis basi pinnato dentatis apice subrotundis. Fl. lapp. Dalib. paris.

E. foliis pinnatis, pinna extrema maxima, subrotunda. Hall. helu. n. 479. R.

Barbarea femina. Tabernæm. 452. R.

Eruca lutea latifolia. f. Barbarea. Bauh. pin. 98.

β. *Sisymbrium, erucæ folio glabro, minus & procerius. Tournef. inst. 226.*

γ. *Sisymbrium orientale, barbareae facie, folio rotundo. Tournef. cor. 16.*

134 TETRADYNAMIA, SILIQUOSA:

Habitat in Europa. 24

3. *ERISYIMUM Alliaria*. E. foliis cordatis. *suec. parif. austr. Hall. helu. n. 480. carn. gallob. pal. sil. Ludw. ed. t. 77. Blakw. t. 372. Kniph. cent. 3. n. 37. Rega. bot. lithu. burg. lugd.*

Alliaria. Bauh. pin. 110. Fuchf. hist. 104. Cam. epit. 589.

Habitat in Europæ sœpibus, cultis, umbrosis. ♂ f. 24

4. *ERYSIMUM repandum*. E. foliis lanceolatis dentatis, racemis oppositifoliis, siliquis racemosis subsessilibus, corollis minutis. *Jacq. austr. t. 22.*

E. (*ramosissimum*) *ramosissimum, patens, diffusum. Crantz. austr. p. 29. n. 5. sec. Jacq. R.*

Habitat in Hispaniæ, Bohemiæ, Austriæ arvis. ☉

Statura Sisymbrii polyceratii, sed racemi oppositifolii longi. Siliquæ filiformes. Caulis angulatus. Folia argentea dentata.

5. *ERYSIMUM cheiranthoides*. E. foliis lanceolatis integerrimis, siliquis patulis. *suec. parif. ged. carn. Hall. helu. n. 477. gallob. pal. sil. Jacq. austr. t. 23. Fl. dan. 731. lugd. litua. burg.*

Turritis foliis integris lanceolatis. Guss. stamp.

Myagræ filiqua longa. Bauh. pin. 109.

Myagro similis planta siliquis longis. Bauh. hist. t. 1. 894. R.

Erysimum III. Tabernæm. p. 449. R.

Camelina, Myagræ alterum thlaspi effigie. Lob. t. 225.

Habitat ubique in Europæ arvis. ☉

Habitus simillimus Cheirantho Erysimoidi præter minores & filiquas patulas. Stigmata in hac parva & divisa, in sequenti quasi ex nodis duobus divaricata constructa.

6. *ERYSIMUM hieracifolium*. E. foliis lanceolatis serratis. *Dalib. parif. suec. pal. sil. Jacq. austr. t. 3. norv. burg. litua.*

E. (*pannonicum*) foliis lanceolatis, dentatis, flore speciei cheiranthi * *Crantz austr.*

Erysimum hieracifolium. Oed. fl. dan. t. 229. hæc præter minere mihi quidem videtur, licet a Linneo ad Cheiranthum erysimoiden allegatur. R.

Keiri sylvestre, foliis hieracii. Rupp. ien. ed. Hall. t. 77. a.

Leucoium luteum sylvestre hieracifolium. Bauh. pin. 102. prodr. 102.

TETRADYNAMIA. SILIQUOSA. 135

Leucoium sylvestre inodorum, flore parvo pallidior, Rai. hist 781.

Habitat in Gallia, Germania, Austria. ♂ 2

879. CHEIRANTHUS.

Germen utrinque denticulo glandulato. Cal. clausus : foliolis duobus basi gibbis. Semina plana.

1. **CHEIRANTHUS erysimoides.** C. foliis lanceolatis dentatis nudis, caule recto simplicissimo, siliquis tetragonis. *Fl. succ. * Allion pedem. t. 8. f. 2. noru. Dærr. nass. delph.*

C. alpinus. Jacq. austr. t. 75. R.

Erysimum cheiranthoides. Crantz austr.

Erysimum foliis integris lanceolatis margine repandodentatis. Hort. upf.

Leucoium luteum sylvestre angustifolium. Bauh. pin. 202.

Leucoium sylvestre. Clus. hist. 1. p. 299. Bauh. hist. 2. p. 873.

Eruca sylvestris angustifolia. Lob. ic. 205. Jacq. R.

Eruca angustifolia. Bauh. pin. 99. Jacq. R.

Habitat in Germania, Helvetia, Hungaria, Gallia, ☉ Radix biennis.

2. **CHEIRANTHUS alpinus.** C foliis linearibus integris subtomentosis, caule ramoso. *delph.*

C. erysimoides. Jacq. austr. t. 74. secundum Linnæum.

Hesperis foliis glabris, linearibus, lanceolatis, obiter dentatis. Hall. helu. n. 449. t. 14. Jacq. R.

Cheiranthus (sylvestris) foliis lanceolatis, asperis, flore specioso odorato. Crantz. austr. secund. Haller & Jacq. R.

Erysimum Hesperis. Scop. carn.

Erysimum foliis linearibus incanis integris. Ger. prou. 363.

Leucoium angustifolium alpinum, flore sulphureo. Tournef. inf. 222. Allion. pedem. 44. t. 9. f. 3.

Habitat in alpinis montibus Italiæ; in rupe Victoriæ Galloprov. in Austria. Jacq. 2

Simillimus C. erysimoidi, distinctus foliis incanis & caule ramoso.

Synonymon Clusii priori loco allegatum huc trahit ill. Jacquin. R.

136 TETRADYNAMIA. SILICULOSA.

3. CHEIRANTHUS *Cheiri*. C. foliis lanceolatis acutis glabris, ramis angulatis, caule fruticoso. *Stamp. parif. gelr. gallob. Dærr. nass. Ludw. ed. s. 157. Blackb. s. 179. Knorr. del. 1. s. V. 1. Kniph. cent. 2. l. 13. Regn. bot. delph. lugd.*

Leucoium foliis glabris, lanceolatis integerrimis. *Helvetu. n. 443.*

Leucoium luteum vulgare. *Bauh. pin. 202.*

Leucoium luteum. *Dod. pempt. 160.*

β. *Leucoium* luteum, magno flore. *Bauh. pin. 202.*

γ. *Leucoium* luteum, serrato folio, flore grandiori. *Bauh. pin. 202. prodr. 102.*

δ. *Leucoium* luteum, pleno flore, maius. *Bauh. pin. 202.*

ε. *Leucoium*, pleno flore, minus. *Bauh. pin. 202.*

Habitat in Angliæ, Helvetiæ, Galliæ, Hispaniæ montibus. ♂ 2 ♀

4. CHEIRANTHUS *fruticulosus*. C. foliis lanceolatis acutis glabris subserratis, caule fruticoso.

Leucoium luteum minus fruticans. *Barr. ic. 1222.*

Habitat in Hispania. Cl. Alstræmer.

Simillimus C. *Cheiri*, sed palmaris. Rami angulati. Folia lanceolata, subpestolata, acuminata, leviuscula, serraturis inferiorum paucis argutis. Flores odores minores, lutei, minus fragrantiores. Petalis profunde rotundis, obsolete emarginatis. Antheræ subcæruleæ. H. V.

5. CHEIRANTHUS *Chius*. C. foliis obovatis emarginatis, siliquis apice subulatis.

Hesperis caule ramosissimo, foliis lanceolatis densis. *Hort. clif.*

Hesperis siliquis hirsutis, flore parvo rubello. *Dill. del. 180. s. 147. f. 178.*

Leucoium thlaspeos facie. *Herm. par. 192 s. 193.*

Habitat in Chio; in Russia. Lepechin. it. 1. p. 208.

6. CHEIRANTHUS *maritimus*. C. foliis ellipticis densius nudis scabriusculis, caule diffusis scabro.

C. foliis lanceolatis acutiusculis, caule diffuso, antheris eminentibus. *Aman. acad.*

Hesperis maritima supina exigua. *Tournef. inst. 224.*

Leucoium maritimum parvum, folio crassiusculo, *Boiss. hist. 2 p. 877. Pluk. phyt. 432. f. 2.*

Leucoium minus, flore violaceo. *Barr. ic. 1227.*

TETRADYNAMIA. SILIQUOSA. 137

Habitat ad littora maris Mediterranei. ☉

Caules *ramosissimi, divaricati, diffusi, rigiduli, pedales, scabri pilis geminis adpressis. Folia ovali lanceolata, obtusa, apice subreflexa, viridia, longius petiolata, rigidula, scabriuscula: superioribus obscure subdentatis. Racemus terminalis. Calyx clausus, laevis. Petala obcordata, rubra, mox violacea: Antheræ in fauce floris.*

8. *CHEIRANTHUS incanus.* C. foliis lanceolatis integerrimis obtusis incanis, siliquis apice truncatis compressis, caule suffruticoso.

Leucoium, incano folio, hortense. Bauh. pin. 200. n. 1. 2. 4. 5. 6. 7.

Viola alba & purpurea. Lob. ic. 329.

Habitat in Hispaniæ maritimis. 24 ♀

Petala integra, Varietas albo flore foliis minus tomentosis & viridibus, sed non, ut Cheiri, rigidulis.

9. *CHEIRANTHUS fenestralis.* C. foliis conferto-capitatis recurvatis undatis, caule indiviso. *Linn. dec. 31. t. 16. Jacq. hort. t. 179.*

Habitat . . . ♂

Radix fibrosa. Caulis semipedalis, crassitie digiti, erectus, nudus. Folia in apice caulis, conferta in capitulum more brassice, lanceolata, recurvata, sessilia, obtusa, undata, incana. Racemi floriferi bracteati, erecti, altitudine caulis. Flores & Fructus Ch. incani, cui nimis affinis, vix tamen varietatem dicerem.

10. *CHEIRANTHUS annuus.* C. foliis lanceolatis subdentatis obtusis incanis, siliquis cylindricis apice acutis, caule herbaceo. *Kniph. cent. 12. n. 26.*

Leucoium incanum minus. Bauh. pin. 200.

Habitat in Europæ australis maritimis. ☉

Petala emarginata.

11. *CHEIRANTHUS littoreus.* C. foliis lanceolatis subdentatis subtomentosis subcarnosis, petalis emarginatis, siliquis tomentosis.

Leucoium maritimum angustifolium. Bauh. pin 201.

Leucoium maritimum minus. Clus. hist. 1. p. 298.

Habitat in Maris Mediterranei littoribus. ☉

Caulis pedalis, alterne ramosus, incanus. Folia alternata, lanceolata, carnosa, canaliculata, obtusa, obtuse dentata, incana. Racemi terminales. Petala obcordata, purpurascencia, basi albida. Stamina longitudine

138 TETRADYNAMIA. SILIQUOSA.

tubi. Siliquæ subulatae, angustae, aequales, folio multo longiores, tomentosae.

12. CHEIRANTHUS tristis. C. foliis linearibus subinnatis, floribus sessilibus: petalis undatis, caule fruticoso. *Læfl.*

C. foliis lanceolato, linearibus acutis glabris, siliquis teretibus. *Hort. ups. 188.*

Leucoium minus; breviori folio, obsoleto flore. *Barr. ic. 999. n. 2.*

Leucoium minus, lavendulae folio, obsoleto flore. *Barr. mus. 148. t. 111. Barr. ic. 302.*

Habitat in Hispania, Italia, Mompelii. H

Flores tristes colore & odore.

13. CHEIRANTHUS trilobus. C. foliis subulatis obtusis, calycibus laevibus, siliquis nodosis mucronatisiformibus laevibus.

Leucoium maritimum minimum hispanicum vernum, foliis erucæ. *Tournef. inst. 221?*

Leucoium maritimum minimum. *Bauh. pin. 201. prob. 103.*

Habitat in Hispania & prope insulas Stoechadum. ☉

Caules ramosi, patuli, spithamei, incani. Folia lanceolata, obtusa, unico vel duobus utrinque dentibus profundioribus. Calyces non pilosi. Corolla purpurea, minuscula. Siliquæ lineares, teretiusculæ; suturis carinatis, lateribus torulis nodosæ, apice longo subulato acuminatæ, glabræ.

16. CHEIRANTHUS sinuatus. C. foliis tomentosis obtusis subsinuatis: ramis integris, siliquis muricatis.

Leucoium maritimum, sinuato folio. *Bauh. pin. 200.*

Leucoium maritimum magnum latifolium. *Bauh. hist. p. 875.*

Habitat in Hispaniæ, Mompelii maritimis. ☉

Caulis rectus, ut tota planta, tomentoso-incanus. Folia Radicalia lato-lanceolata, alternatim dentata, obtusa; Caulina lanceolata, obtusa; canaliculata dentibus paribus duobus obtusis, in medio positis; Rames integra nec dentata. Petala incarnata, obtusa, basi exsertita. Siliquæ longissimæ, muricatæ, tomentosæ.

881. HESPERIS.

Petala oblique flexa. *Glandula* intra stamina breviora.
Siliqua stricta. *Stigma* basi bifurca apice conni-
 vente. *Cal.* clausus.

1. **HESPERIS** *tristis*. H. caule hispido ramoso patente.
austr. s. 202.

Hesperis caule hispido procumbente. *Hort. cliff.*

Hesperis montana pallida odoratissima. *Bauh. pin.* 202.

Hesperis pannonica. *Cam. hort.* s. 18.

Leucoium melancholicum. *Best. cyst.*

Habitat in Hungariæ, Austriæ arvis. ♂

Obs. Leucoium incano folio, hortense. C. B. Hesperis
siliqua articulata. Ei Hesp. peregrina, *siliquis articulatis*.
 Ei. H. *sylvestr. hirsuta*, folio *hieracii*. Ei. Hæ omnes
 a Raio & Pluknetio creduntur synonymæ.

2. **HESPERIS** *matronalis*. H. caule simplici erecto,
 foliis ovato-lanceolatis denticulatis, petalis mucrone
 emarginatis. *Dalib. parls. Kniph. cent.* 7. n. 32.
delph.

H. hortensis. *Bauh. pin.* 202. ♂

Viola matronalis. *Dod. pempt.* 161.

β. Hesperis (*sibirica*) caule simplici, foliis lanceolatis
 dentato-ferratis, petalis obtusissimis integris. *Spec. pl.*
ed. 2.

Habitat in Germania, Helvetia, Sibiria. ♂

3. **HESPERIS** *inodora*. H. caule simplici erecto, foliis
 subhastatis dentatis, petalis obtusis. *sub. Jacq. austr.*
 s. 347.

H. caule hirsuto, foliis ovato-lanceolatis, ferratis, pe-
 tiolatis, subasperis. *Hall. helv. n.* 448.

H. (*sylvestris*) caule subhispido erecto, valvis latitudine
 siliquæ. *Crantz. austr.*

H. sylvestris inodora. *Bauh. pin.* 202. *Rupp. ien. ed.*
Hall. p. 78. s. 1.

Hesperis 3. *Clus. hist.* 1. p. 297.

Habitat Viennæ, Monspelii. ♂

Simillima H. matronali, ut vix nisi a gnaro distingua-
 tur, sed Folia fere hastata basi transversali, argute den-
 tata præsertim ad basin, magis mollia, nec ita scabra.
 Olyces non colorati. Petala obtusa absque mucrone.
 Stamina non extra tubum floris prominentia.

240 TETRADYNAMIA. SILIQUOSA.

Odor *suavis*. Annuam pronuntiat *Cranz*.

5. *HESPERIS verna*. H. caule erecto ramoso, foliis cordatis amplexicaulibus serratis villosis.

Turritis annua verna, flore purpurascens. *Tournef. inst.* 124.

Leucoium minus rotundifolium, flore purpureo. *Bauh. pin.* 876.

Leucoium maritimum latifolium. *Bauh. pin.* 201. *Mor. hist.* 2. p. 231. f. 3. t. 8. f. 5.

Leucoium marinum alterum latifolium purpureo *viscag.* *Lob. ic.* 333.

Rapistrum floribus leucoli marini. *Bauh. pin.* 94. *pin.* 37. *Burf. IV.* 47.

Habitat in Galloprovinciæ maritimis. ☉

6. *HESPERIS laceræ*. H. foliis runcinatis, siliculis cuspidatis.

Cheiranthus (lacerus) foliis lacero-dentatis acuminatis, calycibus pilosis, siliculis nodosis mucronatis. *Spec. ed.* 3. *Gouan.* †

Leucoium lusitanicum purpureum, foliis elegantè dentatis. *Herm. parad.* 193. t. 193.

Habitat in Lusitania, Gallia. ☉

Folia nuda, angusta. Calyces pilosi. Flores nudi colore & odore. Siliculae nodosae tricuspidatae.

Nonne una eademque planta cum *Cheirantho* *lucida* quo supra p. 266? R.

882. A R A B I S.

Glandulae nectariferæ 4, singulae intra calycis fissuram, squamæ instar reflexæ.

1. *ARABIS alpina*. A. foliis amplexicaulibus *Cranz austr. carn. Oed. dan.* t. 62. *delph. lugd.*

A. caule ramoso, folioso, diffuso, foliis amplexicaulibus dentatis incanis. *Hall. helv. n.* 451.

A. foliis radicalibus subovatis denticulatis; caulinis ternis amplexicaulibus erectis. *Neck. gallob.*

A. foliis amplexicaulibus dentatis, caule erecto simplici. *Ry. lugdb.*

A. caule simplici, foliis ovatis utrinque dentatis. *Fl. suec.*

Draba alba siliculosæ repens. *Bauh. pin.* 109.

TETRADYNAMIA. SILIQUOSA. 141

Draba 3, succulento folio. *Clus. hist. 2. p. 125.*

6. **Arabis** foliis amplexicaulibus dentatis, caule diffuso ramoso. *Roy. lugdb.*

Draba alba siliquosa. *Bauh. pin. 109.*

Draba 2. *Clus. hist. 2. p. 24.*

Habitat in Alpibus Helveticis, austriacis, Lapponicis.

Mire varians ex loco planta, a. brevis erecta alpina;

6. *inferne procumbens magna; hinc facie diversissima.*

3. **ARABIS** *italiana*. A. foliis petiolatis lanceolatis integerrimis. *suec. parif. noru. pal. cum. fig. sil. delph. burg. lugd.*

A. foliis radicalibus petiolatis ovatis dentatis, caule subnudo ramoso. *Hall. helu. n. 452.*

A. foliis radicalibus petiolatis lanceolatis, partim integris, partim dentatis. *Crantz austr. 1. 3. f. 2.*

Turritis foliis lanceolatis integris petiolatis, ad exortum ramorum solitariis. *Hort. cliff.*

Bursæ pastoris similis siliquosa major. *Bauh. pin. 108.*

Pilosella siliquosa minor. *Thal. harc. 84. t. 7. f. D.*

Habitat in Europæ septentrionalioris sabulosis. ☉

Nectarium intra foliola calycina non vidit Pollich. R.

4. **ARABIS** *bellidifolia*. A. foliis subdentatis: radicalibus obovatis; caulinis lanceolatis. *Jacq. obs. 1. p. 22. t. 12. Gisech. ic. fasc. 1. t. 15. Jacq. austr. 3. t. 280. delph.*

Leucoium foliis levissimis, rariter dentatis, imis petiolatis ovatis, caulinis amplexicaulibus. *Hall. helu. n. 446.*

Nasturtium alpinum, bellidis folio, majus. *Bauh. pin. 105. prodr. 46. Bauh. hist. 2. p. 870.*

Plantula *Cardamines æmula*. *Clus. hist. 2. p. 129.*

Habitat in Alpibus austriacis, helveticis. 24

Simillima Arabi alpinæ, sed petala erecta angusta. Siliquæ approximata.

7. **ARABIS** *Halleri*. A. foliis caulinis sublyratis, rameis lanceolatis incisis.

Cardamine stolonifera. *Scop. carn. t. 39? R.*

Sisymbrium foliis imis barbæræ, superioribus integris dentatis. *Hall. opusc. 101. t. 1. f. 1. **

Nasturtium barbæræ foliis. *Bauh. pin. 105. prodr. 46.*

Habitat in Harcynia ad Clausthal, locis humidis; in Carniola.

Caulis erectus, angustus. Folia lyrata, glabra. Flores

842 TETRADYNAMIA. SILIQUOSA.

Nullas glandulas nectariferas habere Haller ; eas tamen vidit Scopoli. R.

9. ARABIS *pendula*. A. foliis amplexicaulibus , siliquis ancipitibus linearibus , calycibus subpilosis. *Flor. dan. t. 62.*

A. foliis amplexicaulibus dentatis , caule erecto , simplicissimo , siliquis pendulis. *Gmel. sib. 3. p. 264. t. 60. R.*
Turritis latifolia hirsuta , siliquis pendulis. *Amm. ruth. 58.*
Habitat in Sibiria , Dania. ☉

Folia scabra : Caulina minus , quam sequentis , amplexicaulia , serrata. Caulis scaber pilis raris rigidis. Pedunculi longi , filiformes , laxi. Calyces apicibus hirti. Siliquæ læves , nutantes , laxæ. Flores albi.

10. ARABIS *Turrita*. A. foliis amplexicaulibus , siliquis decurvis planis linearibus , calycibus subrugosis. *Jacq. austr. t. 11. carn. delph. burg. lugd.*

Leucoium foliis acute dentatis , amplexicaulibus , siliquis nutantibus incanis. *Hall. helv. n. 444. R.*

Arabis (*umbrosa*) siliquis compressis latis , uno versu pendulis. *Crantz. austr. p. 39. t. 3. f. 2. Jacq.*

Turritis caule simplici , foliis lanceolatis subdentatis glabris. *Sauv. monsp.*

Turritis perraro ramifera , foliis lanceolatis obscure dentatis. *Guet. stamp. Dalib. parif.*

Leucoium hesperidis folio. *Tournef. inst. 221.*

Brassica sylvestris , albido flore , nutante siliqua. *Bocc. mus. 2. p. 81. t. 72. Barr. ic. 353.*

Turrita major , plateau. *Clus. hist. 2. p. 126.*

Habitat in Helvetiæ , Hungariæ , Galliæ , Siciliæ , sepibus montosis. ☉

Folia Radicalia ovali-oblonga , crassa , incana , margine undata. Caules pedales , vix ramosi. Folia amplexicaulia , dentata : suprema serrata ; minus incana. Siliquæ lineares , compressæ , subsessiles , læves , rigida , alternæ , ad latus inferius flexæ. Flores chroleuci.

A præcedente non differre asserit Hallerus l. c. & de la Chenal in Aÿ. helv. VIII. p. 142. R.

883. TURRITIS.

Siliqua longissima , angulata. Cal. connivens ; erectus. Cor. erecta.

1. TURRITIS *glabra*. T. foliis radicalibus dentatis hispidis , caulinis integerrimis amplexicaulibus glabris. *Fl.*

TETRADYNAMIA. SILIQUOSA. 143

suec. Hall. helu. n. 455. paris. austr. pal. sil. carn. delph. burg. lithu.

T. perfoliata. Nek. gallob.

Dentaria foliis simplicibus, imis petiolatis; superioribus amplexicaulibus. *Scop. carn.*

Brassica sylvestris, foliis circa radicem cichoraceis. *Bauh. pin. 112.*

Habitat in Europæ pascuis siccis apricis. ♂

Monstrum florum viridium habet *Bækmer. Gl. lips. p. 120. R.*

2. *TURRITIS hirsuta. T.* foliis omnibus hispidis, caulinis amplexicaulibus. *suec. paris. Hall. helu. n. 456. austr. gallob. rug. pal. sil. carn. sub. Arabi. delph. burg. lithu. lugd.*

T. foliis lanceolatis, alternis, hispidis. *Scop carn. 1. p. 516. n. 1.*

Erysimo similis hirsuta, non laciniata, alba. *Bauh. pin. 101. prodr. 42. t. 42.*

Habitat in Sueciæ, Germaniæ, Angliæ, pascuis sylvesticis.

3. *TURRITIS alpina. T.* foliis radicalibus dentatis hispidis, caulinis semiamplexicaulibus. *delph.*

Habitat in Austria, Jacq. Gothlandia. ♂ 24

Statura omnino *T. hirsutæ*, sed Caulis lævis & Folia caulina omnino glabra, integerrima, sessilia vel basi parum cordata & obtusa, vixque semiamplexicaulia. Flores majores sunt: Petala alba, obtusissima, obovato-oblonga, calyce duplo longiora. Calyx compressiusculus, erectus.

Flores diversam a *T. hirsuta* dicuntur.

884. B R A S S I C A.

Cal. erectus, connivens. *Sem.* globosa. *Glandula* inter stamina breviora & pistillum, interque longiora & calycem.

* *Stylo obtusiusculo.*

1. *BRASSICA orientalis. B.* foliis cordatis amplexicaulibus glabris: radicalibus scabris integerrimis, filiquis retragonis. (*Pollich. pal. n. 639. nass. Jacq austr. t. 282. R.) delph.*

B. campestris. Mantusch. fil.

144 TETRADYNAMIA. SILIQUOSA.

B. (*Turrita*) foliis oblongis cordato-sagittatis amplexicaulibus, siliquis tetragonis caulem superantibus stric-
tissimis. *Weig. obs. bot. p. 32. R.*

Eruca foliis ovatis integerrimis amplexicaulibus. *Hall. helv. n. 457. R.*

Erysimum campestre. *Scop. carn. non synonymon. R.*

Erysimum perfoliatum. *Crantz. austr. p. 27. R.*

Brassica campestris. 1. *Clus. hist. 2. p. 127.*

B. campestris perfoliata, flore albo. *Bauh. pin 112. Segu-
ver. 1. p. 380.*

B. orientalis perfoliata, flore albo, siliqua quadrangula.
Tournef. cor. 16.

Habitas in Oriente, Monspeli, Germania; &c. ☉

Petala alba. Folia omnia glabra. Syst. Veget. 502.

2. *BRASSICA campestris.* B. radice cauleque tenui,
foliis caulinis uniformibus cordatis sessilibus. *Fl. suec.
Flor. dan. 1. 550. lith. burg. lugd.*

B. foliis oblongis cordato-sagittatis amplexicaulibus siliquis
caule brevioribus patentibus. *Weig. obs. p. 32. R.*

B. austriaca. *Jacq. austr. 1. 283? R.*

B. (*perfoliata*) caule ramisque virgatis, tenuibus, foliis
cordatis amplexicaulibus. *Crantz. austr. p. 37. an?*

B. campestris perfoliata, flore luteo. *Læs. pruss. 29. Fl.
lapp. 265.*

Habitat in agris, non argillosis, Europæ. ☉

*Radix annua. Corolla lutea. Folia radicalia lyrata,
subhispida; caulina lævia. Syst. Veg. p. 502.*

3. *BRASSICA arvensis.* B. foliis amplexicaulibus spatu-
latis repandis: summis cordatis integerrimis.

B. campestris. *Kniph. cent. 5. n. 21. R.*

B. campestris, purpureo flore. *Bauh. pin. 112. Clus. hist.
2. p. 127.*

B. sylvestris, fabariæ foliis. *Bocc. succ. 49. 1. 25. f. 3. 4*

Habitat in Europæ australis arvis humentibus. 2f

*Caulis pedalis, lævis, flexuosus, ramosus, inferne
perennans. Folia glabra, amplexicaulia, spatulata, obtu-
sissima, margine exteriori repanda, subcarnoso-oleracea:
superiora amplexicaulia, cordata, obtusa. Calyx clausus,
lævis, basi bifariam gibbus. Petala laminis patentia,
obovata, emarginata, violacea venis sanguineis. Antheræ
flavæ, eminentes.*

4. *BRASSICA alpina.* B. foliis caulinis cordato-sagittatis
amplexicaulibus, radicalibus ovatis, petalis erectis.
Pollich. pal. delph. Turritus

TETRADYNAMIA. SILIQUOSA. 145

Turritis foliis glaberrimis, radicalibus petiolatis, rotundis, incurvis, caulinis amplexicaulibus, sagittatis, obtusis.
Hall. helv. n. 454.

Turritis Brassica. *Leers fl. herb. n. ** R.

Turritis foliis radicalibus ovatis, caulinis cordato-oblongis integerrimis nitidis. *Grim. Nou. Act. A. N. C. 3. p. 77. R.*

Brassica alpina perennis. *Rupp. ed. Hall. 75. * Mapp. als. p. 42.*

Turritis pauciflora. *Grim. isen. in Nov. Act. A. N. C. III. app. p. 348. R.*

Habitas in Germania, Helvetia. *24 D. Schreber.*

Differt a B. arvensi caule angustiore, foliis tenerioribus, longioribus, radicalibus longe petiolatis; imprimis Petalis erectiusculis, albis, & flore parvo Turritidis. Figuram misit D. Grimm.

5. BRASSICA Napus. B. radice caulescente fusiformi. *Fl. suec. parif. ged. austr. gallob. fil. Ludw. ed. t. 165. Blackw. t. 224. Regn. bot.*

Napus sylvestris. *Bauh. pin. 95.*

Napus. *Dod. pempt. 674.*

β. Napus sativa. *Bauh. pin. 95.*

Habitas in arenosis maritimis Gotlandiæ, Belgii, Angliæ. ♂

Folia radicalia lyrata, glabra, Caulina amplexicaulia, cordato-oblonga, obsolete denticulata. Calyx patens, proximus Sinapios.

6. BRASSICA Rapa. B. radice caulescente orbiculari depressa carnosa. *gallob. Blackw. t. 226.*

Rapa rotunda & oblonga. *Mill. dict. n. 1. & 2. Regn. bot.*

Rapa sylvestris. *Rai. hist. 800.*

Rapa sativa rotunda. *Bauh. pin. 89.*

Rapum. *Cam. epit. 218.*

β. Rapa sativa oblonga s. semina. *Bauh. pin. 89.*

Habitas in arvis Angliæ, Belgii. ☉

7. BRASSICA oleracea. B. radice caulescente tereti carnosa.

B. (oleracea sylvestris) maritima arborea s. procerior ramosa. *Morif. hist. 2. p. 208. Rai. hist. 796. Mill. dict. n. 4.*

β. Brassica (oleracea viridis) alba vel viridis. *Bauh. pin. III.*

γ. Brassica (oleracea rubra) capitata rubra. *Bauh. pin. III.*

Linnaei spec. Plant. Europ. Pars II.

K

146 TETRADYNAMIA SILIQUOSA.

1. *Brassica (oleracea capitata)* capitata alba. *Bauh.* p. 111. *Regn. bot.*

2. *Brassica (oleracea sabauda)* alba crispa. *Bauh.* pin. 111. *Brassica undulata.* *Renealm. spec.* 113.

3. *Brassica (oleracea laciniata)* laciniata rubra. *Bauh.* pin. 2. p. 832.

4. *Brassica (oleracea selenisia)* angusto apii folio. *Bauh.* pin. 112.

5. *Brassica (oleracea fabellica)* fimbriata. *Bauh.* pin. 111. *Brassica fabellica.* *Renealm. spec.* 134. t. 133.

6. *Brassica (oleracea botrytis)* cauliflora. *Bauh.* pin. 111. *Renealm. spec.* 131. t. 133. *Cam. epit.* 253. *Mill. dict.* n. 3. *Kniph. cent.* 12. n. 18.

7. *Brassica (olerac. Napobrassica)* radice napiformi. *Tournef. inst.* 219. *Mill. dict.* n. 2.

Napo Brassica. *Bauh.* pin. 111. *prodr.* 54.

8. *Brassica (olerac. gongylodes)* gongylodes. *Bauh.* pin. 111. *Mill. dict.* n. 8.

Brassica caulo-Rapa. *Bauh.* hist. 2. p. 829.

Caulo-Rapum. *Cam. epit.* 251.

Habitat in maritimis Angliæ. ☉

* * *Eruca siliquis stylo ensiformi.*

10. **BRASSICA** *Erucastrum.* B. foliis runcinatis, caule hispido, siliquis lævibus. *pal. austr. carn. Kniph. cent.* 12. n. 17. *lichu. delph. burg.*

Sisymbrium foliis pinnato-dentatis. *Hort. cliff. par.* *Sisymbrio.*

Eruca caule hirsuto, foliis hirsutis, semipinnatis, subrotunde dentatis. *Hall. helv. n. 459. R.*

Eruca inodora. *Bauh.* hist. 2. p. 862. *Hall. R.*

Eruca sylvestris major lutea, caule aspero. *Bauh.* pin. 111.

Eruca sativa. *Fuchs. hist.* 262.

Habitat in Europæ australioris ruderatis. ☉

11. **BRASSICA** *Eruca.* B. foliis lyratis, caule hirsuto, siliquis glabris. *Blackw. 2. 242. Kniph. cent.* 9. t. 1. *austr. Regn. bot.*

Sisymbrium foliis pinnatifidis: laciniis exterioribus serratis, caule hirsuto. *Hort. cliff.*

Sinapi foliis tetragonis hirsutis, floribus venosis. *Hall. helv. n. 464. R.*

Eruca latifolia alba. *Bauh.* pin. 98.

Sinapi alterum genus. *Fuchs. hist.* 339.

Habitat in Helvetiâ, Austria. ☉

TETRADYNAMIA. SILIQUOSA. 147

1. *BRASSICA vesicaria*. B. foliis runcinatis, siliquis hispidis tectis calyce tumido.

B. foliis lyratis, caule hirsuto, siliquis retrorsum hispidis. Hort. urf.

Eruca chalapensis, caulibus & siliquis hirsutis. Moris. hist. 2. p. 228.

Habitat in Hispania. Læsting. H

Radix fusiformis. Folia lanceolata, pinnatifido-dentata, supra glabra. Caulis pilosus, ramosus. Racemi erecti, terminales. Corollæ flavæ venis saturationibus. Calyx floris cylindricus, in fructu vero ventricosus inflatus persistens evadit. Siliquæ venter calyce non longius pistillo remanente lanceolato. Differt itaque ab *Eruca* calyce, sub fructu, inflato vesicario; Foliis non lyratis, sed lanceolatis; Siliquis hispidis retrorsum nec glabris, Calycibus persistentibus, ad maturitatem fructus, nec deciduis.

885. SINAPIS.

cal. patens. Cor. ungues recti. Glandula inter stamina breviora & pistillum, interque longiora & calycem.

1. *SINAPIS arvensis*. S. siliquis multangulis toroso-turgidis rostro ancipite longioribus. Fl. suec. Hall. helv. 467. parif. ged. austr. carn. gallob. pol. sil. Flor. dan. t. 678. & t. 753. burg. lugd. lithu.

Rapistrum flore luteo. Bauh. pin. 95.

Irion. Fuchs. hist. 257.

Habitat in agris Europæ. ☉

Siliquæ læves.

4. *SINAPIS alba*. S. siliquis hispidis: rostro obliquo longissimo ensiformi. Dalib. parif. Hall. helv. n. 466. Blakw. t. 29. burg.

Sinapi album siliqua hirsuta, semine albo & rufo. Bauh. hist. 2. p. 858. R.

Sinapi apii folio. Bauh. pin. 99.

Sinapi sativum alterum. Dod. pempt. 707.

Habitat in agris Belgii, Angliæ, Galliæ. ♀

Racemi & Pedunculi striati. Siliquæ nodis 3, subcarinatis, pilis patentibus; Rostrium ensiforme, carinatum, etiam per ultimum articulum.

5. *SINAPIS nigra*. S. siliquis glabris racemo appressis,

148 TETRADYNAMIA. SILIQUOSA:

S. filiquis glabris apice tetragonis. *Fl. suec. paris. pal. Blackw. t. 446. Regn. bot. burg. lugd.*

Sinapi filiqua latiuscula glabra, semine rufo s. vulgare. *Bauh. hist. 2. p. 856.*

Sinapi rapi folio. *Bauh. pin. 99.*

Habitat in aggeribus ruderatis Europæ septentrionalioris. 24

6. SINAPIS pyrenaica. S. filiquis striatis scabris, foliis runcinatis lævibus *delph.*

Erysimum, dentis leonis folio, perenne pyrenaicum. *Tournef. inst. 228.*

Habitat in Pyrenæis D. Monnier. 24

Folia runcinata, lævia, costa albida, radicalium laciniis triangulis denticulatis: caulinarum lanceolatis. Caulis læviusculus. Flores racemosi, parvi, lutei. Siliquæ rectiusculæ, striatæ, scabræ, pilis sursum spectantibus obsoletis.

7. SINAPIS pubescens. S. filiquis pubescentibus erectis: rostro compresso, foliis lyratis villosis.

S. hirsuta, caulibus frutescentibus, foliis inferioribus lyrato-pinnatis: superioribus sessilibus. *Ard. spec. 1. p. 21. t. 9.*

Habitat in Siciliæ monte Buffambarenfi. H. Arduin.

Caulis perennans. Folia lyrata obtusa, villosa, ut tota planta. Flores lutei Petalis integris. Siliquæ erectæ, villosæ.

10. SINAPIS eruroides. S. filiquis lævibus æqualibus, foliis lyratis oblongis glabris, caule scabro. *Jacq. hort. t. 170. delph.*

Sinapi hispanicum pumilum album. *Tournef. inst. 227.*

Eruca sylvestris, flore albo, italica. *Barr. ic. 132.*

Habitat in Italia, Hispania ad vias, vineas. ☉

Caulis purpurascens, subangulatus, scaber. Folia lyrato-runcinata, acutiuscula, serrulata. Racemi corymbosi. Calyx semipatulus, pubescens. Petala alba, obtusa, vix emarginata. Filamenta purpurascencia.

11. SINAPIS hispanica. S. foliis duplicato-pinnatis: laciniis linearibus.

Sinapi hispanicum, nasturtii folio. *Tournef. inst. 227.*

Habitat in Hispania.

12. SINAPIS incana. S. filiquis racemo appressis lævibus, foliis inferioribus lyratis scabris; summis lanceolatis, caule scabro. *Jacq. hort. t. 169.*

Sinapi foliis radicalibus pinnatis, hirsutis, incanis, cotu sua filiqua breviori. *Hall. helu. n. 463. R.*

TETRADYNAMIA. SILIQUOSA. 149

Erysimum foliis subincanis, siliquis brevissimis. *Herm. prad.* 115. t. 115. *Vaill. paris.* 51.

Habitat in Gallia, Lusitania, Hispania, Helvetia. ☉

13. *SINAPIS lævigata*. S. siliquis lævibus patulis, foliis lyratis glabris; summis lanceolatis, caule lævi.

Erysimi varietas. *Herm. paris.*

Habitat in Lusitania, Hispania. ☉ ♂

886. R A P H A N U S.

Cal. clausus. *Siliqua* torosa, subarticulata, teres
Glandulæ melliferæ 2. inter stamina breviora &
pistillum, totidem inter stamina longiora &
calycem.

1. *RAPHANUS sativus*. R. siliquis teretibus torosis bilocularibus.

R. minor oblongus. *Bauh. pin.* 96. *Blackw. t.* 81.

R. major orbicularis s. rotundus. *Bauh. pin.* 96.

β. R. niger *Bauh. pin.* 66.

γ. R. chinensis annuus oleiferus.

Habitat in China. ☉ ♂

3. *RAPHANUS Raphanistrum*. R. siliquis teretibus articulatatis lævibus unilocularibus. *suec. Hall. kelu. n.* 468. *paris. monsp. Linn. diss. de Raphania. ged. austr. gallob. pal. sil. tub. lugd. burg. litua.*

Rapistrum flore albo, siliqua articulata. *Bauh. pin.* 95.

Rapistrum flore albo striato. *Bauh. hist. 2. p.* 851.

Rapistrum flore albo lineis nigris depicto. *Bauh. pin.* 95.

Rapistrum purpureum. *Tabernæm.* 407.

δ. *Lampfana* flore melino. *Tabern. p.* 408. R.

Rapistrum flore luteo. *Bauh. pin.* 95.

Habitat inter segetes Europæ.

887. B U N I A S.

Silicula decidua, tetraëdra, angulis inæqualibus
acuminatis muricata.

2. *BUNIAS Erucago*. B. siliculis tetragonis: angulis breviter cristatis. *Jacq. austr. t.* 340. *delph.*

330 TETRADYNAMIA. SILIQUOSA.

B. filiqua tetragonia, quatuor alis emarginatis. *Hall.* 526.

B. foliis antorsum sinuatis. *Hort. ups. Savu. monsp.*
Sinapi echinatum. Bauh. hist. 2. p. 236. R.

Bruca monspeliaca, filiqua quadrangula echinata. *R.*
pin. 99. prodr. 41. t. 41.

Habitat Monspeli in apris humidiusculis. ☉

4. *BUNIAS orientalis*. B. filiculis ovatis gibbis verru-
Kniph. cens. 9. A. 18.

B. filiculis obtuse triangularis, foliis inferioribus ex
nudo-natis, superioribus lanceolatis dentatis. C.
fig. 3. p. 236. n. 16 t. 57. R.

B. foliis retrosum sinuatis. *Hort ups.*

Crambe foliis pinnato-hastatis. Roy. lugdb.

Crambe orientalis, dentis leonis folio, erucaginis
Tournes. cor. 41.

Habitat in Russia. 2c

5. *BUNIAS Cakile*. C. filiculis ovatis laevibus angu-
bus. noru. Kniph. cont. 8. n. 18.

Crambe foliis succulentis linearibus pinnato-dent-
Savu. monsp.

Cakile maritima. Scop. carn.

Raphanus siliquis ovatis angulatis monospermis. R.
Savu. monsp.

Bruca maritima italica, filiqua hastata cuspidi
Bauh. pin. 99.

Cakile serapionis. Lob. ic. 223.

6. *Cakile* L. *Bruca maritima latifolia. Bauh. hist.*
868.

7. Foliis pinnatifidis linearibus carnosiss.

8. Foliis lanceolatis incisiss carnosiss.

Habitat in Europa, Africa, America maritima.
Siliquae deciduae relicta basi bifida. Varietas
stantes quidem, sed folia carnosissima, fructificationes
miles.

9. *BUNIAS balearica*. B. filiculis hispidis foliis pin-
foliolis subdentatis. Jacq. hort. t. 144. R.

B. foliis lyrate-pinnatis, foliolis crosso-pinnatifidis,
culis hispidis, rostro subulato brevioribus. C.
illustr. 45. t. 20. B.

Habitat in Balearibus. ☉ *Dau. Royen.*

Caulis pedalis, ramosissimus, pauculus, angulatus
ria. Folia petiolata, pinnata: pinnis lanceolatis,

TETRADYNAMIA. SILIQUOSA. 191
tis, obtusis, laevibus. Pedunculi racemosi, longissimi, erecti. Pedicelli floribus breviores. Calyx clausus, laevis, lanceolatus, concavus, testaceus. Corolla lutea laminis oblongis, obtusis, longitudine calycis. Siliculæ globosæ, undique echinatae spinis intermixtis confertissimis: rostro subulato, striato, silicula longiore. Cotyledones magnæ, subbipartitæ (Convolvuli) lobis subrotundis.

Similis B. myagroidi, sed Siliculæ statura B. erucaginis, verum omnino, excepto rostro, echinata.

888. I S A T I S.

Siliqua lanceolata, unilocularis, i. sperma decidua, bivalvis: valvulis navicularibus.

1. *ISATIS tinctoria*. I. foliis radicalibus crenatis: caulinis sagittatis, siliculis oblongis. *suec. paris. austr. pal. sub. Kniph. cent. 12. n. 64. Regn. bot. delph. burg.*

I. foliis radicalibus, petiolatis, ovatis, caulinis amplexicaulibus. *Halt. helv. n. 523.*

I. sylvestris f. angustifolia. *Bauh. pin. 113.*

I. sylvestris. *Fuchs. 330.*

β. *Isatis sativa* f. latifolia. *Bauh. pin. 113. Blakw. t. 246.*

I. (*tinctoria*) foliis radicalibus oblongo-ovatis obtusis integerrimis, caulinis sagittatis, siliculis oblongis. *Mill. dict. n. 1. R.*

Habitat ad litora maris Baltici & oceani Europæ, ad vias Helvetiæ, &c. ☞

2. *ISATIS lusitanica*. I. foliis radicalibus crenatis: caulinis sagittatis, pedunculis subtomentosis.

I. sylvestris. minor lusitanica. *Herm. lugdb. 678.*

I. orientalis maritima canescens. *Tournef. cor. 14. Buxb. cent. 1. p. 4. t. 5.*

Habitat in Hispania & Oriente. ☉

Vix a præcedente sufficienter distincta, sed annua & minor. (Flores albi. Gmel. l c. R.)

889. C R A M B E.

Filamenta 4 longiora apice bifurca: altero antherifero. Bacca ficca, globosa, decidua.

1. *CRAMBE maritima*. C. foliis cauleque glabris. *Fl. suec. dan. 316. Kniph. cent. 10. n. 30. K. 4.*

352 TETRADYNAMIA. SILEQUOSA.

C. foliis cordatis crispis carnosis. *Hort. cliff.*

Brassica maritima monospermos. Bank. pin. 112.

Brassica marina sylvestris multiflora monospermos. L. i. 245.

Habitat ad littora Oceani septentrionalis. 24

3. CRAMBE *hispanica*. C. foliis cauleque scabris. *Kniph. cent. 10. n. 29.*

C. foliis subcordatis crenatis scabris. *Hort. cliff. 34. Roy. lugdb.*

Myagrum sphaerocarpum. Jacq. obs. 2. p. 20. t. 41.

Rapistrum maximum rotundifolium monospermum. Can. canad. 147. t. 148. Moris. hist. 2. p. 266. f. 3. 13. f. 1.

Habitat in Hispania. ☉

890. CLEOME.

Glandulae nectariferae 3, ad singulum sinum calycis singulae, excepto infimo. Petala omnia adflectentia. Siliqua unifocularis, bivalvis.

14. CLEOME *violacea*. C. floribus hexandris, terminalibus solitariisque: foliolis lanceolato-linearibus integerrimis.

C. floribus hexandris, genitalibus declinatis, filis subulatis. *Hort. cliff. 341. Hall. gen. 258.*

Trifolium siliquosum; flore violaceo, lusitanicum. Bauh. ic. 866.

β. *Sinapistrum peregrinum alterum, flore luteo. Bauh. ic. 867. melior.*

Habitat in Lusitania. ☉

Planta viscoso-pubescent. Folia subaequalia: Folia simplicia. Pedunculi laeves. Calyx luteus apicibus purpurascentibus. Petala 2. superiora atropurpurascentia, emmis flavis; Lateralia 2 cordata, unguiculata, crenata, concoloria. Antherae purpurascentes. Raceptaculum super flamina glandulis 3, flavis, globosis. Stamina declinata. Pistillum declinatum.

 CLASSIS XVI.

MONADELPHIA.

TRIANDRIA.

DECANDRIA.

897. GERANIUM. Monogyna. *Caps.* 5-coccarof-trata.
902. SIDA. Submonogyna. *Cal.* simplex, angulatus. *Caps.* locularis, 1-sperma.
910. GOSSYPIMUM. Monogyna. *Cal.* exter. 3-fidus. *Caps.* locularis, polysperma, coadunata.
907. LAVATERA. Polygyna. *Cal.* exter. 3-fidus. *Arilli* 1-spermi, verticillati.
906. MALVA. Polygyna. *Cal.* exter. 3-phyl-lus. *Arilli* 1-spermi, verti-cillati, plures.
908. MALOPE. Polygyna. *Cal.* exter. 3-phyl-lus. *Arilli* 1-spermi, conglo-merati.
905. ALCEA. Polygyna. *Cal.* exter. 6-fidus. *Arilli* 1-spermi, plures.
911. HIBISCUS. Monogyna. *Cal.* exter. 8-fidus. *Caps.* loculis polyspermis, coadunatis.
904. ALTHÆA. Polygyna. *Cal.* exter. 9-fidus. *Arilli* 1-spermi, verticillati.

MONADELPHIA

DECANDRIA.

897. GERANIUM

Monogyna. *Stigmata* 5. *Fructus* rostratus, *coccus*.

* *Africana*. *Staminibus* 7 *antheriferis*; *Foliis* alternis
Pedunculis multifloris; *Fruticosa*.

1. GERANIUM *fulgidum*. G. calycibus monophyllis
foliis tripartitis incis: intermedia majore, umbel
geminis, caule fruticoso cainoso. *Hort. cliff.* 498. *Rac*
ger. 52.

G. africanum, folio alceæ, flore coccineo fulgidiss
Dill. elsh. 156. t. 130. f. 137.

G. furiparvula, obliuata folio, flore coccineo, p
equalibus. *Till. pis.* 68. t. 16.

Habuit in Æthiopia. ¶

2. GERANIUM *inquinans*. G. calycibus monophyllis
foliis orbiculato-reniformibus tomentosis crenatis
griusculis, caule fruticoso. *Burm. ger.* 46. n. 24. *Ac*
cent. 3. n. 42. *Knorr. del.* 1. t. S. 19. a.

G. calycibus monophyllis, floribus florentibus ent
foliis subcordatis. *Hort. cliff.*

G. africanum arborescens, malvæ folio pingui, florib
cineo. *Dill. elsh.* 151. t. 125. f. 151. 152.

G. africanum arborescens, malvæ folio plano lucido,
elegantissimo hermineo. *Mart. cent.* 3. f. 3. p. 19.

Habitet in Africa. ¶

Folia digiti tacta inquinant colore ferrugineo.

4. GERANIUM *acetosum*. G. calycibus monophyllis
foliis glabris obovatis carnosus crenatis, caule frut
laxo. *Burm. ger.* 47. *Kniph. cent.* 11. n. 28.

G. africanum frutescens, folio crasso & glauco, ac
sapore. *Comm. præl.* 54. t. 4.

MONADELPHIA. DECANDRIA. 157

Habitat in Africa. 7

5. GERANIUM *papilionaceum*. G. calycibus monophyllis, corollis papilionaceis: alis carinaque minutis, foliis angulatis, caule fruticoso. *Burm. ger. 49. Kniph. cent. 7. n. 27.*

G. africanum arborescens, flore veluti dipetalo eleganter variegato. *Dill. elth. 124. t. 1. 8. f. 155.*

G. africanum arborescens, malvæ folio mucronato, petalis florum inferioribus vix conspicuis. *Mart. cent. 15. t. 15.*

Habitat in Africa. 7

Petala 3 inferiora calycinis foliolis minora sunt; superiora retusa.

10. GERANIUM *cucullatum*. G. calycibus monophyllis, foliis cucullatis dentatis, caule fruticoso. *Burm. ger. 42.*

G. africanum arborescens, foliis cucullatis angulosis. *Dill. elth. 155. t. 129. f. 156.*

G. africanum arborescens, ibisci folio rotundo, carlinæ odore. *Herm. lugdb. 274. t. 375. Seb. mus. 1. t. 26. f. 2.*

G. africanum arborescens, ibisci folio anguloso, floribus amplis purpureis. *Mart. cent. 28. t. 28.*

G. africanum maximum. *Riv. pent. 325.*

Habitat ad Cap. b. Spel. 7

Planta tota pubescens, reliquis altior, rigidior, erectior. Folia alterna, cucullata s. basi contracta, absque lobis manifestis, undulata, crenata. Calycis tubus limbo brevior. Petala incarnata venis rubris: 2 superiora venis atropurpureis.

11. GERANIUM *carnosum*. G. calycibus monophyllis, caule fruticoso: articulari carnosogibbosis, foliis pinnatifidis laciniatis, petalis linearibus. *Burm. ger. 51.*

G. africanum carnosum, petalis angustis albicantibus. *Dill. elth. 153. t. 127. f. 154.*

Habitat in Æthiopia. 7

12. GERANIUM *gibbosum*. G. calycibus monophyllis, caule fruticoso geniculis carnosigibbosis, foliis subpinnatis. *Burm. ger. 50. Kniph. cent. 8. n. 46.*

G. calycibus monophyllis, geniculis nodosis, foliis duplicato pinnatifidis. *Hort. ups.*

G. calycibus monophyllis, caule carnosonodoso: internodiis filiformibus, foliis lobato-pinnatifidis. *Roy. Lugdb.*

334.

G. africanum noctu olens tuberosum & nodosum, equi-

256 MONADELPHIA. DECAENDRIA.

legiæ foliis. *Herm. lugdb.* 284. t. 285. *Stiff. bot.* III. t. 111. *Burm. afric.* t. 37. f. 2.

G. foliis aquilegiæ. *Riv. pent.* 327.

Habitat ad Cap. b. Spei. \bar{h}

13. GERANIUM *peltatum*. *G. calycibus monophyllis*, foliis quinquelobis integerrimis glabris subpeltatis, caule fruticoso. *Burm. ger.* 48. *Mill. dict. n.* 32. & *ic.* t. 142. *Kniph. cent.* 6. n. 43.

G. africanum, foliis inferioribus asari, superioribus sphididis agrizæ maculatis splendentibus & acerosæ sapore. *Comm. præl.* 52. t. 2.

Habitat in Africa. \bar{h}

Planta procumbens.

14. GERANIUM *Zonale*. *G. calycibus monophyllis*, foliis cordato-orbiculatis incisus zona notatis, caule fruticoso. *Hort. upf. ger.* 43.

G. africanum arborescens, alchemillæ ~~hirsuta folia~~, floribus rubicundis. *Comm. præl.* 51. t. 1.

Habitat in Africa. \bar{h}

Folia disco zona rotundata. Varietas ~~foliorum limbo~~ albo f. *flavescente*.

15. GERANIUM *vitifolium*. *G. calycibus monophyllis*, foliis adscendentibus lobatis pubescentibus floribus capitatis; caule fruticoso. *Burm. ger.* 40.

G. africanum arborescens, vitis folio, odore ~~malvæ~~. *Dil. elth.* 152. t. 126. f. 153.

Habitat in Africa. \bar{h}

16. GERANIUM *capitatum*. *G. calycibus monophyllis*, foliis lobatis undatis villosis floribus capitatis, caule fruticoso diffuso. *Burm. ger.* 41. *Kniph. cent.* 9. n. 41.

G. calycibus monophyllis, floribus capitatis, foliis cordatis lobatis crenatis pilosis. *Hort. cliff.* 345. *Rey. lugdb.* 353.

G. africanum frutescens, malvæ folio laciniato, odore. *Herm. lugdb.* 277. t. 278.

G. malvæ folio. *Riv. pent.* 326.

Habitat in Africa. \bar{h}

* * Stemibus 7 antheriferis. Foliis oppositis. Herbacea.

19. GERANIUM *alchimilloides*. *Geranium calycibus monophyllis*, foliis orbiculatis palmatis incisus pilosis, caule herbaceo decumbente. *Burm. ger.* 55. *Kniph. cent.* 9. n. 40.

G. calycibus monophyllis longissimis sessilibus, fructu assurgente foliis palmatis crenatis. *Hort. cliff.*

MONADELPHIA. DECANDRIA. 139

G. africanum, alchimillæ hirsuto folio, floribus albidis
Herm. lugdb. 282. t. 483. *Stiff. bot.* 107. t. 107.

G. africanum, folio alchimillæ maculato. *Riu. pent.* 331.
Habitas in Africa. 24

21. GERANIUM. *odoratissimum*. *G.* calycibus monophyllis, caule carnosissimo brevissimo, ramis herbaceis longis, foliis cordatis mollissimis. *Burm. ger.* 45. *Kniph. cent.* 5. n. 34.

G. africanum humile, folio fragrantissimo molli. *Dill. elth.* 157. t. 131. f. 138.

Habitas in Africa. 24

26. GERANIUM *myrrhifolium*. *G.* calycibus monophyllis, foliis bipinnatis: inferioribus cordatis lobatis, caule herbaceo, calycibus strigosis. *L. Burm. ger.* 59. *Kniph. cent.* 10. n. 47.

G. africanum, betonicæ folio laciniato & maculato, floribus incarnatis. *Herm. lugdb.* 279. t. 281.

G. africanum tuberosum, anemones folio, flore incarnato
Herm. par. 179. t. 178.

G. æthiopicum, myrrhidis folio 3, flore magno striato
Breyn. cent. 129. t. 59.

Habitas ad Cap. b. Spei. 24

Antheræ 5, alternis sterilibus, ut in subdivisione tertia. Caulis herbaceus, patulus, subpubescens. Folia ternata, incisa, subserrata, lineata, adspersa pilis erectiusculis, disco occupato macula lata fusca: foliolo intermedio latiore semipinnatifido. Calyx laciniis 2 trinerviis; 2 uninerviis; 1 binervi. Petala 4 (absque infimo) pallide incarnata: superioribus venis purpureis.

30. GERANIUM *triste*. *G.* calycibus monophyllis sessilibus, scapis bifidis monophyllis. *Knorr. del. hort.* 1. t. S. 19. *Kniph. cent.* 7. n. 29.

G. calycibus monophyllis: tubis longissimis subsessilibus, radice subrotunda. *Hort. ups.* foliis pinnatifidis. *Roy. lugdb.* *Burm. ger.* 57.

G. triste. *corn. can.* 109. t. 110.

G. noctu olens æthiopicum, radice tuberosa, foliis myrrhidis latioribus at angustioribus. *Breyn. cent.* 126. t. 58.

Habitas ad Cap. b. Spei. 24

Folia ad radicem diversa latiora & angustiora. Petala tristi colore viridi-luteo-pallida, omnibus macula rufa excepto solo margine eorum. Corollæ & odore tristes.

118 MONADELPHIA. DECANDRIA

*** Myrrhina. Stamina 5 antheriferis. Cal. 5-p.
Fructus declinati.

31. GERANIUM *romanum*. G. pedunculis multis
floribus pentandris, foliis pinnatis incisis, scapis
calibus. L. *Burm. ger.* 30.

G. myrrhinum tenuifolium, amplo flore purpureo.
rar. 368. t. 1245.

Habitas Romæ in plateis. 24

Simillimum G. cicutaria, sed Caules nulli; nulla
culca in basi petalorum.

Conf. Geran. petraum. *Magnol monsp.* 109. Ger-
ridum *Lob. illustr.* 134. *Moris. hist.* 2. p. 516. f. 3.
f. 27. forte sola varietas.

Non est varietas, sed Geran. (*petraum*) pedun-
culis multis: floribus pentandris, foliis bipinnatis, f.
incisis, scapis radicalibus. *Gouan. Monsp.* 274. *illust.*
t. 21. f. 1. & G. romano abunde diversum. R.

32. GERANIUM *cicutarium*. G. pedunculis multis
floribus pentandris, foliis pinnatis incisis obtusis, G.
ramoso. *succ. paris. Burm. ger.* 33. *ged. gallob. car-*
herb. fil. Kniph. cent. 11. n. 50. *Reg. bot. lugd.*
lihu.

G. caule procumbente, petiolis multifloris, foliis
cato-pinnatis, pinnis acute incisis. *Hall. helu. n.*

G. robertianum. *Riu. pent.* t. 112. & foetidum. *Ei. t.*
Hall. R.

G. cicutæ folio, minus & supinum. *Bauh. pin.* 319.
giff. p. 48.

G. supinum. *Dod. pempt.* 63.

A. G. pimpinellæ folio. *Dillen. giesf.* 69. *Hall. R.*

G. minus, G. arvense. *Tabern.* 57. *Hall. R.*

Habitas in Europæ sterilibus cultis. ☉

33. GERANIUM *moschatum*. G. pedunculis multis
floribus pentandris, foliis pinnatis incisis, corylæ-
pinnatifidis. *Burm. ger.* 29. *Jacq. hort.* t. 55.
Blackw. t. 150.

G. petiolis multifloris, caule diffuso, foliis ovatis p-
tifidis. *Hall. helu. n.* 945. R.

G. moschatum. *Riu. pent. irreg.* t. 110. R.

G. cicutæ folio, moschatum. *Bauh. pin.* 319.

Habitas in Helvetia, Sibiria, Carniola. ☉

Simillimum G. cicutaria & caulescens; herba an-
fiaca.

MONADELPHIA. DECANDRIA. 159

34. GERANIUM *chium*. G. pedunculis multifloris, floribus pentandris, foliis cordatis incis: superioribus lyrato-pinnatifidis. L. *ger.* 35.

G. *chium* vernal, caryophyllatæ folio. *Tournef. cor.* 20. *Mart. cent.* 4. t. 4.

Habitat in Chio. ☉

Cotyledones cordati, integri, retusi, regulares, nec oblique cordati ut G. *gruino* & *ciconio*. Caules G. *ciouta-*torio majores. Macula nulla ad basin petalorum superiorum.

35. GERANIUM *malachoides*. G. pedunculis multifloris, floribus pentandris, foliis cordatis sublobatis. *monsp.* *Burm. ger.* 34. *Kniph. cent.* 11. n. 53.

G. folio *altheæ*. *Bauh. pin.* 318.

G. *malacoides*. *Lob. ic.* 662.

G. pedunculis bifloris, foliis ovatis hirsutis crenato incis obtusis, caulibus procumbentibus. *Roy. lugdb.* 391.

Habitat in Italiæ, G. *narbonensis*, Angliæ *maritimis*. ☉
Petioli oppositi: altero brevior.

36. GERANIUM *maritimum*. G. pedunculis subtrifloris, floribus pentandris, foliis cordatis incis crenatis scabris, caulibus depressis. *gallob.*

G. pedunculis bifloris, foliis ovatis hirsutis crenato-incis obtusis, caulibus procumbentibus. *Roy. lugdb.*

G. *minimum* procumbens, folio *betonicæ*. *Morif. hist.* 2. p. 512. f. 5. t. 35. f. 10.

G. *pusillum* *maritimum*, *altheæ* s. *betonicæ* folio. *Pluk. phys.* 31. f. 4.

Habitat in Angliæ, Gallo-belgiæ *maritimis*.

Pedunculi sæpe 2-flori. Petala integra. Flores sæpe petalis mutilati; Umbella sæpe foliosa.

39. GERANIUM *gruinum*. G. pedunculis submultifloris, floribus pentandris, foliis ternatis lobatis. *Burm. ger.* 32. *Kniph. cent.* 7. n. 26.

G. pedunculis multifloris pentandris. *Hort. ups.* 198, 2.

G. pedunculis bifloris pentandris, radice annua. *Vir. cliff.* 66. *Roy. lugdb.* 352.

G. *latifolium*, acu longissima. *Bauh. pin.* 319.

G. *speciosum* annuum, longissimis rostris, creticum. *Bauh. hist.* 3. p. 479.

Habitat in Creta. ☉

40. GERANIUM *ciconium*. G. pedunculis multifloris, floribus pentandris, foliis pinnatifidis obtusis. *Burm. ger.* 28. *Jacq. hort.* t. 18. *Kniph. cent.* 11. n. 49. *delph.*

260 MONADELPHIA. DECANDRIA.

G. petiolis multifloris, foliis pinnatis, pinnis late serratis, alterne maximis. Hall. helv. n. 946.

G. cicutæ folio, acu longissima. Bauh. pin. 136. 138.

G. apulum coriandrifolium. Col. ecphr. 1. p. 136.

Habitat Monspelli, inque Vallesia, Italia, Helvetia.

41. *GERANIUM pyrenaicum. G. pedunculis bifloris, floribus pentandris, petalis bilobis, calycibus glandulosis, antheris exterioribus sterilibus. delph.*

G. pedunculis bifloris, foliis multifidis: laciniis oblongis, petalis bifidis. Burm. ger. 27. G. prou. 434 t. 16. f. 2.

G. pedunculis bifloris, foliis inferioribus quinquepartitis multifidis rotundatis; superioribus trilobis, caule erecto. Hudson. angl.

G. colombinum pyrenaicum perenne maximum. Tausch. inst. 268.

Habitat in Pyrenæis, Anglia. 24.

Cules villosi. Folia quinquesida, incisa, obtusiuscula, margine rubro. Calycis foliola apice glandula rubra. Petala calyce duplo longiora, purpurascens. Stamina quinque, exteriora breviora antheris abortientibus. Gerard.

† *Geranium alceoides, coriandrifolium, myrrhifolium, auritum & pusillum.*

*** *Batrachia. Stamina decem antheriferis.*

42. *GERANIUM tuberosum. G. pedunculis bifloris, multifidis: laciniis linearibus subdivisis obtusis. Burm. ger. 9.*

Geranium tuberosum majus. Bauh. pin. 318.

Geranium bulbosum. Lob. hist. 377.

Habitat in Anglia, Italia. 24.

43. *GERANIUM macrorrhizum. G. pedunculis bifloris, calycibus inflatis, petalis integris, pistillo longissimo, scapo dichotomo.*

*G. pedunculis bifloris, calycibus inflatis, pistillo longissimo. Hort. cliff. * Burm. ger. 10. Kniph. cent. 1. n. 52.*

G. batrachioides odoratum. Bauh. pin. 318. Mansf. 2. p. 514. f. 4. & 16. f. 15.

G. batrachioides longius radicans odoratum. Bauh. 3. p. 477.

Habitat in Italia. 24.

Folia lobata, incisa, obtusiuscula, supra mollia. Rob.

MONADELPHIA. DECANDRIA. 16t
expens. Scapi aphylli, erecti, dichotomi. Calyces
aristati, globosi, angulati: foliolis 2 exterioribus ma-
ioribus. Petala orbiculata, integra, plana, patentia,
purpurea. Stamina nutantia. Pistillum declinatum, in
fructu longius.

44. GERANIUM phæum. G. pedunculis solitariis op-
positifoliis, calycibus subaristatis, caule erecto, petala-
lis undulatis. *delph. lugd.*

G. pedunculis bifloris foliisque alternis, calycibus sub-
aristatis, caule erecto, petalis undulatis. *Burn. ger.*
11. Jacq. vind. sil. Kniph. cent. 5. n. 35.

G. foliis rugosis hirsutis, semiseptilobis, lobis dentatis,
petalis serratis, acuminatis. *Hall. helv. n. 934.*

G. pedunculis bifloris alternatim cauli integro superne
nudiusculo insidentibus. *

A. Geranium batrachioides hirsutum, flore atro-rubente.
Bauh. pin. 318.

Geranium i. pullo flore. *Clus. hist. 2. p. 99.*

Habitat in Alpibus Pannonicis, Helveticis, Styriacis. **24**

45. GERANIUM fuscum. G. pedunculis bifloris oppo-
sitifoliis geminis, caule patulo, petalis integerrimis.

G. pedunculis bifloris, foliis quinquelobis incis, petalis
reflexis. *Mill. dict. n. 7.*

G. phæum s. fuscum petalis reflexis. *Moriss. hist. 2. p.*
515.

G. montanum fuscum. *Bauh. pin. 318.*

Habitat in Europa australi. **24**

Invitus hanc separe a G. phæo, cum quo convenit
multis, etiam Corolla tota expansa atra: sed differt Fo-
liis rigidis, Corolla magis reflexa, minor; petalis sub-
orbiculatis mucronatis, integerrimis, planis; Pedunculis
non solitariis bifloris, sed 2 distinctis. *H. V.*

46. GERANIUM reflexum. G. pedunculis bifloris foliis-
que alternis, petalis reflexis laciniatis longitudine caly-
cis mutici.

G. flore purpureo reflexo, italicum. *Barr. ic. 39.*

Habitat in Italia. **24**

Affine G. phæo & fusco. Caulis pedalis, dichotomus
s. bifidus, patulus, pubescens. Folia superiora alterna,
quinqueloba, rugosa, tomentosa, non maculata. Flores
cernui: pedunculis oppositifoliis, cernuis. Calyx muticus.
Petala rubra, subincisa, calyce non longiora, reflexa,
Linnaei spec. Plant. Europ. Pars II. **L**

162 MONADELPHIA. DECANDRIA.

bas inter se hiantia. Stamina corollis fere longiora, saccata. Antheræ flavæ margine fusco.

47. GERANIUM nodosum. G. pedunculis bifloris, petalis emarginatis, foliis caulinis trilobis integerratis, subtus lucidis. *delph.*

G. pedunculis bifloris, foliis caulinis trilobis integerratis, summis subsessilibus. *Burm. ger. 7.*

G. nodosum. *Bauh. pin. 318.*

G. 5. 6. plateau. *Clus. hist. 2 p. 101.*

Habitat in Delphinatu. 2.

Caules diffusi, compressi magis quam in G. striatum. Petala emarginata, incarnata striis tribus purpureis.

48. GERANIUM striatum. G. pedunculis bifloris, foliis quinquelobis lobis medio dilatatis, petalis bilobis venoso-reticulatis. *Burm. ger. 6.*

G. foliis septilobatis, rugosis, acute dentatis, petalis venosis, emarginatis. *Hell. helv. n. 936.*

G. romanum versicolor s. striatum. *Part. p. 229. Moris. hist. 2. d. 516. f. 5 t. 10. f. 2. Rai hist. 1063.*

Habitat in Italia. 2.

Folia macula fusca ad sinus in pagina superiore.

49. GERANIUM sylvaticum. pedunculis bifloris, foliis subpeltatis quinquelobis inciso-serratis, caule erecto, petalis emarginatis. *succ. Burm. ger. 12. carn. ph. herb. noru. Flor. dan. t. 124. litua.*

G. caule erecto, foliis rugosis septilobis, lobis serratis, petalis emarginatis. *Hall. helv. n. 932.*

G. batrachioides, folio aconiti. *Bauh. pin. 317.*

Geranium 2. batrachioides minus. *Clus. hist. 2. p. 9.*

Habitat in Europæ borealis sylvis. 2.

Albo flore & flore albo lineis purpureis picto varietas.

50. GERANIUM palustre. G. pedunculis bifloris, foliis simis declinatis, foliis quinquelobis incis, petalis integerratis. *Burm ger. 13. ingr. ged. pal. herb. sil. Kniph. ont. n. 54. Flor. dan. n. 596.*

G. caule diffuso, foliis rugosis semiquinquelobis, lobis trifidis, floribus sparsis, petalis integerrimis. *Hall. n. 933.*

G. sanguineum majus. *Best. cyst. vern. 1. t. 9. f. 1.*

G. batrachioides palustre, folio sanguineo. *Dill. lith. 160. t. 134. f. 161.*

Habitat in Russia, Germaniâ.

MONADELPHIA. DECANDRIA. 163

Rami *divaricatissimi*, *bisariam angulati angulis obtusis*
 Petala nervis 3. *fuscis*, *basi villosa*.

51. GERANIUM *pratense*. G. pedunculis bifloris foliis
 subpeltatis multipartitis rugosis acutis, petalis integris.
delph. lithu.

G. pedunculis bifloris, foliis subpeltatis multipartitis pin-
 nato-laciniatis rugosis acutis, petalis integris. *suec.* *
Burm. ger. carn. gallob. pal. Gmel. tub. p. 208. Leers.
herb. n. 537. Kniph. cent. 5. n. 36.

G. caule erecto, foliis rugosis, hirsutis, multilobis, lobis
 trifidis, lobulis semipinnatis, floribus umbellatis. *Hall.*
helu. n. 931.

G. batrachioides, gratia dei germanorum. *Bauh. pin.*
 318.

Geranium 3. batrachioides majus. *Clus. hist. 2. p. 100.*
Habitat in Europæ borealis pratis. 2

52. GERANIUM *argenteum*. G. pedunculis bifloris, foliis
 subpeltatis septempartitis trifidis tomentoso-sericeis,
 petalis emarginatis. *Burm. ger. 8. delph.*

G. argenteum alpinum. *Bauh. pin. 318. Pluk. phyt. 186.*
f. 3. longius radicum. Pon. bald. 341. t. 342. Segu.
ver. 1. p. 471. t. 10.

G. argenteum montis baldi. *Bauh. hist. 3. p. 474.*

Habitat in Baldi summis rupibus.

**** Staminibus 10. antheriferis. Pedunculis bi-
 floris. ☉

54. GERANIUM *bohemicum*. G. pedunculis bifloris,
 petalis emarginatis, arillis hirtis, cotyledonibus trifidis:
 medio truncato. *Amæn. acad. 4. * Burm. ger. 24.*
cent. 176.

G. batrachioides bohemicum, capsulis nigris, hirsutis.
Dill. elth. 159. t. 133. f. 160.

G. bohemicum batrachioides annuum. *Rai. hist. 1063.*

G. batrachioides minus annuum purpureo-cæruleum. *Morif.*
præl. 267. hist. p. 511. f. 5. t. 15. f. 1. mala.

Habitat in Bohemia? ☉

Planta pubescens viscida. Pedicelli fructus recti. Calyx
 trifidus nervis 3. f. 5. Corolla cærulea, calyce major.
 Arilli hirsuti, atrii.

55. GERANIUM *roberianum*. G. pedunculis bifloris,
 calycibus pilosis decemangulatis. *Burm. ger. ged. carn.*
gallob. pal. fil. tub. Fl. dan. 694. Blackw. t. 480.
Regn. bot. lithu. burg. lugdb.

164 MONADELPHIA. DECANDRIA.

G. pedunculis bifloris, foliis quinque-^{ve} partitis, bipinnatifidis, calycibus hirsutis. Mor. cliff. 344. Fl. parif.

G. foliis duplicato pinnatis, pinnis ultimis confertis calycibus striatis, hirsutis. Hall. helv. n. 943.

G. robertianum primum. Bauh. pin. 319.

Geranium 3. Fusch. hist. 206.

Habitat in Europæ borealis rupibus; etiam in helv. felici. ♂

Calyx pilosus, decem-carinatus, ex foliolis 1 & 4 anterioribus tricarinatis; 2 & 5 unicarinatis; 4 linearibus. Petala integra. Odor hircosus.

Geranium 3. Fusch. ad G. dissectum amandat Hall.

56. GERANIUM *lucidum*. *Geranium pedunculis bifloris calycibus pyramidalis angulatis elevato-rugosis, lobis quinquelobis rotundatis. * succ. parif. monsp. Burm. ger. 19. ingr. pal. Flor. dan. t. 218. Kniph. con. n. 51. delph.*

G. foliis subrotundis, semiquinquelobis, lobis obtusis calycibus transversim rugosis. Hall. helv. n. 942.

Geranium lucidum saxatile. Bauh. pin. 318.

Geranium saxatile. Thal. herc. 44. t. 5.

Geranium rotundifolium saxatile montanum. Col. europ. p. 138. t. 137.

Habitat in Europæ rupibus umbrosis. ☉

57. GERANIUM *molle*. *G. pedunculis bifloris foliis floralibus alternis, petalis bifidis, calycibus muticis caule erectiusculo.*

G. pedunculis bifloris foliisque floralibus alternis, caule ramoso, erectiusculo, calycibus muticis, arillis linearibus. Burm. ger. carn. pal. herb. Fl. dan. t. 679. Burg. 2. delph.

G. foliis mollissimis, hirsutis, reniformibus semipartitis, lobis semitripartitis obtusis. Hall. helv. n. 941.

G. foliis infimis semimultipartitis: laciniis linearibus bifidis, calycibus exaristatis. Neck. gallob.

G. pedunculis bifloris, foliis quinquepartitis rotundatis: laciniis obtusiusculis, capsulis bivalvibus. succ. parif.

G. pedunculis bifloris, foliis sparsis reniformibus quinquesidis mollibus & latissimis. Sauv. monsp.

G. columbinum minus, majori flore & foliis bifidis. monsp.

MONADELPHIA. DECANDRIA. 165

1. columbinum villosum, petalis bifidis purpureis. *Vaill. parif. 79. t. 15. f. 3.*

Habitat in Europa ad plateas. ☉

Petala semibifida. Calyces submutici, corollæ breviores. edicelli declinati. Stipulæ bifidæ.

9. GERANIUM columbinum m. G. pedunculis bifloris foliis longioribus, foliis quinquepartito-multifidis, arillis glabris, calycibus aristatis. *Fl. succ. parif. Burm. ger. ged. carn. pal. herb. fil. Dær. nass. p. 120. burg. lithu. lugd. delph.*

caule procumbente, foliis quinquepartitis, lobis trilobis, lobulis trifidis. *Hall. helu. 938.*

columbinum, foliis dissectis, pediculis florum longissimis. *Vaill. parif. 79. 15. f. 4.*

Habitat in Gallia, Helvetia, Germania.

Pedunculi foliis longiores. Calyces maxime aristati, magni. Petala emarginata. *Hall.*

1. GERANIUM dissectum. G. pedunculis bifloris, foliis quinquepartito-trifidis, petalis emarginatis longitudine calycis, arillis villosis. *Amæn. acad. 4. * Burm. ger. Fl. succ. gallob. carn. pal. herb. delph.*

caule erecto, foliis quinquelobis, lobis trifidis, acutis. *Hall. helu. n. 937.*

majus, foliis imis longis ad usque pediculum divisis. *Morif. hist. 2. p. 311. f. 3. t. 3. f. 3. Vaill. parif. t. 15. f. 2.*

Habitat in Europa australiore. ☉

Calyces aristati, Pistillum villosum. Petala pallido-carnata, calyce non longiora. Stipulæ coloratæ. Antheræ æruleæ, (Petala potius sanguinea. *Læers & mihi. R.*)

2. GERANIUM rotundifolium. G. pedunculis bifloris, petalis subintegris longitudine calycis, caule prostrato, foliis reniformibus incis. *Burm. ger. carn. gallob. herb. fil. Reg. bot. burg. lithu. delph.*

viscidum, caule decumbente, foliis semiquinquelobis, lobis tridentatis & quinquelobatis. *Hall. helu. n. 941. R.*

pedunculis bifloris, foliis subrotundis multifidis, caule procumbente. *Fl. succ. parif.*

folio malvæ rotundo. *Bauh. pin. 328.*

alterum. *Fuchs. hist. 105.*

columbinum majus, flore minore cæruleo. *Vaill. parif. 79. t. 15. f. 1.*

Habitat in Europa cultis. ☉

366 MONADELPHIA. DECANDRIA.

Caulis *divaricatus teres, pubescenti - viscidus*. Petal. *rufi*. Folia *reniformia, rotundata, pubescentia, mollia, subviscosa præsertim subtus, lobata sinibus puncto rubra*. Calyx *aristatus, patulus, rugis 3 longitudinalibus*. Petal. *cuneata, obtusissima, integra, basi striata 3 rufis*. Antheræ *flavae*.

Synonymon. *Vaill. huc non pertinet; monente Haller & Scopolio. R.*

63. GERANIUM *pusillum*. G. pedunculis *bifloris, petalis emarginatis, caule depresso, foliis reniformibus palmatis linearibus acutis*. *Burm. ger. herb. Reich. f. delph.*

G. foliis *hirsutis, semiseptilobis, lobis semitrilobis obtusis*. *Hall. helv. n. 940. R.*

G. columbinum *humile, flore cæruleo minimo*. *Rai. angl. p. 359. t. 16. f. 2.*

G. columbinum *tenuius laciniatum*. *Bauh. pin. 318. p. 138.*

Habitat in *Anglia, Gallia, Germania, Helvetia.* ☉

Caulis *prostratus, vix pubescens*. Folia *magis dissecta laciniis quibusdam ad basin usque sectis, quod non in Geraniis rotundifolio simillimo*. Petala *vere emarginata, purpurea*. Antheræ *cæruleæ*. Stamina *5. alterna castrata sunt*.

Geranium Halleri adductum alii pro varietate G. præcedentis habent, alii huc pertinere perhibent. *R.*

***** *Staminibus 10 antheriferis. Pedunculis 1-floris.*

64. GERANIUM *sibiricum* G. pedunculis *subunifloris, quinquepartitis acutis: foliolis pinnatifidis*. *Burm. germ. Gmel. sib. 3. t. 67. Jacq. hort. t. 19. Kniph. cent. 10. n. 1.*

Habitat in *Siberia.* 24

Caules *pedales, diffusi, divaricati*. Folia *quinquepartita: lobis pinnatifidis*. Pedunculus *e singula bifurcatus*. Caulis *longus, uniflorus, in medio geniculatus bracteis oppositis, subulatis*. Corolla *alba striis purpureis, petalis vix emarginatis*. Calyx *5. phyllus, aristatus*. Antheris *villosis*.

65. GERANIUM *janguineum*. G. pedunculis *unifloris* foliis *quinquepartitis trifidis orbiculatis*. *Burm. ger. pal. fil. Kniph. cent. 7. n. 28. burg. litua. lugd. delph.*

G. pedunculis *unifloris; foliis septilobis, petalis bifidis*. *Hall. helv. n. 930.*

G. pedunculis *simplicibus unifloris*. *Fl. suec. paris.*

G. *janguineum, maximo flore*. *Bauh. pin. 318.*

MONADELPHIA. DECANDRIA. 159

34. GERANIUM *chium*. G. pedunculis multifloris, floribus pentandris, foliis cordatis incis: superioribus lyrato-pinnatifidis. L. *ger.* 35.

G. *chium* vernal, caryophyllatæ folio. *Tournef. cor.* 20. *Mart. cent.* 4. t. 4.

Habitat in Chio. ☉

Cotyledones cordati, integri, retusi, regulares, nec oblique cordati ut G. gruino & ciconio. Caules G. cicuta-

35. GERANIUM *malachoides*. G. pedunculis multifloris, floribus pentandris, foliis cordatis sublobatis. *monsp. Burm. ger.* 34. *Kniph. cent.* 11. n. 53.

G. folio *altheæ*. *Bauh. pin.* 318.

G. *malacoides*. *Lob. ic.* 662.

G. pedunculis bifloris, foliis ovatis hirsutis crenato incis obtusis, caulibus procumbentibus. *Roy. lugdb.* 391.

Habitat in Italiæ, G. *narbonensis*, Angliæ *maritimis*. ☉

Petoli *oppositi: altero brevior.*

36. GERANIUM *maritimum*. G. pedunculis subtrifloris, floribus pentandris, foliis cordatis incis crenatis scabris, caulibus depressis. *gallub.*

G. pedunculis bifloris, foliis ovatis hirsutis crenato-incis obtusis, caulibus procumbentibus. *Roy. lugdb.*

G. *minimum* procumbens, folio *betonicæ*. *Morif. hist.* 2. p. 512. f. 5. t. 35. f. 10.

G. *pusillum* *maritimum*, *altheæ* s. *betonicæ* folio. *Pluk. phyt.* 31. f. 4.

Habitat in Angliæ, Gallo-belgiæ *maritimis*.

Pedunculi *sæpe 2-flori*. Petala *integra*. Flores *sæpe petalis mutilati; Umbella sæpe foliosa*.

39. GERANIUM *gruinum*. G. pedunculis submultifloris, floribus pentandris, foliis ternatis lobatis. *Burm. ger.* 32. *Kniph. cent.* 7. n. 26.

G. pedunculis multifloris pentandris. *Hort. upf.* 198, 2.

G. pedunculis bifloris pentandris, radice annua. *Vir. cliff.* 66. *Roy. lugdb.* 352.

G. *latifolium*, acu longissima. *Bauh. pin.* 319.

G. *speciosum* annuum, longissimis rostris, creticum. *Bauh. hist.* 3. p. 479.

Habitat in Creta. ☉

40. GERANIUM *ciconium*. G. pedunculis multifloris, floribus pentandris, foliis pinnatifidis obtusis. *Burm. ger.* 28. *Jacq. hort.* t. 18. *Kniph. cent.* 11. n. 49. *delph.*

168 MONADELPHIA. POLYANDRIA.

Abutilon indicum. Comm. hort. 2. t. 1. Bauh. hist. 2.

959. Moris. hist. 2. p. 531. f. 5. t. 19. f. 4.

Althææ. theophrasti similis. Bauh pin. 316.

Habitat in India. ☉

Pedunculi erecti. Capsulæ terminatæ angula recta.

21. *SIDA cristata.* S. foliis angulatis: inferioribus crenatis, superioribus panduriformibus, capsulis multiseptatis. Kniph. cent. 2. n. 85.

S. foliis crenatis: inferioribus dentatis obrusis, superioribus acuminatis. Roy lugdb. 349.

S. foliis cordatis obsolete serratis. Hort. cliff. 346.

Abutilon lavateræ flore, fructu cristato. Dill. elch. 3. f. 2.

β. *Sida* foliis inferioribus cordato angulatis, superioribus hastatis, petalis integerrimis. Hort. ups. 199. Kniph. cent. 7. n. 86.

Habitat in Mexico. ☉

904. ALTHÆA.

Cal. duplex: exterior 9. fidus. *Arilli* plures monospermi.

1. *ALTHÆA officinalis.* A. foliis simplicibus tenuibus. parif. gelr. austr. carn. gallob. pal. nass. Flor. 530. Ludw. ed. 1. 1. Kniph. cent. 6. n. 3. Ray. Burg. Lugd.

A. discoridis & plinii. Bauh. pin. 315.

Althæa. Fusch. hist. 15. Cam. epit 667. Blachn. t.

A. tomentosa, herbacea, caule erecto, foliis lanceolatis obsolete trifidis. Hall. helv. n. 1074.

β. *Althæa laciniato folio.* Bauh. pin. 316.

Habitat in Hollandiæ, Angliæ, Galliæ, Sibiria, sub-humidis. 26

2. *ALTHÆA cannabina.* A. foliis inferioribus palmatis superioribus digitatis. monsp. Jacq. austr. 1. 102. burg.

A. foliis compositis scabris. Hort. cliff.

Alcea cannabina. Bauh. pin. 316.

Alcea fruticosa, cannabino folio. Clus. hist. 2. p. 25.

Habitat in Hungaria, Italia, G. Narbonensi, ad frum margines. 26

MONADELPHIA. DECANDRIA. 16t
repens. Scapi aphylli, erecti, dichotomi. Calyces
aristati, globosi, angulati: foliolis 2 exterioribus ma-
ioribus. Petala orbiculata, integra, plana, patentia,
purpurea. Stamina nutantia. Pistillum declinatum, in
fructu longius.

44. **GERANIUM phæum.** G. pedunculis solitariis op-
 positifoliis, calycibus subaristatis, caule erecto, peta-
 lis undulatis. *delph. lugd.*

G. pedunculis bifloris foliisque alternis, calycibus sub-
 aristatis, caule erecto, petalis undulatis. *Burn. ger.*
II. Jacq. vind. sil. Kniph. cent. 5. n. 35.

G. foliis rugosis hirsutis, semiseptilobis, lobis dentatis,
 petalis serratis, acuminatis. *Hall. helv. n. 934.*

G. pedunculis bifloris alternatim cauli integro superne
 nudiusculo insidentibus. *

β. *Geranium batrachioides hirsutum, flore atro-rubente.*
Bauh. pin. 318.

Geranium l. pullo flore. Clus. hist. 2. p. 99.

Habitat in Alpibus Pannonicis, Helveticis, Styriacis. 24

45. **GERANIUM fuscum.** G. pedunculis bifloris oppo-
 sitifoliis geminis, caule patulo, petalis integerrimis.

G. pedunculis bifloris, foliis quinquelobis incis, petalis
 reflexis. *Mill. dict. n. 7.*

G. phæum f. fuscum petalis reflexis. *Moriss. hist. 2. p.*
515.

G. montanum fuscum. *Bauh. pin. 318.*

Habitat in Europa australi. 25

Invitus hanc separe a G. phæo, cum quo convenit
multis, etiam Corollæ tota expansa atra: sed differt Fo-
liis rigidis, Corollæ magis reflexa, minor; petalis sub-
orbiculatis mucronatis, integerrimis, planis; Pedunculis
non solitariis bifloris, sed 2 distinctis. H. V.

46. **GERANIUM reflexum.** G. pedunculis bifloris foliis-
 que alternis, petalis reflexis laciniatis longitudine caly-
 cis mutici.

G. flore purpureo reflexo, italicum. *Barr. ic. 39.*

Habitat in Italia. 26

Affine G. phæo & fusco. Caulis pedalis, dichotomus
f. bifidus, patulus, pubescens. Folia superiora alterna,
quinqueloba, rugosa, tomentosa, non maculata. Flores
cernui: pedunculis oppositifoliis, cernuis. Calyx muticus.
Petala rubra, subincisa, calyce non longiora, reflexa,
Linnaei spec. Plant. Europ. Pars II. L

170 MONADELPHIA. POLYANDRIA.

Malva rosea, folio ficus. *Bauh. pin.* 315.

Malva rosea simplex peregrina. *Tabern. hist.* 315.

Malva arborea. *Blakw. t.* 54.

Habitat in Sibiria. ♂

Vix sufficienter a precedente distincta.

906. MALVA.

Cal. duplex : exterior 3. phyllis. Arilli plures, monospermi.

* Foliis indivisis.

4. MALVA coromandeliana. M. foliis ovato-oblongis acutis, floribus glomeratis, arillis denis tricuspidatis. Althæa coromandeliana angulis prælongis foliis semibicorni. *Pluk. mant.* 10.

Habitat in America ? ☉

Flores axillares glomerati. Fructus truncatus apice limbo

* * Foliis angulatis.

6. MALVA peruviana. M. caule erecto herbaceo, foliis palmatis, spicis secundis axillaribus denticulatis. *hort. t.* 156. *Kniph. cent.* 7. n. 52.

Habitat in Peru. B. Jussieu. ☉

Caulis erectus, tripedalis, pilis rarioribus serpiens pilis. Folia septemlobata, plicata glabra, venosa, serrata; petiolis longitudine folii. Stipulae ovato-lanceolatae. Pedunculus axillaris, nudus, longus. Spica secunda sursum versa, recurvata ante florescentiam. Corollae purpureae. Semina 10, supra exasperata dentibus denis plurimis.

Folia 5-l. 3-loba. Flores rubri. *Jacq. l. c.* R.

8. MALVA bryonifolia. M. caule fruticoso tomentoso foliis pinnatis scabris, pedunculis multifloris.

M. foliis palmatis scabris, caule tomentoso fruticoso, pedunculis multifloris. *Spec. pl.* 2. p. 968. *Roy. bot.* 356. *Mill. dict.* n. 12.

Althæa frutescens, bryoniæ folio. *Bauh. pin.* 316. 138.

Althæa profunde serrato s. dentato folio. *Bauh. hist.* 955.

Habitat in Hispania. ♀

Planta belle echinata per discum foliorum

MONADIPHEIA TITANICA

Ran. diaphanum. *Diastylis* *diastylis* *diastylis*

Petal. dent. 3. fimb. 10.

51. GERANIUM *pratense* *G. pratense*

Subsp. diastylis *diastylis* *diastylis*

diastylis

G. pratense *diastylis* *diastylis* *diastylis*

diastylis *diastylis* *diastylis* *diastylis*

diastylis *diastylis* *diastylis* *diastylis*

diastylis *diastylis* *diastylis* *diastylis*

G. pratense *diastylis* *diastylis* *diastylis*

diastylis *diastylis* *diastylis* *diastylis*

diastylis *diastylis* *diastylis* *diastylis*

G. pratense *diastylis* *diastylis* *diastylis*

diastylis *diastylis* *diastylis* *diastylis*

Geranium *diastylis* *diastylis* *diastylis*

Habitus *diastylis* *diastylis* *diastylis*

52. GERANIUM *argenteum* *G. argenteum*

Subsp. diastylis *diastylis* *diastylis*

petalis emarginatis *Burm. f. 1. 176.*

G. argenteum *diastylis* *diastylis* *diastylis*

f. 3. long. 176. *diastylis* *diastylis* *diastylis*

ver. 1. 176. *diastylis* *diastylis* *diastylis*

G. argenteum *diastylis* *diastylis* *diastylis*

Habitus *diastylis* *diastylis* *diastylis*

**** *diastylis* *diastylis* *diastylis*

54. GERANIUM *bohemium* *G. bohemium*

petalis emarginatis *diastylis* *diastylis*

medio truncatis *Amer. 176.* *diastylis* *diastylis*

cent. 176. *diastylis* *diastylis* *diastylis*

G. bohemium *diastylis* *diastylis* *diastylis*

Dill. elin. 176. *diastylis* *diastylis* *diastylis*

G. bohemium *diastylis* *diastylis* *diastylis*

G. bohemium *diastylis* *diastylis* *diastylis*

pral. 267. *diastylis* *diastylis* *diastylis*

Habitus *diastylis* *diastylis* *diastylis*

Plant. pubescens *diastylis* *diastylis* *diastylis*

aristatus *nervis* *3. f. 3.* *diastylis* *diastylis*

Arilli hirsuti *diastylis* *diastylis* *diastylis*

55. GERANIUM *robertianum* *G. robertianum*

calycibus pilosis *diastylis* *diastylis*

gallab. fol. sub. Fl. dan. 694. *diastylis* *diastylis*

Regn. bot. luth. burg. lugd. *diastylis* *diastylis*

172 MONADELPHIA. POLYANDRIA.

M. cretica minor hirsuta. *Bazb. cent. app. 46. f. 3.*

M. humifusa minima. *Till. pif. 108. t. 33. f. 2.*

Habitas in Bitynia. Arduini. 22

Pedunculi sub florescentia archani, foliis bivalves. *lata rubra, rotundata cum mucronat. Capsula quam 5, in orbem posita aut sparsa.*

14. MALVA sylvestris. M. caule erecto herbaceo, septemlobatis acutis, pedunculis petiolisque pilosis. *Corn. pal. fil. Blackw. t. 22. Regn. bot. burg. luyd. t.*

M. caule erecto, foliis multipartitis. *Pl. succ. parif.*

M. caule erecto, foliis lobatis, lobis serratis, quatuor septentis. *Hall. helv. t. 1064.*

M. caule erecto herbaceo, foliis lobatis obtusis, pedunculis petiolisque pilosis. *Cronq. austr.*

M. foliis quinquelobis serratis, obtusis: calycem exterioribus subtrifidis, *Neck. gallob.*

M. sylvestris, folio sinuato. *Bauh. pin. 314.*

M. sylvestris recta. *Cord. hist. 114.*

Habitas in Europae campestribus.

Caulis & petioli hujus villosi; at sequentis folia latus superius villosum. Albo flore variat.

15. MALVA mauritiana. M. caule erecto herbaceo, quinquelobatis obtusis, pedunculis petiolisque pilosis.

M. hederaceo folio. *Bauh. pin. 315.*

M. major tertia. *Dalech. hist. 586. Kniph. cent. 7. t. 1.*

β. M. folio ficus altera. *Bauh. pin. 315. prod. 17.*

Habitas in Italia, Lusitania, Hispania. ☉

Similis precedenti. Huic folia costis 5; illi 7. Huic exterior lanceolatus, illi ovatus. Huic corolla fere magis, illi caerulea.

16. MALVA hispanica. M. caule erecto, foliis bicalatis crenatis, calyce exteriore diphylo. *Lam.*

M. rotundifolia glabra & hispanica, flore amplo.

Pluk. alm. 238. t. 44. f. 3.

Habitas in Hispania. ☉

Caules diffusi, adscendentes, hirti. Folia rotunda, vix lobata, crenata. Stipulae lanceolatae. Petioli solitarii, uniflori, folio longiores. Calyx exterior phyllus, deficiente tertio interiore foliolo. C. interior fidus lobis cordatis s. sinibus prominentibus. Capsula nata.

17. MALVA verticillata. M. caule erecto folis

MONADELPHIA. POLYANDRIA. 167

Geranium VII. hæmatodes. *Clus. hist. 2. p. 202.*

G. columbinum erectum, tenuius laciniatum, flore magno.

Læf. pruss. 103. n. 18. R.

β. Geranium hæmatodes lancastrienſe, flore eleganter
ſtriato. *Dill. elth. 163. t. 136. f. 163. Barrel. ic. 67.*

Habitat in Europæ pratis ficcis umbroſis. 24

POLYANDRIA.

902. SIDA.

*Cal. ſimplex, angulatus. Stylus multipartitus. Caps.
plures, 1. ſpermæ.*

1. SIDA *ſpiroſa*. S. foliis cordato-oblongis ſerratis, ſti-
pulis ſetaceis, axillis ſubſpinolis.

S. foliis cordato-ovatis ſerratis, capsulis quinque bicuspidatis. *Hort. cliff.*

Alcea, carpinifolia, americana fruteſcens, floſculis lu-
teis, ſemine duplici roſtro. *Comm. hort. 1. p. 3. t. 2.*

Althæa carpinifolia, floſculis luteis. *Bœc. ſic. 11. t. 2.*

Althæa virginiana bidens, pimpinellæ majoris acutiore ſo-
lio, floſculis minimis luteis. *Pluk. alm. 25. t. 9. f. 6.*

Habitat in India utraque. ☉

3. SIDA *rhombifolia*. S. foliis lanceolato-rhomboidibus
ſerratis, axillis ſubſpinolis. *Kœst. cent. 8. n. 85.*

Malvinda unicornis, folio rhomboidi, perennis. *Dill.
elth. 216. t. 172. f. 212.*

Habitat in India utraque. ☉

17. SIDA *Abutilon*. S. foliis ſubrotundo-cordatis indi-
viſis, pedunculis folio brevioribus, capsulis multilocu-
laribus: corniculis bifidis. *Kœst. cent. 4. n. 79.*

Abutilon foliis tomentolis, cordatis, petiolatis, ſiliquis bi-
cornibus. *Hall. helv. n. 1075. R.*

Althæa theophrasti, flore luteo. *Bauh. pin. 316.*

Althæa altera c. Abutilon. *Cam. epit. 668.*

Habitat in Indiis; Helvetia, Sibiria. ☉

*Fruſtus nigri piloli. Calycis foliola complicata ſ. cana-
liculata.*

19. SIDA *indica*. S. foliis cordatis ſublobatis, ſtipulia
reflexis, pedunculis petiolo longioribus, capsulis multi-
locularibus ſcabris calyce longioribus. *Amœn. acad. 4.*

*p. 324.**

174 MONADELPHIA. POLYANDRIA

quepartitis trilobis linearibus, pedunculis folio ca-
longioribus, caule decumbente. *Aman. acad. 4.*

*Malva maritima galloprovincialis, geranii folio. Ta-
nef. inst. 98.*

*Alcea minor maritima tenuifolia procumbens. Herm. par-
9. t. 9.*

*Alcea tenuifolia humilis maritima galloprovincialis, f-
inferioribus ad geranium accendentibus. Pluk. del.
t. 44. f. 4.*

Habitat in Galloprovinciæ, Hispaniæ maritimis. O-

Folia obtusiora, ciliata, ad apices supra setis adsp-

Caules ad basin valde ramosi, prostrati, effigentes. O-

rollæ calyce quadruplo majores.

907. LAVATERA.

*Cal. duplex : exterior 3, fidus. Arilli pl-
monospermi.*

** Caule fruticoso.*

1. *LAVATERA arborea. L. caule arboreo, foliis*
renangularibus tomentosis plicatis, pedunculis cor-
unifloris axillaribus.

L. foliis septemangularibus obtusis plicatis villosis,
fruticoso, floribus ad alas confertis. Hort. cliff.

Malva arborea venera dicta, parvo flore. Bauh. pin-

Malva arborecens. Dod. pempt. 653. Camer. hort.

Habitat inter Pisas & Liburnum. ♂

2. *LAVATERA micans. L. caule arboreo, foliis*
renangularibus acutis crenatis plicatis tomentosis,
mis terminalibus.

Malva foliis mollibus undulatis : in margine
micis sulphureis ad solem splendentibus donatis.

hist. p. 1. 523. f. 5. t. 17. f. 9.

Malva hispanica. Moris. præl. 285.

Habitat in Hispania, Lusitania. ♀

3. *LAVATERA olbia, L. caule fruticoso, foliis*
quelobo-hastatis, floribus solitariis. Jacq. hort. bot.

L. foliis quinquangularibus acutiscrenatis : laciniis
productiore. Hort. cliff.

Althæa frutescens, folio acuto, parvo flore. Bauh.

316.

MONADELPHIA POLYANDRIA. 169

3. *ALTHÆA hirsuta*. A. foliis trifidis piloso-hispidis, supra glabris, pedunculis solitariis unifloris. *paris. monsp. pal. carn. Jacq. austr. t. 170. Delph. burg.*

A. foliis radicalibus reniformibus, caulinis palmatis, trilobis & quinquelobis, calicibus hispidis. *Hall. helv. n. 1073.*

A. foliis imis reniformibus, semiquinquelobis, summis quinquepartitis, serratis, hirsutis. *Grm. isen. in Nov.*

Ad. A. N. C. 3. app. p. 353.

Alcea hirsuta. Bauh. pin. 317.

Alcea hirsuta minima, flore caesio, hispanica. Barr. ic. 1169.

Alcea villosa. Dalech. hist. 394. Bauh. hist. 2. p. 1167.

Habitas in Gallia, Italia, Hispania, Austria sepibus.

Caules diffus, ut tota planta, hirti pilis patulis, sed Folia supra glabra. Calyx exterior 8 phyllus; interior longitudine corollae, acuminatus. Corolla incarnato alba, crenulata.

4. *ALTHÆA Ludwigii*. A. foliis lobatis utrinque nudis, pedunculis congestis unifloris. *Murray prodr. 169. **

Alcea minor siliculata, flore minore. Rai. hist. 605?

Habitat in Sicilia. D. Schreber.

Herba tota facie refert Foliis Malvam Alceam, ut ovum ovo. Pedunculi axillares, 205, uniflori, longitudine petiolorum. Calyx exterior 8-phyllus foliolis lanceolatis; Interior fere brevior, 5-fidus, albis villis hirsutissimus, minor A. hirsutae.

Alcea hirsuta differt ab hac Foliis minus divisis acutioribus, Caulis Pedunculis Petiollisque pilosis; Calicibus strigosis (nec lanatis,) quadruplo majoribus: interiore non hispido, nec lanato-albo.

905. A L C E A.

Cal. duplex: exterior 6. fidus. Arilli plures; monospermi.

1. *ALCEA rosea*. A. foliis sinuato angulosis. *Ludw. ed. t. 41. Knorr. del. 1. t. R. 15. Kniph. cent 6. n. 2. Regn. bot.*

Malva rosea, folio subrotundo. Bauh. pin. 315.

Malva hortensis. Dod. pempt. 652.

Habitat in Oriente. ♂

2. *ALCEA ficifolia*. A. foliis palmatis.

176 MONADELPHIA. POLYANDRIA.

Althæa thuringiaca grandiflora. *Dill. elch. 9. t. 8. f. 4.*

Althæa flore majore. *Bauh. pin. 316.*

Habitat in Pannonia, Tartaria, Suecia, Germania, & ad sepes. ☿

8. *LAVATERA cretica.* L. caule erecto, ramis inferioribus diffusis, pedunculis confertis unifloris, foliis latis : superioribus acutis. *Hort. ups. * Jacq. hort. 1.*

Malva cretica annua altissima, flore parvo ad apicem bellato. *Tournef. cor. 2.*

Malva annua hirsuta : foliis hederæ instar angulosis. *Morist. hist. 2. p. 521. f. 5. t. 7. f. 5.*

Habitat in Creta. ☉

9. *LAVATERA trimestris.* L. caule scabro, herbaceo foliis lanceolatis, pedunculis unifloris, fructibus instructis. *Jacq. hort. 1. 72. Kniph. cent. 8. n. 56.*

L. foliis trilobis : lacinia media productione, caule herbaceo. *Hort. cliff.*

L. foliis ovato-lanceolatis : inferne angulatis, pedunculis unifloris, caule herbaceo. *Sauv. monsp.*

Malva folio vario. *Bauh. pin. 315. prodr. 137. t. 1.*

β. *Lavatera foliis subcordatis obtusis angulatis* caule herbaceo. *Roy. lugdb.*

Habitat in Syria, Hispania, G. Narbonensi.

908. MALOPE.

Cal. duplex : exterior 3 phyllus. Arilli glomerati monospermi.

1. *MALOPE malacoides.* M. foliis ovatis crenatis glabris. *Sabb. hort. 1. t. 50.*

Malva betonicæ folio. *Morist. hist. 2. p. 522. f. 5. t. 11.*

II. *Bocc. sic. 15. t. 8. f. 2.*

Alcea betonicæ folio, flore purpureo violaceo. *Morist. 1189.*

Habitat in Hetruriæ pratis & Mauritania.

909 URENA.

Cal. duplex : exterior 5 fidus. Caps. 5 lobis echinata, monosperma.

1. *URENA lobata.* V. foliis angulatis.

MONADELPHIA. POLYANDRIA. 171

9. MALVA *capensis*. M. caule arborescente, foliis cordatis quinquelobis.

M. foliis subcordatis laciniatis glabris, caule arborescente.

Spec. pl. 2. Kniph. cent. 6. n. 59.

M. *capensis* frutescens, grossularæ folio minori glabro.

Dill. elth. 208. t. 169. f. 206.

10. M. *capensis* frutescens grossulariæ folio majore hirsuto.

Dill. elth. 209. t. 169. f. 207.

Habitat in Æthiopia. ☿

10. MALVA *caroliniana*. M. caule repente, foliis multifidis. *Willich. obs. n. 55.*

Abutilon repens, alceæ foliis, flore helvulo. *Dill. elth.*

5. t. 4. f. 4.

Abutilon carolinianum repens, alceæ foliis, giluo flore.

Mant. 34. t. 34.

Habitat in Carolina. ☉

Fructus *trepani* figura supra tuberculatus ambitu aristis 2 e singulis loculis.

11. MALVA *parviflora*. M. caule patulo, foliis angulatis, floribus axillaribus sessilibus glomeratis, calycibus glabris patentibus. *Amœn. acad. 3. p. 416. Jacq. hort. t. 39.*

M. annua rotundifolia, floribus omnium minimis albis pentapetalis verticillatim genicula ambientibus *Morif. 2. p. 521. Rai. hist. 598.*

M. *tingitana*, flore cæruleo, parva. *Pluk. phyt. 44. f. 2.*

Habitat in Barbaria. ☉

Differt a M. rotundifolia: Caule erectiusculo triplo majore; Foliis acutioribus apice petioli adscendente. Calycibus glabris, inferiore lineari nec lanceolato. Floribus sessilibus. Calycibus maturo fructu purpurascentibus. Arillis supra rugosis & ad latera dentatis.

12. MALVA *rotundifolia*. M. caule prostrato, foliis cordato-orbiculatis obsolete quinquelobis, pedunculis fructiferis declinatis. *Fl. suec. Paris. Hall. helu. n. 1070. austr. carn. gallob. pal. sil. nass. Knorr. del. 2. t. M. 1. Flor. dan. t. 721.*

Malva sylvestris, folio subrotundo. *Bauh. pin. 314.*

Malva sylvestris pumila. *Fuchf. hist. 508.*

Habitat in Europæ ruderatis, viis, plateis. ☉

Albo flore ludit. R.

13. MALVA *sherardiana*. M. caulibus prostratis, foliis orbiculatis plicatis tomentosis crenatis, pedunculis solitariis unifloris arcuatis,

911. H I B I S C U S .

Cal. duplex : exterior polyphyllus. *Caps.* 5 loculi
polyisperma.

1. HIBISCUS *Moscheutos*. H. foliis ovatis acutis
serratis : caule simplicissimo, petiolis floriferis.

H. foliis ovatis crenatis : angulis lateralibus obtusis.
Hort. cliff.

Alcea rosea peregrina, forte *Rosa moscheutos* R.
Corn. canad. 144. t. 145. *Morif. hist.* 2. p. 532. f. 19. f. 6.

Habitat in Canada, Virginia. 24

Flos petiolo insidet uti *Turneræ ulmifoliae*.

2. HIBISCUS *palustris*. H. caule herbaceo simplicissimo
foliis ovatis subtrilobis subtus tomentosis, floribus
axillaribus.

Althæa palustris. *Bauh. pin.* 316. (ex horto C. B. R.
serus.)

Althæa hortensis. s. *peregrina*. *Dod. pempt.* 655.

Ketmia palustris flore purpureo. *Tournef. inst.* 100. f. 1.

Habitat in Virginia. *Gronou. Canada. Kalm.* 24.

Habitus H. *Moscheutos*. Caules sesquipedales
ramosi, annui. Folia lato-ovata, obtuse serrata, vix
acuminata, subtus tomentosa. Pedunculi ex axillis
superiorum solitarii, petiolo longiores, uniflori
e petiolo enati, geniculati. Flos maximus.

4. HIBISCUS *tiliaceus*. H. foliis cordatis subrotundis
divisis acuminatis crenatis, caule arboreo, calyce
interiore decem dentato.

Malva arborea maritima, folio subrotundo minoribus
minuto subtus candido, cortice in futes ducto.
jam. 93. *hist.* 1. p. 215. t. 134. f. 4.

Alcea malabarica, abutili folio, flore minore
flavescente exterius aspero. *Rai. hist.* 1070.

Alcea indica sinarum, flore luteo malvaceo. *Pluk.*
6. p. 355. f. 5.

Althæa maritima arborescens diffusa, foliis ovatis
cordatis crenatis subtus cinereis. *Brown. jam.* 110.

Ketmia zeylanica sempervirens & florens, flore
Burm. zeyl. 136. ind. 150. R.

Ketmia indica, tiliæ folio. *Tournef. inst.* 100. P.

2. R.

MONADELPHIA, POLYANDRIA. 173

latis, floribus axillaribus glomeratis sessilibus, calycibus scabris. *Jacq. hort. t. 40.*

M. sinensis erecta, flosculis albis minimis. *Berth. lugdb. p. 268.*

M. anpua rotundifolia, floribus omnium minimis albis pentaphyllis verticillatim genicula ambientibus. *Morif. hist. 2. p. 521. Rai. hist. 598.*

Habitat in China. ☉

18. *MALVA crispa*. *M. caule erecto*, foliis angulatis crispis, floribus axillaribus glomeratis. *herb. litua.*

M. foliis angulatis crispis, floribus axillaribus glomeratis. *Hort. ups. 200.*

M. foliis crispis. *Bauh. pin. 315.*

M. crispa. *Ded. pempt. 653.*

Habitat in Syria; in Germania quasi sponte. ☉

19. *MALVA Alcea*. *M. caule erecto*, foliis multipartitis scabriusculis. *Succ. parif. ged. carn. pal. Mattusch. fil. Kniph. cent. n. 50. Sabbati hort. 1. t. 48. litua. burg. lugd.*

M. caule erecto, foliis scabris, trilobatis, lobis lateralibus bipartitis, medio tripartito & quinquepartito. *Hall. helu. n. 1071.*

M. caule recto: foliis subpeltatis multipartito-pinnatifidis. *Neck. gallob.*

Alcea vulgaris major. *Bauh. pin. 316.*

Alcea vulgaris. *Clus. hist. 2. p. 25. Blackw. t. 309.*

Habitat in Germania, Anglia, Gallia. 24

Caulis scaber pilis fasciculatis expansis.

20. *MALVA moschata*. *M. caule erecto*, foliis radicalibus reniformibus incis, caulinis quinquepartitis pinnato-multifidis. *Fl. suec. carn. pal. herb. burg. lugd. litua.*

M. foliis radicalibus reniformibus incis, caulinis quinquepartitis pinnatis, pinnis dentatis. *Hall. helu. n. 1072.*

M. montana f. *Alcea rotundifolia laciniata.* *Col. ecphr. 1. p. 148. t. 147.*

Alcea folio rotundo laciniato. *Bauh. pin. 316.*

β. *Alcea vulgaris minor.* *Bauh. pin. 316.*

Habitat in India, Gallia, Germania, &c. 24

Differt a precedente; Floribus ambrosiacis; Caule humiliore pilis solitaris erectis insidentibus puncto prominulo.

21. *MALVA tournefortiana*. *M. foliis radicalibus quin-*

180 MONADELPHIA. DECAENDRIA.

bus palmatis quinquepartitis subtus uniglandulosis, aculeato, floribus sessilibus.

H. vitifolius. *Mill. dict. n.* 18. R.

Alcea benghalensis spinosissima, acetosæ sapore, luteo pallido : umbone purpurascente. *Comm. bot. p.* 35. t. 18.

Ketmia indica, foliis digitatis, flore magno subrotundo, umbone atro-purpureo, petiolis spinosis. *Ehret. t.* 68.

Ketmia indica, *Vitis* folio, magno flore. *Tournef. inf. t.* 100. R.

Ketmia indica, cannabinis foliis, *Bangue* dicta. *Burm. f. ind.* 134. ind. 152. R.

Habitat in India. ☉

Caulis quinquepedalis, adpersus aculeis sursum et inferum ; Ramis paucis, simplicibus. Petioli patentes, longitudine foliorum, adpersi subtus aculeis rectis. Folia inferiora indivisa ovata ; media 3 loba ; superiora 5. palmate digitata, inermia nervo medio subtus basi glandulosa. Exterior subulatus, erectus, aculeatus ; interior glandula in singulo foliolo. Fructus subrotundus, pericarpio calyce tectus.

Convenit cum *H. Sabdariffa* : Structura, facie et odoribus, glandula sub foliis.

17. HIBISCUS *Manihot*. *H.* foliis palmato-digitatis quinquepartitis, caule petiolisque inermibus. *Sabb. de med. t.* 56. *Kniph. cent.* 9. n. 47.

Ketmia folio manihot serrato, flore amplo subrotundo. *Dill. elth.* 189. t. 156. f. 189.

Alcea sinica, *manihot* stellato folio, capsula longiuscula pyramidata quinquesariam divisa. *Pluk. amalt. t.* 355. f. 2.

Habitat in Indiis ☿

18. HIBISCUS *Abelmoschus*. *H.* foliis subpeltatis septemangularibus serratis, caule hispido, ind. 153.

Alcea hirsuta, flore flavo, semine moschato. *bras.* 45. t. 45. *Brown. jam.* 285. n. 4.

Alcea ægyptiaca villosa. *Bauh. pin.* 317.

Flos moschatus. *Mer. surin.* 42. t. 42.

Granum moschatum. *Rumph. amb.* 4. p. 38. t. 1.

Cattu-gasturi. *Reed. mal.* 2. p. 71. t. 38.

Ketmia ægyptiaca, semine moschato. *Tournef. inf. t.* 100. *Burm. zeyl.* 134. R.

MONADELPHIA. POLYANDRIA. 173

Althæa frutescens, folio acuto virente molli, flore specioso. *Pruk. phyt.* 8. f. 1.

Althæa arborea olbia in galloprovincia. *Leb. ic.* 653.

Habitat in Olbia insula Galloprovinciæ. ♀

Variat *Foliis primi anni maximis*, insequentium vero longe minoribus, quod affinis commune. Similima *L. trilobe*, sed huic flores solitarii pedunculo brevissimo; huic *Calyx exterior acuminatus*, fere tripartitus, longitudine interioris.

4 *LAVATERA triloba*. *L.* caule fruticoso, foliis subcordatis subtrilobis rotundatis crenatis, stipulis cordatis, pedunculis aggregatis unifloris. *Jacq. hort.* 1. 74. *Kniph. cent.* 7. n. 41.

L. foliis subcordatis emarginatis obsolete trilobis crenatis petiolo brevioribus, stipulis cordatis. *Roy. lugdb.*

Malva foliis subcordatis trilobis obtusis serratis villosis. *Hort. cliff.*

Althæa frutescens, folio rotundiore incano. *Bauh pin.* 316.

Althæa fruticans hispanica, aceris monspessulani incanis foliis, grandiflora saponem spirans. *Pluk. alm.* 24. t. 8. f. 3.

Habitat in Hispania, Gallia. ♀

Planta tomentosa: tota pilis minimis glutiniferis, eisque immixtis aliis pilis majoribus apice stellatis. Pedunculi tres pluresve ex aliis petiolorum, uniflori Flos magnus, pallide purpurascens. *Calyx interior in fructu duplo longior.* *Ste.*

5. *LAVATERA lusitanica*. *L.* caule fruticoso, foliis septemangularibus tomentosis plicatis, racemis terminalibus. *Kniph. cent.* 9. n. 55.

Althæa frutescens lusitanica, folio ampliore minus incano. *Tournef. inst.* 97.

Habitat in Lusitania, Cap. b. spei. ♀

* * *Caule herbaceo.*

7. *LAVATERA thuringiaca*. *L.* caule herbaceo, fructibus denudatis, calycibus incis. *Scholl. barb. Jacq. austr.* 1. 311. *Kniph. cent.* 6. n. 53.

L. caule herbaceo, foliis summis hastatis calycibusque acutis, pedunculis solitariis unifloris. *Schreb. spicil. flor. lips.*

L. foliis inferioribus cordato-subrotundis superioribus lobatis, crenatis calycibus utrinque incis, caule herbaceo. *Roy. lugdb.*

Althæa thuringiaca. *Cam. hort.* 1. t. 6.

CLASSIS XVII.

DIADELPHIA

HEXANDRIA.

920. FUMARIA.

Cal. 2-phyllus. Cor. 4-
gens, basi gibbosa
stamiferis. Filamentis
theris 3.

OCTANDRIA.

921. POLYGALA.

Cal. 2. laciniae albae
Cor. vexillum cylindricum
Stam. connatis
Carp. obcordatis,
locularis.

DECANDRIA.

* Stamina omnia connata.

929. SPARTIUM.

Filam. adherentia
Stigma adnatum,
sum.

930. GENISTA.

Pisillum deprimens
nam. Stigm. involuta

939. LUPINUS.

Antherae alternae rotundae
alternae oblongae
gum. coriaceum.

936. ANTHYLLIS.

Cal. turgidus, inclausus
siliquam.

932. ULEX.

Cal. diphyllus. Legum.
calyce longius.

935. ONONIS.

Legum. rhombum, falcatum
Vexillum fringed.

MONADELPHIA. POLYANDRIA. 177

V. finica, xanthii facie. *Dill. elsh.* 340. t. 319. f. 412.
Trifolio affinis indiae orientalis, xanthii facie. *Breyn.*
cent. 82. t. 35.

Lappago ambonica. *Rumph. amb.* 6. p. 59. t. 25. *Burm.*
ind. p. 149. R.

Habitat in China ♀

Porus glandulosus subtus in costis 3 intermediis foliorum.

910. GOSSYPIUM.

Cal. duplex exterior 3 fidus. Caps. 4. locularis. Sem.
lana obvoluta.

1. *GOSSYPIUM herbaceum.* G. foliis quinquelobis,
 subtus eglandulosis, caule herbaceo lævi. *Black.* t. 354.
Kniph. cent. 8. n. 47.

G. caule decumbente *Hort. luglb.*

G. frutescens, semine albo. *Bauh. pin.* 430.

Gossypium. *Cam. epit.* 203. *Rumph. amb.* 4. p. 33. t. 12.

Habitat in America. ☉

2. *GOSSYPIUM arboreum.* G. foliis palmatis: lobis
 lanceolatis, caule fruticoso. *Mill. dict. n.* 3. *Forskæhl.*
ægypt. p. 125.

G. caule erecto. *Hort. cliff.*

G. latifolium. *Rumph. amb.* 4. p. 37. t. 13?

G. arboreum caule lævi. *Bauh. pin.* 430.

G. herbaceum s. Xylon maderaspatense, rubicundo flore,
 pentaphyllum, *Pluk. alm.* 172. t. 188. f. 3.

Cudupariti. *Rheed. mal.* 55. t. 31.

Xylon arboreum. *Tournef. inst.* 101. *Burm. ind.* 150. R.

Gossypium arboreum s. segiar. *Alp. exot.* 38. t.
 38. R.

Habitat in India. ♂

Inter foliorum sinus sæpe lacinula, unde omnes sinus obtusi.

3. *GOSSYPIUM hirsutum.* G. foliis quinquelobis subtus
 uniglandulosis, ramulis petiolisque pubescentibus. *Sabb.*
hort. 1. t. 57.

Xylon americanum præstantissimum, semine virescente.
Tournef. inst. 101.

Gossypium frutescens pentaphyllos barbadense. *Pluk.*
alm. 172. t. 299. f. 1.

Habitat America. ☉ ♂

Variat foliis obtusis.

Linnaei spec. Plant. Europ. Pars II.

M

956. CORONILLA. *Legum.* isthmis inter-
tum rectum.
957. ORNITHOPUS. *Legum.* articulatam, r-
cuarum.
959. SCORPIURUS. *Legum.* isthmis inter-
tum, teretiuculum, r-
volum.
958. HIPPOCREPIS. *Legum.* compresso-
br anaceum : altera f-
emarginaturis ad me-
excisa.
971. MEDICAGO. *Legum.* spirale, men-
naceo-compressum. *Fi-*
tilium carinam de-
tens.

* * * * * *Legumen uniloculare polysperum*
(nec priorum.)

970. TRIGONELLA. *Vexillum* *Alaeque* p-
quasi tripetala. *Co-*
minuta.
953. ROBINIA. *Vexillum* reflexo-
subrotundum.
949. CICER. *Calycis* 4 laciniae f-
res vexillo incumb-
948. ERUUM. *Cal.* 5-partitus, f-
lis, longitudine f-
rollae.
951. CYTISUS. *Legumen* pedicellum
lyx bilabiatum.
963. GALEGA. *Legum.* lineare f-
que transversis.
969. LOTUS. *Legum.* teres, f-
minibus cylindricis

MONADELPHIA. DECANDRIA. 179

Habitat in India ad rivos. \bar{h}

Folia juniora subtriloba, subtus poris 3. Stipulae magnae, rotundatae.

9. **HIBISCUS mutabilis.** H. foliis cordato-quinquangularibus obsolete serratis, caule arboreo.

Althaea arborea Rosa sinensis. *Morif. hist.* 2. p. 530 f. 5. t. 18. f. 2.

Rosa sinensis. *Ferr. flor.* 493. t. 497. *Ger. sur.* 31. t. 31.

Habitat in India. \bar{h}

10. **HIBISCUS Galvaviscus.** H. foliis cordatis crenatis: angulis lateralibus extimis parvis caule arboreo. *Sabb. hort.* 1. t. 54. *Kniph. cent.* 1. n. 31.

H. frutescens, foliis angulatis cordatis acuminatis, petalis ab uno latere auritis. *Brown. jam.* 284.

Malvaviscus arboreus, flore miniato clauso. *Dill. elth.* 210. t. 170. f. 208.

Malva folio hederaceo, flore coccineo. *Plum. spec.* 2. ic. 169. f. 2.

Alcea indica arborea, folio molli, flore amplo, elegantef coccineo. *Pluk. alm.* 14. t. 257. f. 2.

Habitat in Mexico. \bar{h}

Corolla convoluta, erecta, coccinea, clausa fauce lacinia laterali connivente. Fructus Baccæ sphaerica: loculis 5, monospermis.

12. **HIBISCUS Syriacus.** H. foliis cuneiformi-ovatis superne inciso-serratis, caule arboreo.

H. foliis cuneiformi-ovatis: superne inciso-dentatis, caule arboreo. *Spec. pl.* 1. *Kniph. cent.* 2. n. 32. *Knorr. del.* 1. t. K. 2. a & tab. K. 2. b.

Ketmia syriaca. *Scop. carn.*

Alcea arboreus syriaca. *Bauh. pin* 316.

Alcea arboreus. *Cam. hort.* t. 3. 4.

Syrischer Hibiscus. *Linn. Pfl. Syst.* 4. p. 157.

Habitat in Syria, Carniolia. \bar{h}

Calyx exterior 6-7-8-fidus. R.

13. **HIBISCUS ficulneus.** H. foliis quinquefido palmatis, caule aculeato, floribus pedunculatis.

Ketmia zeylanica, fici folio, perianthio oblongo integro.

Dill. elth. 190. t. 157. f. 190 *Burm. zeyl.* 137.

Habitat in Zeylona.

Aculei insident verrucis rubris; saepe reflexi, quod non in H. Sabdariffa.

15. **HIBISCUS cannabinus.** H. foliis serratis: superiori-

286 DIADELPHIA. HEXANDRIA.

F. radice bulbosa, cava, caule simplici bifolio, bracteis ovato-lanceolatis. *Hall. helv. n. 348.*

F. (bulbosa caule simplici subangulato, pedunculis brevioribus racemisque solitariis multifloris. *N. gallob.*

F. bulbosa major. *Reich francof. herb. Kniph. cent. n. 31. & 32. Flor. dan. t. 605. Blackw. t. 534.*

F. bulbosa, radice cava, major. *Bauh. pin. 543.*

Pistolochia. Fuchs. hist. 91.

β. *Fumaria bulbosa, intermedia.*

F. radice bulbosa, solida, caule simplici multifolio, bracteis digitatis. *Hall. helv. n. 349.*

F. bulbosa, minor. *Reich. st. manofr. herb. born. pl.*

F. bulbosa, radice non cava, minor. *Bauh. pin. 144.*

γ. F. bulbosa, radice non cava major. *Bauh. pin. 144. Knorr. del. 1. t. H. 9.*

Habitat in Europæ nemoribus & umbrosis. 24

Conjungunt has plantas speciei: Radix carnea, simplex. Varietas β. Plantæ primi anni, producentis bracteas, quæ subsequentibus fissæ evadunt. Gemmæ renales duæ, trivalves. Caulis simplex, subangulatus, in ramo; Foliorum structura decomposita; Racemus immixtus, terminalis, solitarius; Corolla, Calyx, Semina, Stamina, Siliquæ structura; Tempus. Species reliquæ propriis notis distinctissimæ.

Plantæ α & β duæ distinctæ species sunt cultu mutabiles. Ad var. α pertinet varietas γ, dum modo cava, modo solida invenitur. Variat etiam in flore albo. R.

6. FUMARIA lutea. F. filiquis teretibus, caulibus diffusis: angulis obtusis.

F. capnoides. *Mill. dict.*

F. lutea. *Bauh. pin. 143.*

F. lutea montana. *Dalech. hist. 1294.*

F. tingitana, radice fibrosa, perennis, flore albo fructucente, filiquis curtis. *Pluk. alm. 262. t. 90. f. 1.*

Habitat in Mauritania. 24

Simillima F. capnoidi, sed perennis. Corollæ rotundatum, tubo dimidio brevior. Racemi absque involucris.

7. FUMARIA capnoides. F. filiquis linearibus teretibus, caulibus diffusis acutangulis. *Carn. Kniph. cent. 11. 2.*

F. caule ramoso diffuso, filiquis oblongis, radice perennante. *Hort. cliff.*

MONADELPHIA. POLYANDRIA. 131

Mosch f. *Bammia moschata*. *Alp. exot.* 197. R.

Habitat in Indiis. ☞

24. *HIBISCUS pentacarpus*. H. foliis inferioribus cordatis angulatis, superioribus subhastatis, floribus subnutantibus, pistillo cernuo.

H. foliis cordatis angulatis, ferratis, stipulis fetaceis divaricatis. *Hort. cliff.*

Ketmia palustris, foliis lobatis subrotundis inferna parte molli subcinerea lanugine, flore purpureo magno. *Michx. flor.* 54. *Tourtes it. p.* 309.

Ketmia palustris minor, folio angusto, flore parvo purpurascente, fructu depresso pentagono. *Zannich. venet.* 155. t. 91.

Habitat in paludosis Venetiæ, circa Astracan. Lerche. 24

Planta antecedenti nimis affinis; sed fructu parum diverso; forte a precedenti orta? Caulis tripedalis, angustus. Folia caulina cordato-oblonga; superiora hastata, viridia, subtrinervia, subtus tomentosa, tenuissime serrata. Pedunculi axillares in summitate plantæ solitarii, nudæ, petiolo longiores, uniflori. Calyx inferior patens, radiis subulatis, vix longitudine interioris. Corolla patens, dilute rubra, subnutans. Stamina lutea tota; Antheræ saturatius. Stylus cernuus cum staminibus, albus; Stigmata 5, capitata, incarnata.

26. *HIBISCUS Trionum*. H. foliis tripartitis incis, calycibus inflatis. *Sabb. hort.* 1. t. 55. *Kniph. cent.* 5. n. 43.

Ketmia Trionum. Scop. carn.

Alcea vesicaria. Bauh. pin. 317.

Alcea peregrina solisequa. Lob. ic. 650.

β. *Ketmia vesicaria africana. Tournef. inst.* 101.

Hibiscus africanus. Mill. dict. n. 20.

Alcea vesicaria capitis b. spei. Moris. præl. 227. *hist.* 2. p. 533.

Habitat in Italia, Africa, Carnioliæ. ☉

Variatio β Caulis erectior, simplicior. Folia simplicia; semitrifida, s. triloba: supremis 5-lobis, Pedunculi petiolis longiores. Folia *Trionii* e *Mulva capensi* scabrosâ patre.

118 DIADELPHIA. OCTANDRIA.

8. *F. foliis tenuissimis, floribus albis, circa Malvaceum. Bauh. pin. 143. Segu. ver.*

F. tenuifolia erecta hispanica purpurea. Barr. Soc. 41.

F. minor tenuifolia præcox, semine lini. Marif. hist. p. 262. f. 3. t. 12. f. 13.

Capnos tenuifolia. Clus. hist. 2. p. 208.

Habitas in Hispania, G. Narbonensi, Verone alage, &c.

Varietates qui statuas hanc & sequentem esse hanc officinalis viz erret, diditante fructu. An hybrida? ex strobilus arde congestis est. Folia glauca. Flores males.

12. *FUMARIA claviculata. F. filiquis Enciclis, cirrhiferis. Flor. dan. 1. 340.*

F. foliis cirrhiferis, floribus spicatis. Flara. off.

F. clavicularis donata. Bauh. pin. 143. Marif. hist. 266. f. 3. t. 12. f. 3.

F. alba latifolia. Rai. angl. 3. p. 335.

Habitas in Angliæ locis uliginosis saxosis; in Ital. Cirrus foliorum terminalis.

OCTANDRIA.

921. POLYGALA

Cal. 5. phyllus: foliolis duobus albis coloratis. Legumen obcordatum, bilobis.

** Cristata (floribus appendice paniculatis.)*

5. *POLYGALA amara. P. floribus cristatis, caulibus erectiusculis foliis radicalibus obcordatis. Jacq. vind. carn. tub. delph. linæ.*

P. procumbens, foliis imis subrotundis, spicatis. Hall. helv. n. 343.

P. (amarella) foliis circa radicem spicatis obcordatis. Crantz austr.

P. vulgaris, foliis circa radicem oppositis, cæruleo, sapore admodum amaro. Bauh. pin. 143.

P. buxi minoris folio. Vaill. parif. 161. a. 32. f. 1.

Habitas in Gallia, Austria subalpina montibus.

Filia P. vulgaris, sed Folia majora: quibusdam pro majora, cæterum nimis officina.

* * *Stigma pubescens.* (Nec priorum notæ.)

954. COLUTEA. *Legum.* inflatum, supra basin dehiscens
 940. PHASEOLUS. *Carina Stylusque* spirales.
 945. OROBUS. *Stylus* linearis teretiusculus, supra villosus.
 944. PISUM. *Stylus* supra carinatus villosusque.
 946. LATHYRUS. *Stylus* supra planus villosusque.
 947. VICIA *Stylus* sub stigmate barbatus.

* * * *Legumen subbiloculare* (nec priorum.)

965. ASTRAGALUS. *Legum.* 2-loculare, rotundatum.
 966. BISSERRULA. *Legum.* 2-loculare, planum, dentatum.
 964. PHACA. *Legum.* femibiloculare.

* * * * *Legumina submonosperma* (nec priorum.)

967. PSORALEA. *Cal.* punctis glandulosis.
 968. TRIFOLIUM. *Legum.* vix calyce longius; 1-f. 2-spermum *Flores* capitati.
 955. GLYCYRRHIZA. *Cal.* 2-labiatus: superiore 3-fido.

* * * * * *Legumen subarticulatum.* (*).

961. HEDYSARUM. *Legum.* articulis subrotundis, compressis. *Carina* obfusissima.

(*) Umbellatæ *Lotus*, *Coronilla*, *Ornithopus*, *Hippocrepis*, *Scorpiurus*.

90 DIADELPHIA. DECANDRIA.
 Anonymos flore coluteæ. *Clus. hist. I. p. 105. boni*
Habitat in Austriæ, Helvetiæ, Alsatia, montanis. Linn
in Germania. Nou. Ad. A. N. C. IV. app. p. 17

DECANDRIA.

929. SPARTIUM.

Stigma longitudinale, supra villosum. *Filamentum*
 germini adhærentia. *Cal.* deorsum productus.

* *Foliis simplicibus.*

3. SPARTIUM *junceum*. S. ramis oppositis teretibus
 apice floriferis, foliis lanceolatis. *Sauv. monsp. Anst.*
cent. I. n. 83. Regn. bot. burg.

S. arborescens, seminibus lenti similibus. *Bauh. pin. 396.*

S. macrolobium. *Renealm. spec. 34. t. 33.*

Genista juncea. *Bauh. hist. I. p. 995. Scop. carn. ed.*
n. 870. Duham. arb. p. 257. t. 103.

Habitat in G. Narbonensi, Italia, Sicilia, Turcia, C
niola. H

4. SPARTIUM *monospermum*. S. ramis striatis, racemosis
 lateralibus, foliis lanceolatis.

S. ramis angulatis, racemis lateralibus, foliis lanceolatis.
Sauv. monsp.

S. tertium flore albo. *Bauh. pin. 369.*

S. tertium hispanicum. *Clus. hist. I. p. 103.*

Habitat in Hispaniæ sterilioribus. H

Flores albi

5. SPARTIUM *sphærocarpon*. S. ramis teretibus, foliis
 lanceolatis sessilibus subtus pubescentibus.

S. alterum monospermum, semine reni simili. *Bauh. pin. 396.*

S. hyposphærolobium. *Reneal. spec. 35. t. 33.*

Spartium 1. *Clus. hisp. 203. R.*

Spartium 2. *Ejusd. hist. I. p. 109. R.*

Habitat in Europa australi. H

Caules erecti, ramis teretibus, vix striatis. Flores
 lutei; adeoque a præcedenti distinctissimum.

6. SPARTIUM *purgans*. S. ramis teretibus striatis, foliis
 lanceolatis subsessilibus pubescentibus. *prou. monsp.*

Genista purgans. *Spec. plant.*

DIADELPHIA.

HEXANDRIA.

920. FUMARIA.

Cal. diphyllus. *Cor.* ringens. *Filamenta* 2. membranacea, singula *Antheris* 3.

* *Corollis bicalcaratis.*

1. **FUMARIA** *Cucullaria*. F. scapo nudo.

F. tuberosa insipida. *Corn. canad.* 127.

F. siliquosa, radice grumosa, flore bicorporeo ad labia conjuncto, virginiana. *Pluk. alm.* 162. t. 90. f. 3.
Rai. suppl. 475

Capnorchis americana. *Bærh. lugdb.* 1. p. 309.

Bicucullata canadensis, radice tuberosa squamata. *Az. parif.* 1733.

Habitas in Virginia, Canada. 2

Folia radicalia, bina, trisernata, incisa, lævia, tenuia. Petioli rubicundi. Scapus simplex, teres, longitudine folii, rufus. Racemus terminalis, simplex floribus pendulis : Pedicelli uniflori : Bracteis binis, oppositis, ovatis, acutis, rubicundis, parvis, flori approximatis. Calyx ovatus, appressus, parvus, albus. Corolla alba : Limbus flavus, bilabiatus : Labiis æqualibus, concavis, reflexis, ovatis, integris. Faux clausa, flavens, lateribus margine dilatatis, neclare humentibus. Calcaria duo, æqualia, distantia, compressiuscula, longitudine fere corollæ. Stamina : Filamenta utrinque tria distincta.

* * *Corollis unicalcaratis.*

4. **FUMARIA** *bulbosa*. F. caule simplici, bracteis longitudine florum. *ged. austr. carn. sil. Ludw. ed.* t. 151.
litua. delph. burg.

F. caule simplicissimo diphylo, floribus calyce destitutis.
Fl. succ. *

a. *Fumaria bulbosa, cava.*

192 MONADELPHIA. DECANDRIA

Legumina nigra, aspera setis truncatis, sparsis, sis, viscidis.

11. SPARTIUM *scoparium*. S. foliis ternatis solitariis ramis inermibus angulatis. *Fl. succ. gelr. ged. pal. tab. fil. Knorr. del. 1. tab. S. Oeder. den. burg. lugd.*

S. foliis inferioribus ternatis hirsutis, superioribus simplicibus. *Hall. herb. n. 354.*

Cytiso-genista scoparia vulgaris, flore luteo. *Ta inf. 649 Du Ham. arb. 1. t. 84. R.*

Genista angulosa & scoparia. *Bauh. pin. 395.*

Genista. *Dod. pempt. 761. Riv. 1. 65. Blackw. 1.*

Habitat in Europæ australioris arenosis. ¶

12. SPARTIUM *radiatum*. S. foliis ternatis linearifolius, petiolis persistentibus, ramis oppositis latis. *delph.*

S. æquicolorum minimum montanum triphyllum. *ecphr. 1. p. 294. t. 295.*

Genista radiata. *Scop. carn.*

Genista radiata s. stellaris. *Bauh. hist. 1. p. 399.*

Habitat in Italia, Carniolia. ¶

Caules humiles: Ramis oppositis, tetragonis. Foliis posita, ternata, subsessilia: Foliolis sessilibus, linearibus, subpubescentibus. Petioli omnium brevissimi, persistentes, trigoni, gibbi, obtusissimi, ramulo fulciantes flores. Flores terminales, terni, sessiles, Bractea decidua suffulti.

13. SPARTIUM *spinosum*. S. foliis ternatis, angulatis spinosis. *Sauv. monsp.*

Acacia trifolia. *Bauh. pin. 392.*

Aspalathus secunda trifolia. *Bauh. hist. 1. p. 375.*

Habitat in Europæ australis asperis maritimis. ¶

930. GENISTA.

Cal. bilabiatus: 2/3. Vexillum oblongum a staminibusque deorsum reflexum.

* *Inermes.*

1. GENISTA *canariensis*. G. foliis ternatis unguiculatis, ramis angulatis.

DIADELPHIA. HEXANDRIA. 187

F. alba. *Mill. dict. n. 3.*

F. foliis pinnatis, pinnis pinnatis, pinnulis lobatis, ovatis, filiquis brevibus incrispatis. *Hall. helu. n. 347.*

F. sempervirens & florens, flore albo. *Herm. bat.*

Habitat in Helvetia, Gallia, Italia, Carniolia, in muris Goettingæ. *Weber. spicil. p. 20. ☉*

Semper diandram, multoties examinatam reperit *Weber. l. c. R.*

8. FUMARIA enneaphylla. **F. foliis triternatis**, foliolis cordatis.

F. hispanica saxatilis, foliis amplioribus cordiformibus, semine compresso. *Tournef. inst. 422.*

F. enneaphyllos hispanica saxatilis. *Bocc. mus. 2. p. 83. z. 73. Rai. suppl. 475. Barr. ic. 42.*

F. radice fibrosa, foliis ad petiolum sinuatis crassioribus. *Pluk. alm. 162.*

Habitat in Hispaniæ, Siciliæ saxosis.

9. FUMARIA officinalis. **F. pericarpis monospermis racemosis**, caule diffuso. *Litua. burg. lugd. ged. gallob. carn. Regn. bot. pal. sil. Ludw. ed. t. 83. Blackw. t. 237. Kniph. ceut. 1. n. 33.*

F. pericarpis monospermis. *Fl. suéc.*

F. foliis multifidis, subrotunde lanceolatis, fructibus monospermis. *Hall. helu. n. 346.*

F. officinarum & dioscoridis. *Bauh. pin. 143.*

Fumaria. *Fuchf. hist. 338. Cam. epit. 490. Riv. tetrap. t. 1.*

Habitat in Europæ agris cultis. ☉

10. FUMARIA capreolata. **F. pericarpis monospermis racemosis**, foliis scandentibus subcirrhosis. *Flor. dan. t. 340. delph.*

F. viticulis & capreolis plantis vicinis adhærens. *Bauh. pin. 143.*

F. major scandens, flore pallidiore. *Rai. hist. 405.*

Habitat in G. Narbonensi, Anglia, Dania. ☉

Pedunculi partiales recurvati sunt, & **Foliola extrema in cirrhos transeunt.**

Varietas *Fum.* officinalis *Gerad. at singulare*, quod non occurrat in *Sueciæ agris. s. frutetis*, ubi *F. officinalis frequens.*

11. FUMARIA spicata. **F. pericarpis monospermis spicatis**, caule erecto, foliolis filiformibus. *Sauv. monf. delph.*

F. minor tenuifolia, caulibus procumbentibus & caducis. *Bauh. pin. 194.*

394 DIADELPHIA. DECANDRIA.

G. foliis adpressis, ellipticis, sine lanceolato-patente. Hall. helu. n. 353.

G. herbacea l. Chamæ-sparrum. Bauh. hist. 1. p. 393.

Chamæ-Genista sagittalis. Bauh. pin. 395.

Chamæ-Genista sagittalis pannonica. Cam. hort. 1. 13.

Habitat in Germaniæ, Galliæ arenosis sterilibus. 24

Siliqua tetrasperma. R.

5. *GENISTA tridentata. G. ramis triquetris membranaceis subarticulatis, foliis tricuspидatis.*

Genistella fructicosa lusitanica latifolia. Tournef. inst. 646.

Chamæ-Genista peregrina. Dalechamp. hist. 177.

Chamæ-Genista caule foliato. Bauh. pin. 396.

β. *Genista fructicosa lusitanica angustifolia. Tournef. 646.*

Habitat in Lusitania. Læst. 7

Præcedenti valde affinis, at majora omnia & arborum in folium laterale terminati, non integri, tricuspидati. Legumina albo-lanata.

6. *GENISTA tinctoria. G. foliis lanceolatis glabris, ramis striatis teretibus erectis. Fl. suec. gelr. Cræustr. galob. carn. tub. pal. sil. Oed. fl. dan. 1. 50. Kniph. cens. 6. n. 42. burg. lithu. lugd.*

G. inermis, foliis glabris, confertis, ellipticis, lanceolatis floribus sessilibus, spicatis. Hall. helu. n. 350.

G. tinctoria germanica. Bauh. pin. 395.

G. tinctoria vulgaris. Clus. hist. 1. p. 101.

Genistella. Riv. 1. 67. R.

Habitat in Germania, Angliæ. 7

Stamina 8 & 10 habet. Scop. du Roi. R.

7. *GENISTA sibirica. G. foliis lanceolatis glabris, ramis æqualibus teretibus erectis. (Jacq. hort. 1. 190. R.)*

Habitat in Sibiria. 7

Similis antecedenti, sed Caules non striati aut angulati, vix lineati, neque villositas, ulla in caule aut foliis. Folia angustiora nonnihil. Flores multum remoti, paniculatis.

8. *GENISTA florida. G. foliis lanceolatis sericeis, ramis striatis teretibus, racemis secundis.*

G. tinctoria hispanica. Clus. hist. 1. p. 101.

G. tinctoria frutescens, foliis incanis. Bauh. pin. 395.

Habitat in Hispania. 7

9. *GENISTA pilosa. G. foliis lanceolatis obrutis,*

DIADELPHIA. OCTANDRIA. 189

Amaritie mox distinguitur a P. vulgari simili.

6. **POLYGALA vulgaris.** P. floribus cristatis racemosis , caulibus herbaceis simplicibus procumbentibus , foliis lineari-lanceolatis. *Ged. austr. carn. pal. sil. Flor. dan. t. 516. Kniph. cent. II. n. 84. burg. lithua. lugd.*

P. foliis lineari-lanceolatis , caulibus diffusis herbaceis. *Fl. suec.*

P. caule erecto , foliis linearibus acutis. *Hall. helu. n. 344.*

P. caulibus fructescentibus declinatis , floribus racemosis ; cristatis , subsecundis : capsulis obcordatis. *De Neckgallob.*

P. major. *Bauh. pin. 215.*

P. vulgaris major. *Clus. hist. 1. p. 324. Vaill. paris. 161. t. 32. f. 1.*

P. vulgaris. *Bauh. pin. 215.*

Habitat in Europæ pratis , & pascuis siccis. 2f

Apud nos nunquam fere erecta , sed procumbens.

7. **POLYGALA monspeliaca.** P. floribus cristatis racemosis , caule erecto , foliis lanceolato-linearibus acutis. *Sauv. monsp. delph.*

P. annua erecta angustifolia , floribus coloris obsoleti carneis lineis virgatis. *Magn. monsp. 208.*

P. acutioribus foliis monspeliaca. *Bauh. pin. 207.*

P. vulgaris coloris obsoleti , foliis angustissimis. *Bauh. hist. 3. p. 318. Rai. hist. 1336.*

Habitat Monspeli in collibus sterilibus. ☉

Varietas vulgaris secundum Gerard.

9. **POLYGALA sibirica.** P. floribus cristatis , racemo laterali nudo , caulibus herbaceis , foliis lanceolatis. *Gmel. sib. 4. p. 64. t. 32.*

P. multicaulis. *Amm. ruth. n. 43 , 44 , 45.*

Habitat in Sibiria. D. Gmelin.

Habitus antecedentis. Folia inferiora ovata ; superiora lanceolata. Racemus unus (raro duo) , lateralis , latus , absque bracteis , ad exortum pedicellorum partialium.

* * *Imberbes (flores absque penicillo carinali) frutescentes.*

20. **POLYGALA Chamæbuxus.** P. floribus imberbibus sparsis : carina apice subrotundo , caule fruticoso , foliis lanceolatis. *Jacq. austr. t. 233. delph.*

Polygaloides procumbens , foliis duris , ovatis , nervo aristato. Hall. helu. n. 345.

Chamæbuxus flore coluteæ. Bauh. pin. 471.

296 **DIADELPHIA. DECA NDRIA.**

Conf. Genista sylvestris. Scop. carn. ed. 2. n. 875.

24. **GENISTA lusitanica.** G. caule aphylo, spinis decussatis.

Genista spartium spinosum minus. Bauh. pin. 394

Scorpius secundus. Clus. hist. 1. p. 107. Bauh. hist. 1. p. 401.

Habitat in Lusitania, Hispania. H

932. **U L E X.**

Cal. 2. phyllus. Legumen vix calyce longius.

1. **ULEX europæus.** U. foliis villosis acutis, spinis sparsis.
Neck. gallob. Fl. dan. t. 608. lugd. burg.

U. folio sub singulis spinis subulato plano acuto. *Hort. ups.*

Genista spinosa major longioribus aculeis. Bauh. pin. 394

Genista spinosa. Dod. pemps. 659.

2. *Genista spinosa major, brevioribus aculeis. Bauh. pin. 394.*

Habitat in Anglia, Gallia, Dalmatia, Brabantia. H

933. **A M O R P H A.**

Cor. vexillum ovatum, concavum. Alae o. Carinae

1. **AMORPHA fruticosa.** Amorpha. 353. t. 19. Mill. & ic. t. 27. *Duham. arb. 1. t. 16. Kniph. cent. 10. t. 2.*

Barba Jovis americana, pseudo-acaciae flosculis purpureis minimis. Anglic. hort. 11. t. 4.

Habitat in Carolina. H

935. **O N O N I S.**

Cal. 5. partitus: laciniis linearibus. Vexillum fructum. Legumen turgidum, sessile. Filamentum connata absque fissura.

** Floribus subsessilibus.*

1. **ONONIS antiquorum.** O. floribus solitariis foliolis majoribus, foliis inferioribus ternatis; ramis læviusculis spinosis. *delph.*

Anonis legitima antiquorum. Tournef. cor. 28.

Habitat in Europa australi. 24

Similis O. spinosa, sed brevior, rigidior, majoris fructibus.

MONADELPHIA. DECANDRIA. 198

Genista s. **spartium purgans**. *Bauh. hist. 1. p. 404. Tournef. hist. 644.*

Habitat Monspeli. $\bar{\eta}$

Rami inermes. Legumina ovata, pendula, hinc sinuata.

7. SPARTIUM scorpius. *S. ramis spinosis patentibus; foliis ovatis. monsp. prou. delph.*

Genista spartium spinosum majus, flore luteo. *Bauh. pin. 394.*

Aspalathus secunda monspeliensis. *Bauh. hist. 1. p. 402.*

8. Genista spartium spinosum majus secundum, flore pallido. *Bauh. pin. 394.*

7. Genista spartium spinosum majus tertium hirsutum. *Bauh. pin. 394.*

Aspalathus alter III. *Clus. hist. 1. p. 106.*

Habitat in Hispania. *G. Narbonensi.* $\bar{\eta}$

Frutex totus ex spinis alternis, quibus flores insident.

* * *Foliis ternatis.*

9. SPARTIUM patens. *S. foliis ternatis, ramis virgatis, floribus lateralibus geminis cernuis.*

Cytisus lusitanicus, medicæ folio, floribus in foliorum alis. *Tournef. inst. 648.*

Habitat in Lusitania. $\bar{\eta}$

Frutex Ramis virgatis, patentibus, striatis. Folia ternata, parva: foliolis obovatis, subæqualibus, subtus subpubescentibus. Flores laterales, magni, tumidi, flavissimi, penduli deorsum, bini: vexilli dorso macula fulva. Semen hirsutum.

10. SPARTIUM complicatum. *S. foliis ternatis: foliolis conduplicatis, caulibus inermibus prostratis glabris, leguminibus scabris.*

Cytisus ramis hamifusis albidis, floribus capitatis terminalibus, foliolis ovalibus glabris aggestis. *Sauv. monsp.*

Cytisus foliis viridibus angustis quasi complicatis, monspeliensis. *Bauh. 391. Pluk. alm. 529. t. 86. f. 1.*

Cytisus foliis incanis angustis quasi complicatis. *Bauh. pin. 390. Guett. stamp.*

Cytisus montis calcaris. *Bauh. hist. 1. p. 370.*

Cytisus II. *Clus. hist. 1. p. 94. hisp. 190. ic. 192.*

Habitat in Hispania, Gallia australiore. $\bar{\eta}$

Caules pubescentes. Foliola lanceolata, nunquam explicata. Stamina alterna antheris magnis; alterna parvis-

THE UNIVERSITY OF CHICAGO
LIBRARY

1961
JAN 10 1961

1961

1961

1961

1961

1961

1961

1961

DIABELPHIA. DECANDRIA. 194

Habitat in Europæ australis maritimis ; ex Hispania. Alfræmer.

Facies *Euphrasie*, *palmaris*, *ramosa*, *diffusa*, *viscoso-pubescentis*. Stipulæ *cordatæ*, *latissimæ*, *plicatæ*, *dentatæ*, *emarginatæ*, *foliis latiores*. Folia *simplicia*, *cuneiformi-obovata*, *plicata*, *argute denticulata*. Flores *axillares*, *solitarii*, *subpedunculati*. Calyces *corolla breviores*.

* * *Floribus pedunculatis : pedunculo mutico.*

3. *ONONIS pubescens*. O. pedunculis *muticis brevissimis*, foliis *superioribus simplicibus*, stipulis *ovato-lanceolatis integerrimis*.

O. (*Morissoni*) pedunculis *unifloris nudis*, foliis *ovatis : imis ternatis*, *summis simplicibus*, calycibus *maximis*. *Gouan. illustr. R.*

Anonis purpurea procumbens verna præcox fruticosa annua viscosa, siliquis *crassioribus lentiformibus rarius dispositis*. *Morif. hist. 2. p. 168.*

Habitat in Europa australi. ☉ Ex Insulis balearicis. Gouan.

Caulis *pedalis*, *ramosus*, *diffusus*, *teretes*, *uti totæ plantæ pubescenti-viscidus*, Ramorum *apicibus congestis foliis floribusque*, qui *sensim remotiores evadunt*. Folia *inferiora ternata : reliqua simplicia*. Foliola *ovata s. ovalia*, *obtusæ*, *acute serrata*. Stipulæ *ovato-lanceolatæ*, *petiolo adnatæ ejusque longitudinis*, *integerrimæ*. Petioli *longitudine foliorum*. Pedunculi *axillares*, *mutici*, *longitudine petiolorum*. Calyces *lanceolati*, *obtusiusculi*, *longitudine corollæ*.

14. *ONONIS reclinata*. O. pedunculis *muticis unifloris*, foliis *ternatis subrotundis crenatis*, leguminibus *cernuis*, *Anonis pusilla hirsuta*, flore *ex albo & rubro colore mixto*. *Morif. hist. 2. p. 171.*

Anonis purpurea non spinosa minor. Barr. ic. 354. mala.
Anonis annua pumila, flore *purpurascente*. *Tournef. inst. 408.*

Habitat in Delphinatu, Hispania, Italia. ☉

Planta *viscoso-pubescentis*, *parva diffusa*, *patula*. Folia *ovato-lanceolata*, *ternata*, *orbiculata*, *crassiuscula*, *plicata*, *crenatis*, *viscoso-pubescentia*. Stipulæ *ovales*, *patulæ*, *obtusæ*, *adnatæ*. Pedunculi *axillares*, *uniflori*, *mutici*, *longitudine foliorum*. Calyces *longitudine pedunculi obtusi*. Corollæ *albida*, *perillo subius purpureo*. Calyx *deffloratus reclinatus ad caulem*. Legumen *calyce sesquiflongius*.

1444. **ORCANDRIA**

Orcandria L. pedunculata uniflora
 foliis oppositis, stipulis teretibus. Calibus pro-

longis, vaginis lanceolatis cristatis
 distans pedunculata 47. t. 12.

Orcandria pedunculata multiflora *Hort.* 6.

Orcandria *Cenifu*, *Lamourea* 22.

Orcandria *apicalis*, *subramosa*, *arcta*, *P.*
arcta, *cristata*, *apice mucronata*, *P.*

Orcandria *Sagittata lanceolata*, *terrestris*
arcta, *arcta*, *arcta*, *foliis* *aristatis*
arcta, *arcta*; *Corallia* *Fraxilla* *sp.*

*** *Pedunculus ariflorus*.
Orcandria *aristata* *O. pedunculata* *uniflora* *ar-*
istata, *stipulis* *teretibus*. *Ger. p. 104*

Orcandria, *calycibus* *ad* *aristam* *foliis* *aristatis* *ar-*
istata *terrestris* *aristata*. *Saxr. monsp.*

Orcandria *aristata* *viscosa* *et* *viscosa*, *parviflora*
aristata. *inf. 404*

Orcandria *aristata* *aristata* *aristata* *aristata*. *Hort.*
aristata

Orcandria *G. Narbonne*, *Hispania*, *Italia*.
Orcandria *O. multiflora*, *fol. Peta* *foliis* *aristatis*

Orcandria, *aristata*. *Sagittata* *divergens*. *aristata*, *ar-*
istata, *aristata*, *aristata*, *aristata* *f. aristata*. *Orcandria*

Orcandria, *aristata* *aristata*; *aristata*, *ar-*
istata.

17. **ONONIS** *viscosa*. *O. pedunculata* *uniflora* *ar-*
istata *aristata*: *aristata* *aristata*. *Ger. p. 104*

O. pedunculata *uniflora* *aristata* *aristata*, *foliis* *ar-*
istata. *Saxr. monsp.*

Ononis *lutea* *viscosa* *aristata* *aristata*, *aristata* *aristata*
aristata. *Sp. 4. 1239*

Ononis *viscosa*, *stipulis* *aristatis* *lutea* *Mag.* *aristata*
Ononis *aristata* *aristata* *aristata* *aristata*

aristata. *inf. 409*
Ononis *aristata*, *O. in Hispania* *O*
aristata *aristata* *O. Calyces* *aristata* *aristata*

18. **ONONIS** *omnipedunculata*. *O. pedunculata* *ar-*
istata, *leguminibus* *aristatis* *aristata*.

DIADELPHIA. DECANDRIA. 198

tuberculato decumbente. *Fl. succ. Jacq. austr. t. 208, carn. pal. Kniph. cent. 5. n. 33. burg. lugd. delph.*

G. inermis, procumbens, foliis duris, subhirsutis, spiciis brevibus, foliosis. *Hall. helu. n. 351.*

G. ramosa, foliis hyperici. *Bauh. pin. 395.*

Chamæ-Genista foliis genistæ vulgaris. *Bauh. pin. 395.*

Chamæ-Genista montana hispida. *Bauh. pin. 396.*

Genistella pilosa. *Bauh. hist. 1. P. 2. p. 393. Hall. R.*

Chamæ-Genista prima. *Clus. hist. 1. p. 103. pann. 49.*

Habitat in Pannonia. G. Narbonensi, Germania. H

Siliquæ poly. & disperma. R.

10. GENISTA. *humifusa.* **G.** foliis lanceolatis ciliatis, ramis prostratis striatis villosis. *delph.*

G. orientalis minima humifusa, foliis subrotundis ad hores pilosis. *Tournef. cor. 44.*

Habitat in Oriente D. Gerard.

** * Spinozæ.*

11. GENISTA *anglica.* **G.** spinis simplicibus, ramis fl. a-
riferis inermibus, foliis lanceolatis. *Flor. dan. t. 619, delph.*

G. minor asphaltoides. *Bauh. pin. 395. prodr. 157.*

Genistella aculeata. *Lob. ic. 2. t. 93.*

Genistella. *Dod. pempt. 670.*

Genista-Spartium minus anglicum. *Tournef. inst. 645. R.*

Habitat in Angliæ ericetis humidiusculis. H

12. GENISTA *germanica.* **G.** spinis compositis, ramis flo-
riferis inermibus, foliis lanceolatis. *Gelr. Crantz. austr. carn. tub. pal. sil. Kniph. cent. 11. n. 49. burg. delph. lugd.*

G. foliis ovato-lanceolatis, hirsutis, spinis ramosis, in ramis senescentibus. *Hall. helu. n. 352.*

G. spinosa major germanica. *Bauh. pin. 395.*

Genistella spinosa. *Riv. tetr. 67. aculeata. Tabern. 1101.*

Habitat in Germania. H

Siliqua trisperma. Hall. R.

13. GENISTA *hispanica.* **G.** spinis decompositis, ramis flo-
riferis inermibus, foliis linearibus pilosis. *monsp. delph.*

Genista spinosa minor hispanica villosissima. *Bauh. pin. 395.*

Genistella monspeliaca spinosa. *Bauh. prodr. 157.*

Genistella, montis ventosi, spinosa. *Bauh. hist. 1. p. 400.*

Habitat in Hispania, G. Narbonensi. H

Spinæ innocuæ, G. germanicæ adfinis.

N. R.

800 **DIADELPHIA. DECAENDRIA.**

15. **ONONIS** *cenisia*. O. pedunculis muticis unifloris, foliis ternis cuneatis, stipulis serratis, caulibus prostratis. *delph.*

Anonis isermis glabra, vaginis lanceolatis cristatis, perianthiis longis unifloris. *Allion. pedem. 47. t. 10. f. 1. bona.*

Anonis luteo-purpurea minima angustifolia. *Barr. ic. 83. t. 1104.*

Habitat ad pedem montis Cenisi. Latourette. 24

Caulis prostratus, digitales, subramosi, laeves. Folia ternata. Foliola laevia, cuneata, apice rotundata, argenteo-minimeque serrata. Stipulae lanceolatae, serratae. Pedunculi laterales, uniflori, mutici, foliis multo longiores. Flores Calyce brevi, lavi; corollae Vexillo purpureo striato.

*** *Pedunculis aristatis.*

16. **ONONIS** *cherleri*. O. pedunculis unifloris aristatis, foliis ternatis, stipulis serratis. *Ger. prou.*

Ononis trifolia, calycibus adscissis, lobis sessilibus erectis, stipulis longas setas exferentibus. *Sauv. monsp.*

Anonis pusilla villosa & viscosa, purpurascens flore. *Tournef. inst. 408.*

Anonis trifolius minima hirculensis *cherleri. Bauh. hist. 3. p. 394.*

Habitat in G. Narbonensi, Hispania, Italia. 24

Affinis O. minutissima, sed Folia subtus villosa-viscida, nec glabra. Stipulae divergentes, serratae, apice lanceolatae, nec vaginantes, integerrimae s. setaceae. Calyx pubescentes, longitudine corollae; nec glabri, corollae longiores.

17. **ONONIS** *viscosa*. O. pedunculis unifloris aristatis, foliis simplicibus: inferioribus ternatis. *Ger. prou. delph.*

O. pedunculis unifloris filis terminatis, foliis ternatis simplicibusque. *Sauv. monsp.*

Anonis lutea viscosa latifolia minor, flore pallido. *Barr. rar. 840. t. 1239.*

Anonis viscosa, spinis carens lutea. *Magn. monsp. 211*

Anonis annua erectior latifolia glutinosa lusitanica. *Tournef. inst. 409.*

Haec ut Monspeli, & in Hispania. O

Varietas B major & Calyces corolla longiores.

18. **ONONIS** *ornithopodioides*. O pedunculis bifloris multifloris, leguminibus linearibus cernuis.

DIADELPHIA. DECANDRIA. 199

fere glabra : pedunculis solitariis , bractea duplo longioribus.

2. *ONONIS arvensis*. O. floribus racemosis geminatis, foliis ternatis superioribus solitariis, ramis inermibus subvillosis. *gallob. ged. austr. pal. sil. sub. Blackw. t. 301. f. 1. 2. burg. lugd. lithu.*

Anonis caule decumbente , ramis senescentibus spinosis. Hall. helu. n. 316.

Ononis floribus lateralibus , solitariis : spinis rameis terminalibus foliosis. Scop. carn.

Anonis spinosa. Scop. carn. ed.

3. *Ononis (mitis)* floribus subsessilibus solitariis lateralibus; ramis inermibus. *Fl. suec.*

Anonis spinis carens purpurea. Bauh. pin. 329.

Anonis mitior. L. Clus. hist. 99.

4. *Ononis (spinosa)* floribus subsessilibus lateralibus, caule spinoso. *Hort. cliff. Kniph. cent. 5. n. 61. Regn. bot.*

Anonis spinosa , flore purpureo. Bauh. pin. 329.

Anonis. Fuchs. hist. 60. Ruz. t. 69.

Habitat in aridis, Europæ. 24

Folia in caule ternata ; Floralia vero simplicia , Flores duo propriis pedunculis exeunt simul. Planta pubescenti-subviscida ; prima æstate inermis ; adultior spinosa evadit , ut sæpe observavit Læf. pruss. 17.

Floribus albis variat. R.

3. *ONONIS repens*. O. caulibus diffusis, ramis erectis; foliis superioribus solitariis, stipulis ovatis.

Anonis maritima procumbens, foliis hirsutis pubescentibus. Pluk. alm. 33. Dill. elth. 29. t. 25. f. 18.

Habitat in Angliæ litoribus maris. Oriente. 24

Affinis admodum præcedenti , differt caulibus procumbentibus undique diffusis , & quod minor ; Flores ex alis solitarii.

4. *ONONIS minutissima*. O. floribus subsessilibus lateralibus, foliis ternatis glabris, stipulis ensiformibus, calycibus scariosis corolla longioribus. *Jacq. austr. t. 240. burg.*

Anonis foliis ternatis, ovatis serratis, hirsutis, floribus sessilibus, calycibus hiantibus. Hall. helu. n. 359. R.

Ononis trifolia, calycibus ad axillas sessilibus erectis & stipulis longas setas exercentibus. Sauv. monsp. 190.

O. floribus lateralibus sessilibus solitariis, petiolis longissimis. Roy. lugd.

O. lutea sylvestris minima. Coh. aphr. t. p. 304. t. 301.

DIADREPETA DECANDRIA

forma lutea minor. Benth. p. 349.

uirg. dentata non spinosa minima. *Barr. ic. 1107.*

Monspeliu, Helvetia, Austria.

Siliques subformes, diffusi, glabri, suffruticosi. Stipulae
~~cuneatae~~, subbipartitae, integerrimae. Folia ternata,
~~pinnatis~~, petiolata : Foliolis oblongiusculis, rigidulis, ser-
~~vatis~~, recurvatis, viz mucronatis. Petioli brevissimi. Flores
 axillares, foliarii. Calyces glabri, subcarinosi, profunde
 quinquefidi, recti, subulati, corollae longiores. Corollae
 limbo. Legumen ovatum, fuscum, calyce brevius. Folia
 floralia superiora simpliciora.

♂. *ONONIS mitissima*, floribus sessilibus spicatis, bracteis stipularibus ovatis ventricosis scariosis imbricatis.

O. stipulis floralibus ovatis membranaceis integerrimis. Roy.
ludb.

Anonis alopecuroides mitis annua purpurascens, *Dill. elth.*
23, 4. 24. f. 27.

Anonis purpurea spiceta erecta annua latifolia, filiquis rectis
lentiformibus. *Meris. hist.* 2. p. 169, f. 2. t. 17. f. 4.

Woburn in Barbados, Lusitania. ☉

Folia caulina cernua; floralia simplicia. Siliquæ & Bracteæ amplexicaules, membranaceo-carilagineæ, læves, ovatae, apicem ad flores imbricatae in spicam, calyces obtegentes. Calycis laciniæ ovatae, lateribus reflexæ, intus adpersæ setis viscosis ut filiqua.

ONONIS alopecuroides. O. spicis foliosis foliis simplicibus
ovatis obtusis, stipulis dilataris. *Kniph. cent. 8. n. 75.*

Q. floribus spicatis. Mort. cliff.

Anonis ficula alopecuroides. *Tournef. inst.* 408.

Anonis spicata L. alopecuroides hufianica non spinosa. Rai.
suppl. 468.

Anonis purpurea spicata. *alopecuroides* major. *Berk. lugd.*
2. p. 22.

Nabices in Sicilia, Lusitania, Hispania. ☉

7. *ONONIS variegata*. O. stipulis bracteisque densatis, foliis simplicibus striatis, floribus subpedunculatis. Roy. Lugdb. ⁴

Anonis lutea, trixaginis folio, maritima. Barr. ic. 776.
mala.

Anonis maritima mitis, folio lucido ferrato. Boer. sic. 7.
s. 32. f. 3.

Anonis non spinosa, flore luteo variegato, angustifolia
maritima. *Tournef. inf.* 409.

DIADELPHIA. DECANDRIA. 194

Habitat in Europæ australis maritimis ; ex Hispania. Alfræmer.

Facies Euphrasie , palmaris , ramosa , diffusa , viscoso-pubescent. Stipulæ cordatæ , latissimæ , plicatæ , dentatæ , emarginatæ , foliis latiores. Folia simplicia , cuneiformi-obovata , plicata , argute denticulata. Flores axillares , solitarii , subpedunculati. Calyces corolla breviores.

* * *Floribus pedunculatis : pedunculo mutico.*

8. **ONONIS pubescens.** O. pedunculis muticis brevissimis , foliis superioribus simplicibus , stipulis ovato-lanceolatis integerrimis.

O. (*Morissoni*) pedunculis unifloris nudis , foliis ovatis : imis ternatis , summis simplicibus , calycibus maximis. *Gouan. illustr. R.*

Anonis purpurea procumbens verna præcox fruticosa annua viscosa , siliquis crassioribus lentiformibus rarius dispositis.

Morif. hist. 2. p. 168.

Habitat in Europa australi. ☉ Ex Insulis balearicis. Gouan.

Caulis pedalis , ramosus , diffusus , teretes , uti tota planta pubescenti - viscidus , Ramorum apicibus congestis foliis floribusque , qui sensim remotiores evadunt. Folia inferiora ternata : reliqua simplicia. Foliola ovata s. ovalia , obtusa , acute serrata. Stipulæ ovato - lanceolatæ , petiolo adnatæ ejusque longitudinis , integerrimæ. Petioli longitudine foliorum. Pedunculi axillares , mutici , longitudine petiolorum. Calyces lanceolati , obtusiusculi , longitudine corollæ.

14. **ONONIS reclinata.** O. pedunculis muticis unifloris , foliis ternatis subrotundis crenatis , leguminibus cernuis ,

Anonis pusilla hirsuta , flore ex albo & rubro colore , mixto. *Morif. hist. 2. p. 171.*

Anonis purpurea non spinosa minor. *Barr. ix. 354. mala.*

Anonis annua pumila , flore purpurascente. *Tournef. inst. 408.*

Habitat in Delphinatu , Hispania , Italia. ☉

Planta viscoso-pubescent , parva diffusa , patula. Folia petiolota , ternata , orbiculata , crassiuscula , plicata , crenata , viscoso-pubescentia. Stipulæ ovales , patulæ , obtusæ , serratæ. Pedunculi axillares , uniflori , mutici , longitudine vix foliorum. Calyces longitudine pedunculi obtusi. Corollæ albidæ vexillo subtus purpureo. Calyx defloratus reflectitur apice ad caulem. Legumen calyce sesquilongius villosum.

100 DIADELPHIA. DECANDRIA.

15. *ONONIS cenisia*. O. pedunculis muticis unifloris, foliis ternis cuneatis, stipulis serratis, caulibus prostratis. *delph.*

Anonis inermis glabra, vaginis lanceolatis cristatis, petiolis longis unifloris. *Allion. pedem. 47. t. 10. f. 3. bona.*

Anonis luteo-purpurea minima angustifolia. *Barr. ic. 833. t. 1104.*

Habitat ad pedem montis Cenicii. Latourette. 24

Caules prostrati, digitales, subramosi, laeves. Folia ternata. Foliola laevia, cuneata, apice rotundata, argute tenuissimeque serrata. Stipulae lanceolatae, serratae. Pedunculi laterales, uniflori, mutici, foliis multo longiores. Flores Calyce brevi, laevi; Corollis Vexillo purpureo, striato.

* * * *Pedunculis aristatis.*

16. *ONONIS cherleri*. O. pedunculis unifloris aristatis, foliis ternatis, stipulis serratis. *Ger. prou.*

Ononis trifolia, calycibus ad axillas sessilibus erectis, stipulis longas setas exferentibus. *Sauv. monsp.*

Anonis pusilla villosa & viscosa, purpurascente flore. *Tournef. inst. 408.*

Anonis trifoliis minima hirsuta cherleri. *Bauh. hist. 21 P. 334.*

Habitat in G. Narbonensi, Hispania, Italia. 24

Affinis Q. minutissima, sed Folia subtus villosa-viscida, nec glabra. Stipulae divergentes, serratae, apice lanceolatae, nec vaginantes, integerrimae s. setaceae. Calyces pubescentes, longitudine corollae; nec glabri, corolla longiores.

17. *ONONIS viscosa*. O. pedunculis unifloris aristatis, foliis simplicibus: infimis ternatis. *Ger. prou. delph.*

O. pedunculis unifloris filo terminatis, foliis ternatis simplicibusque. *Sauv. monsp.*

Anonis lutea viscosa latifolia minor, flore pallido. *Barr. rar. 840. t. 1239.*

Anonis viscosa, spinis carens lutea. *Magn. monsp. 21?*

18. *ONONIS annua erectior latifolia glutinosa lusitanica*. *Tournef. inst. 409.*

Habitat Monspelii, & in Hispania. ☉

Varietas β major & Calyces corolla longiores.

18. *ONONIS ornithopodioides*. O pedunculis bifloris aristatis, leguminibus linearibus cernuis.

DIADELPHIA. DECANDRIA. 201

Anonis filiquis ornithopodii. *Bærh. lugub. 2. p. 34.*

Foenum græcum ficulum, filiquis ornithopodii. *Tournef. inst. 409.*

Habitat in Sicilia ad Capo passaro.

Caulis erectus; flexuosus: Ramis alternis, brevibus. Folia petiolis oblongis, viscosa: foliolis; obovatis serratis: intermedio duplo majore. Stipulae semicordatae. Pedunculi petiolo longiores. Legumina linearia compressa, torulosa, cernua. Calycis foliola filiformia.

19. *ONONIS pinguis*. O. pedunculis unifloris aristatis, foliis ternatis lanceolatis, stipulis integerrimis.

Anonis non spinosa, flore luteo variegato. *Bauh. pin. 389.*

Anonis lutea non spinosa, *Natrix* Plinii. *Dalech. hist. 449.*

β *Anonis* lutea mitis oxytriphylia ad foliorum petiolos capreolata. *Pluk alm. 33. t. 135. f. 5.*

Habitat in Europa australi. ♀

Similissima O. *Natrix*, sed Caulis succosior, magis angulatus; Folia majora, lanceolato-oblonga; Arista pedunculorum longitudine floris, quam in precedenti breviores.

Varietas O. *Natrix* secundum Gerard.

20. *ONONIS Natrix* O. pedunculis unifloris aristatis, foliis ternatis viscosis, stipulis integerrimis, caule fruticoso. * *monsp. burgd. lugd.*

Anonis viscida, foliis petiolatis ternatis, petiolis unifloris.

Hall. helv. n. 358.

Anonis Natrix. *Scop. carn.*

Anonis viscosa spinis carens lutea major. *Bauh. pin. 389.*

Ononis lutea. *Cam. epit. 445.*

Natrix. *Riu. tetr. 68. Lob: ic. 2. p. 28.*

Habitat in G. Narbonensi, Hispania: inter segetes. ♀

Planta graveolens, odore theriacæ, parum viscida. Folia ovalia; planiuscula, apice tantum serrata, non carnosæ. Pedunculus aristæ longitudine. Calyces visciidi.

* * * * Fruticosa.

21. *ONONIS tridentata*. O fruticosa, foliis ternatis carnosis sublinearibus tridentatis, pedunculis bifloris.

Anonis hispanica frutescens, folio tridentato carnosio. *Tournef. inst. 408 Magn. hort. 16. t. 16. Rai. suppl. 466.*

Habitat in Hispania. ♀

Caulis erectus, paniculatus. Stipulae subulatae minutæ. Foliola linearia, carnosæ, subcanaliculata, apice tridentata, ad latera denticulis obsoletis. Racemus terminalis,

102. DIADELPHIA. DECANDRIA.

parrus hispidoglutinosus : pedunculis bifloris. Flos *purpureus*.

22. *ONONIS crispa*. O fruticosa, foliis ternatis subrotundis undulatis dentatis viscoso, pubescentibus, pedunculis, unifloris muticis. *delph.*

Anonis hispanica frutescens, folio rotundiore. *Magn. monsp.* 17. t. 17. *Rai. suppl.* 466.

β *Anonis hispanica frutescens*, folio rosæ sylvestris. *Tournef. inst.* 409.

Habitat in Hispaniæ monte Mariola prope Valentiam. ἥ

Caulis erectus, paniculatus, pubescenti-viscosus. Foliola obovata, obtusa, recurvata, canaliculata, plicata, subdentata, crispa. Stipulæ bifidæ, amplexicaules, dentatæ, uti tota planta piloso-viscidæ. Flores laterales, axillares, solitarii, pedunculo brevissimo. Calyx quinquepartitus: laciniis corolla dimidio brevioribus, subæqualibus, tridentatis. Corolla flava, magnitudine foliorum: vexillo extus sanguineo-striato, ut in *O. Natrice*, cui simillima, sed crispa, magis viscosa & pedunculis muticis, quosque deflorata valde recurvat, ut fructus caute approximetur.

Magnolius flores spicatos pingit.

23. *ONONIS fruticosa*. O fruticosa, foliis sessilibus ternatis lanceolatis serratis, stipulis vaginalibus, pedunculis subtrifloris. *Ger. prou. delph.*

O. fruticosa, floribus paniculatis, pedunculis subtrifloris, stipulis vaginalibus, foliis ternatis lanceolatis serratis. *Hort. cliff.* *

Anonis purpurea frutescens non spinosa. *Dodart. mem.* 57. t. 57.

Anonis purpurea verna præcox frutescens, flore rubro amplo. *Morif. hist.* 2. p. 170. *Mill. dict.* 1. 36. *Duham. arb.* 1. t. 58.

Habitat in montibus Delphinatus. ἥ

Stipulæ vaginantes quadriaristatæ, aridæ.

24. *ONONIS rotundifolia*. O fruticosa, foliis ternatis ovatis dentatis, calycibus triphyлло - bracteatis, pedunculis subtrifloris. *Ger. prou. delph.*

O. fruticosa, pedunculis trifloris, calycibus triphyлло-bracteatis, foliis ternatis subrotundis. *Hort. cliff.* *

Anonis foliis subrotundis serratis, ternatis, petiolis multifloris. *Hall. helu. n.* 357.

Cicer sylvestre latifolium triphyllum. *Bauh. pin.* 347.

DIADELPHIA. DECANDRIA. 205

Cicer sylvestre tertium. Dod. pempt. 525.

Cicer sylvestre verius. Lob. ic. 2. p. 73.

Habitat in Alpibus Helveticis. H

*Vexillum pubescens. Alæ carina breviores. Calyceis
sine bracteis 3, cordatis, approximatis, brevioribus.
Supulæ virentes.*

936. ANTHYLLIS.

Cal. ventricosus. Legumen subrotundum, rectum.

* *Herbaceæ.*

1. **ANTHYLLIS tetraphylla.** A. herbacea, foliis quaternopinnatis, floribus lateralibus. *Hort. ups. Sauv. monsp.*

A. foliis quinto-pinnatis inæqualibus, floribus lateralibus. *Hort. cliff.*

Lotus pentaphyllos vesicaria. Bauh. pin. 331.

Trifolium alicacabum. Cam. hort. 171. t. 47.

Habitat in Italia, Sicilia. ☉

Folium pinnatum tetraphyllum loco pentaphylli, cum foliolum infimum opposito careat. Stamina diadelphea.

2. **ANTHYLLIS Vulneraria.** A. Herbacea, foliis pinnatis inæqualibus, capitulo duplicato. *aust. ged. gallob. pal. sil. Kniph. cent. 6. n. 8. lugd. litu. burg.*

Anthyllis foliis pinnatis, foliolis pluribus: terminali majore. Fl. suec.

Vulneraria foliis radicalibus ovatis, caulinis pinnatis, umbellulis transversis. Hall. helv. n. 398.

Vulneraria anthyllis. Scop. carn.

Anthyllis. Rtu. t. 18. R.

Lotus affinis Vulneraria pratensis. Bauh. pin. 332.

Lagopodium flore luteo. Tabern. ic. 925.

3. *Vulneraria lupina, flore coccineo. Dill. elik. 431. t. 310. f. 413.*

Lotus affinis hirsuta, flore rubente. Bauh. pin. 332.

Vulneraria rustica, flore albo. Tournef. inst. 291.

Habitat in pratis Europæ borealioris. 24

3. **ANTHYLLIS montana.** A. herbacea, foliis pinnatis æqualibus, capitulo terminali secundo, floribus obli-
manis. *Jacq. austr. t. 334. burg.*

939. LUPINUS.

Cal. 2. labiatus. *Antheræ* 5. oblongæ, 5 subrotundæ.
Legumen coriaceum.

1. LUPINUS *albus*. L. calycibus alternis inappendiculatis : labio superiore integro ; inferiore tridentato. *Ludw. ed.* : s. 191. *Blackw. t.* 282. *Kniph. cent.* 4. n. 43. *Regn. bot.* L. caule composito. *Hort. cliff.*

L. sativus, flore albo. *Bauh. pin.* 347. *Clus. hist.* 2. p. 228.

L. flore albo. *Riu. tetr.*

Habitas . . . ☉

Flores albi. Semina compressa, orbiculata.

3. LUPINUS *varius*. L. calycibus semiverticillatis appendiculatis : labio superiore bifido ; inferiore subtridentato. *Kniph. cent.* 7. n. 47.

L. sylvestris, flore cæruleo. *Bauh. pin.* 348.

L. sylvestris, flore purpureo, semine rotundo vario. *Bauh. hist.* 2. p. 291.

Habitas Messanæ, Monspeli inter segetes. ☉

Flores rubri. Foliola linearia.

4. LUPINUS *hirsutus*. L. calycibus alternis appendiculatis : labio superiore bipartito ; inferiore tridentato.

L. lanuginosus latifolius humilis, flore cæruleo purpurascente, stoloniferus. *Shav. afr.* 393.

L. peregrinus major s. villosus cæruleus major. *Bauh. pin.* 348. *prodr.* 148.

L. exoticus hirsutissimus. *Bauh. hist.* 2. p. 289.

Habitas in Arabia, Archipelagi insulis, Hispania. ☉

Sylvestris foliis quinatis, longe petiolatis. Foliola spatulata. Flores cærulei. Legumina, uti tota planta, vestita, pilis ferrugineis densis, longis. Flores huic sæpe laterales.

6. LUPINUS *angustifolius*. L. calycibus alternis appendiculatis : labio superiore bipartito ; inferiore integro. *cent.* 4. n. 46. *Knorr. del.* 2. s. L. 7.

L. angustifolius cæruleus elatior. *Rai. hist.* 908.

L. flore cæruleo minore. *Riu. tetr.*

Habitas in Hispania, Messanæ inter segetes. ☉

Foliola linearia. Flores cærulei,

DIADELPHIA. DECANDRIA. 207

7. **LUPINUS luteus**, L. calycibus verticillatis appendicularis : labio superiore bipartito , inferiore tridentato. *Sauv. monsp. Kniph. cent. 4. n. 47.*

L. sylvestris, flore luteo. *Bauh. pin. 348. Riv. tetr.*

L. flore luteo , semine compresso vario. *Bauh. hist. 24 p. 291.*

Habitat in Sicilia arenosis. ☉

Huic flores lutei , subsessiles , odorati , bracteis ovatis.

940. PHASEOLUS.

Carina cum staminibus styloque spiraliter tortis.]

** Scandentes.*

1. **PHASEOLUS vulgaris**. P. caule volubili , floribus racemosis geminis , bracteis calyce minoribus , leguminibus pendulis.

P. radice annua , caule volubili , leguminibus pendulis compressis torosis. *Hart. upf.*

P. vulgaris. *Lob. ic. 59.*

Smilax hortensis f. **phaseolus major**. *Bauh. pin. 339.*

β. **Phaseolus (coccineus)** volubilis , floribus racemosis geminis , bracteis calyce brevioribus , leguminibus pendulis. *Kniph. cent. 12. n. 75.*

P. puniceo flore. *Cornut. canad. 184.*

P. indicus , flore coccineo f. puniceo. *Morif. hist. 2. p. 69.*

Habitat in Indiis. ☉

γ. **PHASEOLUS nanus**. P. caule erecto lævi , bracteis calyce majoribus , leguminibus pendulis compressis rugosis.

P. vulgaris italicus humilis f. minor , albus cum orbita nigricante. *Bauh. hist. 2. p. 358. Rai. hist. 885.*

Smilax filiqua sursum rigente f. **Phaseolus parvus italicus**. *Bauh. pin. 339.*

Habitat in India.

941. DOLICHOS.

Vexilli basis callis 2. parallelis , oblongis , alas subtus comprimantibus.

1. **DOLICHOS lablab**. D. volubilis , leguminibus ovato

ADELPHIA. DECANDRIA.

— na
te
minibus, seminibus ovatis hilo arcuato vix
trematatem. *Kniph. cent. 6. n. 37.*

Phal. ægyptius nigro semine. *Bauh. pin. 341.*

Phaseolus niger Lablab. *Alp. ægypt. 74. t. 77.*
ægypt. 27.

Habitat in Ægypto. ☉

Legumina dorso scabra. Caules ramique teretes, ret
scabri. Pedunculi semiverticillati.

4. DOLICHOS unguiculatus. D. volubilis, legum
capitatis subcylindricis : apice recurvo concavo.
hort. t. 23.

Cacara nigra. *Rumph. amb. 5. p. 381. t. 138. f.*

Habitat in Barbados.

9. DOLICHOS pruriens. D. volubilis, leguminibus
mosis : valvulis subcarinatis hirtis, pedunculis ternis.
amer. 201. t. 121.

Phaseolus americanus, solli lanugine obsito,
quis pungentibus, se usco punctato. *Pluk. p.*
214. f. 1.

Phaseolus utriusque Indiæ lobi villosis pungentibus
Sloan.

Strizolobium spicis multis pennis alaribus, floribus
Brown. jam. 190.

Cacara pruriens. *Rumph. ai 6. p. 393. t. 142.*

Nai corana. *Rheed. mal. 8. p. 1. Fl. zeyl. 539.*

Habitat in Indiis.

Foliola subtus hirta. Calyx campanulatus, 5 dent
medio brevior. Corollæ vexillum brevissimum, reliqu
satis triplo minus, ovatum. Carina linearis, acuta,
naviculari. Stamina diadelphæ. Filamenta 4. alterna ju
vata, majora. Legumen compressum, basi inflexum,
reflexum, (adeo liseram s. flexura refert.) Hujus & sup
legumina hirta urunt.

944. P I S U M.

Stylus triangulus, supra carinatus pubescens. Cal
laciniæ superiores 2 breviores.

1. PISUM sativum. P. petiolis teretibus, stipulis
rotundatis crenatis, pedunculis multifloris. *Ladv.*
t. 169.

P. stipulis crenatis. *Hort. cliff.*

Pisum. Cam. epit. 213. Blackw. t. 83.

6. *Pisum hortense majus. Bauh. pin. 342.*

7. *Pisum sine cortice duriore. Bauh. pin. 343.*

8. *Pisum umbellatum. Bauh. pin. 344.*

P. (*umbellatum*) stipulis quadrifidis acutis, pedunculis multifloris terminalibus. *Mill. dict. n. 3.*

9. *Pisum (quadratum) majus quadratum. Bauh. pin. 342.*

Habitas in Europæ agris. ☉

Foliola conduplicata.

2. *PISUM arvense. P. petiolis tetraphyllis, stipulis crenatis, pedunculis unifloris. Fl. suec. norv. delph.*

P. pulchrum, folio anguloso. *Bauh. hist. 2. p. 197. Moris. hist. 2. p. 47. f. 2. t. 1. f. 4.*

Habitas inter Europæ segetes. ☉

3. *PISUM maritimum. P. petiolis supra planiusculis, caule angulato, stipulis sagittatis: pedunculis multifloris. Fl. suec. Flor. dan. t. 338.*

P. stipulis integerrimis. *Hort. cliff.*

P. marinum. *Rai. hist. 892.*

P. spontaneum perenne repens humile. *Moris. hist. 2. p. 47. f. 2. t. 1. f. 5.*

Habitas in Europæ, Canada borealis littoribus maris arenosis. 4

Europea planta tota, excepta foliorum pagina superiore, leviter pubescens. Canadensis vero tota glabra & duplo major. Utraque foliolis convolutis.

4. *PISUM ochrus. P. petiolis decurrentibus membranaceis diphyllis pedunculis unifloris. Kniph. cent. 10. n. 71.*

Ochrus folio integro capreolos emittente. Bauh. pin. 343.

Ervilia sylvestris. Dod. pempt. 522.

Habitas inter segetes Cretæ, Italiæ. ☉

Foliola convoluta.

945. OROBUS.

Stylus linearis. Cal. basi obtusus: laciniis superioribus profundioribus, brevioribus.

3. *OROBUS luteus. O. foliis pinnatis ovato-oblongis, stipulis rotundato-lunatis dentatis, caule simplici. delph.*

O. foliis ovato-lanceolatis, stipulis triangulis, caule ramoso herbaceo. Gmel. sib. 4. p. 13. t. 4.

Linnaei spec. Plant. Europ. Pars II.

☉

[illegible]

DIADELPHIA. DECANDRIA. 211

O. orientalis, foliis angustissimis costæ brevissimæ. *Tournef. cor.* 29.

Araci s. Apios leguminosæ species rossano carolo. *Bauh. hist.* 2. p. 326.

Habitat in Sibiria, Gallia. 24

Habitus O. tuberosi, sed Folia ensiformi-lanceolata & pinnata ex 2. f. 3. oppositionibus absque cirrho. Racemi floribus paucis luteis.

7. OROBUS niger. O. caule ramoso, foliis sexjugis ovato-oblongis. *suæc. ged. austr. carn. pom. pal. Kniph. cent.* 3. n. 69. *delph. lith. burg. lugd.*

O. caule ramoso, foliis ovatis duodenis. *Hall. helu. n.* 418.

O. sylvaticus, viciæ foliis. *Bauh. pin.* 352. *Riu. t.* 60.

O. pannonicus 2. *Clus. hist.* 2. p. 230.

Habitat in Europæ borealis montosis. 24

Foliola convoluta.

8. OROBUS pyrenaicus. O. caule ramoso, foliis bijugis lanceolatis nervosis, stipulis subspinosis. *carn.*

O. caule ramoso, foliis trijugis lanceolatis nervosis. *Sauv. monsp.*

O. pyrenaicus, foliis nervosis. *Tournef. inst.* 235.

O. pyrenaicus latifolius nervosus. *Pluk. phyt.* 210. f. 2.

Habitat in Pyrenæis. 24

9. OROBUS sylvaticus. O. caulibus decumbentibus hirsutis ramosis.

O. caule ramoso hirsuto decumbente, foliis subseptemjugis.

O. sylvaticus nostras. *Rai. angl.* 3. p. 324. *hist.* 1892. *Ad. paris.* 1706. p. 87. t. 90.

Habitat in Anglia, Gallia. 24

946. LATHYRUS.

Stylus planus, supra villosus, superne latior.

Cal. lacinia superiores 2 breviores.

* *Pedunculis unifloris.*

1. LATHYRUS Aphaca. L. pedunculis unifloris, cirrhis aphyllis, stipulis sagittato-cordatis. *carn. pal. gallob. lugd. burg. delph.*

L. aphyllus stipulis sagittatis latissimis. *Hall. helu. n.* 442.

L. cirrhys aphyllis. *Hort. upf. paris.*

Vicia lutea, foliis convolvuli minoris. *Bauh. pin.* 345.

210 DIADELPHIA. DECANDRIA.

O. caule ramoso erecto , foliis ovato - lanceolatis. *Hall. helu. n. 419.*

O. foliis quinatis : foliolis ovato-oblongis , stipulis semi-fagittatis , pedunculis longissimis , caule simplici. *Ger. prou.*

O. alpinus latifolius. *Bauh. pin. 351. prodr. 149.*

O. sylvaticus pallido flore. *Bauh. pin. 351.*

Galega montana Dalechampi. Bauh. hist. 2. p. 343.

Habitat in Sibiria, Helvetia, Verona, Pyrenæorum, alpinis. 24

Caulis bicubitalis, simplex s. ramosus. Folia subtus glauca. Stipulae semisagittatae, integerrimae, minime subrotundo-lunatae, aut dentatae. Flores lutei. Turra.

4. *OROBUS vernus.* *O.* foliis pinnatis ovatis , stipulis semisagittatis integerrimis , caule simplici. *ged. gallob. austr. carn. norv. pom. pal. sil. Riu. t. 58. delph. litua. burg. lugd.*

O. caule simplicissimo , foliolis pluribus ovatis acutis. *Fl. suec. Hall. helu. n. 416.*

O. sylvaticus purpureus vernus. *Bauh. pin. 351. Thal. herc. 80. t. 6. Blackw. t. 208. f. 1. 2.*

Habitat in Europæ borealis nemoribus. 24

5. *OROBUS tuberosus.* *O.* foliis pinnatis lanceolatis , stipulis semisagittatis integerrimis , caule simplici. *carn. pom. pal. burg. delph.*

O. caule simplicissimo , foliolis pluribus lanceolatis integris. *Fl. suec.*

O. caule simplici , foliis senis , ellipticis , radice tuberosa. *Hall. helu. n. 417.*

O. foliis imis ovatis , superioribus lanceolatis , caule alato simplici. *Crantz. austr.*

O. radice tuberosa. *Riu. t. 59. R.*

Astragalus sylvaticus , foliis oblongis glabris. Bauh. pin. 351.

Astragalus sylvaticus. Thal. herc. 7. t. 1.

Lathyrus angustifolius , radice tuberosa, Læs. pruss. 138. t. 37.

Habitat in Europæ borealis pratis & sylvis. 24

6. *OROBUS angustifolius.* *O.* foliis bijugis ensiformibus , stipulis subulatis , caule simplici. *Gmel. sib. 4. t. 14. f. 5. delph.*

β. *O.* caule ramoso , foliis quaterno pinnatis linearibus , stipulis semisagittatis subulatis. *Ger. prou.*

DIADELPHIA. DECANDRIA. 211

O. orientalis, foliis angustissimis costæ brevissimæ. *Tournef. cor.* 29.

Araci f. Apios leguminosæ species tossano carolo. *Bauh. hist.* 2. p. 326.

Habitat in Sibiria, Gallia. 24

Habitus O. tuberosi, sed *Folia ensiformi-lanceolata & pinnata ex 2. f. 3. oppositionibus absque cirrho. Racemi floribus paucis luteis.*

7. OROBUS niger. O. caule ramoso, foliis sexjugis ovato-oblongis. *suec. ged. austr. carn. pom. pal. Kniph. cent.* 3. n. 69. *delph. lith. burg. lugd.*

O. caule ramoso, foliis ovatis duodenis. *Hall. helu. n.* 418.

O. sylvaticus, viciæ foliis. *Bauh. pin.* 352. *Riv. t.* 60.

O. pannonicus 2. *Clus. hist.* 2. p. 230.

Habitat in Europæ borealis montosis. 24

Foliola convoluta.

8. OROBUS pyrenaicus. O. caule ramoso, foliis bijugis lanceolatis nervosis, stipulis subspinosis. *carn.*

O. caule ramoso, foliis trijugis lanceolatis nervosis. *Sauv. monsp.*

O. pyrenaicus, foliis nervosis. *Tournef. inst.* 235.

O. pyrenaicus latifolius nervosus. *Pluk. phyt.* 210. f. 2.

Habitat in Pyrenæis. 24

9. OROBUS sylvaticus. O. caulibus decumbentibus hirsutis ramosis.

O. caule ramoso hirsuto decumbente, foliis subseptemjugis.

O. sylvaticus nostras. *Rai. angl.* 3. p. 324. *hist.* 1892. *Ad. paris.* 1706. p. 87. t. 90.

Habitat in Anglia, Gallia. 24

946. LATHYRUS.

Stylus planus, supra villosus, superne latior.

Cal. lacinia superiores 2 breviores.

* *Pedunculis unifloris.*

1. LATHYRUS Aphaca. L. pedunculis unifloris, cirrhis aphyllis, stipulis sagittato-cordatis. *carn. pal. gallob. lugd. burg. delph.*

L. aphyllus stipulis sagittatis latissimis. *Hall. helu. n.* 442.

L. cirrhys aphyllis. *Hort. upf. paris.*

Vicia lutea, foliis convolvuli minoris. *Bauh. pin.* 345.

212 DIADELPHIA DECANDRIA

Aphaca. *Lob. ic. 2. p. 70.*

Elatine 3. *Tabernaem. p. 116. R.*

Habitat in Italia, Gallia, Anglia, Helvetia, Germania, inter segetes. ☉

Cirrhus interdum aliquis gerit Foliola conjugata, 2. lanceolata, reliquis Lathyris simillima, ac hoc rarissime.

2. **LATHYRUS Nissolia.** L. pedunculis unifloris, foliis simplicibus, stipulis subulatis. *gallob. carn. Scholl. barb. n. 1013. pal. delph.*

L. pedunculis unifloris, foliis gramineis, stipulis minimis. *Hall. helu. n. 441.*

L. foliis solitariis lentiformibus. *Hort. cliff. paris.*

L. sylvestris minor. *Bauh. pin. 344.*

L. angustifolius erectus, folio singulari, sine capreolis, nissoli. *Magn. hort. 112. t. 112.*

Nissolia parva, flore purpureo. *Buxb. cent. 3. p. 84. t. 45. f. 1.*

Nissolia vulgaris. *Lindern. hort. 73. cum. ic.*

Habitat in Gallia, Anglia, Germania, &c. ☉

Cirrhii nulli. Caulis erectus, Stipulae minimae, subulatae.

3. **LATHYRUS amphicarpos.** L. pedunculis unifloris calyce longioribus, cirrhis diphyllis simplicissimis.

L. amphicarpos f. supra infraque terram filiquas gerens. *Morif. hist. 2. p. 51. f. 2. t. 23. f. 1.*

L. pedunculis unifloris, cirrhis diphyllis, radicibus etiam sub terra fructificantibus. *Hort. cliff. Sauv. monsp.*

Viciae similis supra & infra terram filiquas gerens. *Bauh. pin. 345.*

Arachidna f. Arachioides honorii belli. *Bauh. hist. 2. p. 323.*

Habitat in Syria. ☉

Conf. Vicia amphicarpa. Aracus homoion. Clus. exot. 87. t. 88.

4. **LATHYRUS cicera.** L. pedunculis unifloris, cirrhis diphyllis, leguminibus ovatis compressis: dorso canaliculatis. *Scop. carn. delph.*

L. foliis lanceolatis, stipulis latissimis, scapis brevibus, unifloris, filiquis brevissime alatis. *Hall. helu. n. 339.*

L. pedunculis unifloris, cirrhis diphyllis, foliis lanceolato-linearibus, internodiis membranaceis. *Sauv. monsp.*

L. sativus flore purpureo. *Bauh. pin. 344.*

L. flore rubro. *Bauh. hist. 2. p. 507. R.*

Aracus f. Cicera. *Dod. pempt. 523.*

Habitat in Hispania. ☉

DIADELPHIA. DECANDRIA. 213

Medius inter antecedentem & sequentem magnitudine. Penoli latere non membranacei. Corollæ rubra. Legumina dorso canaliculata, at non marginibus acutis, uti antecodens & insequens. Legumen latum, ovali-oblongum.

5. LATHYRUS *sativus*. L. pedunculis unifloris, cirrhis diphyllis tetraphyllisque, leguminibus ovatis compressis dorso bimarginatis. *parif. meth. carn. gallob. Kniph.*

sem. 5. n. 49.

L. foliis geminis, gramineis, stipulis folii latitudine, petiolis unifloris, siliquis alatis. *Hall. helv. n. 438.*

L. siliqua latiori. *Riv. 1. 47. Hall. R.*

L. anguloso semine. *Bauh. hist. 2. p. 306. Hall. R.*

L. annuus, flore caruleo, ochri siliqua. *Herm. lugd. 357.*

6. L. *sativus*, flore fructuque albo. *Bauh. pin. 343.*

Lathyrus f. *Gicercula*. *Dod. pempt. 522.*

Habitas in Hispania, Gallia, Helvetia. &c. ☉

7. LATHYRUS *setifolius*. L. pedunculis unifloris, cirrhis diphyllis, foliolis semiceo-linearibus. *Alben. nicæns. Ger. prov. delph.*

L. pedunculis unifloris, cirrhis diphyllis: foliolis linearibus, stipulis subulatis, caulibus nodis. *Sauv. mansp.*

L. foliis angustis, floribus singularibus coccineis. *Sauv. 2.*

L. *baldi*. *Riv. setr.*

L. folio tenuiore, floribus rubris. *Bauh. hist. 2. p. 308.*

L. sylvestris major, angustissimo folio. *Bauh. pin. 344.*

L. angustissimo f. capillaceo folio. *Bauh. prodr. 148.*

Habitas Montpelii, in Baldo, Italia.

Caules subfolitarii. Foliola subavenia. Stipulæ appendiculatæ. Petiolus trifidus, stipula triplo longior. Legumen ovatum, turgidum. Gerard.

8. LATHYRUS *angulatus*. L. pedunculis unifloris aristatis, cirrhis diphyllis, simplicissimis, foliolis linearibus. *Ger. prov. Sep. carn.*

L. foliis gramineis, petiolis unifloris setigeris, siliquis longis erectis. *Hall. helv. n. 440.*

L. pedunculis unifloris cirrhosis, foliolis linearibus, seminibus angularis. *Dalib. parif.*

Lathyrus pedunculis unifloris, cirrhis diphyllis, foliolis capillaceis. *Sauv. mansp.*

L. gramineus, floribus coccineis. *Rai. exot. 160. R.*

Lathyrus angustissimo folio, semine anguloso. *Tournef. infl.*

214 **DIADELPHIA DECANDRIA**

Lathyrus folio tenuiore, floribus rubris. Barb. cens. 3. p. 23. t. 42. f. 2.

Lathyrus angustifolius leptolobus, semine hexaedro, flore caeruleo. Moris. blas. 279.

Habitat in Gallia, Hispania, Oriente. &c. ☉

Caules multi. Foliola nervosa Stipulae lanceolata. Petiolus viz stipula longior. Legumen oblongum, compressum. Ger.

9. **LATHYRUS articulatus.** L. pedunculis subnatis, cirrhis polyphyllis : foliolis alternis.

L. hispanicus. Mill. dict. n. 3. & ic. t. 96.

L. angustissimo folio, americanus variegatus. Bauh. pin. 344.

Clymenum hispanicum, flore vario, filiqua articulata. Tournef. inst. 396.

10. *Lathyrus angustifolius humilior. Bauh. pin. 344.*

Habitat in Bactica, Monspeli. ☉

Folia sapinis tetraphylla ; inferiores petioli lanceolati, aphylli. Stipulae utrinque mucronatae. Florum carina pisl, lateribus undulatis, nec obliqua Lathyri. Vexillum purpureum alis & carina albis.

** * Pedunculis bifloris.*

10. **LATHYRUS odoratus.** L. pedunculis bifloris, cirrhis diphyllis, foliolis ovato - oblongis, leguminibus infutis.

a. Lathyrus ficulus.

Lathyrus ficulus. Rupp. ic. 210.

L. distoplatyphyllos hirsutus mollis, magno & perennino flore odoro. Comm. hors. 2. p. 219. t. 80.

β. Lathyrus zeylanicus.

L. zeylanicus, odorato flore amaro ex albo & rubro vario. Burm. zeyl. 138. Kniph. cens. 4. n. 37.

Habitat α. in Sicilia, β. in Zeylona. ☉

Foliola involuta.

11. **LATHYRUS annuus.** L. pedunculis bifloris, cirrhis diphyllis : foliolis ensiformibus, leguminibus glabris, stipulis bipartitis.

L. hispanicus. Riv. tetrap. 158.

L. hispanicus flore luteo. Herm. lugdb. 357.

L. luteus latifolius. Magn. monsp. 130.

Lathyri species lutea. Bauh. hist. 2. p. 304.

Habitat in Hispania, Monspeli. ☉

13. **LATHYRUS Clymenum.** L. pedunculis bifloris, cirrhis polyphyllis, stipulis dentatis.

DIADELPHIA. DECANDRIA. 215

L. vicioides, vexillo rubro, labialibus petalis rostrum ambicantibus caeruleis. *Marif. hist. 2. p. 46. Pluk. phyt. 114. f. 6.*

Clymeum hispanicum, flore vario, siliqua plana. *Toarnes. inst. 395.*

Habitas in Mauritania, Oriente. ☉

* * * *Pedunculis multifloris.*

14. LATHYRUS hirsutus. L. pedunculis subtrifloris, cirrhis diphyllis, foliolis lanceolatis, leguminibus hirsutis, seminibus scabris. *Pollich. pal. burg. delph.*

L. foliis geminis, ellipticis, floribus solitariis & geminis; siliquis hirsutis. *Hall. helu. n. 437.*

L. pedunculis bifloris, cirrhis diphyllis, foliolis linearilanceolatis, leguminibus hirsutis, seminibus scabris. *Spec. pl. paris. monsp.*

L. angustifolius, siliqua hirsuta. *Bauh. pin. 344.*

L. siliqua hirsuta. *Bauh. hist. 2. p. 305. Riu. tetr. t. 41.*

Habitat inter Angliæ, Galliæ, Germaniæ &c. segetes. ☉

Pedunculi sæpius triflori sunt. Flores purpurei.

15. LATHYRUS tuberosus. L. pedunculis multifloris; cirrhis diphyllis: foliolis ovalibus, internodiis nudis. *paris. pal. sil. Kniph. cent. 5. n. 51. burg. delph.*

L. scapis multifloris, foliis ovatis, capreolis trifidis. *Hall. helu. n. 435.*

L. foliis binatis ovatis, radice tuberosa; pedunculis multifloris. *Crantz. austr.*

L. arvensis. *Riu. tetr. t. 42. Hall. R.*

L. arvensis repens tuberosus. *Bauh. pin. 344.*

Arachidna Theophrasti. *Column. ecphr. 301. 304. R.*

Terræ glandes. *Dodon. cer. 168. Hall. R.*

Apios. *Fuchf. hist. 131.*

Habitat inter Belgii, Genevæ, Germaniæ, Tartariæ, segetes. 24

Foliola involuta. Corolla tota rubra.

16. LATHYRUS pratensis. L. pedunculis multifloris; cirrhis diphyllis simplicissimis: foliolis lanceolatis. *suec. paris. ged. carn. gallob. pal. sil. Flor. dan. 527. Kniph. cent. 11. n. 63. burg. lithu. delph. lugd.*

L. scapis multifloris, foliis lanceolatis, capreolis simplicibus. *Hall. helu. n. 436.*

L. foliis binatis utrinque acutis, pedunculis multifloris, floribus luteis. *Crantz. austr.*

L. pratensis. *Riu. tetr. t. 43.*

16 DIADELPHIA. DECANDRIA.

L. sylvestris luteus, foliis viciæ, *Bauh. pin.* 344.

L. luteus sylvestris dumetorum. *Bauh. hist.* 2. p. 304. t. 304.

Habitat in Europæ pratis. 24

Cirrhii quidem interdum trifidi sunt, at sæpius simplicissimi.
Foliola convoluta involuta.

17. LATHYRUS sylvestris. *L. pedunculis multifloris*,
cirrhis diphyllis : foliolis ensiformibus, internodiis mem-
branaceis. *suec. parif. ged. austr. gallob. pal. sil. Oed. fl.*
dan. t. 325. burg. litu. delph. lugd.

L. caule alato, foliis geminis ensiformibus. *Hall. helu. n.*
434.

L. sylvestris major. *Bauh. pin.* 433.

L. sylvestris. *Clus. hist.* 2. p. 129. *Dod. pempt.* 523. *Rin.*
tetr. t. 39.

Habitat in Europæ pratis montosis. 24

18. LATHYRUS latifolius. *L. pedunculis multifloris*,
cirrhis diphyllis : foliolis lanceolatis, internodiis mem-
branaceis. *suec. sil. Kniph. cent. 7. n. 40. Mill. illustr.*
burg. delph. litu. lugd.

L. caule alato, foliis ovato-lanceolatis, rigidis, scapis
multifloris. *Hall. helu. n.* 433.

L. pedunculis multifloris, caule alato, foliis & floribus
maximis. *Crantz. austr. R.*

L. narbonensis. *Rin. tetr. t. 40. R.*

L. latifolius. *Bauh. pin.* 344. *Garid. prou. 271. t. 271.*

L. major latifolius, flore purpureo, speciosior. *Bauh. hist.*
2. p. 302.

Habitat in Europæ sepibus. 24

Foliola involuta, elliptica.

19. LATHYRUS heterophyllus. *L. pedunculis multifloris*,
cirrhis diphyllis tetraphyllisque : foliolis lanceolatis,
internodiis membranaceis. *Fl. suec. ged.*

L. caule alato, foliis quaternis & binis, lanceolatis, scapis
multifloris. *Hall. helu. n.* 432.

L. pedunculis multifloris, cirrhis diphyllis : foliolis grami-
neis trinerviis, internodiis membranaceis. *Sauv. monsp.*

L. major narbonensis angustifolius. *Bauh. hist.* 2. p. 304.

Habitat ad radices montium Europæ. 24

20. LATHYRUS palustris. *L. pedunculis multifloris*,
cirrhis polyphyllis, stipulis lanceolatis. *Fl. suec. parif.*
ged. pal. sil. Fl. dan. 399. burg. lithu.

L. foliis senis, floribus racemosis. *Hall. helu. n.* 431.

L. pedunculis multifloris, cirrhis 3-4-phyllis, stipulis lan-
ceolatis; internodiis membranaceis. *Crantz. austr.*

DIADELPHIA: DECANDRIA: 217

L. foliis pinnatis, pedunculis multifloris. Fl. lepp.

L. palustris, flore orobi mucronatis verni. Rupp. bot. 210. t. 1.

L. peregrinus, foliis viciae, flore subcaeruleo pallidove purpurascente flore. Bauh. pin. 344.

Vicia lathyroides f. Lathyrus viciformis. Pluk. aln. 387. t. 71. f. 2.

Habitas in Europæ borealis pascuis paludosis. 2.

21. LATHYRUS pisiformis. *L. pedunculis multifloris, cirrhis polyphyllis, stipulis ovatis foliolo latioribus.*

Lian. dec. 39. t. 20. Leers herb. Kniph. cent. 11. n. 62.

Vicia pedunculis multifloris, cirrhis polyphyllis, stipulis ovatis, basi acutis. Sp. ed.

L. pedunculis multifloris, cirrhis heterophyllis, foliolis ovatis, stipulis maximis. Gmel. fib. 4. p. 7. n. 4. t. 1.

Vicia pedunculis multifloris, foliis ovatis, stipulis maximis. Hall. germ. 295.

Habitas in Sibiria, Germania. 2.

Planta ut Pisum crescens. Caules acute angulati. Petioli cirrhosi, sæpe hexaphylli: foliolis ovato-oblongis, acutis. Stipulae foliis latiores, ovatae, basi acuminatae, viz angulatae. Pedunculi longitudine foliorum, multiflori. Flores vexillo & alis albidis venis purpureis. Vexillum parvum, carina Pisi, nec obliqua. Stylus supra longitudinaliter barbatus. Foliola convoluta ut in Piso maritimo & Ochro, nec involuta ut in plerisque Lathyris; minus conduplicata ut in Piso sativo.

947. VICIA.

Stigma latere inferiore transverse barbatum.

** Pedunculis elongatis.*

1. VICIA pisiformis. *V. pedunculis multifloris, petiolis polyphyllis, foliolis ovatis: infimis sessilibus. Jacq. austr. t. 364. fil. Kniph. cent. 10. n. 99. burg.*

V. foliis ovatis maximis, filiquis racemosis pendulis. Hall. helv. n. 1428.

Pisum sylvestre perenne. Bauh. pin. 343.

Pisum sylvestre. Clus. hist. 2. p. 119.

Czeczca flore ochroleuco. Riv. tetr. t. 52.

Habitas in Pannoniæ Austriæ, Germaniæ sylvis. 2.

218 DIADELPHIA. DECANDRIA.

Foliola petiolorum infima sunt cauli approximata, sessilia, stipulas minimas occultantia. Flores ochroleuci.

2. *VICIA dumetorum*. V. pedunculis multifloris, foliolis reflexis ovatis mucronatis, stipulis subdentatis. *Ged. aust. tub. burg. delph.*

V. pedunculis multifloris, petiolis polyphyllis cirrhosis: foliolis alternis ovatis, stipulis dentatis. *Fl. suec. paris. monsp.*

V. foliis ovatis oblongis, stipulis ferratis, siliquis racemosis pendulis. *Hall. helu. n. 427.*

V. maxima dumetorum. *Bauh. pin. 345.*

V. sylvatica maxima piso similis. *Bauh. hist. 2. p. 315.*

Cracca sylvatica. Riu. tetr. 50.

Habitat in Gallia, Germania, &c. 24

Differt a præcedente Corollis non flavis, Foliolis longioribus, nec infimis cauli approximatis.

3. *VICIA sylvatica*. V. pedunculis multifloris, foliolis ovalibus, stipulis denticulatis. *Ged. carn. norv. herb. fil. Flor. dan. t. 277. delph. lithu.*

V. pedunculis multifloris, stipulis crenatis. *Fl. suec. monsp.*

V. foliis ovatis, stipulis argute dentatis, siliquis racemosis pendulis. *Hall. helu. n. 426. t. 12. R.*

V. multiflora maxima perennis, tetro odore, floribus albescentibus: lineis cæruleis notatis. *Pluk. alm. 387. t. 71. f. 1.*

Habitat in Sueciæ, Germaniæ, Galliæ, &c. sylvis. 24

4. *VICIA cassubica*. V. pedunculis subsexfloris, foliolis denis ovatis, stipulis integris. *Fl. suec. 3. Oed. dan. 98.*

V. pedunculis multifloris, caule fruticoso. *Hort. cliff.*

V. multiflora cassubica frutescens, siliqua lentis. *Pluk. alm. 387. t. 72. f. 2.*

Habitat in Germania? Dania.

5. *VICIA Cracca*. V. pedunculis multifloris, floribus imbricatis, foliolis lanceolatis pubescentibus, stipulis integris. *Pal. gallob. tub. fil. kniph. cent. 10. n. 98. burg. delph. lugd. lithu.*

V. pedunculis multifloris, stipulis utrinque acutis integris. *Fl. suec. paris. monsp.*

V. foliis linearibus sericeis, racemis, multifloris reflexis, stipulis integerrimis. *Hall. helu. n. 424.*

V. racemis folio brevioribus; floribus imbricatis, pendulis: foliolis lanceolatis, pubescentibus. *Scop. carn.*

V. foliis linearibus hirsutis, pedunculis multifloris, siliquis subhirsutis longis. *Crantz. austr.*

DIADELPHIA. DECAENDRIA. 219

V. sylvestris spicata. *Bauh. pin.* 345.

V. multiflora. *Bauh. pin.* 343.

Cracca. Riv. tetr. 49.

Habitat in Europæ pratis, agris. 2.

Foliola involuta. Variat floribus albis.

6. *VICIA onobrychioides*. *V. pedunculis, floribus, distantibus, foliolis linearibus, stipulis inferne denticulatis, delph.*

V. foliis ellipticis, floribus racemosis erectis, stipulis argute dentatis. Hall. helv. n. 425. *R.*

V. pedunculis multifloris, stipulis crenatis, foliolis linearibus. Sauv. monsp.

T. onobrychidis flore. Bauh. pin. 345. *prodr.* 149.

Habitat in Gallia. D. Sauvages. Helvetia. Haller.

Statura Viciæ Craccæ. Folia pinnata, cirrhosa: foliolis sæpius 14, linearibus, angustissimis: oppositionibus distantibus. Pedunculi axillares, foliis triplo longiores, striati. Floribus 10. s. 12, subsessilibus, remotissimis, Craccæ triplo majoribus, purpureis. Legumina latissima.

3. *VICIA biennis*. *V. pedunculis multifloris, petiolis sulcatis subdodecaphyllis: foliolis lanceolatis glabris. Mill. dict. n.* 5. *Gmel. sib.* 4. t. 2. *delph.*

*V. pedunculis multifloris, petiolis polyphyllis: foliolis lanceolatis glabris. Hort. upf. **

V. perennis multiflora, flore cœruleo ex albo mixto.

Magn. monsp. 300? sec. Sauv. monsp. 234.

Habitat in Sibiria. ♂

Foliola involuta. Planta biennis: maxima ultra 12 pedes alta.

10. *VICIA sativa*. *V. leguminibus sessilibus subbinatis erectis, foliis retusis, stipulis notatis. Ged. austr. gallob. herb. sil. Flor. dan. t.* 522. *Kniph. cent. I. n.* 96.

V. leguminibus erectis, petiolis polyphyllis: foliolis acumine emarginatis, stipulis denticulatis. Fl. succ. parisi. burg. delph. lithu. lugd.

V. sativa vulgaris, semine nigro. Bauh. pin. 344.

V. sativa alba. Bauh. pin. 344.

β. *Vicia (nigra) semine rotundo nigro. Bauh. pin.* 345.

V. foliis imis ovatis, superioribus linearibus, scapis brevissimis, bifloris. Hall. helv. n. 430. *pal.*

Vicia sativa. Scopoli carn.

V. vulgaris, acutiore folio, semine parvo nigro. Bauh. pin. 345.

220 **DIADELPHIA. DECANDRIA.**

V. folio angustiore, flore rubro. Dill. app. 47.

V. angustifolia. Riv. tetr. tab. 54.

Habitat inter Europæ segetes hodie. ☉

Stipulæ subtus macula quasi inusta notatæ.

Varietas β foliis angustioribus sublinearibus.

11. VICIA lathyroides. *V. leguminibus sessilibus solitariis erectis glabris, foliolis senis: inferioribus obcordatis. Ged. austr. pal. herb. fil. Fl. dan. t. 58. Kniph. cent. 3. n. 96.*

V. lathyroides, purpureo-cœruleis floribus. Herm. par. 242. t. 242.

V. montana minima, flosculo purpureo. Mænn. acad. 1.

Vicia minima. Riv. tetr. 169.

Habitat in Scotia, Lusatia, Norvegia, &c. 2c

12. VICIA lutea. *V. leguminibus sessilibus reflexis pilosis solitariis penta-spermis, corollæ vexillis glabris. pal. tugd. delph.*

V. foliis subsessilibus obverse ovatis apice emarginatis, leguminibus solitariis villosis. Dalib. paris.

V. leguminibus solitariis deorsum fixis hirsutis. Sauv. monsp.

V. sylvestris lutea, siliqua hirsuta. Bauh. pin. 345.

V. flore luteo pallido, siliquis hirsutis crassis brevibus pro-pendentibus. Moris. hist. 2. p. 62. f. 2. t. 21. f. 5.

Habitat in Gallia, Germania, Hispania, Italia, Oriente. ☉

Variat foliis lanceolatis & retusis, Leguminibus hirsutis & subglabris. Stipulæ alternatim integerrimæ & tri-dentatæ.

13. VICIA hybrida. *V. leguminibus sessilibus reflexis pilosis pentaspermis, corollæ vexillis villosis. Jacq. aust. t. 146.*

Habitat Monspeliū, Massilia. ☉

Affinis R. luteæ. Foliola obovata, retusa, parva, pilis adspersa. Stipulæ non ad caulem fuscae, sed totæ virides. Flores flavi, supra subgrisei, pilosi. Cætera V. luteæ.

14. VICIA peregrina. *V. leguminibus subsessilibus pendulis glabris tetraspermis, foliolis linearibus emarginatis. burg.*

V. peregrina angustissimis foliis, siliqua lata glabra. Pluk. alm. 386. t. 233. f. 6.

V. angustifolia purpureo-violacea, siliqua lata glabra. Magn. monsp. 276.

Habitat in Gallia. D. Sauvages. ☉

15. VICIA sepium. *V. leguminibus pedicellatis subqua-ternis erectis, foliolis ovatis integerrimis: exterioribus*

DIADELPHIA. DECANDRIA. 418

decreſcentibus. *Auſtr. gallob. ped. ſil. Flor. Jan. 1. 699.*
Kniph. cent. 5. n. 100. delph. Italia.

V. foliis ovatis oblongis, ſcapis brevibus quadrifloris.
Hall. helv. n. 428.

V. axillis ſubquadrifloris; floribus erectis, pedunculatis;
 foliis ovatis, ſtipulis dentatis. *Jacq. carn.*

V. leguminibus adſcendentibus, petiolis polyphyllis, folio-
 lis ovatis acutis integerrimis. *Fl. ſucc. pariſ. monſp.*

Cracca major. Tabern. p. 506.

Vicia ſepium perennis. Bauh. hiſt. 2. p. 313. R.

V. ſepium, folio rotundiore acuto. *Bauh. pin. 345.*

V. ſepium. *Riu. tetr. 56.*

Habitat in Europæ ſepibus. 24

Folia conduplicata. Racemi foliis breviores, ſubquadriflori. Floribus albis variis.

16. VICIA biſkynica. V. leguminibus pedunculatis foliariis erectis, foliolis quaternis ovali-lanceolatis, ſtipulis dentatis. *Jacq. hort. 1. 147.*

Lathyrus pedunculis uniſloris, cirrhis tetraphyllis, ſtipulis dentatis. *Roy. lugdb.*

Lathyrus pedunculis uniſloris, cirrhis tetraphyllis, ſtipulis ſetaceis, caule ancipiti. *Amoen. acad. 1.*

Cracca floribus albis, foliis circa radicem denticulatis.
Bauh. denſ. 3. p. 25. t. 45. f. 2.

Clymenum biſkynicum, ſiliqua ſingulari, flore minore.
Bark. lugdb. 2. p. 43.

Habitat in Italiæ, Bævariæ arvis. 24

Caules angulati. Cirrhi foliis bijugis, lato-lanceolatis. Stipule valde dentate. Flores axillares, ſolitarii, Calyce longitudine pedunculi; Corolla vexillo violaceo, alis albidis. Legumen compreſſum, ovato-oblongum, villoſum. Semina circiter 6. Viciam eſſe Stigma extus, barbatum & Stipulas ſubans poro glanduloſo evincunt.

17. VICIA narbonenſis. V. leguminibus ſubſeſſilibus ſubternatis erectis, foliolis ſenis ſubovatis, ſtipulis denticulatis. *Savv. monſp. Kniph. cent. 4. n. 98. Kærr. dell. 2. t. L. 1.*

V. petiolis inferioribus tetraphyllis: ſuperioribus polyphyllis, foliolis ovatis cirrhiferis. *Roy. lugdb.*

V. petiolis tetraphyllis cirrhosis. *Hort. cliff.*

V. narbonenſis. *Riu. tetr. 1. 40.*

Faba ſylveſtris, fructu rotundo atro. *Bauh. pin. 338.*

Bona ſylveſtris. *Dod. pempt. 516.*

121. DIADELPHIA. DECANDRIA.

Habitat in Gallia, Anglia, Sibiria. ☉

Folia conduplicata ut in *V. sepium*, nec involuta ut in plerisque vicis. Pedunculi uniflori in macris, triflori in fertilibus. Corollæ atropurpurea.

18. VICIA Faba. *V.* caule erecto, petiolis absque cirrhis.

Paris. Ludw. ed. 1. 159. Regn. bot.

Faba. *Bauh. pin. 338. Mill. dict. Blackw. t. 19.*

Bona f. Phaseolus. *Dod. pempt. 513.*

β Faba minor f. equina. *Bauh. pin. 338.*

Habitat non procul a mari Caspio in confinis Persia.

D. Lerche. ☉

Corollarum alæ in medio Macula atra holosericea. Foliola convoluta.

948. ERVUM.

Cal. 5. partitus, longitudine corollæ.

1. ERVUM *Lens.* *E.* pedunculis subbifloris, seminibus compressis convexis. *Carn. herb. Ludw. ed. t. 141. Kniph. cent. 9. n. 35.*

Cicer pedunculis bifloris, foliolis integerrimis, stipulis indivisis. *Sauv. monsp.*

Cicer pedunculis bifloris, seminibus compressis. *Dalib. Paris. Lens vulgaris. Bauh. pin. 346.*

Lens. Dod. pempt. 526. Hall. helu. n. 421. Rin. tetr. t. 35.

Habitat inter Gallia segetes, in pratis Carniolia. ☉

2. ERVUM *tetraspermum.* *Ervum* pedunculis subbifloris, seminibus globosis quaternis. *Fl. succ. paris. carn. pat. herb. fil. lith. delph. lugd.*

V. foliis linearibus, siliquis gemellis glabris. *Hall. helu. n. 923.*

V. (gemella) foliis linearibus, siliquis obtusis gemellis, seminibus globosis quaternis. *Crantz. austr.*

V. foliolis quadri-quinquejugis linearibus utrinque attenuatis: pedunculis filiformibus, subquinquefloris. *Neck. gallob.*

Cracca minor cum siliquis gemellis. *Rin. tetr. 167.*

Vicia segetum, singularibus siliquis glabris. *Bauh. pin. 345.*

Vicia minor segetum, cum siliquis paucis glabris. *Morist. hist. 2. p. 64. f. 2. t. 4. f. 16.*

Habitat inter Europæ segetes. ☉

Gaulis subtetragonus, anceps. Folia foliolis sæpius 10.

DIADELPHIA DECANDRIA. 209

plerumque alacris. Pedunculis forte capitis sessilibus, unifloris s. bifloris. Flos natus, saepe sanguineus, albus violaceus. Legumina ovato-oblonga, lora.

3. ERVUM hirsutum. E. pedunculis multifloris, foliis lobis globosis ovatis. Ager. pol. get. 1202. herb. M. M. den. 1. 639. burg. luth. delph. lugh.

Cicer pedunculis multifloris, seminibus globosis. Fl. salt. parif. monsp.

Vicia foliis lanceatis, stipulis racemosis, dispersis, hirsutis. Hall. helv. 2. 421.

Vicia racemes folio longioribus; floribus intricatis, secundis, pendulis; foliis lanceatis, hirsutis villatis. Scop. carn.

Ervum foliis integerrimis lanceatis apice lobulatis; pedunculis subsessilibus. Herb. polifol.

Vicia legumina cum stipulis planis hirsuta. Bauh. pin. 345.

Cracca minor. Tabern. ic. 905. Rer. rar. 1. 53.

Habitat in Europe apic, Oriente. ☉

4. ERVUM solonense. E. pedunculis subsessilibus nullis; petalis simplicibus, foliis ovatis. Amer. acad. 1. burg.

Vicia pedunculis uni-florisve, petalis diphyllis brevifloris circatis. Germ. flup.

Vicia praecox verna minima solonensis, semine hexedro. Loxf. blaf. 321. lig. 1. p. 63. Rei. lig. 901.

Vicia minima praecox perithensium. Turres. lig. 397.

Habitat in Anglia, Parisiis, Montpelii.

Affinis Lathyrus aspalatus?

5. ERVUM monanthos. E. pedunculis unifloris. Less. herb.

Vicia pedunculis unifloris, foliis integerrimis, stipulis alternis-dentatis. Horv. upf.

Lenis monanthos. Herm. legib. 360.

Habitat in Asia ruthenica; circa Herborn quass sponte. ☉

Foliola convoluta.

6. ERVUM Ervilia. E. germinibus undato-plicatis, foliis impari-pinnatis. Monsp. Leers. herb.

E. leguminibus pendulis. Horv. cliff.

Ervum. Cam. epis. 215. Hall. helv. 2. 420. Rer. rar. 1. 6.

Blakw. 1. 208. f. 3.

Orobis siliquis articulatis, flore majori. Bauh. pin. 346. /

Lenis minor. Camer. epis. p. 211.

Habitat in Gallia, Italia, Oriente. ☉

Folia cirrhis destituta. Pedunculi subsessilibus.

224 DIADELPHIA. DECANDRIA:

949. C I C E R.

Cal. 3. partitus, longitudine corollæ: laciniis 4. superioribus vexillo incumbentibus. *Legumen* rhombenum, turgidum, dispermum.

1. *CICER arietinum*. C. foliolis serratis. *Hall. helv. n. 399.*
Blackw. t. 557. Ludw. ed. t. 53. Kniph. cent. 7. n. 11.
Rosa. bot.

C. sativum. *Bauh. pin. 347. Camer. epis. p. 104.*

C. arietinum. *Dod. pempt. 325. Riv. t. 19.*

Habitas inter Hispaniam, Italiam, Orientis septem. ☉

Folia pinnata cum impari. Pedunculus uniflorus, suberectus.

951. C Y T I S U S.

Cal. 2. labiatus: $\frac{1}{2}$. *Legumen* basi attenuatum.

1. *CYTISUS Laburnum*. C. racemis simplicibus pendulis
 foliolis ovato-oblongis. *Hall. helv. n. 360. carn. Jacq.*
ausp. t. 306. delph. lugd.

Anagyris non ferida major alpina. Bauh. pin. 391.

Anagyris non ferida minor. Bauh. pin. 391.

Laburnum arbor trifolia anagyridi similis. Bauh. pin. t. 2.
p. 361. Nov. Act. A. N. C. vol. 1. t. 2. f. 2.

Der Bohnenbaum. Linn. Pfl. Syst. 4. p. 242.

Habitas in Helvetia, Sabaudia. ☿

2. *CYTISUS nigricans*. C. racemis simplicibus
 foliolis ovato-oblongis. *Savv. monsp. carn. Jacq.*
t. 387. tab. Kniph. cent. 10. p. 37. lila.

C. foliis ovatis nitidis, floribus spicatis, cernuis, calycibus & filiquis sericeis. *Hall. helv. n. 361.*

C. glaber nigricans. *Bauh. pin. 390. Tournes. inst. 648.*

Cytisus IV. Clus. hist. 1. p. 95.

Habitas in Austria, Pannonia, Bohemia, Italia, Germania, &c. ☿

Folia subtus pilis canescentia, supra glabra, impari mutua.

3. *CYTISUS sessilifolius*. D. racemis erectis, calycibus
 bractea triplici, foliis floralibus sessilibus. *delph.*

C. foliis saepius sessilibus, calycibus squamula triplici
Hort. cliff. monsp.

DIAPDELPHIA. DECANDRIA. 227

C. glabris foliis subrotundis , pediculis brevissimis. *Bauh. pin. 390.*

C. glaber , filiqua lata. *Bauh. hist. 1. p. 373.*

Habitat in Italia , Galloprovincia. H

Folia longius petiolata , at in ramulis sessilia. Legumina nigricantia.

5. CYTISUS patens. **C.** floribus pedunculatis subbinatis lateralibus nutantibus.

C. lusitanicus , medicæ folio , floribus in foliorum alis. *Tournef. inst. 648. Vandelli.*

Habitat in Lusitania. H

Rami patentissimi , virgati. Pedunculi gemini , uniflori. Flores penduli , lutei.

6. CYTISUS hirsutus. **C.** pedunculis simplicibus lateralibus , calycibus hirsutis trifidis obtusis ventricoso-oblongis. *Jacq. obs. 4. t. 96.*

C. calycibus hirsutis sessilibus , pedunculis simplicibus brevissimis. *Hort. cliff. Sa. monsp. Gm. fib. 4. d. 17. t. 6. f. 2.*

C. (*supinus*) floribus axillaribus , ramis decumbentibus , foliolis ovatis , subtus incanis. *Crantz. austr.*

C. hirsutus & prostratus. *Scop. carn.*

C. incanus , filiqua longiore. *Bauh. pin. 390.*

C. foliis subrufa lanugine hirsutis. *Bauh. pin. 390.*

Cytisus. Clus. hist. 1. p. 94.

Habitat in Hispania , Sibiria , Austria , Italia. H

Rami decumbentes , ut in sequenti. Folia subtus hirsuta. Calyces hirsutissimi.

In Cytis. Laburno , alpino , sessilifolio & hirsuto Stamina omnia connata cum cel. Crantzio observat ill. Medicus in obs. Soc. æcon. Lutr. 1774. p. 192. R.

7. CYTISUS supinus. **C.** floribus umbellatis terminalibus , rarius (ramis) erectis , foliolis ovatis. *fil. delph.*

C. capitatus. *Jacq. austr. t. 33. secund. Linn. ergo & Scop. carn. R.*

C. floribus umbellatis terminalibus , ramis decumbentibus , foliolis ovatis. *Spec. 2.*

C. (*hirsutus*) foliis ovatis , ramis decumbentibus , floribus capitato-umbellatis. *Crantz. austr.*

C. supinus , foliis infra & filiquis molli lanugine pubescentibus. *Bauh. pin. 390.*

Cytisus VII. species altera. Clus. hist. 1. p. 96.

Habitat in Austria , Sibiria , Italia , Sicilia , Galloprovincia. H

Linnaei spec. Plant. Europ. Pars II, R

226 DIADELPHIA. DECANDRIA.

Caulis petiolique pubescens, folia læviuscula. Calyx tubulosus, ventricosus, pubescens, bilabiatus.

8. CYTISUS *austriacus*. C. floribus umbellatis terminalibus, caulibus erectis, foliolis lanceolatis. *Jacq. austr. t. 21.*

C. floribus capitatis, foliolis ovato-oblongis, caule fruticoso. *Mill. dict. t. 107. f. 2. ed. germ. t. 117. f. 2.*

C. incanus, folio oblongo, austriacus. *Bauh. pin. 390.*

Cytisus V. *Clus. hist. 1. p. 195.*

Habitat in Sibiria, Austria, Italia. ☞

Similis C. *Jupino*, sed Caules erecti, ramosissimi, paniculati. Foliola lanceolata, nec ovata.

Stamina etiam omnia connata sunt. *Crantz. l. c. R.*

9. CYTISUS *argenteus*. C. floribus subsessilibus subbinatis, foliis tomentosis, caulibus decumbentibus, stipulis minutis. *prou. carn. delph.*

C. acaulis, floribus solitariis: foliolis sericeis lanceolatis. *Sauv. monsp.*

Lotus fruticosus incanus siliquosus. *Bauh. pin. 332.*

Lotus asperior fruticosa narbonensis incana. *Lob. ic. 2. p. 41.*

Trifolium argenteum, floribus luteis. *Bauh. hist. 2. p. 359.*

Habitat in G. Narbonensi, Carniolia. ☞

Planta incana, nitida. Rami versus radicem, fruticantes, superne herbacei. Bractæ 2. f. 3. calycis basi adnatæ. Calycis laciniae longiores, quam in congeneribus. Legumina hirsuta.

953. ROBINIA.

Cal. 4. fidus. Legumen gibbum, elongatum.

1. ROBINIA *Pseudacacia*. R. racemis pedicellis unifloris, foliis impari-pinnatis, stipulis spinosis. *Kniph. cens. 3. n. 76.*

R. aculeis geminatis. *Hort. cliff. Du ham. arb. 2. t. 42.*

Acacia americana, siliquis glabris. *Rai. hist. 1719. Seb. mus. 1. t. 15. f. 1.*

Acaciæ affinis virginiana spinosa, siliqua membranacea plana. *Pluk. alm. 9. p. 73. t. 4.*

Habitat in Virginia. ☞

Racemi penduli albi.

5. ROBINIA *Caragana*. R. pedunculis simplicibus, foliis abrupte pinnatis, petiolis inermibus. *Kniph. cens. 5. n. 76.*

Aspalathus arborescens, pinis foliorum crebrioribus oblongis. *Amm. ruth. 285.*

DIADELPHIA. DECAEDRIA. 227

Pseudo-Acacia foliorum parvis crebris. *Dalman. arb. 30*

Caragana fibrica. Roy. lugdb.

Habitas in Sibiria. H

Folia quinque vel sexiuga. Stipulae & petioli apice subsistentes, ut in sequentibus. Flores sex, ad gemmam praecursum anni, propriis pedunculis. Color florum luteus.

954. COLUTEA.

Cal. 5. fidus. Legumen inflatum, basi superiore dehiscens.

1. *COLUTEA arborescens. C. arborea, foliolis cuneatis, carn. Kniph. cent. 5. n. 24. Regn. bot. delph.*

C. foliis ovatis emarginatis, floribus racemosis. Hall. bot. n. 400. R.

C. vesicaria. Bauh. pin. 396.

Colutea. Dod. pempt. 784. Riv. t. 25.

β. *C. africana, sennae foliis, flore sanguinea. Comm. rar. II. t. II.*

γ. *C. foliolis ovatis integerrimis, caule fruticoso. Mill. dict. t. 100.*

Habitas in Anglia, G. Napolitani, Italia, &c. egypte ad Vesuvium. H

Arbor foliolis fere obovatis, floribus luteis.

2. *COLUTEA frutescens. C. fruticosa, baculis ovatis, longis Kniph. cent. 5. n. 25.*

C. aethiopica, flore purpureo. Breyn. cent. 79. t. 25. Mill. ic. 99.

Habitas in Aethiopia, Sibiria. ♂ H

Frutex incanus, foliolis tomentosis supra glabris, floribus rubris: vexillis carina brevioribus; alio minutissimo, ut fere nullis. Legumina inflata, dorsalto sutura hians.

955. GLICYRRHIZA.

Cal. bilabiatus: 5. Legumen ovatum compressum.

1. *GLICYRRHIZA echinata. G. leguminibus echinatis, foliis stipulatis, foliolo impari sessili. Jacq. hort. t. 95. Kniph. cent. 5. n. 37.*

G. capite echinato. Bauh. pin. 352.

228 DIADELPHIA DECANDRIA.

Dulcis radix. *Cam. epit.* 423.

Habitat in Gargano Apuliæ, in deserto Nagico Tartariæ. 24

Petiolus communis marginatus margine distincto mediante nervo. Foliola lanceolato-ovata: impari sessili. Stipulæ ensiformes. Spica subrotundo-capitata.

Stipulæ parvæ, oblongæ, non ensiformes. *Gmel. it. 1. p. 155. R.*

2. GLYCYRRHIZA glabra. G. leguminibus glabris, stipulis nullis, foliolo impari petiolato. *monsp. Ludw. ed. 1. 64. Kniph. cent. 4. n. 29. Re. n. bot. delph.*

G. lævis. *Pall it. 1. app. n. 120. * R.*

G. siliquosa & germanica. *Bauh. pin. 352.*

G. vulgaris. *Dod. pempt. 341.*

Habitat in Franconia, Gallia, Hispania, Italia. 24

Petiolus communis ultra ultimum foliorum par elongatus. Foliola ovata: impari petiolato. Stipulæ nullæ. Spicæ oblongæ floribus distantibus.

956. CORONILLA.

Cal. 2. labiatus: 3. dentibus superioribus connatis. Vexillum vix alis longius. Legumen isthmis interceptum.

1. CORONILLA Emerus. C. fruticosa, pedunculis subtrifloris, corollarum unguibus calyce triplo longioribus caule angulato. *Fl. succ. carn. austr. Kniph. cent. 3. n. 33. lugd. delph.*

C. frutescens, foliis retusis, septenis, siliquis longissimis, obiter articulatis *Hall. helv. n. 389.*

C. major. *Riv. t. 97. R.*

Emerus major. *Mill. dict. n. 1. & ic. t. 132. f. 1.*

Colutea siliquosa s. scorpioides major. *Bauh. pin. 397.*

Colutea scorpioides. *Cam. epit. 541.*

β. Colutea siliquosa minor. *Bauh. pin. 397.*

Emerus minor. *Mill. dict. n. 2. & ic. t. 132. f. 2.*

Habitat Genevæ, Monspëlii, Salerni, Viennæ. 7

2. CORONILLA juncea. C. fruticosa, foliis quinatis ternatisque lineari-lanceolatis subcarnosis obtusis.

Dorychnium luteum hispanicum carnosius. *Barr. ic. 133.*

Colutea caule genistæ fungoso. *Bauh. hist. 1. p. 383.*

Polygala major massiliotica. *Bauh. pin. 349.*

Habitat Massiliæ, Monspëlii. 7

Caulis Spartii. Folia parva, subcoriacea.

3. **CORONILLA** *valentiana* C. fruticosa, foliis oblongis, stipulis suborbiculatis. *Caup. monsp. p. 10. t. 1. stip.*

Coronilla fruticosa, leguminibus crassioribus. *Caup. monsp. p. 10. t. 1. stip.*

multifloris, caule fruticoso. Fl. 1. t. 1. p. 10.

C. leguminibus crassioribus, stipulis suborbiculatis, caule fruticoso. Fl. 1. t. 1. p. 10.

C. leguminibus crassioribus, stipulis suborbiculatis, caule fruticoso. Fl. 1. t. 1. p. 10.

Polygala altera. Bauh. pin. 349. Riv. petr. 207.

Polygala valentiana. Clus. hist. 1. p. 98.

Coronilla s. Colutea monsp. Lab. ic. 2. p. 72. 2.

Habitat in Hispania, Italia. H

Odor florum graveolens, radu magis. Umbellæ vexilla continent in capitulum florum, quæ inter radu cingunt

fulvæ Carinæ cum alis. Folia prout vere colore Ruta: Foliola mucrone reflexo, unde retusa apparent. Florens stipulas dejicit.

4. **CORONILLA** *glauca*. C. fruticosa, foliis septenis oblongis, stipulis lanceolatis. *Ar. can. acad. 4.*

C. fruticosa, leguminibus crassioribus. Mill. ic. 1. 289. f. 1.

C. maritima, glauco folio. Tournef. inst. 650.

Colutea scorpioides maritima, glauco folio. Bauh. pin. 397. prodr. 157.

Colutea scorpioides altera. Clus. pann. ic. 46. Gouan. monsp. 378. R.

Habitat in Gallia Narbonensi. H

Flores nōdu inodori, die suavissime fragantes. Caulis erectus, ramis rubris. Foliola 5. f. 7, cuneiformia, obtusa cum mucrone. Stipulæ lanceolatae, patulae.

5. **CORONILLA** *coronata* C. fruticosa, foliolis novenis obovatis: internis cauli approximatis, stipula oppositifolia bipartita. *Jacq. austr. 1. 95. delph.*

C. monsp. Riv. petr. 207. Scop. carn. 1. 44.

C. caule erecto, foliis undenis ovatis, floribus umbellatis, siliquis articulatis pendulis. Hall. helv. n. 388.

Colutea scorpioides minor coronata. Bauh. pin. 397.

Colutea scorpioides II. Clus. hist. 1. p. 98.

Habitat in Europa australi. H

Foliola obovata, distantia: infimo petiolorum pari articulo cauli approximato, ut in Vicia pisiiformi, tamen stipulae non sunt hæc foliola.

Caulis fruticosus, sesquipedalis, erectus.

230 **DIADELPHIA. DECANDRIA.**

6. **CORONILLA minima.** C. suffruticosa procumbens; foliolis novenis ovatis, stipula oppositifolia emarginata, leguminibus, angulatis nodosis. *Jacq. austr. t. 271. burg. delph.*

C. caulibus lignosis procumbentibus, foliis ovatis, floribus umbellatis, siliquis alatis. *Hall. helu. n. 390.*

C. procumbens, foliolis novenis lanceolatis, stipulis oppositifoliis emarginatis, leguminibus angulatis nodosis. *Amœn. acad. 4. **

Ferrum equinum, siliquis in summitate. *Bauh. pin. 349.*

Polygalum Cörtusi. *Bauh. hist. 2. p. 351.*

Lotus enneaphyllos. *Dalech. hist. 510.*

Habitat in Gallia australi, Helvetia, Italia, Hispania. 24

Caulis suffruticosus, prostratus, teres, statura Loti corniculati. Foliola infima petiolorum cauli approximata (uti C. coronatæ s. variæ) Stipularum instar, patentia, in plano expansa, reliquis non minora. Foliola ovata, acutiuscula.

Obs. Cur varietas C. valentinæ dicta fuit Tournefortio, Gerardo? anne C. coronatam pro C. valentina assumere?

8. **CORONILLA Securidaca.** C. herbacea, leguminibus falcato-gladiatis, foliolis plurimis. *Sauv. monsp.*

Securidaca lutea. *Mill. dict.*

Securidaca lutea major. *Bauh. pin. 398.*

Hedysarum primum. *Dod. pempt. 546.*

Habitat inter Hispaniæ segetes. ☉

Dormis foliolis supra petiolum conniventibus, versusque petioli basem flexis.

9. **CORONILLA varia.** C. herbacea, leguminibus erectis teretibus torosis numerosis, foliolis plurimis glabris. *Paris. monsp. ged. austr. t. 432. carn. pal. fil. Kniph. cent. 7. n. 16. burg. delph. lithu. lugd.*

C. caule brachiato angulato, foliis vigenis, aristatis, floribus umbellatis. *Hall. helu. n. 387.*

C. flore vario. *Riu. tetr. t. 94. R.*

Hedysarum purpureum. *Tabern.*

Securidaca dumetorum major, flore vario, siliquis articulat. *Bauh. pin. 349.*

Securidaca 2. altera species. *Clus. hist. 2. p. 237.*

Habitat in Lusatia, Bohemia, Dania, Gallia, Germania, &c. ☉

Foliola in petiolo communi infima cauli, uti stipulæ, approximata sæpius sunt, quemadmodum in C. coronata,

DIADELPHIA. DECANDRIA. 118
est minima. Caules prostrati. Pedunculi longitudine foliorum.

957. ORNITHOPUS.

Legumen articulatum, teres, arcuatum.

1. ORNITHOPUS perpusillus. O. foliis pinnatis, leguminibus incurvatis. *Monsp. ged. gallob. pal. Rich fl. franc. Flor. dan. 730 lugd. burg. delph.*

O. foliis pinnatis, articulis leguminum subrotundis latis, *Hort. cliff. paris. gelr.*

Ornithopodium caule prostrato, foliis pinnatis, siliculis umbellatis, articulatis. *Hall. helu. n. 393. R.*

Ornithopodium minus. *Bauh. pin. 350.*

Ornithopodium perpusillum. *Dalech. hist. 487.*

β Ornithopodium majus. *Bauh. pin. 350.*

γ Ornithopodium radice tuberculis nodosa. *Bauh. pin. 250. Kniph. cent. 7. n. 66.*

Ornithopus (nodosus) foliis pinnatis, leguminibus confertis pedunculatis. *Mill. dict. n. 2. R.*

Habitat in Angliæ, Belgii, Galliæ, Hispaniæ, &c. arenosis. ☉

Planta glabra, procumbens. Folia petiolata. Pedunculi longitudine foliorum. Flores 3. s. 4. lutei, vexillo rubro reticulato.

Planta tota hirsuta. Legumina incurvata. *R.*

2. ORNITHOPUS compressus. O foliis pinnatis, leguminibus recurvatis compressis rugosis, bractea pinnata.

O. foliis pinnatis, leguminibus compressis rugosis. *Spreo. 2. monsp.*

Ornithopodio affinis hirsuta scorpioides. *Bauh. pin. 350.*

Scorpioides leguminosa. *Dalech. hist. 493.*

Habitat in Italia, Sicilia. ☉

Planta tomentosa. Foliola sessilia. Pedunculi folio breviores: Bractea pinnata. Flores sæpius 2, flavi. Legumina recurva.

3. ORNITHOPUS scorpioides. O. foliis ternatis subsessilibus: impari maximo. *Scop. carn. Kniph. cent. 10. n. 67. delph.*

Telephium dioscoridis s. Scorpioides. *Bauh. pin. 287.*

Scorpioides portulacæ folio. *Bauh. pin. 287.*

Scorpioides marthioli. *Dod. pempt. 71. Riv. tetr. 210.*

Habitat in G. Narbonensi, Hispania, Italia inter segetes. ☉

232 DIADELPHIA. DECANDRIA.

Folia ternata s. potius simplicia, petiolata, stipulæ subrotundis. Planta erecta. Pedunculi aphylli, subquadristlori.

958. HIPPOCREPIS.

Legumen compressum, altera futura pluries emarginatum, curvum.

1. **HIPPOCREPIS unifiliquosa.** H. leguminibus sessilibus solitariis erectis. *Monsp.*

Ferrum equinum siliquis solitariis lunatim excisis. *Hall. helu. n. 392.*

Ferrum equinum siliqua singulari. *Bauh. pin. 349. Garid. t. 114.*

Ferrum equinum vulgare. *Col. ecphr. I. p. 302, t. 300. Riu. tetr. t. 98.*

Habitat in Italia, Helvetia. ☉

2. **HIPPOCREPIS multifiliquosa.** H. leguminibus pedunculari confertis circularibus: margine altero lobatis. *Monsp. lugd.*

Ferrum equinum siliqua multiplici. *Bauh. pin. 346.*

Ferrum equinum alterum polyceraton, *Col. ecphr. I. t. 300.*

Habitat in G. Narbonensis, Hispaniæ, Italiæ cretaceis.

3. **HIPPOCREPIS comosa.** H. leguminibus pedunculatis confertis arcuatis: margine exteriori repandis. *Parif. austr. carn. gallob. pal. delph. burg.*

H. leguminibus pedunculatis confertis: margine exteriori lobatis. *Hort. cliff.*

Ferrum equinum siliquis umbellatis, undulatis. *Hall. helu. n. 391.*

Ferrum equinum germanicum, siliquis in summitate. *Bauh. pin. 346.*

Ferrum equinum comosum, s. capitatum. *Col. ecphr. I. p. 302. t. 301. Riu. tetr. 97.*

Habitat in Germania, Italia, Gallia, Angliæ cretaceis. 2.

Caules plures, adscendentes, spithamei, laeves. Stipulæ quasi decurrentes. Foliola 15. ovali-linearia, patentia, consistentia. Pedunculi angulato-striati, foliis longiores, e summis alis 2. s. 3. Umbella orbicularis, 8-flora: Vexillum subforficatum, subtus striatum, lateribus non reflexum, præter basim, quæ retusa, unde quasi porus utrinque.

959. SCORPIURUS.

Legumen isthmis interceptum, revolutum teres.

1. SCORPIURUS *vermiculata*. S. pedunculis unifloris, leguminibus tectis undique squamis obtusis. *Carn. riu. tetr.* 210. *delph.*

Scorpioides filiqua crassa boelii. *Tournef. inst.* 402.

Scorpioides bupleuri folio, corniculis crassioribus & magis spongiosis instar litui contortis & in se convolutis.

Morif. hist. 2. p. 127. f. 2. t. II. f. III.

Habitat in Europa australi. ☉

2. SCORPIURUS *muricata*. S. pedunculis bifloris, leguminibus extrorsum obtuse aculeatis. *Carn. Kniph. cent.* 8. n. 82.

Scorpioides bupleuri folio corniculis asperis & rugosis f. rigidis, striatis f. sulcatis litui instar contortis & in se convolutis. *Morif. hist.* 2. p. 127. f. 2. t. II. f. IV.

Habitat in Europa australi. ☉

3. SCORPIURUS *fulcata*. S. pedunculis subtrifloris, leguminibus extrorsum spinis distinctis acutis.

Scorpioides bupleuri folio. *Bauh. pin.* 287.

Scorpioides prius. *Dod. pempt.* 71.

Habitat in Europa australi. ☉

4. SCORPIURUS *subvillosa*. S. pedunculis subquadrifloris, leguminibus extrorsum spinis confertis acutis. *Kniph. cent.* 9. n. 96.

Scorpioides bupleuri folio, corniculis asperis magis in se contortis & convolutis. *Morif. hist.* 2. p. 127. f. 2. t. II. f. II.

Habitat in Europa australi. ☉

Species hasce omnes olim ex una specie ortas esse dubium non est, nec sufficit locus harum generationi, qui tum mutatus easdem redderet; quæ itaque mixtura harum produxerit constantes plantas? Qui has omnes aut conjungat aut distinguat, videtur argumentis inniti.

961. HEDYSARUM.

Cor. carina tranverse obtusa. *Legumen* articulis
1. spermis.

40. HEDYSARUM *alpinum*. H. foliis pinnatis, legumi-

234. DIADELPHIA. DECANDRIA.

nibus articulatis glabris pendulis , caule erecto. *Gmel. fib. 4. p. 26. n. 35. t. 10. delph.*

H. saxatile , filiqua lævi , floribus purpureis , inodorum. *Amm. ruth.*

c. Onobrychis semine clypeato lævi. Bauh. pin. 350. prodr. 149.

Habitat α . in Sibiria β in Helvetia. $2\frac{1}{2}$

Caulis *rectus* ; stipulæ *breves*. Foliola 17 , ovato-lanceolata. Planta β differt caule humiliore , floribus majoribus paucioribus , calyce pubescente , at attributa specifica tot communia , ut vix sufficienter distinguatur.

41. HEDYSARUM obscurum. *H. foliis pinnatis , stipulis vaginalibus , caule erecto flexuoso , floribus racemosis pendulis. Gmel. fib. 4. p. 29. t. 12. Jacq. austr. t. 168.*

H. caule erecto , ramoso , foliis ovatis , filiquis pendulis levissimis. Hall. helu. ed. 2. n. 395. t. 12.

H. (controversum) foliis pinnatis ; stipulis vaginalibus ; floribus racemosis & leguminibus articulatis glabris , pendulis. Crantz. austr. p. 425. t. 2. f. 3. Legumen Ex Jacq. R.

Astragalus caulescens erectus , stipulis vaginalibus. Sp. pl.

Astragalus caule erecto-ramoso , spica purpurea nitente terminato. Hall. helu. ed. 1. 567. t. 14.

Habitat in Sibiria , Helvetia. $2\frac{1}{2}$

Caulis *flexuosus*. Stipulæ *amplexicaules* , pollicares. Foliola *exac̄te elliptica* , mucronata setula. Fructus mihi *ignotus*.

Ill. Hallerus *eandem plantam* , cum *H. alpino* , suam facit , at meum specimen ex Helvetia optime representatur figura Halleriana ; & est certe species distinctissima ab *H. alpino meo sibirico*.

42. HEDYSARUM coronarium. *H. foliis pinnatis , leguminibus articulatis aculeatis nudis rectis , caule diffuso. Kniph. cen. 3. n. 45.*

Onobrychis , semine clypeato aspero , major. *Bauh. pin. 350.*

Onobrychis altera. Dod. pempt. 549.

Habitat in Italiæ pratis. $2\frac{1}{2}$

44. HEDYSARUM humile. *H. foliis pinnatis , leguminibus articulatis asperis , corollæ alis obsoletis , spicis hirsutis , caulibus depressis.*

H. clypeatum minus , flore purpureo. Rai. hist. 929.

Onobrychis semine clypeato aspero , minor. Bauh. pin. 350. prodr. 149.

DIADELPHIA. DECANDRIA. 439

Polygalo gesneri affine caput gallinaceum. Bauh. hist. 4. p. 336.

Habitat in G. Narbonensi, Hispania. 2.

Folia pinnata: foliolis obovato-oblongis, subinae v. llysto. Caules semipedales folio ramoque saepius unico. Spica erecta pubescens. Corollae similes H. coronario, sed minores alarum solis rudimentis. Legumina saepius biarticulata.

Radix perennis. Vexillum longitudine carinae. Alae vix ullae. Calyx hirsutus, corolla dimidio brevior, in seorsum terminatus.

Gerard vult varietatem H. pumili, sed perperam.

Alae non obsoletae, sed vexillo tantum triplo minores. Gouan. l. c. R.

45. HEDYSARUM spinosissimum. H. foliis pinnatis, leguminibus articulatis aculeatis tomentosis, caule diffuso. Hort. upf. *

H. hispanicum supinum annuum angustifolium, floribus parvis ex albo purpurascens. Bærh. lugdb.

Onobrychis minor, foliolis cordatis, siliquis magnis asperis compressis. Pluk. phys. 30. f. 2.

Habitat in Hispania. ☉

Legumen saepius articulis 2, orbiculatis, aequaliter aculeatis.

47. HEDYSARUM pumilum. H. foliis pinnatis, caule suffruticoso, corollarum alis vexillo vexilloque carina brevioribus, leguminibus monospermis.

Habitat in Hispania. ☿

Fruticulus palmaris. Folia pinnata, longitudine caulis: pinnis ovali-oblongis. Spica glabra. Florum vexillum carina dimidio brevius. Carina maxima, latissima obtusissima. Calyx laevis densibus subulatis, brevissimis.

48. HEDYSARUM Onobrychis. H. foliis pinnatis, leguminibus monospermis aculeatis, corollarum alis calycem aequantibus, caule elongato. Jacq. austr. 1. 352. pul. tub. lugd. delph.

H. foliis pinnatis, leguminibus subrotundis aculeatis. Hort. upf. monsp.

Onobrychis caule erecto, ramoso, floribus spicatis. Hall. helu. n. 396.

Onobrychis. Riu. t. 2. viciaefolia. Scop. carn.

Onobrychis folio viciae, fructu echinato, major. Bauh. pin. 350.

Caput gallinaceum belgarum. Lob. ic. 2. p. 81.

236 DIADELPHIA. DECANDRIA.

β. *Onobrychis incana*, foliis longioribus. *C. B.*

γ. *Onobrychis foliis viciæ*, fructu echinato minimo. *Bauh. pin. 350. ex Gouan. R.*

Habitat in Sibiria, Gallia, Angliæ, Bohemia aprica, cretaceis. 24

49. *HEDYSARUM saxatile*. *H.* foliis pinnatis, leguminibus monospermis sulcatis muticis, corollarum alis brevissimis, scapis subradicatis. *delph.*

H. foliis pinnatis, foliolis linearibus, leguminibus monospermis sulcatis lævibus, petalis inæqualibus. *Ger. prou.*

Onobrychis saxatilis, foliis viciæ angustioribus & longioribus aquisextiensis. *Tournef. inst. 390. Garid. aix. Allion. nicæns. 124.*

Habitat in Galloprovincia & in agro Nicæensi, Sibiria. 24

Caules sub terra lignosi. Caulis brevissimus, ut fere nullus.

Scapi itaque fere radicati. Spica onobrychidis, sed albida.

Corollæ alæ calyce vix longiores, s. vexillo carinaque breviores. Folia linearia.

50. *HEDYSARUM Caput galli*. *H.* foliis pinnatis, leguminibus monospermis cristæ dentibus subulatis, alis brevissimis, caule diffuso. *Kniph. cent. 9. n. 44.*

H. foliis pinnatis, leguminibus aculeatis, corollarum alis minimis calyce brevioribus. *Ger. prou.*

Onobrychis fructu echinato, minor. Bauh. pin. 350. prodr. 149. Burf. XIX. 135.

Habitat in Galloprovincia. 24

Legumen carina spinis æqualibus, acutis. Radix perennis.

Synonymon Bauhini ad H. Onobrychidis var. γ. refero. Gouan. R.

262. INDIGOFERA.

Cal. patens. Corollæ carina utrinque calcari subulato patulo. Legumen lineare.

II. *INDIGOFERA Anil*. *I.* foliis pinnatis lanceolatis, racemis brevibus, caule suffruticoso.

I. foliis pinnatis: foliolis ellipticis acutiusculis, leguminibus declinatis recurvis, caule erecto. *Mant. 2. p. 272.*

Habitat in India H

Habitus & facies. I. tinctoriæ. Caulis suffruticosus. Folia pinnata, palmaria: foliola oblonga, obtusiuscula (recentioribus).

DIADELPHIA. DECANDRIA.

flora acutiuscula) , *supra nuda* , *subius incana* , omni
æqualia. Racemi laterales , subspicati , foliis brevibus. l. 4.
gumina declinata , *recurvata* , *teretiuscula* , *poly sperma*
sutura utraque gibbiflora (*dehiscencia sutura superne recedente*) , *mucronata*.

12. INDIGOFERA *indica*. I. foliis pinnatis *obovatis* ;
racemis brevibus , *caule suffruticoso*.

I. leguminibus *arcuatis incanis* , *racemis foliis brevibus*.
Mat. med. 174. *Mill. dict. n.* 1. & *ic.* t. 34. *Regn. bot.*
Blackw. t. 596.

Indigofera foliis nudis. Hort. cliff. 487.

Isatis indica , *foliis rosmarini* , *glasse affinis. Bauh. pin.* 113.

Anil. f. Nil inodorum color. Bauh. hist. 2. n. 947.

Colutea indica humilis , *exqua Indigo* , *folio viridi. Burma*
zeyl. 69: R.

Anil f. Indigo americana. March. paris. 1718. R.

Habitat in India. H

Differt a præcedente. Foliolis obovatis , *obtusis* , *utrinque*
nudis. Leguminibus teretibus , *rectis* , *etiam sutura gibbo-*
flora , *subtorulosa. Racemis laxis* , *minusis.*

963. G A L E G A.

Cal. dentibus subulatis , *subæqualibus. Legumen*
striis obliquis , *feminibus interjectis.*

I. GALEGA *officinalis*. Galega leguminibus strictis erectis ,
foliolis lanceolatis striatis nudis. Regn. bot.

G. foliis ellipticis , spinis & siliquis acutis, *Hall. helv.*
n. 394.

Galega. Riv. t. 72. *Ludw. ed.* t. 161.

G. vulgaris. *Bauh. pin.* 352. *Moris, hist.* 2. p. 91. f. 2. t. 7.
f. 9. *Blackw. t.* 92.

Habitat in Hispania , *Italia* , *Africa ad versuras.* 24

964. P H A C A.

Legumen semibiloculare.

I. PHACA *bætica*. P. caulescens erecta pilosa , leguminibus
tereti - cymbiformibus.

Phaca leguminibus rectis. Roy. lugdb.

Astragalus bæticus lanuginosus , *radice amplissima. Bauh.*
pin. 351.

DIADELPHIA. DECANDRIA:

Psoralea baticus. Clus. hist. 2. p. 234.

in Hispania, Lusitania.

PHACA alpina. P. caulescens erecta glabra, leguminibus oblongis inflatis subpilosis. *delph.*

P. leguminibus pendulis semiovatis. *Gmel. sib. 5. p. 35. t. 14. bona.*

P. frigida. *Fl. suec.*

Astragalus caule erecto ramosissimo, foliis ellipticis, hirsutis, siliquis vesicariis. *Hall. helv. n. 401. R.*

Astragaloides elatior erecta, viciae foliis, floribus luteis, siliquis pendulis. *Amm. ruth.*

Hab. in Alpibus Sibiriae, Lapponiae, Helvetiae umbrosis. 2.

Sibirica Gmel. & Lapponica nostra omnino eadem. Syst.

Veg. 365. Ergo etiam Astragalus Hall. l. c. sed cur glabram dicit plantam b. Linnaeus, quam hirsutam facit Hall. R.

3. PHACA australis. P. caule ramoso prostrato, foliolis lanceolatis, florum alis semibifidis. *Hall. helv. n. 403.*

P. caulescens diffusa, pedunculis longissimis, leguminibus semiovatis. *Ger. prou. **

Astragaloides alpina supina glabra, foliis acutioribus. *Tilepif. 19. t. 14. f. 1.*

Habitat in alpinis Helvetiae, Italiae, Galloprovinciae. 2.

Folia sexjuga circiter, cinerea, pilosa. Flores albido-flavicantes. Alae semibifidae, quod singulare. Legumina glabra. Carina corollae violacea, nec ad apicem 2 maculis violaceis, ut in Ph. alpina, a qua parum differt.

Junior hirsuta, adultior glabra evadit, quasi distincta planta.

963. ASTRAGALUS.

Legumen biloculare, gibbum.

** Caulibus foliosis erectis; nec prostratis.*

1. ASTRAGALUS alopecuroides. A. caulescens, spicis cylindricis subsessilibus, calycibus leguminibusque lanatis. *delph.*

A capitulis oblongis sessilibus, calycibus leguminibus lanatis. *Hort. cliff.*

A. alpinus procerior alopecuroides. *Tournef. inf. 416. Mill. dict. t. 58.*

Habitat in Sibiria, Hispania.

DIADELPHIA. DECANDRIA.

239

stora acutiuscula) , *supra nuda* , *subtus incana* , om.
æqualia. Racemi laterales , subspicati , foliis breviores. Legu-
 gmina declinata , recurvata , teretiuscula , polysperma
sutura utraque gibbiflora (*dehiscencia futura superiore rece-*
dente) , mucronata.

12. INDIGOFFERA *indica*. I. foliis pinnatis obovatis &
 racemis brevibus , caule suffruticoso.

I. leguminibus arcuatis incanis , racemis folio brevioribus.
Mat. med. 174. *Mill. dict. n.* 1. & *ic.* t. 34. *Regn. bot.*
Blackw. t. 596.

Indigofera foliis nudis. *Hort. cliff.* 487.

Isatis indica , foliis rosmarini , glasti affinis. *Bauh. pin.* 113.

Anil. f. Nil inodorum color. *Bauh. hist.* 2. n. 945.

Colutea indica humilis , exqua Indigo , folio viridi. *Burmæ*
zeyl. 69: R.

Anil f. Indigo americana. *March. paris.* 1718. R.

Habitat in India. H

Differt a præcedente. Foliolis obovatis , obtusis , utrinque
 nudis. Leguminibus teretibus , rectis , etiam futura gibbo-
 flore , subtorulosa. Racemis laxis , minutis.

963. G A L E G A.

Cal. dentibus subulatis , subæqualibus. *Legumen*
 striis obliquis , seminibus interjectis.

1. GALEGA *officinalis*, Galega leguminibus strictis erectis ,
 foliolis lanceolatis striatis nudis. *Regn. bot.*

G. foliis ellipticis , spinis & siliquis acutis, *Hall. helv.*
 n. 394.

Galega. *Riu.* t. 72. *Ludw. ed.* t. 161.

G. vulgaris. *Bauh. pin.* 352. *Moris, hist.* 2. p. 91. f. 2. t. 7.
 f. 9. *Blackw.* t. 92.

Habitat in Hispania , Italia , Africa ad versuras. 24

964. P H A C A.

Legumen semibiloculare.

1. PHACA *bætica*. P. caulescens erecta pilosa , leguminibus
 tereti - cymbiformibus.

Phaca leguminibus rectis. *Roy. lugdb.*

Astragalus bæticus lanuginosus , radice amplissima. *Bauh.*
pin. 351.

240 DIADELPHIA. DECANDRIA:

penduli, flavo-albicantes. Legumina glabra, brevia, triquetra, acuta, intra calycem pedicellata. Semina utrinque 2.

8. **ASTRAGALUS** *Onobrychis*. A. caulescens erectus pubescens, floribus spicatis, vexillis duplo longioribus, stipulis solitariis. *Gmel. sib. 4. p. 43. n. 57. t. 21. Jacq. austr. t. 38. delph.*

A. caule subrecto, ramoso, foliis linearibus, sericeis vexillis longissimis. *Hall. helv. n. 412.*

Phaca caulescens erecta subhirsuta, leguminibus cordato-oblongis inflatis subpilosus. *Willich. obs. 115.*

Onobrychis spicata, flore purpureo. *Bauh. pin. 350.*

Onobrychis I. *Clus. hist. 2. p. 238.*

β. *Astragalus (tenuifolius)* caulescens erectus glaber, foliis linearibus, pedunculis elongatis triquetris. *Spec. 2. p. 1065. n. 6.*

Habitat in Austria, Sibiria, Helvetia. 2

* * *Caulibus foliosis, diffusis.*

12. **ASTRAGALUS** *Cicer*. A. caulescens prostratus, leguminibus subglobosis inflatis mucronatis pilosis. *Jacq. austr. t. 251. Kniph. cent. 8. n. 10. delph.*

A. leguminibus subglobosis, floribus spicatis, caule erecto. *Hort. cliff.*

A. caule ramoso, diffuso, spicis erectis, siliquis inflatis, rotundis, unguiculatis. *Hall. helv. n. 409.*

Cicer sylvestre, foliis oblongis hispidis, majus. *Bauh. pin. 247.*

Cicer sylvestre. Cam. epit. 205.

Glaux. Riv. t. 108. Hall. R.

Habitat in Austria, Helvetia, Italia, Germania. 2

Foliola oblonga, sublanceolata sunt.

13. **ASTRAGALUS** *microphyllus*. A. caulescens erecto patulus: foliolis ovalibus, calycibus tumidiusculis, leguminibus subrotundis.

A. caulibus procumbentibus, pedunculis longioribus, spicis ovatis glaberrimis. *Gmel. sib. 4. p. 41. t. 19.*

Cicer, foliis oblongis hispidis, minus. *Bauh. pin. 347.*

Habitat in Sibiria, Hareynia, Thuringia. 2

Habitus A. *Ciceris dicti*, at quadruplo minor. Caulis erectus, flexuosus, pedalis, non glaber. Rami patentes, breviores. Folia pinnata: foliolis 13. f. 15. ovalibus, obtusis, interdum emarginatis. Stipulae subulatae, minimae. Pedunculi solitarii, foliis longiores, superne floribus horizontalibus instructi. Calyces glabri, subrotundi, parum ventricosi, pilis

DIADILPHEIA DECANDRIA

pl. rar. v. 1. p. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

24. **ASTRAGALUS** *p. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.*

A. luteus *caulescens* *procumbens* *leguminibus* *subulatis* *recurvatis* *glabris* *foliis* *obcordatis* *subtus* *villosis* *manfr. Kniph. cens. 11. n. 13. p. 15.*

A. repens *caulescens* *procumbens* *leguminibus* *subulatis* *recurvatis* *glabris* *foliis* *obcordatis* *subtus* *villosis* *manfr. Kniph. cens. 11. n. 13. p. 15.*

A. luteus *caulescens* *procumbens* *leguminibus* *subulatis* *recurvatis* *glabris* *foliis* *obcordatis* *subtus* *villosis* *manfr. Kniph. cens. 11. n. 13. p. 15.*

Gymnocarpus *lychnis* *caulescens* *procumbens* *leguminibus* *subulatis* *recurvatis* *glabris* *foliis* *obcordatis* *subtus* *villosis* *manfr. Kniph. cens. 11. n. 13. p. 15.*

Habitat in Europa *temeribus.*

25. **ASTRAGALUS** *luteus*. *A. caulescens* *procumbens* *leguminibus* *subulatis* *recurvatis* *glabris* *foliis* *obcordatis* *subtus* *villosis* *manfr. Kniph. cens. 11. n. 13. p. 15.*

A. leguminibus *pendulis* *recurvis* *filiculis* *Hort. cliff.*

A. filiqua *curva* *Riz. terr. 221.*

A. monspeliacus *Clus. hif. 2. p. 234.*

Securidaca *lutea* *minor* *corniculis* *recurvis* *Bauh. pin. 349.*

Habitat *Messanæ* *Monspeli.* ☉

Caulis *sæpius* *decumbens* *rarius* *erectus*. *Foliola* *obcordata* *subtus* *villosa*. *Pedunculi* *foliis* *æquales* *aut* *longiores*. *Spicæ* *paucifloræ* *4. f. 5. floribus*. *Flores* *leucophæi* *remoti*. *Legumina* *declinato-hamosa* *acutissima* *apice* *et* *basi* *distincto*.

17. **ASTRAGALUS** *boeticus*. *A. caulescens* *procumbens* *spicis* *pedunculatis* *leguminibus* *prismaticis* *rectis* *tri-quetris* *apice* *uncinatis*. *Kniph. cens. 11. n. 14.*

A. caule *erecto* *floribus* *spicatis* *leguminibus* *oblongis* *acinaciformibus* *apice* *recurvo*. *Roy. lugdb.*

A. annuus *foliis* *&* *filiquis* *longioribus* *glabris* *ad* *folio* *un-*
orum *communi* *pediculo* *innixis*. *Pluk. alm. 60.*

A. luteus *annuus* *boeticus* *maritimus* *rectus* *alter*. *Morif. hif. 2. p. 108. f. 2. t. 9. f. 11.*

Securidaca *ficula* *filiquis* *foliaceis*. *Boc. sic. - - -*
Rai. hif. 935.

Habitat in *Sicilia* *Lusitania* *Hispania.* (-)

18. **ASTRAGALUS** *stella*. *A. caulescens* *diffusus* *capitulis* *pedunculatis* *lateralibus* *leguminibus* *rectis* *subulatis* *unc-*
ronatis.

Linnaei spec. Plant. Europ. Pars II.

Q

242 DIADELPHIA. DECANDRIA.

A. annuus maritimus procumbens latifolius, floribus pediculo infidentibus. Tournef. inst. 416.

A. hirsutus, corniculis plurimis uno pediculo stelliformiter adaptis. Pluk. phys. 79. f. 4.

Astrag. annuus angustifolius flosculis pedunculis oblongis infidentibus. Tournef. 416.

Glauci leguminosæ affinis. Bauh. prodr. 150.

Stella leguminosa. Bauh. hist. 2. p. 350. Tabern. ic. 512.

Habitat Monspeli.

Differt ab A. sesameo Pedunculis elongatis leguminibusque plurimis.

19. *ASTRAGALUS sesameus. A. caulescens diffusus, capitulis sessilibus lateralibus, leguminibus erectis subulatis acumine reflexis.*

A. annuus angustifolius, flosculis subæruleis. Garid. aiz. 50. t. 12.

A. annuus, foliis & siliquis hirsutis: plurimis in foliorum alis sessilibus. Pluk. alm. 60. t. 79. f. 3.

Ornithopodio affinis hirsuta, fructu stellato. Bauh. pin. 350.

Vicia sesamea apula. Col. ecphr. 1. p. 303. t. 301.

Habitat in G. Narbonensi, Italia. ☉

Flores minuti, cærulescentes, axillares, sessiles, calyce non longiores. Legumina subterna, erecta, incurva, hispida, apice patula.

Flores luteos pronunciat. Gouan. illustr. p. 50. n. 2. in nota. R.

20. *ASTRAGALUS pentaglottis. A. caulescens procumbens, leguminibus capitatis replicatis compressis conniventibus cristatis acumine reflexo.*

*A. (echinatus) caulescens procumbens, leguminibus capitatis ovatis triquetris echinatis, apice hamoso. Murray. prodr. 222. **

A. (cristatus) caule diffuso, leguminibus capitatis hirsutis squamoso-strigosis ovato-triquetris. Gouan. illustr.

A. supinus siliquis villosis glomeratis. Tournef. inst. 416. Gouan. R.

A. hispanicus, siliqua epiglottidi simili, flore purpureo, major. Herm. lugdb. 74. t. 75.

Glaux hispanica. Riu. tetr. 223. f. 2.

Habitat in Hispania. ☉

Caules procumbentes, palmares. Folia pinnata: 15, oblonga, emarginata, subtus pubescentia. Petioli. Stipulæ ovato-lanceolatae. Pedunculi axillares.

tenes, foliis oppositis, alternis, ovatis, serratis, petalis, calice, legumine, siliqua, stipite, radice, etc.

21. ASTRAGALUS *leguminibus, stipite, radice, etc.*

A. *Stipite, radice, etc.*
Hort. cist.

A. *Stipite, radice, etc.*
Hern. lugub. -

Glaux minima. Rau. etc.
Habitat in Orientis, A. etc. *Corollae albr.*

22. ASTRAGALUS *leguminibus, stipite, radice, etc.*
acumine reflexo.

A. *villosus procumbens*
Rai. app. 151.

Habitat in Hispania
Similitudinem gerit

lescens. Car. etc.
Flores plures

caerulei, etc.
subulata

cum, etc.
in A. etc.

23. A. *etc.*
car. etc.

A. *etc.*
A. *etc.*

Habitat *etc.*
in

244 DIADELPHIA. DECANDRIA.

bens, floribus subracemosis, erectis, foliolis tomentosis.
Fl. suec. Scholl. barb. Flor. dan. t. 614. Kniph. cent. 9. n. 15. lithu.

A. pedunculis folia villosa æquantibus, floribus erectis, caulibus procumbentibus. delph.

A. incanus parvus nostras. Pluk. alm. 59. Rai. angl. 3. p. 326. t. 12. f. 3. mala.

Glaux montana purpurea nostras. Rai. hist. 939.

Habitat in Angliæ, Scanizæ arena mobili. 24

25. ASTRALAGUS Glaux. *A. caulescens diffusus, capitulis pedunculatis imbricatis, ovatis, floribus erectis, leguminibus ovatis callosis inflatis. delph.*

Ciceri sylvestri minori affinis. Bauh. pin. 347.

Glaux dioscoridis. Clus. hist. 2. p. 240. t. 241.

Habitat in Hispania, Sibiria. 24

Radix lignosa. Caules spithamæi, plurimi, superne magis subvillosi. Folia pinnata: foliolis 21 f. 23. minimis, ovali-oblongis, subtus pilis albis adspersis. Pedunculi ex summis foliorum alis, foliis sæpius breviores, pilosi. Capitulum Trifolii facie, ovatum, calycibus ante & post florescentiam apicibus nigris. Corollæ erectæ, acuminatæ, vexillo fere duplo longiores.

27. ASTRAGALUS alpinus. *A. caulescens procumbens, floribus pendulis racemosis, leguminibus utrinque acutis pilosis. Flor. dan. t. 51. delph.*

A. pedunculis folio longioribus, floribus laxè spicatis pendulis. Fl. suec.

A. caule procumbente ramoso, foliis ovatis, siliquis inflatis, pendulis, hirsutis. Hall. helv. n. 404.

A. alpinus minimus. Fl. lapp. 267. t. 9. f. 1.

A. alpinus, folio viciæ. Scheuch. alp. 509. f. 7.

Habitat in Alpibus Lapponicis, Helveticis. 24

** * * Scapo nudo, absque caule foliis.*

30. ASTRAGALUS montanus. *A. subacaulos, scapis folio longioribus, floribus laxè spicatis erectis, leguminibus ovatis acumine inflexo.* Jac. austr. t. 167. carn. 45. Pall. it. 2. p. 568. delph.*

A. foliis ovato lanceolatis, sericeis, siliquis erectis, inflatis, subhirsutis. Hall. helv. n. 408.

Phaca (montana) subacaulis procumbens hirsuta, scapo floribus erectis, leguminibus ovato-lanceolatis. Crantz, austr.

Astragaloides alpina purpurea villosa pumila, foliis brevioribus acuminatis. Till. pis. p. 19. t. 14. f. 3. R.

646 DIADELPHIA. DECANDRIA.

A. incurvus, *siliqua incurva*. *Boiss. monsp.* 33. *affinis*
nicaens. 158.

Onobrychis incana. *Bauh. pin.* 149. *foliis longioribus.*
Bauh. pin. 340. *Roem. hist.* 928.

Habitas in Galloprovincia. 2.

Scapi scabri, floribus saepe ciliatis. Calyx florum non retrorsus.
Vexillum triplicatum, alis albis longius. Folia lanceolata.
Legumina subincurva, angusta.

Differt ab Ononispistulano foliis rotundioribus, legumine;
Leguminibus brevioribus angustioribus.

37. ASTRAGALUS campestris. *A. acaulis, calycibus leguminibusque villosis; foliis lanceolatis acutis, scapo decumbente.* *Fl. haec. delph.*

A. spicis aphyllis, foliis lanceolatis hirsutis, siliquis erectis, inflatis, semibilocularibus. *Hall. helv. ed. 2. n. 406. s. 13.*

A. acaulos, foliis peracutis, calyce & fructu villosis. *Hall. helv. ed. 1. p. 567. s. 13.*

A. perennis supinus, foliis & siliquis hispidis, flore luteo. *Bux. Hall.* 32.

Habitas in Oelandia, Germania, Helvetia. 2.

Corolla flavae Carina acuminata basi purpurascens.

38. ASTRAGALUS depressus. *A. acaulis, scapis foliis brevioribus, leguminibus cernuis, foliolis subemarginatis nudis.* *Amoen. acad. 4. * delph.*

A. supinus minor, flosculus albis. *Berth. lugdb. 2. p. 54.*

Habitas in alpibus Europae: ad mare caspium. *Pall. it. 2. p. 329. 2.*

39. ASTRAGALUS exscapus. *A. acaulis exscapus, leguminibus lanatis, foliis villosis.*

A. perennis supinus, foliis & siliquis hispidis. *Buxb. hall.* 32.

A. acaulos flore luteo. *Buxb. cent. 3. p. 21. s. 38. f. 2.*

Glux lanuginosa montana acaulos. *Rupp. ien. 2. p. 270.*

Glux montana acaulon. *Bauh. pin.* 347. *prodr.* 147.

Habitat in Thuringia. 2. *D. Schreber.*

Caulis aut Scapus nullus. Folia pinnata; spichamena, villosa. Foliola 21, 33, ovata, obtusa, sessilis. Flores numerosi, radicales, subsessiles, lutei, demum flavi. Calyx ovatus, venricosus, albo-villosus, dentibus albis.

Obf. Lob. ic. 2. p. 79. hanc refert, sed flores lutei.

* * * *Caulis lignoso.*

40. ASTRAGALUS tragacanthoides. *A. subacaulis, floribus radicebusque numerosis subsessilibus.* *Kni. cent. 3.*

** 12.*

THE UNIVERSITY OF CHICAGO

~~SECRET~~

THE UNIVERSITY OF CHICAGO

1. DATE 10/10/1964 BY 100749 100749

151. - - - - -

 DEPARTMENT OF THE INTERIOR
 BUREAU OF LAND MANAGEMENT
 WASHINGTON, D. C. 20080

2000 1000 500

~~Index : 200 : 1000 : 10000~~

~~SECRET~~ ~~CONFIDENTIAL~~ ~~RESTRICTED~~ ~~NOFORN~~

7-11-1944

THE UNIVERSITY OF CHICAGO LIBRARY

512

۱۰۰

1. The first part of the document is a list of names and titles, including "The Hon. Mr. Justice" and "The Hon. Mr. Justice".

3. ~~THE~~ ~~REASONING~~ ~~PROCESSES~~ ~~OF~~ ~~THE~~ ~~MIND~~

127-100-112. 127-100-112. 127-100-112. 127-100-112. 127-100-112.

294 פתח עיני מלך

ACTY. = "

Business in Indian Territory **July 1, 1900 to Dec. 31, 1900.**

班主 王 二 三 四 五

Leggieri, Giuseppe, 1898-1974, 1974

10

I. BISSER, 14. Februar, 1914, 10. 11. 1914, 10. 11. 1914

4. 10

At the time of the investigation, the following persons were interviewed:

1000

Source: *Journal of the American Medical Association*, 1940, 115: 1000-1001.

SECRET

Наименование: _____

367 3 2 2 2 1 1 1

Casey Sullivan, 24, 1000 1st St., N.E., Wash., D.C.

1. *Chlorophyll a* (Chl *a*)
 2. *Chlorophyll b* (Chl *b*)
 3. *Chlorophyll c* (Chl *c*)
 4. *Chlorophyll d* (Chl *d*)
 5. *Chlorophyll e* (Chl *e*)
 6. *Chlorophyll f* (Chl *f*)
 7. *Chlorophyll g* (Chl *g*)
 8. *Chlorophyll h* (Chl *h*)
 9. *Chlorophyll i* (Chl *i*)
 10. *Chlorophyll j* (Chl *j*)
 11. *Chlorophyll k* (Chl *k*)
 12. *Chlorophyll l* (Chl *l*)
 13. *Chlorophyll m* (Chl *m*)
 14. *Chlorophyll n* (Chl *n*)
 15. *Chlorophyll o* (Chl *o*)
 16. *Chlorophyll p* (Chl *p*)
 17. *Chlorophyll q* (Chl *q*)
 18. *Chlorophyll r* (Chl *r*)
 19. *Chlorophyll s* (Chl *s*)
 20. *Chlorophyll t* (Chl *t*)
 21. *Chlorophyll u* (Chl *u*)
 22. *Chlorophyll v* (Chl *v*)
 23. *Chlorophyll w* (Chl *w*)
 24. *Chlorophyll x* (Chl *x*)
 25. *Chlorophyll y* (Chl *y*)
 26. *Chlorophyll z* (Chl *z*)
 27. *Chlorophyll aa* (Chl *aa*)
 28. *Chlorophyll ab* (Chl *ab*)
 29. *Chlorophyll ac* (Chl *ac*)
 30. *Chlorophyll ad* (Chl *ad*)
 31. *Chlorophyll ae* (Chl *ae*)
 32. *Chlorophyll af* (Chl *af*)
 33. *Chlorophyll ag* (Chl *ag*)
 34. *Chlorophyll ah* (Chl *ah*)
 35. *Chlorophyll ai* (Chl *ai*)
 36. *Chlorophyll aj* (Chl *aj*)
 37. *Chlorophyll ak* (Chl *ak*)
 38. *Chlorophyll al* (Chl *al*)
 39. *Chlorophyll am* (Chl *am*)
 40. *Chlorophyll an* (Chl *an*)
 41. *Chlorophyll ao* (Chl *ao*)
 42. *Chlorophyll ap* (Chl *ap*)
 43. *Chlorophyll aq* (Chl *aq*)
 44. *Chlorophyll ar* (Chl *ar*)
 45. *Chlorophyll as* (Chl *as*)
 46. *Chlorophyll at* (Chl *at*)
 47. *Chlorophyll au* (Chl *au*)
 48. *Chlorophyll av* (Chl *av*)
 49. *Chlorophyll aw* (Chl *aw*)
 50. *Chlorophyll ax* (Chl *ax*)
 51. *Chlorophyll ay* (Chl *ay*)
 52. *Chlorophyll az* (Chl *az*)
 53. *Chlorophyll aza* (Chl *aza*)
 54. *Chlorophyll abz* (Chl *abz*)
 55. *Chlorophyll acz* (Chl *acz*)
 56. *Chlorophyll adz* (Chl *adz*)
 57. *Chlorophyll aez* (Chl *aez*)
 58. *Chlorophyll afz* (Chl *afz*)
 59. *Chlorophyll agz* (Chl *agz*)
 60. *Chlorophyll ahz* (Chl *ahz*)
 61. *Chlorophyll aiz* (Chl *aiz*)
 62. *Chlorophyll ajz* (Chl *ajz*)
 63. *Chlorophyll akz* (Chl *akz*)
 64. *Chlorophyll alz* (Chl *alz*)
 65. *Chlorophyll amz* (Chl *amz*)
 66. *Chlorophyll anz* (Chl *anz*)
 67. *Chlorophyll aoz* (Chl *aoz*)
 68. *Chlorophyll apz* (Chl *apz*)
 69. *Chlorophyll aqz* (Chl *aqz*)
 70. *Chlorophyll arz* (Chl *arz*)
 71. *Chlorophyll asz* (Chl *asz*)
 72. *Chlorophyll atz* (Chl *atz*)
 73. *Chlorophyll auz* (Chl *auz*)
 74. *Chlorophyll avz* (Chl *avz*)
 75. *Chlorophyll awz* (Chl *awz*)
 76. *Chlorophyll axz* (Chl *axz*)
 77. *Chlorophyll ayz* (Chl *ayz*)
 78. *Chlorophyll azz* (Chl *azz*)
 79. *Chlorophyll azaa* (Chl *aza*)
 80. *Chlorophyll abz* (Chl *abz*)
 81. *Chlorophyll acz* (Chl *acz*)
 82. *Chlorophyll adz* (Chl *adz*)
 83. *Chlorophyll aez* (Chl *aez*)
 84. *Chlorophyll afz* (Chl *afz*)
 85. *Chlorophyll agz* (Chl *agz*)
 86. *Chlorophyll ahz* (Chl *ahz*)
 87. *Chlorophyll aiz* (Chl *aiz*)
 88. *Chlorophyll ajz* (Chl *ajz*)
 89. *Chlorophyll akz* (Chl *akz*)
 90. *Chlorophyll alz* (Chl *alz*)
 91. *Chlorophyll amz* (Chl *amz*)
 92. *Chlorophyll anz* (Chl *anz*)
 93. *Chlorophyll aoz* (Chl *aoz*)
 94. *Chlorophyll apz* (Chl *apz*)
 95. *Chlorophyll aqz* (Chl *aqz*)
 96. *Chlorophyll arz* (Chl *arz*)
 97. *Chlorophyll asz* (Chl *asz*)
 98. *Chlorophyll atz* (Chl *atz*)
 99. *Chlorophyll auz* (Chl *auz*)
 100. *Chlorophyll avz* (Chl *avz*)
 101. *Chlorophyll awz* (Chl *awz*)
 102. *Chlorophyll axz* (Chl *axz*)
 103. *Chlorophyll ayz* (Chl *ayz*)
 104. *Chlorophyll azz* (Chl *azz*)
 105. *Chlorophyll azaa* (Chl *aza*)
 106. *Chlorophyll abz* (Chl *abz*)
 107. *Chlorophyll acz* (Chl *acz*)
 108. *Chlorophyll adz* (Chl *adz*)
 109. *Chlorophyll aez* (Chl *aez*)
 110. *Chlorophyll afz* (Chl *afz*)
 111. *Chlorophyll agz* (Chl *agz*)
 112. *Chlorophyll ahz* (Chl *ahz*)
 113. *Chlorophyll aiz* (Chl *aiz*)
 114. *Chlorophyll ajz* (Chl *ajz*)
 115. *Chlorophyll akz* (Chl *akz*)
 116. *Chlorophyll alz* (Chl *alz*)
 117. *Chlorophyll amz* (Chl *amz*)
 118. *Chlorophyll anz* (Chl *anz*)
 119. *Chlorophyll aoz* (Chl *aoz*)
 120. *Chlorophyll apz* (Chl *apz*)
 121. *Chlorophyll aqz* (Chl *aqz*)
 122. *Chlorophyll arz* (Chl *arz*)
 123. *Chlorophyll asz* (Chl *asz*)
 124. *Chlorophyll atz* (Chl *atz*)
 125. *Chlorophyll auz* (Chl *auz*)
 126. *Chlorophyll avz* (Chl *avz*)
 127. *Chlorophyll awz* (Chl *awz*)
 128. *Chlorophyll axz* (Chl *axz*)
 129. *Chlorophyll ayz* (Chl *ayz*)
 130. *Chlorophyll azz* (Chl *azz*)
 131. *Chlorophyll azaa* (Chl *aza*)
 132. *Chlorophyll abz* (Chl *abz*)
 133.

9. PSOR 4734

foliis lanceolatis ovatis, apice mucronatis, nervis

Клірн. сент. 2. 7. 63, 101, 14

P. foliis omatibus ternatis *perfoliatus* *cap. 4-5* *Spec. 12*

4

248 DIADELPHIA. DECANDRIA.

Trifolium capitulis subrotundis , calycibus ventricosus , caule
arborescente. *Hort. cliff.*

Trifolium bitumen redolens. *Bauh. pin.* 327.

Trifolium bituminosum. *Dod. pempt.* 566.

β. Trifolium bitumen redolens angustifolium. *Bærh. lugdb.*
2. p. 32.

Trifolium bituminosum inodorum. *Best. cyst. æst.* t. II. f. 2.

Habitat in Siciliae , Italiae , Narbonæ collibus maritimis. H

Variat foliolis lanceolatis & ovatis. Spicæ subrotundæ.
Caulis suffruticosus.

Stamen decimum distinctum , ut P. glandulosæ & ameri-
canæ , an itaque distincti generis.

10. PSORALEA glandulosa. P. foliis omnibus ternatis : fo-
liolis lanceolatis , petiolis scabris , floribus spicatis. *Gouan.*
illustr. Regn. bot.

Psoralea foliis omnibus ternatis , caule erecto arborescente ,
pedunculis spicatis , bracteis minimis ciliatis. *Arduin. spec.*
24. t. 8.

Barba Jovis triphylla , flore ex albo & cæruleo. *Fewill.*
per. 7. t. 3.

Habitat in Peru , Hispania & insulis balearicis. Gou. H

Similis P. americanæ : sed Spicæ elongatæ fere racemosæ.
Bracteis minimis , nec calycum longitudine. Odor ruræ.
Caulis suffruticosus.

968. TRIFOLIUM.

Flores subcapitati. Legumen vix calyce longius non
dehiscens , deciduum.

** Meliloti leguminibus nudis polyspermis.*

1. TRIFOLIUM M. cærulea. T. racemis ovatis ; legumi-
nibus seminudis mucronatis , caule erecto , spicis oblongis.
Darr. nass. Blackw. t. 284. *Kniph. cent.* 5. n. 92. *Regn.*
bot.

T. spicis oblongis , leguminibus seminudis mucronatis , caule
erecto. *Spec.* 2.

Lotus hortensis odorata. *Bauh.* 331.

Lotus sylvestris. *Cam. epit.* 724.

Habitat in Bohemia , Lybia.

2. TRIFOLIUM M. indica. T. leguminibus racemosis nudis
monospermis , caule erecto. *Kniph. cent.* 10. n. 90.

DIADELPHIA. DECANDRIA. 249

T. erectum, caule teretiusculo, pedunculis compresso-angulatis, pericarpis racemosis rotundis rugosis nudiusculis monospermis. *Fl. zeyl.*

Melilotus lutea Indiae orientalis erecta, folliculis rotundis parvis. *Pluk. alm. 246. t. 45. f. 4.*

β. Melilotus humilis erecta, floribus exiguis odoratis albis. *Pluk. alm. 247. t. 45. f. 5.*

γ. Melilotus vulgari nostrati affinis, folio minore & angustiore. *Bauh. hist. 2. p. 371.*

Melilotus angustifolia repens, folliculis rotundis. *Bauh. pin. 331. prodr. 144.*

δ. Melilotus lutea minor, floribus & siliculis minoribus spicatum & dense dispositis. *Morif. hist. 2. p. 161.*

Melilotus nova Berardi s. erecta, folliculis rotundis minor. *Rai. hist. 951.*

Habitat in India, Africa. ☉

Varietates datas vix notae sufficientes distinguunt.

3. TRIFOLIUM M. polonica. T. leguminibus racemosis nudis dispermis lanceolatis, caule erecto.

Habitat in Polonia. ☉

Differt a T. Meliloto officinali Caule omnino tereti. Folii minoribus, acuminatis, superne acute serratis. Racemo floribus remotis longiusque pedicellatis. Pedunculis teretibus, nec sulcatis. Florum vexillis replicatis, nec alis extrorsum marginibus longitudinaliter conniventibus, sed oblique divaricatis. Leguminibus dispermis, parum rugosis, lanceolato acuminatis, longioribus. Statura & Odor conveniunt. Flores pallidissime flavescentes.

4. TRIFOLIUM M. officinalis. T. leguminibus racemosis nudis dispermis rugosis acutis, caule erecto. *suec. paris. sib. austr. gallob. pal. Kniph. cent. 7. n. 95. Blakw. t. 80. Riu. t. 6. Regn. bot. burg. lithu. lugd.*

Melilotus filiquis racemosis, pendulis, acutis, rugosis, dispermis. *Hall. helv. n. 362.*

Trifolium corollis polypetalis; racemis; filiquis monospermis rugosis. *Scop. carn.*

T. odoratum s. Melilotus. *Dod. pempt. 567.*

Melilotus officinarum germaniae. *Bauh. pin. 331.*

β. Melilotus officinarum germaniae, flore albo. *Tournef. inst. 407. lugd. lith.*

Lorus sylvestris, flore albo. *Tabern. hist. 893.*

γ. Melilotus vulgaris altissima frutescens, flore albo s. luteo. *T. Rai. suppl. 407.*

Habitat in Europae campestribus. ☉ ♂

150 DIADELPHIA. DECANDRIA.

Varietas frutescens γ. radice bienni, caule altiori, floribus minoribus, vexillo lateribus deflexo.

5. TRIFOLIUM *M. italica*. T. leguminibus racemosis nudis dispermis rugosis obtusis, caule erecto, foliolis integris.

Melilotus italica, folliculis rotundis. *Bauh. pin.* 331.

Melilotus italica. Cam. hort. 99. t. 29.

Habitat in Italia. ☉

7. TRIFOLIUM *M. ornithopodioides*. T. leguminibus nudis octospermis subternis calyce duplo longioribus, caulibus declinatis. *Flor. dan.* t. 368.

Medicago leguminibus ternatis erectis recurvis descendentibus, pedunculo communi. *Hort. cliff.*

Trifolium siliquosum, loto affine, siliquis ornithopodii. *Pluk. alm.* 375. t. 68. f. 1.

Fœnum græcum humile repens, ornithopodii siliquis brevibus erectis. *Rai. angl.* 3. p. 331. t. 14. f. 1.

Habitat in Anglia, Gallia, Dania. ☉

Racemus ovatus. Media inter Trifolia & Trigonellas.

* * *Lotoidea leguminibus tectis, polyspermis.*

8. TRIFOLIUM *Lupinaster*. T. capitulis dimidiatis, foliis quinatis sessilibus, leguminibus polyspermis. *Kniph. cent.* 5. n. 91.

T. leguminibus polyspermis, foliis pluribus. *Gmel. sib.* 4. p. 19. t. 6. f. 1.

Lupinaster. Buxb. ac. 2. p. 345. t. 20.

Lupinaster floribus purpureis & albis, siliquis minoribus. *Amm. ruth.* 143. 144.

Habitat in Sibiria. 24

10. TRIFOLIUM *strictum*. T. capitulis globosis, leguminibus dispermis; calycibus longitudine corollæ, foliolis ferrulatis, stipulis rhombeis.

T. pratense annum minimum, foliis longis angustis venosis tenuissime serratis, floribus albis congestis, siliquis minoribus dispermis. *Mich. gen.* 29. t. 25. f. 7.

Habitat in Italiae, Hispaniae pratis, pascuis. ☉

Caulis inferne ramosus, patulus, lævis. Folia longe petiolata, ternata, glabra, denticulata, striata, inferiora obovata, superiora elliptica, demum lanceolata. Stipulæ latissimæ, rhombeæ, acumine subulato Pedunculi sulcati, axillares, pedunculo breviores. Spicæ rachi oblonga s. subglobosa. Calyx nudus dentibus subulatis: 2 superioribus corolla longioribus, post anthesin reflexis dentibus patulis.

DIADELPHIA. DECANDRIA. 251

Corolla valde parva. Bractæ inter flores, lineares, acuminatæ, vix corollæ longitudine.

11. TRIFOLIUM *hybridum*. T. capitulis umbellaribus, leguminibus tetraspermis, caule adscendente. *Fl. suec. pal. Dærr. nass. burg.*

Trifolium pratense corymbiferum. *Mich. gen. 28. t. 25. f. 2. 6.*

β. Trifolium caule fistuloso erecto, capitulis subglobosis; foliolis obverse ovatis serratis. *Dalib. parif. 226.*

T. orientale altissimum, caule fistuloso, flore albo. *Vaill. parif. 195. t. 22. f. 5.*

T. flore albo. *Riu. tetr. t. 127. f. 2.*

Habitat in Europæ cultis.

12. TRIFOLIUM *repens*. T. capitulis umbellaribus, leguminibus tetraspermis, caule repente. *Fl. suec. ged. austr. gal. pal. sil. Kniph. cent. 7. n. 96. burg. delph. lugd. lithu.*

T. capitulis subrotundis, flosculis pedunculatis, leguminibus tetraspermis, caule procumbente. *Fl. suec. parif.*

T. caule repente, spicis depressis, siliquis tetraspermis. *Hall. helu. n. 367.*

T. corollis polypetalis; scapis axillaribus; siliquis tetraspermis; caule repente. *Scop. carn.*

T. pratense album. *Bauh. pin. 327.*

T. repens. *Riu. tetr. 17. Fl. lapp. 274.*

Habitat in Europæ pascuis. 24

14. TRIFOLIUM *alpinum*. T. capitulis umbellaribus; scapo nudo, leguminibus dispermis pendulis, foliis lineari-lanceolatis. *delph.*

T. scapis radicatis, floribus racemosis, foliis lanceolatis nervosis. *Hall. helu. n. 369.*

T. scapo nudo simplicissimo, foliis lineari-lanceolatis. *Sauv. monsp.*

T. alpinum, flore magno, radice dulci. *Bauh. pin. 328. prodr. 143.*

T. angustifolium alpinum. *Pon. bald. 340. t. 340.*

Anonis acaulis, floribus longissimis, foliis non serratis. *Hall. helu. 588.*

Habitat in Alpibus Italicis, Helveticis, Pyrenæis, Baldo. 24

Variat albo flore. R.

* * * *Lagopoda calycibus villosis.*

15. TRIFOLIUM *subterraneum*. T. capitulis villosis quin-

252 DIADELPHIA. DECANDRIA.

quefloris, coma centrali reflexa rigida fructum obvolvente.

*Hort. cliff. * Dalib. paris. burg.*

T. subterraneum. Riu. tetr.

T. pumilum supinum, flosculis longis albis. Rai. angl. 3. p. 317. & 478. t. 3. f. 2.

T. album tricoccum subterraneum gastonium reticulatum. Moris. hist. 2. p. 132. f. 2. t. 14. f. 5.

T. pratense supinum cathobleps. Barr. ic. 881.

Habitat in Gallia, Italia. ☉

Fructus capitula globosa terram penetrantia.

17. TRIFOLIUM Cherleri. *T. capitulis villosis globosis terminalibus solitariis, calycibus omnibus fertilibus, caulibus procumbentibus. foliis obcordatis. delph.*

T. capitulis globosis villosis, stipulis floralibus dilatatis cordatis, caulibus procumbentibus. Ger. prou.

T. capitulis villosis, hemisphericis terminalibus bractea subcalyculatis calycibus omnibus fertilibus. Amæ. acad. 4.

T. capitulis villosis globosis, calycibus omnibus fertilibus tomento connexis. Roy. lugdb.

T. capitulis villosis globosis terminalibus bractea orbiculata terminatis. Sauv. monsp.

T. globosum repens. Bauh. pin. 329. prodr. 143.

T. glomerulis personatæ cherleri. Bauh. hist. 2. p. 277.

Lagopus minor supinus, molli & compresso capite. Barr. ic. 859.

Habitat Monspeli. ☉

Affine T. lappaceo, sed Calyces toti pilosi. Cauliculi tomentosi. Bracteæ 3, maximæ, cordatæ, ad margines coloratæ. Stipulæ magnitudine foliorum. Ger.

18. TRIFOLIUM lappaceum. *T. spicis subovatis, calycinis dentibus setaceis hispidis caule patulo, foliis ovatis. delph.*

T. globosum s. capitulo lagopi rotundiorè. Bauh. pin. 329. prodr. 143.

T. capitulo glomerato rigido. Bauh. hist. 2. p. 377.

Habitat Monspeli.

Caulis divaricatus. Calyx lævis dentibus absoluto flore rigescentibus. Corollæ 1. petalæ, pallidæ s. albæ.

19. TRIFOLIUM rubens. *T. spicis villosis longis, corollis monopetalis, caule erecto, foliis serrulatis. monsp. Jacq. austr. t. 385. pal. sub. Kniph. cent. 5. n. 93. burg. lishu. lugd. delph.*

DIADELPHIA. DECANDRIA. 253

T. foliis nervosis, ciliatis, spicis ovatis, obesis, calycis denticulis plumosis, imo longissimo. *Hall. helu. n. 375.*

T. corollis monopetalis æqualibus, calycino imo dente nihilo longioribus, spicis oblongis; foliis ferrularis. *Hall. helu. n. 925.*

T. montanum, spica longissima rubente. *Bauh. pin. 328.*

Lagopus major alter. *Dod. pempt. 578.*

β. **T.** spica oblonga rubra. *Bauh. pin. 328.*

Habitat in Italia, G. Narbonensi, Helvetia, &c. ☉

Dens calycis infimus longitudine corollæ.

20. TRIFOLIUM pratense. **T.** spicis globosis subvillosis; cinctis stipulis oppositis membranaceis, corollis monopetalis. *austr. gallob. tub. herb. pal. sil. Knorr. del. 2. t. T. 3. Kniph. cent. 1. n. 91. Blackw. t. 20. Regn. bot. burg. lithu. lugd.*

T. spicis villosis, caule diffuso, foliolis integerrimis. *Fl. suec. paris.*

T. caule obliquo, foliis ovatis hirsutis, supremis conjugatis, vaginis aristatis. *Hall. helu. n. 377.*

T. corollis monospermis, inæqualibus, spicis subrotundis; stipulis setaceis; foliis integerrimis. *Scop. carn.*

T. pratense, flore monopetalo. *T. Fl. lapp. 273.*

T. pratense purpureum. *Bauh. pin. 327.*

T. pratense. *Cam. epit. 582. Riu. t. 11.*

Habitat in Europæ graminosis. 24

Spica sessilis inter Folia duo, opposita, sessilia, quorum Stipulæ membranaceæ, dilatatæ, calycem quasi communem constituunt.

21. TRIFOLIUM alpestre. **T.** spicis subglobosis villosis terminalibus, caule erecto, foliis lanceolatis ferrularis. *Jacq. obs. 3. p. 14. t. 64. pal. austr. herb. sil. Fl. dan. t. 662. burg. lithu. delph.*

T. foliis nervosis, supremis conjugatis, vaginis lanceolatis. *Hall. helu. n. 376.*

T. spica longiore, flore purpureo. *Riu. t. 12. R.*

T. spicis villosis sobovatis, caule erecto, foliis ovato-oblongis integerrimis. *Roy. lugdb.*

T. montanum purpureum majus. *Bauh. pin. 328.*

T. majus 2. *Clus. hist. 2. p. 245. non album sed rubrum. Bauh. hist. 2. p. 375.*

Habitas in Europa, etiam in Suecia. 24

Simillimum T. pratensi, herba T. montani. Distinguitur a Tr. pratensi, quod hoc alpestre capitulis terminalibus 2,

254 DIADELPHIA. DECANDRIA.

Illud 1. hoc stipulis viridibus, illud scariosis, venis rubris; hoc stipulis ultimis utrinque lanceolatis illud subovatis.

Non est T. quod sub nomine T. alpestris Linn. describit Gouan. in illustr. p. 52, & cui flores ochroleucos adscribit. Videtur Trifolium, Gouano alpestre dictum, ad T. pannonicum referri posse. R.

22. TRIFOLIUM *pannonicum*. T. spicis villosis longis, corollis monopetalis, foliis integerrimis cauleque erecto villosissimis. *Jacq. obs. 2. p. 21. t. 42.*

Habitat in pratis roridis Hungariæ inferioris.

Statura Tr. pratensis, sed majora omnia, undique albavillosa. Caules sesquipedales, erecti. Folia ternata, ovali-oblonga, in stipula sessilia. Spica ovato-oblonga, densa. Flores pollicares, albi.

23. TRIFOLIUM *squarrosum* T. spicis oblongis subpilosis, calycum infimo dente longissimo reflexo, caule herbaceo erecto.

T. purpureum lagopoides hirsutum annuum rotundifolium, spica dilute rubente. Moris. hist. 2. p. 140. f. 2. t. 13. f. 1.

T. hispanicum angustifolium, spica dilute rubente. Bauh. pin. 328.

Habitat in Hispania. ☉

Habitus Trifolii pratensis. Folia suprema opposita, subpilosa, lanceolata aut ovalia, emarginata. Spicæ terminales, ovatae; Calyces parum villosi, 5. dentati: dentibus nudis, setaceis: 4. erectis; 5. seu infimo reliquis triplo majore, reflexo, unde Spica squarrosa & uncinata valde existit.

Variat foliis ovalibus & lanceolatis.

24. TRIFOLIUM *incarnatum*. T. spicis villosis oblongis obtusis aphyllis, foliolis subrotundis crenatis. *Dalib. parif. lith. lugd.*

T. foliis subrotundis, spicis cylindricis, calycibus plumosis. Hall. helu. 374.

T. latifolium. Riv. t. 77. R.

T. spica subrotunda rubra. Tournef. inst. 405. R.

T. stellatum glabrum. Ei. ib. R.

T. capitulo displaci. Rai. hist. 194. R.

T. spica rotunda rubra. Bauh. pin. 328.

T. album incarnatum spicatum s. Lagopus maximus. Bauh. hist. 2. p. 376.

Lagopus maximum, folio & facie trifolii pratensis. Loh. 16, 39.

96. TRIFOLIUM *M. messanensis*. T. leguminibus arcuatis, rugosis folio brevioribus, caule diffuso.

Melilotus messanensis procumbens, foliis rugosis, sublongis, spicis florum brevioribus. *Rai. hist.* 952.

Habitas in Sicilia. ☉

Caulis *diffusus*. Facies plantæ *Meliloti*. Folia *ternata*, *carnea*, *suserrata*. Flores *lutei*, *parvi*. Legumen *oblongiusculum*, *albicans*, *rugosum* striis *semi-circularibus*. Sem. *Tr. M. officinalis* *majora*.

969. LOTUS:

Legumen cylindricum, strictum. *Alæ* sursum longitudinaliter convinentes. *Cal.* tubulosus.

* *Leguminibus rarioribus, nec capitulum constituentibus.*

1. LOTUS *maritimus*. L. leguminibus solitariis membranaceo-quadrangulis, foliis glabris bracteis lanceolatis. *Fl. succ. mansp. kal. Kniph. cent.* 7. n. 45.

L. *retragonolobus* *maritimus*, flore *luteo*. *Barth. ad.* 2. p. 346.

L. *filiquosa* *lutea* *filiquis* *pinnatis* *strictioribus* & *longioribus*. *Rai. hist.* 967.

Habitas in Europæ maritimis. 2.

Affinis *nimum* *L. filiquoso*, *ut forte varietas*.

2. LOTUS *filiquosus*. L. leguminibus solitariis membranaceo-quadrangulis, caulibus procumbentibus, foliis subtrus pubescentibus. *Jacq. austr.* 1. 361. *p. 11. delph. burg. lugl.*

L. *caule prostrato*, *scapis unifloris*, *filiquis alatis*. *Hall. helv.* n. 386.

Trifolii character, utut genus habitu facillimum, diffillime exiit.

Racemosa, 1. 2. 3. 4. 5. 6. 7. 39. 40. 41. 42. 43. 46.

Umbellata, 8. 9. 11. 12. 13. 14.

Fasciculata, 15. 16. 44.

Capitata, 17. 18. 29. 30. 31. 32. 33. 34. 35. 36. 37.

38. 45.

Spicata, 10. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28.

R 3.

256 **DIADELPHIA. DECANDRIA.**

28. **TRIFOLIUM stellatum.** T. spicis pilosis ovatis, calycibus patentibus, caule diffuso, foliolis obcordatis. *Scop. carn. delph.*

T. spicis ovatis pilosis, denticulis calycinis lanceolatis patulis. *Hort. cliff. monsp.*

T. stellatum. *Bauh. pin. 329. prodr. 143. hist. 2. p. 376.*

Lagopus minor erectus, capite globofo stellato. *Barr. ic. 860 & 755.*

Habitat in Sicilia, Italia, G. Narbonensi, Carniolia. ☉

Sicæ ovata. Calyx æqualis extus pilosus.

30. **TRIFOLIUM scabrum.** T. capitulis sessilibus lateralibus ovatis, calycibus inæqualibus rigidis recurvis. *carn. pal. delph.*

T. spicis sessilibus lateralibus ovatis scabris. *Hort. cliff. * paris. monsp.*

T. caulibus prostratis, capitulis ovatis, in alis sessilibus, calycibus rigidis, segmentis rectis. *Hall. helu. n. 371.*

T. capitulo oblongo aspero. *Bauh. pin. 329. prodr. 140.*

T. cujus caules ex geniculis glomerulos oblongos proferunt. *Bauh. hist. 2. p. 378.*

T. minus, capite subrotundo parvo albo echinato. *Barr. ic. 870.*

T. dilute purpureum, glomerulis florum oblongis sine pediculis caulibus adnatis. *Rai. cant. 168.*

T. flosculis albis in glomerulis asperis cauliculis adnatis. *Vail. paris. 196. t. 33. f. 1.*

Habitat in Anglia, Gallia, Italia, Germania, &c. ☉

Caules minus ramosi. Foliola ovalia, crassiuscula, margine quasi crenata. Capitula axillaria, ovata, sessilia, dura, rigida, persistencia. Calycis dentes recurvi: exteriores longiores, validiores. Corollæ albidæ, angustæ, longitudine calycis.

31. **TRIFOLIUM glomeratum.** T. capitulis sessilibus hemisphæricis rigidis, calycibus striatis patulis æqualibus.

T. capitulis æqualibus rigidis sessilibus, calycibus striatis patulis æqualibus. *Spec. pl. 2.*

T. capitulis sessilibus lateralibus subglobosis. *Hort. cliff. * meth.*

T. cum glomerulis ad caulium nodos rotundis. *Rai. angl. 3. p. 329. Pluk. phyt. 113. f. 5.*

T. arvense supinum verticillatum *Barr. ic. 882.*

Habitat in Anglia, Hispania.

Caules decumbentes. Capi

Calycibus

DIADELPHIA DECANDRIA 259

palchro ordine digestis : laciniis repens perennans interpres
religuarum occupantibus.

32. TRIFOLIUM *bracteatum*. T. cap. ...
libus ovatis, calycibus ...
Leers herb.

T. capitatus ...
tridis villosis ...

T. laxiale ...
Gou. R.

T. capitatus thymi, Tournef. ...

T. parvum ...
merulis mollioribus ...
paris. 196. t. 33. f. 2.

Habitas in Germania, Gallia, Hispania

Calyces 10. striati, extus ...

*** Vesicaria calycibus inflatis ...

35. TRIFOLIUM *spumosum*. T. ...
inflatis glabris quinquedentatis involucriis ...
pentaphyllis. Hort. cliff.

T. capitulo spumoso laevi. Bauh. pin. 329. t. 10

T. caule nudo, glomerulis glabris Bauh. cliff.

Habitas in Gallia, Italia, Apulia

Legumina calyce longiora, ...
plentia; tetrasperma. Calyces ...
rollæ rubræ.

36. TRIFOLIUM ...
resupina ...
tratis. Hort. cliff.

T. pratense

T. pratense

1. 2. 3.

T. ...

2. 3. 4.

Hab. ...

...

perennans

apice ...

latere ...

unde ...

37. Trifolium

...

258 DIADELPHIA. DECANDRA:

T. fragiferum tomentosum. Magn. monsp. 265. t. 264.

T. glomerulis tomentosis per caulium longitudinem. Bauh. hist. 2. p. 379.

Habitat in G. Narbonensi, Hispania, Lusitania, Mauritania. D. Sauvages. Læfling. 24

Caules undique decumbentes, non repentes. Folia obovata, denticulata, obtusa, glabra stipulis acuminatis. Pedunculi axillares, brevissimi. Capitula parva. Calycis dentes 3 superiores elongati, inferiores 2 parvi. Calyx crassissimo tomento involutus, ut vasa minime conspiciantur. Accedit propius T. folliculato, quam fragifero. Involucrum universale dodecaphyllum. Corollæ fere resupinata.

38. TRIFOLIUM fragiferum. *T. capitulis subrotundis, calycibus inflatis bidentatis reflexis, caulibus repentibus.*

** Fl. suæc. parif. gelr. gallob. pal. burg. delph. lugd.*

T. caule repente, spicis glabris, calycibus sericis, ampelascensibus. Hall. helu. n. 370.

T. capitulis subrotundis, calycibus, indeffloratis, inflatis, caule repente. Crantz. austr.

T. corollis polypetalis; spicis subrotundis; calycibus inflatis bidentatis reflexis reticulatis. Scop. carn.

T. fragiferum friscum. Bauh. pin. 329. Clus. cur. app. 38. n. 39.

T. capitulo spumoso aspero minus. Bauh. pin. 329. prodr. 140. sec. Hallerum.

T. fragiferum. Vaill. parif. 195. t. 22. f. 2.

T. caule nudo, glomerulis glabris. Bauh. hist. 2. p. 279. fig. majore.

Habitat in Suecia, Gallia, Anglia, &c. 24

Pedunculi in hoc foliis longiores, & Caules radices agunt. Calyces parum pubescentes. Corollæ alba, aut rubicunda uti apud exteros. Calyces subglobosi, villosi, bidentati, basi inferiore tridentata.

****** Lupulina vexillata, corollæ inflexis.*

39. TRIFOLIUM montanum. *T. spicis subimbricatis subtribus, vexillis subulatis emarcescentibus, calycibus nudis, caule erecto. pal. sil. burg. delph. lithu.*

T. cauliculo paucifloro, foliis elliptico-lanceolatis, nervosis, subtus subhirsutis. Hall. helu. n. 372.

T. (album) foliis cauleque erecto hirsutis; corollis florentibus erectis; vexillis subulatis in dorsum reclinatis. Crantz. austr.

DIADELPHIA. DECA. 21.

- T. corollis purpureis.
foliis lanceolatis
T. folio ovato
T. capite
Cane
T.
T.
Habitus
Cane
m
ac. T.
V
erect
S
T.
bus,
is,
T.
n.
T.
X
T.
J
T.
Habitus
Cane
altera
dicat
41 T.
n
T
T
T
L
L
B
H
D
I
C
l

260 DIADELPHIA. DECANDRIA.

42. **TRIFOLIUM procumbens.** T. spicis ovalibus imbricatis, vexillis deflexis persistentibus, caulibus procumbentibus. *Fl. succ. herb. pal. tub. fil. delph.*

T. caule erecto, spicis strepentibus paucifloris. *Hall. helv. n. 364. R.*

T. caulibus decumbentibus elongatis; florum vexillis deflexis. *Neck gallob.*

T. pratense luteo-croceum. *Vaill. parif. 196.*

T. lupulinum alterum minus. *Rai. angl. 3. p. 330. t. 14. f. 3. Hall. opusc. 193.*

Habitas in Europæ campestribus. 2.

Similis sequenti, sed major floribus sæpe 10. f. 12. Folia non striata. Sata etiam evasit diversa a sequenti.

Semen 1, sæpe 2. Leers. l. c. R.

43. **TRIFOLIUM filiforme.** T. spicis subimbricatis, vexillis deflexis persistentibus, calycibus pedicellatis, caulibus procumbentibus. *Fl. succ. * herb. fil. delph.*

T. caulibus filiformibus, spicis subquinquefloris: flosculis pedicellatis. *Neck. gallob.*

T. lupulinum minimum. *Morif. hist. 2. p. 142. Rai. angl. 3. p. 331. t. 14. f. 4. Hall. opusc. 87.*

Habitat in Anglia, Germania, &c. ☉

Caules filiformes, procumbentes. Pedunculi filiformes, foliis longiores, sustinentes Flores 3 ad 5, in satis sæpe 12 ad 15, deflexos: Calycibus fructum ferentibus induratis manifestis & distinctis pedicellis. Foliola emarginata, striata.

Semen unicum. Leers l. c. Cum præcedente conjungit Hallerus. R.

44. **TRIFOLIUM suffocatum.** T. acaule exscapum floribus subterfilibus radicalibus. *Jacq. hort. 60.*

T. minimum supinum, flosculorum & seminum glomeratis plurimis confertim ad radicem nascentibus. *Rai. hist. 942.*

T. epitymi capitulis, inter genicula, annuum. *Cupan. Cathol.*

T. vernum repens. *Buxb. cent. 3. p. 18. s. 31 f. 2. forte.*

Habitat in Sicilia arenosis. ☉

Caules vix palmares, multum ramosi in caespitem, depressi terræque impressi ut vix ulli. Folia ternata, glabra, obovata, denticulata. Petioli erecti, longitudine minus plantæ s. pollicares. Flores glomerulis axillaribus sessilibus, quasi terræ innatis. Calyces glomerati, glabri, quinquesidi: laciniis rectis, tubum, hyalina.

Habitat in Hispania, Italia. ☉

5. **MEDICAGO sativa.** M. pedunculis racemosis, leguminibus contortis, erecto glabro. *austr. gallob. pal. Kniph. cent. 8. n. 67. Regn. bot. lugd. delph. burg.*

Medica caule erecto, foliis oblongis serratis, racemis erectis, siliquis planis, iterato contortis. *Hall. helu. n. 382.*

Medica legitima. *Clus. hist. 1. p. 242.*

Medica sativa. *Morif. hist. 2. p. 150. f. 2. t. 16. f. Mill. dict. n. 1.*

Pœnum burgundicum. *Lob. ic. 2. p. 36.*

Habitat in Hispaniæ, Galliæ &c. apricis. 24

6. **MEDICAGO falcata.** M. pedunculis racemosis, leguminibus lunatis, caule prostrato. *parif. austr. gallob. pal. Kniph. cent. 11. p. 67. Fl. dan. t. 233. delph. lugd. burg.*

Medica foliis oblongis serratis, siliquis racemosis falcatis. *Hall. helu. n. 381.*

Medicago racemis axillaribus; siliquis polyspermis falcatis, compressis. *Scop. carn.*

Trifolium sylvestre luteum, siliqua cornuta. *Bauh. pin. 330. Falcata. Riu. t. 84.*

Lens major repens. *Tabern. p. 502. Hall. R.*

Medica flavo flore. *Clus. hist. 2. p. 243.*

Habitat in Europæ pratis apricis, siccis. 24

7. **MEDICAGO lupulina.** M. spicis ovalibus, leguminibus reniformibus monospermis, caulibus procumbentibus. *Succ. parif. austr. gallob. tub. pal. sil. Kniph. cent. 11. n. 68. lugd. delph. lithua. burg.*

Trifolium leguminibus spicatis reniformibus nudis monospermis, caule procumbente. *Hort. cliffi*

Medica caule diffuso, capitulis hemisphæricis, siliquis reniformibus. *Hall. hell. n. 380*

M. spicis axillaribus, ovatis, siliquis monospermis, rugosis. *Scop. carn.*

Trifolium pratense luteum, capitulo brevior. *Bauh. pin. 328.*

Trifolium pratense luteum. *Fusch. hist. 819.*

Melilotus minima. *Riu. t. 8. Hall. R.*

Habitat in Europæ pratis. ♂

Legumina striato-rugosa, pilis rigidis subhispida.

8. **MEDICAGO marina.** M. pedunculis racemosis, leguminibus cochleatis spinosis, caule procumbente tomentoso. *Hort. cliff. * monsp. Kniph. cent. 4. n. 46.*

262 DIADELPHIA. DECANBRIA.

Tetragonolobus Scandalida. Scop. carn.

Lotus pratensis filiquosus luteus. Bauh. pin. 332. major & durior. Tournesf. inst. 403. Segu. ver. 2. p. 90.

L. tetragonolobus. Riv. tetr. 79. bona.

Habitas in Europæ australioris pratis subhumidis. 24

Caules humifusi, pedales. Folia & calyces hirsuta.

3. *LOTUS tetragonolobus. L. leguminibus solitariis membranaceo - quadrangulis, bracteis ovatis. Kniph. cent. 5. n. 33.*

L. leguminibus solitariis : angulis quatuor membranaceis. Hort. cliff.

L. ruber, filiqua angulosa. Bauh. pin. 332.

L. pulcherrima tetragonolobus. Comm. hort. 91. t. 26.

Habitas in Sicilia collibus. ☉ anno 1579. Clusio.

4. *LOTUS conjugatus. L. leguminibus conjugatis membranaceo quadrangulis, bracteis oblongo-ovatis.*

L. luteus, filiqua angulosa. Bark. lugdb.

Habitas Montpelii. ☉

Convenit cum 3 habitu, ætate, foliis hirsutis, leguminibus; differt vero Corollis duplo minoribus, Leguminibus duobus, Alis integris minus crispis, foliorum basi caulem non amplexante, sed ab opposito latere dehiscente.

5. *LOTUS edulis. L. leguminibus subsolitariis gibbis incurvis.*

L. pentaphyllos, filiqua cornuta. Bauh. pin. 332.

L. edulis cretica. Rai. hist. 967.

L. oligoceros cretica lutea, filiquis binis curvis propendentibus. Moris. hist. 2. p. 176. f. 2. t. 18. f. 5.

Trifolium corniculatum creticum. Alp. exot. t. 268.

Habitas in Italia, Sicilia, Creta.

Planta fere L. tetragonolobi. Caulis subvillosus. Folia glabra. Supræ ovata. Pedunculi longi: bracteis, 3 ovatis, longitudine floris. Legumina Crotalaria, ovato-oblonga, gibba, sed sutura dorsali recta.

7. *LOTUS peregrinus. L. leguminibus subbinoatis linearibus compressis natantibus Sauv. monsp.*

L. filiquis geminis, peregrina. Bark. lugdb. 2. p. 38.

Habitas in Europa australi.

8. *LOTUS angustissima. L. leguminibus subbinoatis linearibus strictis erectis, caule erecto, pedunculis alternis.*

L. leguminibus septem linearibus strictis erectis. Hort. cliff. monsp.

L. pentaphyllos monsp.

Lotus angustissima, Bauh.

DIADELPHIA. DECANDRIA. 469

Medica magna turbinata. Bauh. hist. 2. p. 385.

Trifolium cochleatum turbinatum, fructu compresso oblongo. Bauh. pin. 329.

4. *M. (intertexta)* pedunculis unifloris leguminibus cochleatis sphaericis, spinis longioribus divaricatis recurvis. *Ger. prov.*

Medica leguminibus cochleatis spinosissimis, aculeis utrinque tendentibus. Mill. dict.

Medica cochleata spinosa major dicarpos, capsula s. spinis longioribus sursum & deorsum tendentibus. Moris. hist. 1. p. 153. f. 2. t. 15. f. 8. 9. 7. Knorr. del. 2. t. M. 1. a.

Medica echinata spinosa, echinis magnis utrinque turbinatis cum spinulis reflexis. Rai. hist. 962.

Fructus 2. f. 4, ovati: anfractibus spinis setaceis reflexis. Stipulae dentatae.

5. *M. (arabica)* pedunculis subtrifloris, leguminibus echinatis, foliolis obcordatis. *Gou.*

Medica cochleata minor polycarpos annua, capsula majore alba, folio cordato macula fusca. Moris. hist. 2. p. 144. f. 2. t. 15. f. 17.

Medica arabica. Cam. hort. 97. t. 27.

Trifolium cochleatum, folio maculato cordato. Bauh. pin. 329.

Folia macula sanguinea lanceolata, quae in nupta dispareret. Flores 4. f. 5. Fructus spinis longe divergentibus.

n. *M. (coronata.)* pedunculis subbifloris, leguminibus compressis subspinosis, stipulis subulatis integerrimis. *Gou. Scop. l. c.*

M. pedunculis multifloris, leguminibus cochleatis compressis spinulis oppositis. Ger. prov.

M. fructu minimo, coronulam utrinque dentatam referente. Sauv. monsp.

Medicago cochleata polycarpos annua, capsula minima coronata. Moris. hist. 2. p. 155. f. 25. t. 15. f. 21.

Medica coronata cherleri parva. Bauh. hist. 2. p. 386.

Trifolium folio obruso, folliculis cordatis. Bauh. pin. 329: prodr. 141. n. 13.

Flores minimi. Legumina spinis brevioribus erectiusculis. Gouan.

3. *M. (ciliaris)* pedunculis multifloris, leguminibus congestis globosis hispidis, stipulis ciliatis. *Gouan. monsp.*

M. pedunculis multifloris legum. cochleatis ciliatis. Ger. prov.

264 DIADELPHIA. DECANDRIA.

L. siliquis umbellatis rectissimis, cylindricis. *Hall. helm.* n. 385.

L. caule herbaceo, florum capitula depresso leguminibus decumbentibus teretibus. *Fl. succ. parif.*

Lotus s. melilotus pentaphyllos minor glabra. *Bauh. pin.* 332. *Riu.* 1. 76.

Trifolium corniculatum. *Dod. pempt.* 573.

β. Lotus pentaphyllos, flore majore luteo splendente. *Bauh. pin.* 332. *Pollich. l. c.*

γ. Lotus (*tenuifolius*) pentaphyllos frutescens, tenuissimis glabris foliis. *Bauh. pin.* 332. *Pollich. l. c.*

Trifolium corniculatum frutescens tenuissimis foliis. *Bauh. prodr.* 144.

Habitat in Europa. 24

Corollæ odoratissima.

Hujus forte varietas etiam γ est, cujus Caulis duplo longiores & angustiores, Folia linearia, & Legumina angustiora. Videtur in umbra nata; Burserus C. Bauhino hanc attulit.

17. LOTUS *cytisoides*. L. capitulis dimidiatis, caule diffuso, ramosissimo, foliis tormentosis.

L. siliquosa maritima lutea, cytisi facie. *Bar. rar.* 71. 1031.

Habitat in maritimis Europæ australis.

18. LOTUS *Dorycnium*. L. capitulis aphyllis, foliis sessilibus quinatis. *austr. Kniph. cent.* 3. n. 60. *delph.*

Dorycnium foliis digitatis sessilibus. *Hort. cliff. monsp.*

Dorycnium pentaphyllum. *Scop. carn.*

Dorycnium hispanicum. *Clus. hisp.* 201, 202. *Hall. herb.* n. 384.

Dorycnium monspeliensium. *Lab. ic.* 51.

Trifolium album angustifolium, floribus veluti in capitulum congestis. *Bauh. pin.* 329.

Habitat in Hispania, G. Narbonensi, Austria, Carniola. 24

970. TRIGONELLA.

Vexillum & alæ subæquales, patentes, forma corollæ tripetalæ.

1. TRIGONELLA *ruthenica*. T. leguminibus pedunculatis congestis pendulis linearibus rectis, foliolis sublanceolatis. *Emel. sib.* 4. p. 24. n. 33. t. 8.

T. leguminibus pedunculatis. Hort. cliff. 378.

Melilotus supina angustifolia, medicæ facie, siliqua compressa. Amm. ruth. 159. & n. 156.

Lorus montana humilior trifolia ad caulem alata lutea, siliqua modo recta, modo medicaginis instar falcata nigra. Amm. ruth. 119.

Habitat in Sibiria,

Foliola lanceolata, obtusissima, superne acute serrata, dentata. Stipulæ integerrimæ. Legumina linearia.

3. *TRIGONELLA polycerata. T. leguminibus sessilibus congestis erectis subrectis longis linearibus, pedunculis muticis.*

Medicago leguminibus confertis sessilibus erectis. Sauv. monsp.

Medicago leguminibus confertis longis rectis parallelis pedunculo communi. Hort. cliff.

Fœnum græcum sylvestre alterum polyceration. Bauh. pin. 348.

Fœnum græcum sylvestre alterum, Dod. pempt. 547.

Habitat in Hispania, Italia, Monspeli. ☉

6. *TRIGONELLA corniculata. T. leguminibus pedunculatis congestis declinatis subfalcatis, pedunculis longis subspinosis, caule erecto.*

Trifolium floribus spicatis, leguminibus nudis mucronatis falcatis declinatis. Hort. cliff. gelr. delph.

Melilotus, corniculis reflexis, major. Bauh. pin. 331.

Melilotus lutea major, corniculis reflexis ex eodem centro ortis. Moris. hist. 2. p. 162. f. 2. t. 16. f. 11.

Trifolium corniculatum 2. Dod. pempt. 573.

Habitat in Europa australi. ☉

Caulis huic erectus. Stipulæ simplices, subdentatæ. Pedunculus tetragonus, folio longior, apice molliusculo-spinoso, sub quo Legumina 10 - 12, falcata, pedunculo breviora, Calyces ante explicationem obscuriores. Flores odoratissimi.

7. *TRIGONELLA monspeliaca. T. leguminibus sessilibus congestis arcuatis divaricatis inclinatis brevibus, pedunculo mucronato inermi. delph. burg.*

Medicago leguminibus confertis sessilibus arcuatis inclinatis. Hort. cliff. paris. monsp.

Fœnum græcum polyceration. Riu. tetr.

Fœnum græcum sylvestre polyceratum monspeliacum. Brey. cent. 80. t. 33. f. 2.

Hedysarum minimum. Dalech. hist. 446.

266 DIADELPHIA. DECANDRIA.

Habitat Monspeli. ☉

Legumina 10 ad 12, *declinata*. Pedunculus *brevissimus*, *molle mucronatus*, folia non longior. Caules *prostrati*, terræ *adpressi*.

9. TRIGONELLA *Fœnum græcum*. T. leguminibus sessilibus strictis erectiusculis subfalcatis acuminatis; caule erecto. *Blakw. t. 38. Ludw. ed. t. 160. Regn. bot.*

T. leguminibus sessilibus. *Dalib. paris. monsp.*

Buceras siliquis sessilibus, arcuatis undique divergentibus. *Hall. helv. n. 379.*

Medicago leguminibus subsolitaris sessilibus erectis reflexo-falcatis acuminatis. *Hort. cliff.*

Fœnum græcum sativum. *Bauh. pin. 348.*

Fœnum græcum. *Fuchs. hist. 798. Cam. epit. 199. Riu. t. 82.*

β. Fœnum græcum sylvestre. *Bauh. pin. 348.*

Habitat Monspeli. ☉

Sylvestre β. differt Stolonibus ad radicem longis, præter erectum caulem, terræ *adpressis*. Foliolis obovatis, nec obtuse lanceolatis. Geniculis foliorum purpurascens. Leguminibus magis hirsutis.

971. M E D I C A G O.

Legumen compressum, cochleatum. Carina corollæ a vexillo deflectens.

1. MEDICAGO *arborea*. M. leguminibus lunatis margine integerrimis, caule arboreo. *Kniph. cent. 5. n. 55.*

Medica arborea. *Mill. dict. n. 7.*

Cytisus incanus, siliquis falcatis. *Bauh. pin. 389.*

Cytisus marante. *Lob. ic. 2. p. 46.*

Habitat in Rhodo, Neapoli. ☿

3. MEDICAGO *radiata*. M. leguminibus reniformibus margine dentatis, foliis ternatis. *Hort. cliff. Viadr. 231. Kniph. cent. 10 n. 58.*

M. annua, siliqua falcata striata. *Morif. hist. 2. p. 158. f. 2. t. 16. f. 3.*

Trifolium siliqua falcata. *Bauh. pin. 330.*

Lunaria radiata italorum. *Lob. ic. 2. p. 38.*

Habitat in Italia. ☉

4. MEDICAGO *circinnata*. M. leguminibus reniformibus margine dentatis, foliis pinnatis. *Hort. cliff.*

Loto affinis siliquis hirsutis circinnatis. *Bauh. pin. 333.*

Falcata foliis Anthylloidis. *Riu. tetr. 201.*

DIADELPHIA DECANDRIA. 167

Habitas in Hispania, Italia. ☉

5. **MEDICAGO** *serena*. M. pedunculis racemosis, leguminibus compressis, caule glabro. *aust. gallob. pal. Kniph. cent. 8. n. 67. Regu. bor. lugd. delph. burg.*

Medica caule erecto, foliis oblongis serratis, racemis, erectis, siliquis planis, immixto canonicis. *Hall. helv. n. 382.*

Medica legumina. Clus. hist. 1. p. 242.

Medica serena. Monf. hist. 2. p. 150. f. 2. n. 16. f. Mill. del. n. 1.

Ficaria burgundica. Lab. ic. 2. p. 36.

Habitas in Hispania, Gallia &c. apricis. ☿

6. **MEDICAGO** *falcata*. M. pedunculis racemosis, leguminibus lanceis, caule quadrato. *par. aust. gallob. pal. Kniph. cent. 12. p. 67. Fl. Aen. 1. 233. delph. lugd. burg.*

Medica foliis oblongis serratis, siliquis racemosis falcatis. *Hall. helv. n. 381.*

Medicago racemis axillaribus; siliquis polyispermis falcatis, compressis. *Scop. carn.*

Trifolium sylvestre luteum, siliqua coriacea. *Bauh. pin. 330.*

Falcata. Riv. s. 84.

Leas major repens. Tabern. p. 502. Hall. R.

Medica serena bore. Clus. hist. 2. p. 243.

Habitas in Europæ pratis apricis, ficcis. ☿

7. **MEDICAGO** *lupulina*. M. spicis ovatis, leguminibus reniformibus monospermis, caudibus procumbentibus. *Succ. par. aust. gallob. tab. pal. fl. Kniph. cent. 11. n. 68. lugd. delph. kithra. burg.*

Trifolium leguminibus spicatis reniformibus nudis monospermis, caule procumbente. *Hort. cliff.*

Medica caule diffuso, capitulis hemisphaericis, siliquis reniformibus. *Hall. helv. n. 380.*

M. spicis axillaribus, ovatis, siliquis monospermis, rugosis. *Scop. carn.*

Trifolium pratense luteum, capitulo brevior. *Bauh. pin. 328.*

Trifolium pratense luteum. *Fusch. hist. 819.*

Melilotus minima. *Riv. s. 8. Hall. R.*

Habitas in Europæ pratis. ♂

Legumina striato-rugosa, pilis rigidis subhispida.

8. **MEDICAGO** *marina*. M. pedunculis racemosis, leguminibus cochleatis spinosis, caule procumbente tomentoso. *Hort. cliff. * monsp. Kniph. cent. 4. n. 46.*

268 DIADELPHIA. DECANDRIA.

Trifolium cochleatum maritimum tomentosum. Bauh. pin.
329.

Medica marina. Clus. hist. 2. p. 243.

Medicago incana. Riv. 205.

Habitat in maris Mediterranei litoribus. 2.

9. *MEDICAGO polymorpha. M. leguminibus cochleatis, stipulis dentatis, caule diffuso Paris. carn.*

Trifolium cochleatum, fructu nigro hispido. Bauh. pin. 329.

α. *Medicago (orbicularis) leguminibus solitariis cochleatis depressis planis, stipulis ciliatis, caule diffuso. Sauv. monsp. Scop. l. c.*

M. pedunculis unifloris aristatis, leguminibus cochleatis compressis rugosis nudis. Ger. Prov.

Medica major dicarpos, capsula compressa orbiculata nigra plana oris crispis. Moris. hist. 2. p. 152. f. 2. t. 15. f. 1.

Medica orbiculata. Bauh. hist. 2. p. 384. Rai. hist. 961.

Trifolium cochleatum s. scutellatum, folio latiore, fructu minuto obtuso. Bauh. pin. 329. prodr. 140.

Fructus solitarius, lentiformis, anfractibus inermibus planis. Pedunculi aristati. Stipulae ciliatae.

β. *M. (scutellata) cochleata major dicarpos, capsula rotunda globosa scutellata. Moris. 2. p. 152. f. 2. t. 15. f. 4.*

M. pedunculis multifloris leguminibus cochleatis globosis inermibus nudis. Ger. prov.

Medica scutellata. Bauh. hist. 3. p. 384.

Trifolium cochleatum, fructu latiore. Bauh. pin. 329.

Fructus solitarius, hemisphaericus: anfractibus inermibus concavis. Pedunculi aristati. Stipulae dentatae.

γ. *M. (tornata) leguminibus cochleatis nudis cylindricis apice basique planiusculis. Ger. prov. lugd. delph.*

Medica leguminibus ternatis inermibus, stipulis acute dentatis, foliolis serratis. Mill. dict.

Medica tornata major & minor lenis. Park, theatr. 1116. Moris. hist. 2. f. 2. t. 15. f. 11.

Fructus plures, subcylindrici: anfractibus inermibus planis. Stipulae dentatae.

δ. *M. (turbinata) leguminibus cochleatis inermibus subcylindricis: extremitatibus camplanatis. Gouan. monsp. 517.*

Medicago fructu turbinato. Sauv. monsp.

Medica dicarpos, fructus capsula turbinata rugosa. Moris. hist. 2. p. 153. Knorr. del. 1. t. S. 20.

DIADELPHIA. DECANI

Medica magna turbinata. *Bauh. hist. 2. p.*

Trifolium cochleatum turbinatum, fructu co
go. *Bauh. pin. 329.*

e. M. (intertexta) pedunculis unifloris legumi
chleatis sphæricis, spinis longioribus divaricatis. *Ger. prov.*

Medica leguminibus cochleatis spinosissimis, aculeis utrin-
que tendentibus. *Mill. dict.*

Medica cochleata spinosa major dicarpos, capsula s. spin
longioribus sursum & deorsum tendentibus. *Morif.*

1. p. 153. f. 2. t. 15. f. 8. 9. 7. Knorr. del. 2. t. M.

Medica echinata spinosa, echinis magnis utrinque
cum spinulis reflexis. *Bauh. pin. 329.*

Fructus 2. f. 4.

Libr. 4

flexis. Stipulae densa

f. M. (arabica.) pe

natis, foliolis ob

Medica cochleata minor polycarpos annua, cap. ma-
jore alba, folio cordato macula fusca. *Morif. hist. 2.*
p. 144. f. 2. t. 15. f. 17.

Medica arabica. *Cam. hort. 97. t. 27.*

Trifolium cochleatum, folio maculato cordato. *Bauh. pin.*
329.

Folia macula sanguinea lanceolata, quæ in nupta dis-
paret. Flores 4. f. 5. Fructus spinis longe divergentibus.

n. M. (coronata.) pedunculis subbifloris, leguminibus
compressis subspinosis, stipulis subulatis integerrimis.
Gou. Scop. l. c.

M. pedunculis multifloris, leguminibus cochleatis compres-
sis spinulis oppositis. *Ger. prov.*

M. fructu minimo, coronulam utrinque dentatam refe-
rente. *Sauv. monsp.*

Medicago cochleata polycarpos annua, capsula minima
coronata. *Morif hist. 2. p. 155. f. 25. t. 15. f. 27.*

Medica coronata cherleri parva. *Bauh. hist. 2. p. 386.*

Trifolium folio obruso, folliculis cordatis. *Bauh. pin. 329.*
prodr. 141. n. 13.

Flores minimi. Legumina spinis brevioribus erectiusculis.
Gouan.

s. M. (ciliaris) pedunculis multifloris, leguminibus con-
gestis globosis hispidis, stipulis ciliatis. *Gouan. monsp.*

M. pedunculis multifloris legum. cochleatis ciliatis. *Ger.*
prov.

276 DIADELPHIA. DECANDRIA.

Medica cochleata polycarpus, capsula spinosa minore
perennis ciliaris. *Morif. hist. 2. p. 154.*

Medica ciliaris Guilandini. Rai hist. 995.

Medicago leguminibus congestis globosis hispidis, stipu-
lis ciliatis. *Sauv. monsp.*

Medica annua, fructu ciliari. *Tournef. inst. 410.*

Trifolium fructu cochleato ciliari. Magn. monsp. 269.

*Fructus plures ovales : anfractibus spinis reclusculis vil-
losis. Stipulae ciliatae.*

l. *M. (hirsuta.)* pedunculis multifloris, legum. cochlea-
tis, spinulis hamatis, stipulis integris. *Ger. 518. Neck.
gallob. Scop. l. e. Hall. helu. n. 383. var. β.*

M. leguminibus solitariis globosis brevissime Chinatis,
stipulis ferratis. *Sauv. monsp.*

Medica echinata hirsuta. Bauh. hist. 2. p. 386.

Trifolium echinatum arvense. Bauh. pin. 329.

u. *M. (rigidula)* pedunculis multifloris, legum. Cochlea-
tis intense aculeatis *Ger. prov.*

M. triphylla, leguminibus spinosis, foliolis interioribus
cuneiformibus retusis, superioribus subrotundis. *Dalib.
paris.*

Medicago hirsuta, echinis rigidioribus. *Bauh. hist. 2. p. 385.*

Trifolium fructu compresso spinis horrido. Bauh. pin. 329.

λ. *M. (minima.)* leguminibus cochleatis subternatis, spi-
nulis uncinatis, stipulis integris. *Gouan. Oed. dan. 211.
Crantz. austr. Pollich. pal. delph. lugd. burg.*

*M. leguminibus cochleatis aculeis uncinatis alternis externe
donatis*, stipulis integris. *Guett. stamp. Dalib. paris.*

Medica foliis emarginatis, ferratis, racemis paucifloris,
siliquis cochleatis globosis. *Hall. helu. n. 383.*

*Cochleata echinata. minima. Riu t. 88. Morif. hist. 2. p.
144. t. 15. n. 20. Hall. R.*

Medica echinata minima. Bauh. hist. 2. p. 386.

Trifolium echinatum, fructu minore. *Bauh. pin. 330.*

*Fructus plures, vix semine Orobi majores : anfractibus
hispidis : spinis lateraliter recurvatis. Stipulae integrae.*

μ. *M. (muricata)* pedunculis multifloris, legum. cochleatis
subrotundis spinosis incanis, foliis villosis *Ger. delph.
prov.*

Medica cochleata dicarpus, capsula spinosa rotunda minore.
Morif. hist. 2. p. 153. f. 2. t. 15. f. 11.

ν. *M. (nigra.)* pedunculis paucifloris, leguminibus cochlea-
tis, hamulis longitudine leguminum. *Ger. prov.*

DIADELPHIA. DECAND.

Medica cochleata polycarpus annua, capsu-
diore. *Mors. hist.* 2. p. 154. f. 2. s. 1.

β. *Medicago (lacinata)*, leguminibus coch-
lipulis contortis, foliis oblongis duplicato-
orient.

M. fructu echinato, foliis linearibus dentatis. *Savv. m.*

Medica leguminibus cochleatis spinosis, foliolis acute den-
tatis tricuspidisque. *Mull. dict.* n. 5.

Trifolium fructu echinato, foliis eleganter dissectis. *monsp.* 271. s. 270.

Trifolium cochleatum spinosum syriacum, foliis lacinia-
Breyn. cent. 81. t. 34.

Medicago syriaca. Riv. tetr. 205.

M. dicarpus lacinata ~~Med. cent.~~ tu spinosis
245.

Trifolium echinato c.

Folia quasi *Trigonel* ^{je non exe-}
sis a Larva. Foliola ^{en.} ^{dentibus utrin-}
que 2 longioribus pinnatipaa. Pedunculi aristati, subbiflori,
folius breviores. Legumina spinosa.

Habitat in Europa australi. ☉

Polymorpha hæc species canis instar produxit numere-
fas varietates, quamvis non in eadem regione proveniunt,

CLASSIS XVIII.

POLYADELPHIA.
ICOSANDRIA.

9 CITRUS.

Cal. 3. dentatus. Cor. 5.
petala. Stam. 10. in cylind-
rum passim connata. Pist.
Bacca locularis pulpe
locularis.

YANDRIA.

911. HYPERICUM.

Cal. 3. pariter, inferus. Cor.
5. petala. Styli 1. 3. l. 3.
Caps. locularis.

POLYADELPHIA.

POLYADELPHI

ICOSANDRIA

974. CITRUS.

Cal. 5. fidus. *Petala* 5, oblonga. *Antheræ* 20.
mentis connatis in varia corpora. *Bac.* 9. loc

1. CITRUS *medica*. C. petiolis linearibus. *Regn. bot.*
Malus Medica. *Bauh.* 1. 361.

β. Citrus. *Limon.*

Limon vulgaris. *Ferr.*
del. 1. 1. C. 1.

Malus Limonia acida. β.

Habitat in Asia, Media

Varietates hujus hodie numerosissimæ una cum sequentibus
mixtae, de quibus Hesperides consulendæ.

Venis in Italiam post Virgilii & Plinii tempora ante Palladii.

2. CITRUS *Aurantium*. C. petiolis alatis, foliis acuminatis.
Mill. illustr. Regn. bot.

Aurantium. *Blakw.* 1. 349. *Kniph. cent.* 9. n. 23. *Knör. del.*
1. 1. P. 4.

Malus Aurantia major. *Bauh. pin.* 436.

Aurantia mala. *Cam. epit.* 150.

β. Citrus *Sinensis*.

Malus aurantia cortice eduli. *Bauh. pin.* 436.

Habitat in India. β

POLYANDRIA.

981. HYPERICUM.

Cal. 5. partitus. *Petala* 5. *Filamenta* multa, in 5
phalanges basi connata. *Capsula*.

* *Pentagyna*.

1. HYPERICUM *balearicum*. H. floribus pentagynis.
Linnaei spec. Plant. Europ. Pars II.

274 POLYADELPHIA. POLYANDRIA.

caule fruticoso, foliis ramisque cicatrifatis. *Mill. dict. n. 1. 54. Kniph. cent. 2. n. 35.*

H. floribus pentagynis, folio & ramis verrucosis. Hort. cliff.

Hypericum f. Ascyrum frutescens, magno flore. Magn. char. 260.

Myrto - Cystus pennæi. Clus. hist. 1. p. 68.

Habitat in Majorca. H

Caules tetragoni, punctis racemoso - glandulosis adspersi. Folia flexuosa, margine glandulis verrucosis. Flores solitarii, terminales.

6. *HYPERICUM Ascyron. H. floribus pentagynis, caule tetragono herbaceo erecto simplici, foliis lævibus integerrimis. Gmel. fib. 4. p. 178. t. 69. Kniph. cent. 9. n. 50.*

H. floribus pentagynis, foliis ovato-oblongis glabris integerrimis. Hort. cliff.

Androsæmum flore & theca quinquecapsulari omnium maximis. Moris. hist. 2. p. 472.

Ascyrum magno flore. Bauh. pin. 280. prodr. 130. Burf. VVI. 25.

Habitat in Sibiria, Canada, Pyrenæis. 24

Folia acuta. Linn. hyper. p. 4. R.

* * *Trigyna.*

8. *HYPERICUM Androsæmum. H. floribus trigynis, fructibus baccatis, caule fruticoso ancipiti. Regn. bot. delph. lugd.*

H. floribus trigynis, fructu baccato, foliis ovatis pedunculo longioribus. Hort. cliff. paris.

Androsæmum maximum frutescens. Bauh. pin. 280.

Androsæmum. Dod. pempt. 78. Blackw. t. 94.

Habitat in Angliæ, G. Narbonensis, Italiæ humentibus. H

9. *HYPERICUM olympicum. H. floribus trigynis, calycibus acutis, staminibus corolla brevioribus, caule fruticoso. §. ic. t. 151. f. 1.*

H. montis olympi. Wheel. itin. 222. Dill. elth. 182. t. 151. f. 183.

H. orientale flore magno. Tournef. cor. 19.

Habitat in Pyrenæis, Olympo. H

12. *HYPERICUM hircinum. H. floribus tryginis, staminibus corolla longioribus, caule fruticoso ancipiti. Kniph. cent. n. 51.*

H. foetidum frutescens minus. Dill. elth. 182.

Androsæmum foetidum, capitulis longissimis filamentis donatis. Bauh. pin. 280.

POLYADELPHIA POLYANDRIA. 273

Tournef. Chyf. diff. 2. p. 205.

Habitat in Sicilia; Calabria; Creta ad rivet. B

17. **HYPERICUM** *ericaides*. H. floribus trigynis, foliis
nerviis serratis.

H. *ericaides* minimum, foliis cinereis. *Pluk. alm. 189. et*
93. f. 5.

H. tomentosum lusitanicum minimum. *Tournef. infl. 256.*

Habitat in Lusitania, Hispania. 24

21. **HYPERICUM** *quadrangulare*. H. floribus trigynis,
caule quadrato herbaceo. *Fl. succ. parif. gallob. herb. pal.*
fl. Flor. dan. 640. delph. lith. burg. lugd.

H. caule quadrangulo, foliis ovatis perforatis, punctatis
Hall. helu. n. 1038.

H. calyce integro, foliis margine punctatis, caule quadrato
herbaceo. *Crantz. austr.*

Hypericum *Ascyron* dictum, caule quadrangulo. *Bauh. Ager*
3. p. 382.

H. vulgare minus, caule quadrangulo, foliis non perfoliatis
Bauh. pin. 279.

Ascyron. Dod. pempt. 78.

Habitat in Europæ pratis. 24

Folia absque poris pellucidis esse perhibet Linnæus in *dis*
de hyper. p. 5. quod vero alii negant. R.

22. **HYPERICUM** *perforatum*. H. floribus trigynis, caule
ancipiti, foliis obtusis pellucido-punctatis. *succ. parif. pal.*
herb. fil. Blackw. t. 15. Ludw. ed. t. 11. Kniph. cent. 30
n. 46. Regn. bot. lugd. lithu. burg.

H. caule terete, alato, ramosissimo, foliis ovatis perforatis
Hall. helu. n. 1037.

H. (*officinarum*) calyce integro, caule ancipiti ramosissimo,
foliis confertis, pellucido-punctatis. *Crantz. austr.*

H. floribus trigynis; petalis uno latere crenatis, caule an-
cipiti. *Scop. carn.*

H. vulgare. *Bauh. pin. 279. Neck. gallob. p. 319.*

Hypericum. Dod. pempt. 76.

Habitat in Europæ pratis. 24

Antheræ puncto atro. Stigmata sanguinea.

23. **HYPERICUM** *humifusum*. H. floribus trigynis axilla-
ribus solitariis, caulibus ancipitibus prostratis filiformibus,
foliis glabris. *Fl. succ. gallob. pal. fl. Flor. dan. t. 141*
delph. lugd. burg. lithu.

H. foliis ovatis, caulibus supinis. *Guss. stamp. Dalib. parif.*

276 POLYADELPHIA POLYANDRIA.

H. caule prostrato, foliis ovatis, calycibus ferratis punctatis. Hall. helv. n. 1039.

H. minimum supinum. Bauh. hist. 3. p. 384. R.

H. minus supinum f. supinum glabrum. Bauh. pin. 279.

H. supinum III. minimum. Clus. hist. 2. p. 181.

Habitat in Europa australiore. 24

Affine nimium antecedenti, sed flores axillares, solitarii. Folia absque punctis pellucidis.

24. *HYPERICUM crispum. H. florib. trigynis, caule tereti, foliis sessilibus lanceolatis basi undulato-dentatis.*

H. triquetrifolium. Turr. fars. 12.

H. crispum, triquetro & cuspidato folio. Bocc. mus. 2. p. 31. t. 12.

H. folio brevior. Bauh. pin. 279.

H. foliis parvis crispis f. sinuatis siculum. Rai. hist. 1019.

Habitat in Calabria, Sicilia, Græcia. 24

Folia minutissima, semiamplexicaulia, basi trippa.

25. *HYPERICUM perfoliatum. H. floribus trigynis, caule subanticipiti, foliis amplexicaulibus ovatis, cyma floribus sessilibus.*

H. panicula subdichotoma. Diff. de hyper. p. 5.

H. (fallax) floribus trigynis, foliis ovato-oblongis, rugosis, calycisque laciniis ovatis obtusis, caule subquadrangulati. Grim. flor. isen. in Nou. Ad. A. N. C. vol. 3. app. 362. R.

H. (dubium) floribus trigynis, caule subquadrato herbaceo, foliis ovatis, calycibus obrusissimis. Leers herbarn. n. 594. R.

Habitat in Germania. 24

Cl. Grimii & Leersii nomina unam eandemque plantam indicant, an vero sit H. perfoliatum Linn. ob brevem Linnae denominationem haud facile dictu. Leersius flores subcorymbosos pedunculatos dicit. Cl. Grimm. pro mera H. perforati varietate suum Hypericum habet. Nou. Ad. A. N. C. Tom. app. p. 130. R.

26. *HYPERICUM elodes. H. floribus trigynis, caule tereti repente foliisque villosis subrotundis. gallob.*

Ascyrum supinum villosum palustre. Bauh. pin. 280.

Ascyrum supinum elodes. Clus. hist. app. alt.

Caryophyllus palustris foliis subrotundis incanis, floribus aureis. Mentz. pug. 1. 7. f. 3. an.

Habitat in Sibiria, Anglia, Gallia paludosis. 24

HYADELPHIA POLYANDRIA 277

Hypericum *simul* *H. montanum*.

27. **HYPERICUM** *barbatum*. *H.* floribus trigynis, caule tereti, foliis calycinis barbatis. *Lin. diff. de hyper.* p. 6. *Jacq. austr.* 1. 259.

Habitat in Austria. 24

* *β. Trigyna*; calycibus *Bracteis*que serrato-glandulosis.

28. **HYPERICUM** *montanum*. *H.* floribus trigynis, calycibus serrato-glandulosis, caule tereti erecto, foliis ovatis glabris. *succ. Fl. dan.* 1. 173. *pal. herb. fil. lithu. lugd. delph.*

H. foliis ovatis, ora punctata, calyceibus lanceolatis, serratis, glanduligeris. *Hall. helv. n.* 1042.

H. calycibus serrato-glandulosis, caule tereti erecto, foliis glabris punctatis sessilibus. *Crantz. austr.*

H. elegantissimum non ramosum, folio lato. *Bank. hist.* 3. p. 383.

H. caule simplici elegans latifolium. *Hill. plant.* 557. t. 14.

Androsæmum campoclarense. *Col. ecphr.* 1. p. 73. t. 74.

Afigrum *L. Hypericum* bifolium glabrum non perforatum. *Bank. pin.* 280.

Ascyrum. *Fuchs. hist.* 74.

Habitat in Europæ montibus. 24.

Bractæ etiam in summo caule glanduloso - ciliatæ, & parvæ, quod non in *H. hirsuto*.

29. **HYPERICUM** *hirsutum*. *H.* floribus trigynis, calycibus serrato glandulosis, caule tereti erecto, foliis ovatis subpubescentibus. *herb. pal. Kniph. cent.* 8. n. 52. *delph. burg. lugd.*

H. foliis ovatis hirsutis, calycibus lanceolatis, serratis, glanduligeris. *Hall. helv. n.* 1043.

H. (*villosum*) calyce serrato glanduloso, foliis pedicellatis, perforatis, caule tereti villosa. *Crantz. austr.*

H. floribus trigynis; calycibus serrato-glandulosis; foliis petiolatis, ovato-lanceolatis; caule tereti. *Scop. carn.*

H. foliis lanceolatis, calycinis laciniis linearibus. *Guett. stamp.*

H. floribus trigynis, calicum serraturis capitatis, caule tereti. *Fl. suec. paris.*

β. H. foliis ovato-oblongis brevi pediculo donatis calycinis laciniis oblongo-acutis. *Guett. stamp.*

Androsæmum *Ascyron* dictum, caule rotundo hirsuto. *Morif. hist.* 2. p. 171. f. 5. t. 6. f. 11.

278 POLYADELPHIA: POLYANDRIA

Androsæmum hirsutum. Bauh. pin. 280. caute.

Hypericum Androsæmum dictum. Bauh. hist. 3. p. 382.

Androsæmum alterum hirsutum. Col. ecphr. 1. p. 745.

Habitat in Europæ collibus & montibus. 24

Flores noctu clauduntur, quod non in H. perforato.

30. HYPERICUM tomentosum. H. floribus trigynis, calycibus serrato-glandulosis, foliis semiamplexicaulibus flexuosis tomentosis, caulibus prostratis.

H. supinum tomentosum minus & monspeliacum. Bauh. pin. 279. Sauv. monsp.

H. supinum tomentosum alterum. Clus. hist. 2. p. 181.

H. supinum tomentosum majus & hispanicum. Bauh. pin. 279.

H. supinum tomentosum hispanicum. Clus. hist. 2. p. 181.

Habitat in G. Narbonensi, Hispania. 24

Racemi terminales bifidi. Folia sessilia. Caulis teres. Stamina phalangibus tribus.

32. HYPERICUM pulchrum. H. floribus trigynis, calycibus serrato-glandulosis, caule tereti, foliis amplexicaulibus cordatis glabris. *Qed. dan. t. 75. pal. delph. burg. lugd.*

H. foliis amplexicaulibus cordatis, calycibus ovatis, serratis glanduligeris. Hall. helv. n. 1041.

H. (elegantissimum) calyce serrato, foliis perfoliatis, margine punctatis, caule tereti glabro. Crantz. austr.

H. foliis cordatis connatis, laciniis calycinis ovatis obtusis. Guett. stamp.

H. calyce serrato, foliis connatis. Hall. gætt. 116.

H. minus erectum. Bauh. pin. 279.

H. minus glabrum erectum pulchrum. Moris. hist. 2. p. 470.

H. pulchrum tragi. Bauh. hist. 3. p. 383.

Habitat in Europa australiore. 4

33. HYPERICUM nummularium. H. floribus trigynis, calycibus serrato-glandulosis, foliis cordato-orbiculatis glabris. *delph.*

H. nummulariæ folio. Bauh. pin. 279. prodr. 130.

H. fere orbiculato folio, floribus amplis pallide luteis petalis in ambitu crenatis. Pluk. alm. 188. t. 93. f. 4.

Androsæmum supinum saxatile, nummulariæ folio averfa parte rubente. Bocc. mus. 2. p. 134. t. 91.

Habitat in Pyrenæis. 24

34. HYPERICUM Coris. H. floribus trigynis: calycibus serrato-glandulosis, foliis subverticillatis. *lugd.*

POLYADELPHIA POLYAD

. foliis quaternis, linearibus, calyce seu glanduligeris. *Hall. helv. n. 1040.*

*N. f. Coris legitima ericæ similis. Moris. hist. 2
1. 6. f. 4.*

Hypericoides Coris quorundam. Bauh. hist. 3. p. 384.

Coris lutea. Bauh. pin. 280.

Habitas in Europæ australis collibus siccis, & in Oriente.

Supulæ utrinque solitariae s. ad singulum folium duæ, magnitudine & figura foliorum, hinc, quum folia sint opposita, quasi sena articulos cingunt, quorum quatuor sunt. Supulæ.

CLASSIS XIX.

SYNGENESIA.

POLYGAMIA ÆQUALIS.

* *Semiflosculosi Tournef. corollis ligulatis omnibus,*

1001. SCOLYMUS.

Recept. paleaceum. Pappus nullus. Cal. imbricatus, spinosus.

1000. CICHORIUM

Recept. subpaleaceum. Pappus sub 5. dentatus. Cal. calyculatus.

999. CATANANCHE.

Recept. paleaceum. Pappus 5. aristatus, sessilis. Cal. imbricatus, scariosus.

996. SERIOLA.

Recept. paleaceum. Pappus subpilosus. Cal. simplex.

997. HYPOCHÆRIS.

Recept. paleaceum. Pappus subplumosus. Cal. imbricatus.

983. GEROPON.

Recept. paleaceum. Pappus plumosus disci. 5. aristatus radii. Cal. simplex.

994. ANDRYALA.

Recept. villosum. Pappus pilosus sessilis. Cal. subæqualis, rotundatus.

984. TRAGOPOGON.

Recept. nudum. Pappus plumosus, stipitatus. Cal. simplex.

986. PICRIS.

Recept. nudum. Pappus plumosus, stipitatus. Cal. calyculatus.

991. LEONTODON.

Recept. nudum
plumosus
Cal. imbricatus
laxis.

985. SCORZONERA.

Recept. nudum
plumosus, et
Cal. imbricatus marg
scarioso.

993. CREPIS.

Recept. nudum. Pappus
pilosus. *Cal. caliculatus*
squamis difformibus.

989. CHONDRILLA.

Recept. nudum Pappus
pilosus, stipitatus. *Cal.*
calicul. multiflorus.

990. PRENANTHES.

Recept. nudum. Pappus
pilosus. *Cal. calicul.*
sub 5. florus.

988. LACTUCA.

Recept. nudum. Pappus
pilosus, stipitatus. *Cal.*
imbricat. margine sca-
rioso.

992. HIERACIUM.

Recept. nudum. Pappus
pilosus, sessilis. *Cal.*
imbric. ovatus.

987. SONCHUS.

Recept. nudum. Pappus
pilosus, sessilis. *Cal.*
imbric. gibbus.

998. LAPSANA.

Recept. nudum. Pappus
nullus. *Cal. calycula-*
tus.

995. HYOSERIS.

Recept. nudum. Pappus
calyculo coronatus. *Cal.*
subaequalis.

* * Capitati.

1009. ATRACTYLIS.

Cor. radiata.

1008. CARLINA.

Cal. radiatus : radiis co-
loratis.

1005. CNICUS.

Cal. bracteis obvallatus.

1002. ARCTIUM.

Cal. squamis apice inae-
vato-hamosis.

- | | |
|-------------------------|--|
| 1010. CARTHAMUS. | <i>Cal.</i> squamis squarrosis foliaceis. |
| 1007. CYNARA. | <i>Cal.</i> squamis squarrosis canaliculatis, spinosis. |
| 1004. CARDUUS. | <i>Cal.</i> squamis spinosis ventricosus. |
| 1000. ONOPORDON. | <i>Cal.</i> squamis ventricosis spinosis. <i>Recept.</i> savorum. |
| 1003. SERRATULA. | <i>Cal.</i> squamis acutiusculis muticis imbricatus, subcylindricus. |

*** *Discoides.*

- | | |
|--------------------------|--|
| 1013. CACALIA. | <i>Recept.</i> nudum. <i>Pappus</i> pilosus. <i>Cal.</i> calyculatus. |
| 1019. CHRYSOCOMA. | <i>Recept.</i> nudum. <i>Pappus</i> pilosus. <i>Cal.</i> imbricatus. <i>Pist.</i> brevissima. |
| 1015. EUPATORIUM. | <i>Recept.</i> nudum. <i>Pappus</i> plumosus. <i>Cal.</i> imbricatus. <i>Pistil.</i> longissima. |
| 1012. BIDENS. | <i>Recept.</i> paleaceum. <i>Papp.</i> aristatus. <i>Cal.</i> imbricatus. |
| 1018. STÆHELINA. | <i>Recept.</i> paleaceum. <i>Papp.</i> plumosus, ramosus. <i>Antheræ</i> candatae. |

* *Tapacetum.*

POLYGAMIA SUPERFLUA.

* *Discoides.*

- | | |
|-------------------------|--|
| 1025. ARTEMISIA. | <i>Recept.</i> subnudum. <i>Papp.</i> nullas. <i>Cor.</i> radii nulli. |
|-------------------------|--|

1013. CARPESIUM.	<i>Recept.</i> i radat. fidz.
1014. TANACETUM.	<i>Recept.</i> . submersum. — r. 3. fidz.
1015. COTULA.	<i>Recept.</i> subnudum, marginatus. Cor 4. fidz.
1019. BACCHARIS.	<i>Recept.</i> nudum. <i>Pappus</i> pilosus. Cor. femineæ hermaphroditis mixtæ.
1020. CONYZA.	<i>Recept.</i> nudum. <i>Pappus</i> pilosus. Cor. radii 3. fidz.
1026. GNAPHALIUM.	<i>Recept.</i> nudum. <i>Pappus</i> plumosus. <i>Cal.</i> scario- sus: squam. concavis.
1027. XERANTHEMUM.	<i>Recept.</i> paleaceum m. <i>Papp.</i> subsetaceus. <i>Cal.</i> scario- sus radio explanato.
1031. ANACYCLUS.	<i>Recept.</i> paleaceum. <i>Pap.</i> nullus. <i>Sem.</i> marginata, emarginata.

**** Semistofculi, subbilabiani.**

***** Radiati.**

1042. BELLIS.	<i>Recept.</i> nudum. <i>Pappus</i> nullus. <i>Cal.</i> squamis æqualibus simplex.
1049. MATRICARIA.	<i>Recept.</i> nudum. <i>Pappus</i> nullus. <i>Cal.</i> squamis imbricatus acutis.
1048. CHRYSANTHEMUM.	<i>Recept.</i> nudum. <i>Pappus</i> nullus. <i>Cal.</i> squamis in- timis scariosis.
1039. DORONICUM.	<i>Recept.</i> nudum. <i>Pappus</i> pilosus. <i>Pappus</i> radii nullus.
1038. ARNICA.	<i>Recept.</i> nudum. <i>Pappus</i> pilosus. <i>Stamina</i> radii castrata.

1037. INULA.	<i>Recept. nudum. Pappus pilosus. Anthera basi bifida.</i>
1031. ERIGERON.	<i>Recept. nudum. Pappus pilosus. Cor. radii capillares.</i>
1035. SOLIDAGO.	<i>Recept. nudum. Pappus pilosus. Cor. radii subleni, remoti.</i>
1036. CINEBARIA.	<i>Recept. nudum. Pappus pilosus. Cal. æqualis simplex.</i>
1033. SENECIO.	<i>Recept. nudum. Pappus pilosus. Cal. squamis apice sphacelatis.</i>
1032. TUSSILAGO.	<i>Recept. nudum. Pappus pilosus. Cal. squamis submembranaceis.</i>
1034. ASTER.	<i>Recept. nudum. Pappus pilosus. Cal. subsquarrosus.</i>
1044. TAGETES.	<i>Recept. nudum. Pappus aristatus. Cal. 1. phyllus. Radius 5 - florus.</i>
1041. HELENIUM.	<i>Recept. seminudum. Pappus 5. aristatus. Cal. multipartitus. Cor. radiis 3. fidis.</i>
1057. SIEGESBECKIA.	<i>Recept. paleaceum. Pappus nullus. Radius dimidiatus.</i>
1052. ANTHEMIS.	<i>Recept. paleaceum. Pappus nullus. Cal. hemisphaericus.</i>
1053. ACHILLEA.	<i>Recept. paleaceum. Pappus nullus. Radius 5. florus. Cal. oblongus.</i>
1059. BUPHTHALMEUM.	<i>Recept. paleaceum. Pappus marginat. Sigma hermaphroditi simplex.</i>
1058. VERBESINA.	<i>Recept. paleaceum. Pappus aristatus. Fasciculi radii circiter 5.</i>

1046. ZINNIA.

Recept. pubescens. Papp.
aristatus. Rad. 55 filis.
pilosus. Cal. imbricatus.
caulis.

POLYGAMIA FRUSTRANEA.

1066. CENTAUREA.

Recept. totum. Papp.
pilosus. Rad. coroll.
tubulosus.

1068. RUDBECKIA.

Recept. pubescens. Papp.
marginatus. Cal. fere
duplex.

1069. COREOPSIS.

Recept. pubescens. Papp.
aristatus. Cal. calycis
latus.

1069. MELANTHUS.

Recept. pubescens. Papp.
aristatus. Cal. squarrosus.

POLYGAMIA NECESSARIA.

1079. FILAGO.

Recept. nudum. Pappus
nullus. Flasc. sem. in-
ter squamis calycis.

1080. MICROPS.

Recept. nudum. Pappus
nullus. Flasc. sem. squa-
mis calycis vaginatis.

1076. OTHONNA.

Recept. nudum. Pappus
pilosus. Cal. monophyl-
lus.

1073. CALENDULA.

Recept. nudum. Pappus
nullus. Sem. membra-
nacea.

1074. ARCTOTIS.

Recept. subpilosum. Papp.
s. phyllis. Sem. co-
mposita.

POLYGAMIA SEGREGATA.

1084. ECHINOPS.

Perianthium i. florum
Papp. pubescens. Pe-
lyg. aequalis.

MONOGAMIA

1090. JASIONE.

Cal. communis. *Cor.* 5.
petala regul. *Caps.* in-
fera, 2. locularis.

1091. LOBELIA.

Cal. 5. dentatus. *Cor.* 2.
petala irrenul. *Caps.* in-
fera. 2. locularis.

1092. VIOLA.

Cal. 5. phyllus. *Cor.* 5.
petala irregularis. *Caps.*
supera, 3. valvis.

1093. IMPATIENS.

Cal. 2. phyllus. *Cor.* 5.
petala irregularis. *Caps.*
supera, 3. valvis.

SYNGENE

POLYGAMIA ÆQUA

983. GEROPOGON.

Receptaculum setoso-paleaceum. *Cal.* simplex. *Semina* disci pappo plumoso : radii 5. aristato.

1. GEROPOGON *glabrum*. G. foliis glabris. *Jacq. hort.* t. 33.

Tragopogon, gramineo folio, glabrum, flore dilute incarnato. *Rai. suppl.* 149.

Tragopogon calycibus corollæ radio longioribus, foliis integris, seminibus lævibus : disci plumosi ; radiis seraceis. *Hort. ups.* *.

Habitat in Italia. ☉

2. GEROPOGON *hirsutum*. G. foliis pilosis.

Tragopogon gramineis foliis hirsutis. *Bauh. pin.* 275.

Tragopogon gramineo folio, suave rubente flore. *Col. æphr.* 1. p. 232. t. 231.

Habitat in Italia. ☉

3. GEROPOGON *calyculatum*. G. calycibus calyculatis. *Jacq. hort.* t. 106.

Habitat.

Radix perennis. Caules plures, ramosi. Flores terminales, ante anthesin nutantes. Calyx calyculatus, flosculis numerosis.

984. TRAGOPOGON.

Receptaculum nudum. *Cal.* simplex Pappus plumosus.

* *Caulescentia.*

1. TRAGOPOGON *pratense*. Tragopogon calycibus corollæ radium æquantibus, foliis integris strictis. *Carr.*

288 SYNGENESIA. POLYGAMIA ÆQUALIS.

*gallob. pal. fil Ludw. ed. t. 49. Knorr. del. 2. t. 1. T. 35.
Kniph. cent. 9. n. 91. lugd. delph. burg. lithua.*

T. calycibus florem superantibus. Fl. suéc. paris.

T. foliis gramineis carinatis, amplexicaulibus, caudatis, subcrispis, calyce flori æquali Hall. helv. n. 8.

T. pratense luteum majus. Bauh. pin. 274.

Tragopogon Fuchf. hist. 827.

Habitat in Europæ pratis apricis. ♂

Corolla calyce sæpe brevior. R.

3. TRAGOPOGON porrifolium. *T. calycibus corollæ radio longioribus, foliis integris strictis, pedunculis superne incrassatis. Kniph. cent. 7. n. 93. delph. burg. lugd.*

T. foliis gramineis amplexicaulibus, calyce florem superante. Hall. helv. n. 9.

T. purpureo cæruleum, porri folio, quod Artefij vulgo Bauh. pin. 274. Moris. hist. 3. p. 8. f. 7. t. 9. f. 5.

T. alterum s. Barba hirci. Dalech. hist. 1079.

Barba hirci altera. Cam. epis. 313.

β. Tragopogon folio oblongo sinuato. Bauh. pin. 274.

Habitat in Helvetia. ♂

4. TRAGOPOGON crocifolium. *T. calycibus corollæ radio longioribus, foliis integris, radicalibus pedunculisque basi villosis. monsp. delph.*

T. purpureo-cæruleum crocifolium. Bauh. pin. 275.

T. croci folio montanum, flore nigro-purpureo. Col. ecphr. I. p. 229. t. 230.

Habitat in Italia, Monspeli. ♂

Calyces pentaphylli. Similis præcedenti, vix pedalis (nec 6-pedalis,) tenera. Folia basi villosa, ætate glabra. Flos violaceus ex seriebus tantum duabus, sed disco magis luteus.

γ. TRAGOPOGON villosum. *T. calycibus corollæ radio sesquilongioribus, caule foliisque villosis.*

T. caule tomentoso. Hall. gatt. 418.

Habitat in Hispania, Sibiria. ♂

Magnitudo T. porrifolii, sed tota planta villis albis pubescens, imprimis caulis, & folia subtus magis copiosa. Pedunculi cylindrici. Calyx enneaphyllus, corolla fere sesquilongior, non imbricatus. Corolla magis reliquis pallide flava radiis 18. Anthæræ fuscescentes. Planta ex omnibus foliorum aliis secundum totam longitudinem ramos demum emittit floridos, unde corymbosa evadit. Flos expansus subnutans est.

SYNGENESIA. POLYGAMIA ÆQUALIS. 289

6. **TRAGOPOGON** *Dalechampii*. T. calycibus monophyllis corolla brevioribus inermibus; foliis runcinatis. *Monsp. delph.*

Hieracium asperum; flore magno dentis leonis. *Bauh. pin.* 127.

Hieracium magnum. *Dalech. hist.* 569.

Hieracium sulphureum, incisus foliis, montanum. *Barr. rar.* 1043. t. 209.

Hedypnois monspessulana. *Bauh. hist.* 2. p. 1036.

Habitat in Hispania, G. Narbonensi. 2.

Caulis brevissimus. Folia tomentoso-scabra; integerrima nec denticulata. Corolla subtus purpurea: flosculis luteis apice s. dentibus purpurascens.

7. **TRAGOPOGON** *picroides*. T. calycibus monophyllis corolla brevioribus aculeatis; foliis runcinatis denticulatis. *Monsp. delph.*

Picris calycibus simplicibus aculeatis, foliis inermibus hastato-sinuatis. *Hort. ups.*

Sonchus asper laciniatus creticus. *Bauh. pin.* 124. *prodr.* 60.

Hieracium majus, folio sonchi, semine incurvo. *Bauh. pin.* 127.

Chondrilla cretica nomine missa, semine crispo. *Bauh. hist.* 2. p. 1022.

Habitat in Creta, *Monspeli.* ☉

8. **TRAGOPOGON** *asperum*. T. calycibus corolla brevioribus hispida, foliis integris: caulinis oblongis.

Sonchus asper, subrotundo folio, major & minor. *Bauh. pin.* 124. *prodr.* 60.

Habitat Monspeli.

985. SCORZONERA.

Receptaculum nudum. Pappus plumosus. Calyx imbricatus squamis margine scariosis.

2. **SCORZONERA** *humilis*. S. caule subnudo unifloro; foliis lato-lanceolatis nervosis planis. *Suec. ingr. carn.* *Jacq. austr.* t. 36. *tub. pal. sil. delph. lithua. burg.*

S. caule nudo unifloro, foliis lanceolatis. *Hall. helv. n.* 10.

S. *humilis latifolia nervosa*. *Bauh. pin.* 275.

S. *humilis latifolia pannonica*. *Clus. hist.* 2. p. 138.

Habitat in Europæ septentrionali.

3. **SCORZONERA** *hispanica*. 1

Linnaei spec. Plant. Europ.

2. 2. folia

290 SYNGENESIA. POLYGAMIA ÆQUALIS.

amplexicaulibus integris serrulatis. *Carn. delph. Blackw.*
t. 406. *Kniph. cent.* 7. n. 83.

S. latifolia sinuata. Bauh. pin. 275.

S. major hispanica. I. Clus. hist. 2. p. 137.

Habitat in Hispania, Sibiria. 24

4. *SCORZONERA graminifolia. S. foliis lineari-ensiformibus integris carinatis. Jacq. obs.* 4. p. 13. t. 100. *delph.*

S. lusitanica gramineo folio, flore pallide luteo. T. Buxb. cent. 2. p. 26. t. 21.

S. caule ramoso, foliis linearibus acuminatis carinatis, calycibus acutis. Gmel. sibir. 2 p. 4.

Habitat in Lusitania, Sibiria.

5. *SCORZONERA purpurea. S. foliis lineari-subulatis integris canaliculato-triquetris, pedunculis cylindricis. Jacq. austr.* t. 35. *carn.* t. 46.

S. caule ramoso tereti, foliis linearibus acuminatis, calycibus obtusiusculis. Gmel. sib. 2. p. 7. t. 2.

S. angustifolia subcærulea. Bauh. pin. 275.

S. sylvestris, foliis angustis, flore cæruleo. Mentz. pug. t. 1. f. 1.

Habitat in Marchia Brandenburgica, Austria, Sibiria.

Corolla cæruleo-carnea. Antheræ albæ. Mygind.

6. *SCORZONERA angustifolia. S. foliis subulatis integris, pedunculo incrassato, caule basi villosa. Vind. monsp. carn. burg.*

S. caule simplici unifloro, foliis lineari-lanceolatis. Gmel. sib. 2. p. 2. t. 1.

S. caule nudo unifloro, foliis linearibus. Hall. helv. n. 11.

S. volkameriana. Fehr. p. 37. c. *icone. Hall. R.*

S. angustifolia prima. Bauh. pin. 275.

S. humilis angustifolia pannonica III. Clus. hist. 2. p. 137. t. 138.

Tragopogon pinifolium hispanicum. Barr. ic. 496.

Habitat in Hispaniæ, Monspeli, Austriæ collibus saxosis.

Caulis vix spithameus, basi ramosus, villosus. Fila, conferta, linearia, subulata, longitudine fere caulis. Pedunculi terminales, versus florem crassiores ipso caule. Flos magnus. Corolla lutea subtus purpurascens, longitudine calycis.

Synonymon Barrelieri non huc quadrat ob caulem ab imo ad summum ramosum. Hall. R.

7. *SCORZONERA hirsuta. S. foliis linearibus cauleque unifloro hirsutis.*

SYNGENESIA. POLYGAMIA ÆQUALIS. 192

Tragopogon calycibus corolla brevioribus, foliis integris strictis ciliatis, caule hitto, pedunculis superne incrassatis. *Gerard. prou.*

Tragopogon hirsutum. *Bauh. pin. 274.*

Tragopogon alter luteus. *Col. ecphr. 1. p. 234. t. 233.*

Habitat in Apulia.

Folia canaliculata. Semina hirsuta, alba, papposa. H. V.

8. SCORZONERA resedifolia. *S. foliis obtuse dentatis, caule divaricato, calycum apicibus tomentosis. delph.*

S. caule diffuso lanato, foliis hastato-pinnatifidis, caulinis linearibus integerrimis, calycibus tomentosis. Gouan. illustr.

S. laciniatis foliis, supina. Barr. rar. 1050. t. 800.

Chondrilla sicula tragopogonoides maritima. *Bocc. sic. 130. t. 7. f. A.*

Habitat in Hispania, Monspelii. 2

Caulis divaricatus. Folia Resedæ s. Coronopi dentibus obtusis.

9. SCORZONERA laciniata. *S. foliis linearibus dentatis acutis, caule erecto, calycum squamis patulo mucronatis. Par. monsp pal. Jacq. austr. t. 356. delph. burg.*

S. caule folioso, ramoso, foliis angustissimis, imis rari-ter semipinnatis, superioribus linearibus lanceolatis. Hall. helu. n. 12.

S. foliis laciniatis. Tournesf. inst. 477. Sauv. monsp.

Tragopogon laciniatum luteum. *Bauh. pin. 274. Rai. hist. 251.*

Habitat in Germania, Gallia. ♂

Caulis erectus. Calycis squamæ infra apicem mucronatæ.

12. SCORZONERA picroides. *S. foliis superioribus amplexicaulibus integerrimis, inferioribus runcinatis, pedunculis squamosis. Delph.*

Crepis foliis radicalibus ovato-oblongis denticulatis, caulinis lanceolatis amplexicaulibus, pedunculis unifloris squamosis. Roy. lugdb.

Sonchus subrotundo folio nostras. *Pluk. phyt. 61. f. 5.*

Sonchus lævis angustifolius. *Bauh. pin. 124?*

Crepis dalechampii. *Dalech. hist. 562. **

Habitat Monspelii. ☉

Semina incurva crenulata: Pappo simplici sessili; erga media inter Scorzoneram & Sonchum.

986. P I C R I S.

*Recept. nudum. Cal. calyculatus. Pappus plumosus.
Sem. tranversim fulcata.*

1. **PICRIS** *echioides*. P. perianthiis exterioribus pentaphyllis interiore aristato majoribus. *Galloba burg.*

P. calycibus simplicibus perianthio maximo obvallatis, *Dalib. paris. Gouan. monsp.*

Hieracium echioides capitulis cardui benedicti. Bauh. pin. 128.

Buglossum echioides luteum hieracio cognatum. Lob. ic. 577.

β. *Hieracium capitulis cardui benedicti. Herm. par. 185, t. 185.*

Habitat in Angliæ, Galliæ, Italiæ sylvis carduis, aggeribus. ☉

Folia indivisa, amplexicaulia. Calyx Exterior florum pentaphyllus, interiore major: foliolis subcordatis, spinosis. Interior octangulatus: foliolis totidem carinatis, infra apicem aristatis; his exterius incumbunt foliola totidem minuta, lævia.

2. **PICRIS** *hieracioides*. Perianthiis laxis foliis integris, pedunculis squamatis in calycem. *Fl. suec. paris. monsp. pal. carn. gallob. sil. delph. lugd. burg.*

Picris aspera, foliis oblongis dentatis, superioribus amplexicaulibus. Hall. helv. n. 24.

Cichorium pratense luteum hirsutie asperum. Bauh. pin. 126.

Hieracium asperum, majore flore, in agrorum limitibus. Bauh. hist. 2. p. 1029.

Habitat in Germaniæ, Angliæ, Belgii, Galliæ versuris agrorum. 24

Folia lanceolata. Pedunculi in calycem usque adspersi squamis subulatis, donec calyx fere imbricatus sit. Pappus vix plumosus.

987 S O N C H U S.

*Receptacul. nudum. Calyx imbricatus, ventricosus.
Pappus pilosus.*

1. **SONCHUS** *maritimus*. S. pedunculo nudo foliis lan-

SYNGENESIA POLYGAMIA ÆQUALIS. 293

ceolatis amplexicaulibus indivisis retrorsum argute dentatis.

S. angustifolius. *Gallob.*

S. angustifolius maritimus. *Bauh. pin. 124. prodr. 61. Pluk. alm. 354. t. 62. f. 5. All. nicæns. 84.*

Chondrilla palustris longifolia sinuata læviter spinosa. *Rai. suppl. 137.*

Habitat in Europa australi. 24.

2. SONCHUS palustris. *S. pedunculis calycibusque hispidis subumbellatis, foliis runcinatis basi sagittatis.* *Gallob. delph. lithua. burg.*

S. pedunculis calycibusque hispidis, floribus distanti-corymbosis. *Hort. ups.*

S. caule erecto, foliis pinnato-hastatis apiceque hastatis, floribus congestis villosis. *Dalib. parif.*

S. asper arborescens. *Bauh. pin. 124.*

S. lævior austriacus s. altissimus. *Clus. hist. 2. p. 147.*

Habitat in Belgii, Angliæ, Germaniæ, Galliæ, Hungariæ pratis paludosis. 24.

Facies a *S. arvensi* diversissima; Structura tamen ita eadem, ac si *S. palustris* ex *S. arvensi* olim ortus fuisset.

3. SONCHUS arvensis. *S. pedunculis calycibusque hispidis subumbellatis, foliis runcinatis basi cordatis.* *Gallob. pal. sil. Flor. dan. t. 606. lithua. burg. lugd.*

S. foliis amplexicaulibus, semipinnatis, serratis, calycibus hispidis. *Hall. helv. n. 23.*

S. pedunculis calycibusque hispidis, floribus congestis. *Fl. suec. parif.*

S. foliis lanceolato-oblongis dentatis, floribus congestis hispidis. *Hort. cliff.*

Hieracium arvense. *Scop. carn.*

Hieracium majus, folio sonchi, s. Hieracium sonchites. *Bauh. pin. 126.*

Hieracium majus. *Fuchs. hist. 319.*

Habitat in Europa agris argillosis. 24.

4. SONCHUS oleraceus. *S. pedunculis tomentosis, calycibus glabris.* *Suec. parif. ged. gallob. pal. sil. Flor. dan. t. 682. carn. lugd. lithua. burg.*

S. annuus ramosus diffusus, foliis laciniatis. *Hort. cliff.*

S. caule ramosissimo diffuso: foliis summis amplexicaulibus, pedunculis inæqualibus. *Roy. lugdb.*

Hieracium oleraceum. *Scop. carn.*

5. Sonchus lævis.

294 SYNGENESIA. POLYGAMIA ÆQUALIS.

S. foliis amplexicaulibus dentatis integris semipinnatis, calycibus levibus. *Hall. helu. n. 21. Blackw. t. 130.*

S. lævis laciniatus latifolius. *Bauh. pin. 124. Gmel. sib. 2. p. 9.*

S. lævis. *Cam. epit. 279.*

β. S. lævis minor, paucioribus laciniis. *Bauh. pin. 124.*

S. lævis latifolius & angustifolius. *Taber. ic. 190. 189.*

γ. Sonchus asper.

S. foliis amplexicaulibus, rigidis integris & semipinnatis, dentatis, calycibus levibus. *Hall. helu. n. 22.*

S. asper. *Blackw. t. 30.*

Sonchus asper laciniatus & non laciniatus. *Bauh. pin. 134.*

Sonchus asper laciniatus latifolius & angustifolius. *Laf. pruf. 157. t. 77. 78.*

δ. Sonchus asper non laciniatus. *Tournef. inst. 474.*

Sonchus aspera. *Fuchf. hist. 674. Dodon. 643.*

Habitat in Europæ cultis. ☉

Folia compresso-amplexicaulia. Pedunculi ætate glabri sunt.

5. *SONCHUS tenerrimus.* *S. pedunculis tomentosis, calycibus pilosis.*

S. lævis in plurimas & tenuissimas lacinias divisus. *Bauh. pin. 124. prodr. 61.*

Hieracium foliis in tenues lacinias profunde sectis flore luteo. *Pluk. phyt. 93. f.*

Chondrilla lutea. *Bauh. hist. 1. p. 120.*

Habitat Monspeli, Florentiæ. ☉

Caulis ramosissimus: pilis sparsis & glutiniferis. Pedicelli albo tomento vestiti. Calyces cylindrici: pilis patentibus adspersi. Folia glabra, tenera, pinnatifida, inermia.

6. *SONCHUS Plumieri.* *S. pedunculis nudis, floribus paniculatis, foliis runcinatis.*

S. caule glaberrimo, foliis runcinatis glabris, racemo laxo glaberrimo, pedunculis nudis. *Gouan. illustr.*

Lactuca alpina glabra, acanthi folio, flore magno ceruleo. *Vaill. act. 1721. p. 200. Monnier. obs. 157.*

Habitat in Pyrenæis: Monte Aureo prope Carthusiam majorem. *Monnier. h*

Simillimus S. alpino, sed Corolla unica flosculorum ordine s. flosculis dimidio parioribus, at quadruplo majoribus. Caulis altitudinis humanæ. Folia bipedalia, Taraxaci runcinata, supra lævia, subtus venis scabra; Caxlinæ angustiora, acutiora. Panicula terminalis: pedunculis

SYNGENESIA. POLYGAMIA ÆQUALIS. 295
nudis. Corolla Cichorii. Calyx exsudat guttulas resinosas, minimas, sparsas. Pappus stipitatus Lactucæ.

7. SONCHUS alpinus. S. pedunculis squamosis floribus racemosis, foliis runcinatis. *Suec. monsp. norv. fil. Oed. dan. t. 182. delph.*

S. caule erecto, foliis pinnato-hastatis : apice cordato-triangularis, floribus racemosis. *Hort. cliff. **

S. foliis amplexicaulibus, petiolis, triangularibus, petiolo pinnato, floribus spicatis. *Hall. helv. n. 20.*

Hieracium cæruleum. Scop. carn.

Sonchus lapponum altissimus, floribus cæruleis. Fl. lapp.

S. lævis laciniatus cæruleus f. *Sonchus alpinus cæruleus. Bauh. pin. 124.*

S. cæruleus. *Cam. epit. 284.*

Habitat ad latera Alpium. Lapponiæ, Helvetiæ, Austriæ. ☉

Albo flore datur. R.

9. SONCHUS sibiricus S. pedunculis squamatis foliis lanceolatis indivisis sessilibus. *Fl. suec. ingr. Kniph. cent. 10. n. 82.*

S. foliis lanceolatis sessilibus plerumque denticulatis, floribus corymbosis, caulibus glabris. *Gmel. sibir. 2. p. 11. t. 3.*

Lactuca salicis folio, flore cæruleo. Amm. ruth. 211. 212.

Habitat in Sibiria, Suecia septentrionali, Finlandia. 24
Pappus simplex, sessilis. Variat albo flore.

988. LACTUCA.

Receptaculum nudum. Calyx imbricatus, cylindricus, margine membranaceo. Pappus simplex, stipitatus. Sem. lævia.

1. LACTUCA quercina. L. foliis runcinatis denticulatis, acutis : subtus lævibus, caule glabro. *Fl. suec. **

L. foliis quernis. *Rai. hist. 221.*

L. sylvestris, angusto laciniato folio. *Magn. monsp. app ?*

Habitat in Insula Carolina Baltici; in Germania. 24

2. LACTUCA sativa. L. foliis rotundatis, caulibus cordatis, caule corymboso.

L. sativa. *Bauh. pin. 122. 242. Blackw. t. 88.*

β. *Lactuca capitata. Bauh. pin. 123. Moris. hist. 3. p. 57. f. 7. t. 2. f. 2.*

γ. *Lactuca crispa. Bauh. pin. 123. Dod. pempt. 644.*

296 SYNGENESIA. POLYGAMIA ÆQUALIS.

Habitat ——— ○

Cultura mangonio forte produca ex proximis.

3. **LACTUCA Scariola.** L. foliis verticalibus carina aculeatis. *Amæn. acad.* 4. * *gallob. pal. sil. Regn. bot. delph. lithua burg.*

L. foliis semipinnatis, nervo spinoso. *Hall. helu. n.* 14.

L. sylvestris costa spinosa. *Bauh. pin.* 123.

L. sylvestris laciniata. *Moris. hist.* 3. p. 58. f. 7. t. 2. f. 17.

Endivia major lactucina spinosa. *Barr. ic.* 135. R.

Lactuca agrestis. *Cord. p.* 157. R.

β. L. sylvestris italica, costa spinosa, sanguineis maculis adspersa. *Herm. parad.* 191. t. 191.

γ. L. sylvestris annua, costa spinosa, folio integro colore cæcio. *Moris. hist.* 3. p. 58.

Habitat in Europa australi. ○

Folia radicalia sinuata. *Rai.*

In γ. Folia inferiora runcinata, obliqua s. verticalia; Caulina superiora amplexicaulia, sagittata, carina latiori margine scabro.

4. **LACTUCA virosa.** L. foliis horizontalibus carina aculeatis dentatis. *Monsp. carn. delph. lugd.*

L. caule foliisque aculeatis. *Hort. cliff. * paris. monsp.*

L. foliis ovatis patulis, nervo spinoso. *Hall. helu. n.* 15.

L. sylvestris, odore viroso. *Bauh. pin.* 123. *Moris. hist.* 3. p. 58. f. 7. t. 2. f. 16.

L. atra. *Cæsalp.* 515. *Hall.* R.

L. sylvestris lato folio, nervo viroso. *Bauh. hist.* 2. p. 1002. *Hall.* R.

Habitat in Europæ australioris aggeribus, sepibus.

Caullis inferne aculeatus. Folia sagittata, sessilia, margine & imprimis carina aculeatis. Varietas prioris ex Gerardo & Hallero.

5. **LACTUCA saligna.** L. foliis hastato-linearibus sessilibus carina aculeatis. *Monsp. carn. pal. Jacq. austr. t.* 259. *delph. burg.*

L. foliis imis semipinnatis, superioribus amplexicaulibus, hamatis, nervo spinoso. *Hall. helu. n.* 13.

L. foliis linearibus hastatis: inferioribus dentatis. *Roy. lugd.*

L. foliis imis pinnatifidis, cæteris ensiformibus basi hastatis, dorso ferratis. *Sauv. monsp.*

L. foliis radicalibus linearibus dentato pinnatis: laciniis sursum dentatis: caulinis alato-lanceolatis, floralibus linearibus sessilibus. *Guet. stamp. Dalib. paris.*

SYNGENESIA. POLYGAMIA ÆQUALIS. 297

- L. sylvestris altissima**, foliis inferioribus dentatis, superioribus acuminatis integris. *Rai. suppl.* 135.
L. sylvestris altera, angusto saligno folio: costa albicante. *Rupp. ien.* 167. *Hall. ien.* 207. t. 4. *optim.*
L. sylvestris, angusto laciniato folio., *Magn. monsp.* 298. *
Chondrilla viscosa humilis. *Bauh. pin.* 130. *prodr.* 68. *Taill.*
Endivia minor lactucina spinosa. *Barr. ic.* 136.
Habitat in Gallia, Lipsiæ.
Folia plano-verticalia, fere ut in L. scariola.
8. LACTUCA perennis. L. foliis linearibus dentato-pinnatis: laciniis sursum dentatis. *Paris. Mill. dict.* 1. 157. *monsp. ged. carn. pal. delph. burg.*
Chondrilla cærulea altera, cichorii sylvestris folio. *Bauh. pin.* 130.
Chondrilla cærulea latifolia laciniata. *Bauh. pin.* 130.
Chondrilla, *Dod. pempt.* 637. cærulea. *Tabern.* 176. *Apate. Dalech. hist.* 561.
Habitat in Germania, Helvetia, Gallia. 24

989. CHONDRILLA.

Receptaculum nudum. Cal. calyculatus. Pappus simplex, stipitatus. Flosculi multicipliserie. Sem. muricata.

- 1. CHONDRILLA juncea**. C. foliis radicalibus runcinatis, caulinis linearibus integris. *Monsp. pal. ged. sil. Hall. helu. n.* 17 *Jacq. austr.* 1. 427. *delph. burg. lugd. lithua.*
C. (juncea) foliis radicalibus eroso-pinnatifidis lyratis, caule basi aculeato, floribus spicatis. *Gouan. illustr.*
C. foliis radicalibus pinnatifidis, caulinis lanceolato-linearibus. *Wach. ultr. Bœhm. lips.*
C. juncea viscosa arvensis. *Bauh. pin.* 130.
C. juncea. *Tabern. hist.* 487.
C. viminea. *Bauh. hist.* 2. p. 1021. *
Habitat in Germania, Helvetia, Gallia ad agrorum margines inque locis glareosis.
Caulis inferne strigosus.
2. CHONDRILLA crepoides. C. foliis sagittatis amplexicaulibus, caule simplici, floribus subsessilibus lateralibus.
Habitat ——— ○

298 SYNGENESIA. POLYGAMIA ÆQUALIS.

Posset Crepidis generi associari. Caulis simplex, sæpiusque simplicissimus, sesquipedalis, basi purpureo striatus, adpersus setis albis raris. Folia (Turritidis) sagittata, amplexicaulia, indivisa, oblonga, scabra margine & imprimis carina albidis pilis; supra lævia, infima denticulata. Flores alterni pedunculo vix calycis longitudine bractea una alterave. Calyx cylindricus, striatus, adpersus tuberculis nigris seta alba. Calyx inferior brevissimus, foliolis subulatis persistentibus. Corolla lutea, subtus purpurascens, longitudine calycis.

990. P R E N A N T H E S.

Recept. nudum. Cal. calyculatus. Pappus simplex subseffilis. Flosculi simplici ferie.

1. **PRENANTHES tenuifolia.** P. foliis linearibus integerrimis. *delph.*

P. angustifolius, flore purpureo. *Vaill. ad. 1721. p. 193.*
Chondrilla angustissimo longissimo integroque folio. *Juss. ad. 1709.*

Habitat in Alpibus. Europæ australis.

2. **PRENANTHES viminea.** P. foliorum ramentis cauli adnatis. *Jacq. austr. t. 9. delph. burg. lugd.*

P. foliorum laciniis extrorsum dentatis, ramentis foliaceis cauli ramisque adnatis. *Sauv. monsp.*

Chondrilla viminea viscosa monspeliaca. *Bauh. pin 131. prodr. 68.*

Chondrilla viscosa, caule foliis obducto. *Bauh. pin. 130.*

Chondrilla saxatilis viscosa, caule inermidopha. *Col. ecphr. 1. p. 238. t. 240.*

Habitat in Austria, Gallia, Lusitania.

3. **PRENANTHES purpurea.** P. flosculis quinis, foliis lanceolatis denticulatis. *Monsp. carn. Jacq. austr. t. 317. pal. sil. delph.*

P. foliis amplexicaulibus, petiolatis, lanceolatis, serratis. *Hall. helv. n. 19.*

Lactuca montana purpureo-cærulea minor. *Bauh. pin. 123.*

Sonchus lævior pannonicus 4, purpureo flore. *Clus. hist. 3. p. 147.*

β. Lactuca montana, purpureo-cærulea, major. *Bauh. pin. 123.*

Sonchus montanus purpureus tetrapetalos. *Col. ecphr. 1. p. 245.*

SYNGENESIA. POLYGAMIA ÆQUALIS. 299

Habitat in Germaniæ, Helvetiæ, Italiæ nemoribus montanis.

Folia inferiora amplexicaulia. Pappus sessilis.

4. **PRENANTHES muralis.** P. flosculis quinis, foliis runcinatis. *Suec. parif. ged. carn. pal. sil. Fl. dan. t. 509. delph. lugd. lithua. burg.*

P. foliis serratis, pinnatis, pinna suprema triangulari, trilobata. *Hall. helu. n. 18.*

Sonchus levis laciniatus muralis, parvis floribus. Bauh. pin. 124.

Sonchus lævis vulgator. 2. Clus. hist. o. p. 136.

Sonchus lævis sylvaticus IV. Tabern. 194. Hall. R.

Habitat in Europæ nemoribus umbrosis.

Pappus simplex, stipitatus.

6. **PRENANTHES chondrioides.** P. flosculis denis, calycibus octofidis, foliis lanceolatis: radicalibus indivisis subdentatis. *Ard. specim. 2. p. 36. t. 17.*

Habitat in Europa australi. 2.

Caulis panniculatus. Folia caulina linearia, ad ramificationes tantum, minuta; radicalia, lanceolata, petiolata, nuda, minute dentata.

991. LEONTODON.

Recept. nudum. Cal. imbricatus squamis laxiusculis,

Pappus plumosus.

1. **LEONTODON Taraxacum.** L. calyce inferne reflexo, foliis runcinatis denticulatis lævibus. *Suec. lugd. burg. lithua. parif. pal. sil. Flor. dan. t. 574. Knorr. del. 1. t. K. 3. Blackw. t. 1. & 501. Regn. bot.*

Taraxacum calycibus glabris, squamis imis reflexis. Hall. helu. n. 56.

Dens leonis, latiore folio. Bauh. pin. 126.

Hedypnois. Fuchs. hist. 680.

β. *Dens leonis angustiore folio. Bauh. pin. 126. Segu. ver. 2. p. 188.*

γ. *Dens leonis latiore & rotundiore folio. Tournef. inst. 410.*

δ. *Dens leonis folio tenuissimo. Bauh. prodr. 62.*

Habitat in Europæ pascuis. 2.

Pappus stipitatus, simplex. Leers herb. n. 605. Scop. l. c. R.

300 SYNGENESIA. POLYGAMIA ÆQUALIS.

2. LEONTODON *bulbosum*. L. foliis oblongo-ovatis subdentatis glabris, calyce lævi, scapo superne hirsuto.

Taraxacum calyce toto erecto subhirsuto, radice bulbosa. *Sauv. monsp.*

Chondrilla bulbosa. *Bauh. pin.* 130.

Chondrilla altera *dioscoridis*. *Col. phys.* 11. t. 4.

Chondrilla pusilla marina lutea bulbosa. *Lob. ic.* 230.

Habitat *Monspeli*, inque Italia.

3. LEONTODON *aureum*. L. foliis runcinatis, caule subunifolio, calyce hispido. *Jacq. austr.* t. 297. *delph.*

Andryala aurea. *Scop. ann.*

Hieracium aureum. *Scop. carn.*

Taraxacum calycibus hirsutis, squamis rectis. *Hall. helv.* ed. 2. n. 57. tab. 1.

Taraxacum foliorum dentibus triangularibus sine subrotundo. *Hall. helv.* 760. t. 24. opus. 276.

Dens leonis foliis extremo subrotundis, floribus aureis. *Rai. eur.* 399.

Hieracium alterum minus IV. *Col. ecphr.* 2. p. 31. t. 29.

Habitat in Alpibus Helvetiæ, Baldi, Italiæ. *Seguier.* 24

Folia *Taraxaci*, sed minora, obtusiora. Scapus nudus striatus, in medio foliola lineari notatus. Calyx fuscus, erectus, basi hirsutus foliorum carinis ciliatis. Corolla fulva, subtus rubra. Receptaculum villosum.

4. LEONTODON *hastile*. L. scapo calyceque lævi, foliis lanceolatis dentatis integerrimis glabris. *Jacq. austr.* t. 164. *delph.*

L. danubiale. *Jacq. vind.* 270. sec. *Jacq. R.*

Picris caule unifloro, foliis glabris, semipinnatis. *Hall. helv.* n. 26. *Jacq. R.*

Dens leonis glaber, angustiore & longiore folio, flore minore. *Morif. hist.* 3. p. 75.

Taraxaconoides *chondrillæ* glabro virenti folio. *Vaill. ad.* 170.

Habitat in Europa australi. ☉.

Folia longa, pedalia dentibus alternis, latis, brevibus, glabra, vix costa hirsuta. Scapus indivisus, lævis, nudus absque squamis. Calyx glaber, sulcatus, basi tantum squamis aliquot minutis auctus, ante anthesin nutans.

5. LEONTODON *tuberosum*. L. foliis runcinatis scabris, calyce acuto hirsuto. *Gouan. monsp.*

Taraxaconoides *chondrillæ* folio hirsuto, asphodeli radice. *Vaill. ad.* 1721. p. 170.

SYNGENESIA. POLYGAMIA ÆQUALIS. 307

Dens Leonis asphodeli bulbulis. *Bauh. pin.* 126.

Chondrilla altera dioscoridis monspeliensium. *Lob. ic.* 232.

Habitat in Etruriæ, Galloprovinciæ, Narbonæ pratis. 24

Folia runcinata, obtusiuscula: laciniis lateralibus integris, nec denticulatis, sed curvatis versus basin. Scapus uniflorus, parum. Calyx vero magis hirtus.

6. **LEONTODON autumnale**. L. caule ramoso, pedunculis squamosis, foliis lanceolatis dentatis integerrimis glabris. *suec. carn. pal. sil. Flor. dan. t. 501. herb. lugd. lith. delph. burg.*

Crepis foliis longis dentatis linearibus, caule declinato nudo. *Hort. cliff.*

Picris caule nudo brachiato, foliis glabris semipinnatis. *Hall. helu. n. 28.*

Hieracium V. *Reyg. ged.*

Hieracium foliis coronopi. *Bauh. pin.* 128. R.

Hieracium nigrum. *Tabern. hist.* 181. R.

Hieracium, chondrillæ folio glabro, radice succisa, majus & minus. *Bauh. pin.* 127.

Hieracium minus. *Fuchf. hist.* 320.

Habitat in Europæ pratis, pascuis, autumn. 24

Pappus substipitatus plumosus.

Pappus sessilis, centralium substipitatus. *Leers. l. c. R.*

7. **LEONTODON hispidum**. L. calyce toto erecto, foliis dentatis integerrimis hispidis: pilis furcatis. *suec. ingr. Jacq. vind. carn. pal. sil. gallob. lith. delph. lugd. burg.*

Picris caule nudo, unifloro, foliis asperis, dentatis. *Hall. helu. n. 25.*

Picris calyce erecto, foliis hispidis dentatis: dentibus integerrimis. *Sauv. monsp.*

β. **Hieracium** foliis lanceolato-linearibus integerrimis villosis. *Ger. prou.*

Hieracium foliis integris subdenticulatis lanceolatis scabris, scapo unifloro. *Sp. pl.* 1.

Hieracium montanum angustifolium nonnihil incanum. *Bauh. pin.* 129.

Hieracium VI. montanum. *Clus. hist.* 2. p. 141. *Bauh. hist.* 3. p. 1038.

γ. **Hieracium** foliis oblongo-dentatis pubescentibus, scapo unifloro. *Ger. prou.*

Hieracium foliis dentatis viscidis hirsutis subasperis, scapo nudo unifloro, calyce hirt. *Sauv. monsp.* 298. *Gouan. monsp.*

302 SYNGENESIA. POLYGAMIA ÆQUALIS.

Hieracium montanum, dentis leonis folio, f. laciniatum & lanuginoso folio. *Bauh. pin.* 128.

Hieracium folio Hedypnoidis. *Clus. hist.* 2. p. 142.

Habitat in Europæ borealis pratis. 24

Bractea linearis infra florem, quod non in *L. hirt.* Pappus sessilis, plumosus. Planta vexata ob synonyma aliter ab aliis assumpta.

Semen ex stylo pappigerum. *Hall. l. c.* R.

Var. *β.* ad *Hieracium incanum* alii referunt. R.

8. **LEONTODON hirtum.** L. calyce toto erecto, foliis dentatis hirtis: setis simplicissimis. *carn. gallab. herb. delph.*

L. foliis cauleque hispidis, pedunculo incrassato striato. *Ger. prou.*

Crepis foliis lanceolatis dentato sinuatis hispidis, setis subulatis, scapo unifloro. *Sp. pl.* 1.

Rhagadiolus foliis semipinnatis asperrimis. *Hall. helu. n.* 7.

Hieracium, dentis leonis folio, hirsutie asperum magis laciniatum. *Bauh. pin.* 127.

Hieracium, dentis leonis folio, hirsutie asperum minus laciniatum. *Bauh. pin.* 127. *prodr.* 63.

Hieracium parvum hirtum, caule aphylo. crispum ubi siccatum. *Bauh. hist.* 2. p. 1038.

Dens leonis foliis minimis hirsutis asperis. *Tournef. inst.* 469.


Hieracium asperum, folio magno dentis leonis. *Bauh. pin.* 127.

Hieracium pumilum saxatile asperum, radice præmorfa. *Bauh. prodr.* 66.

Habitat in Germania, Helvetia, Gallia, Hispania &c. 24

Simillimum L. hispido. Sed diversum: Pedunculi & Calyces minus hispidi & fere glabri. Folia pilis apice indivisis & disco, dum sicca, valde flexuoso. Radius corollæ subtus, etiam luteus nec virescens. Calyx ante florescentiam nutans, maturo fructu canaliculatus. Pappus plumosus est.

Folia rigidula, tactu quasi arida, laciniis obliquatis. Calycis foliola extima basi utrinque dilatata mediante plica. Syst. Veget. 596.

Semina radii calyculo brevi, multidentato, loco pappi; Semina interiora pappo sessili; Sem. centralia pappo subtipitato coronata. *Leers l. c.* R. 

992. HIERACIUM.

Receptaculum nudum. *Cal.* imbricatus, ovatus.
Pappus simplex, sessilis.

* *Scapo nudo unifloro.*

1. **HIERACIUM** *incanum*. H. foliis integerimis subdenticulatis lanceolatis scabris, scapo unifloro. *Jacq. austr. t. 287. excluso Bauhino. pal. delph.*

Apargia incana. Scop. carn.

Leontodon (pyrenaicum) scapo lævi, pedunculis tumidis squamosis, calycibus villosis, foliis ovato-lanceolatis petiolatis subintegerrimis. *Gouan. illustr. p. 55. t. 22. f. 1. 2. Sec. De la Chenal. R.*

Leontodon hispidum. var. β. Linn. huc pertinet. R.

Picridis var. β. Hall. helu. n. 25.

Picris scapo nudo squamoso, foliis ovato-oblongis rariter dentatis. *De la Chenal. Act. helu. VIII. p. 133. R.*

Hieracium montanum VI. Clus. pan. 645. 646. R.

Hieracium montanum angustifolium II. Tabernæm. ic. 147. R.

Habitat in Austria, Helvetia, Palatinatu.

2. **HIERACIUM** *pumilum*. H. foliis ovatis, petiolis dilatatis, scapis subunifloris. *Jacq. austr. 2. t. 189 ? delph.*

*Leontodon dentatum. Mant. 1. p. 107. **

H. foliis ovato-lanceolatis, dentatis, petiolatis, petiolis dentatis. *Hall. helu. n. 42.*

Habitat in alpibus Helvetiæ, Sabaudia. 24

Planta vix palmaris. Folia omnia radicalia, cordata s. ovata, subtus subtomentosa, obtusiuscula, petiolata, denticulata. Petioli longitudine foliorum, superne dilatati, denticulati, sanguinei. Scapi uniflori, longitudine foliorum, adspersi 2 s. 3. foliolis subulatis, minusis; pubescentes, procumbentes, Calyx pubescens, oblongus, incanus squamis acutis. Corolla magnitudine *Pilosellæ*.

Est *Hieracium (prunellæfolium)* caule ramoso multifloro, foliis lyratis tomentosis, pedunculis unifloris. *Gouan. illustr. p. 57. t. 22. f. 3. quocum vero Crepidem pygmæam. Linn. conjungit. R.*

Hieracium sub nomine H. pumili Lin. a Jacquinno descriptum & depictum in flor. austr. t. 189. convenit cum planta

304 SYNGENESIA. POLYGAMIA ÆQUALIS.

in rupibus M. Dornacensis a me lecta & abunde differt ab Hier. Hall. helu. n. 42. De la Chenal. in Ad. Helu. VIII. p. 137. R.

2. **HIERACIUM alpinum.** H. foliis oblongis integris dentatis, scapo subnudo unifloro, calyce piloso. *Monsp. Jacq. austr. t. 191. tub. fil. Fl. dan. t. 27 delph. burg.*

H. caule unifloro, calyce villosa. *Fl. suec.*

H. caule unifloro; foliis lingulatis, dentatis hirsutis, calyce hirsuto. *Hall. helu. n. 49.*

H. scapo unifloro; folioso; foliis lingulatis mollibus; calyce piloso. *Scop. carn.*

H. alpinum pumilum, folio lanuginoso. *Bauh. pin. 129.*

H. villosum alpinum, flore magno singulari. *Rai. angl. 3. p. 169. t. 6. f. 2. **

Pilosella monoclonos non repens alpina minor lanuginosa; amplo flore. Moris. hist. 3. p. 78. f. 7. t. 7. f. 5.

Habitat in Alpibus Lapponicis, Britannicis, Austriacis 24

4. **HIERACIUM Taraxaci.** H. foliis lanceolatis dentatis glabris, scapo subnudo unifloro, calyce hirsuto. *delph.*

Picris caule hirsuto unifloro, foliis glabris semipinnatis, calycibus hirsutissimis. Hall. helu. n. 27.

Dens leonis alp. foliis oblongis, raro dentatis, calyce hispido nigrescente. Seguiet. suppl. 266. Hall. R.

Habitat in alpibus Helveticis, Pithoënsi-Lapponicis. 24

Hybrida an planta vel nova, ne pereat, proponitur.

Folia omnino Leontod. autumnalis; Scapus & Flos absolute Hier. alpini, ut si scapus & Flos hujus adponatur illius foliis, habeat quivis specimen plantæ. Folia lanceolata, glabra, dentata: dentibus aliis brevioribus, aliis oblongis, sæpius antrorsum flexis. Scapus foliorum rudimentis minutis aliquot, pilosus, versus florem crassior & apice uti calyx tumidus, fusco-pilosus. Differt a Leontod. aureo omnino, Calyx enim est valde imbricatus, quod vix in L. aureo, autumnali.

6. **HIERACIUM Pilosella.** H. foliis integerrimis ovatis subtus tomentosis, stolonibus repentibus, scapo unifloro.

Suec. paris. pal. ged. gallob. tub. Ludw. ed. t. 154.

Regn. bot. lugd. lithua. burgd.

H. caule unifloro, stolonibus reptantibus, foliis petiolatis ovatis longe pillosis, subtus tomentosis. *Hall. helu. n. 55.*

Pilosella major repens hirsuta. Bauh. pin. 262.

Pilosella minor. Cam. epis. 709. Fuchs. hist. 605. Blackw. t. 365.

Habitat

Habitat in Europæ pascuis aridis. 24

Corolla extus rubra.

* * *Scapo nudo multifloro.*

7. **HIERACIUM** *dubium*. H. foliis integris ovato-oblongis, stolonibus repentibus, scapo nudo multifloro. *Suéc. parif. monsp. herb. pal. sil. lithua. delph. burg.*

H. caule nudo, paucifloro, staminibus reptantibus, foliis longe pilosis, lanceolatis. *Hall. helu. n. 33.*

Pilosella major repens minus hirsuta. Bauh. pin. 262.

Habitat in Suecia. &c. 24

Caulis hirtus. Folia hirta, præsertim subtus.

8. **HIERACIUM** *Auricula*. H. foliis integerrimis, lanceolatis, stolonibus repentibus, scapo nudo multifloro. *suec. * gallob. tub. ged. lithua. delph. burg. herb. pal. sil.*

H. foliis integerrimis, caule repente, scapo nudo multifloro. *Hort. cliff. 388. **

H. caule nudo, stolonibus reptantibus foliis longe pilosis, lanceolatis, floribus umbellatis. *Hall. helu. n. 52.*

H. murorum angustifolium non sinuatum. *Bauh. prod. 64. R.*

H. alpinum, folio oblongo angustissimo. *Bauh. prodr. 64. Hall. R.*

Pilosella major umbellifera macrocaulos montana & pratensis. Col. ecphr. 248. 249. Hall. R.

Pilosella minor, folio angustiori, minus piloso, repens. Bauh. hist. 2. p. 1040. Hall. R.

Pilosella major erecta altera. Bauh. pin. 262.

Pilosella repens minor, caule pedali, polyanthos, foliis angustis oblongis. Rai. suppl. 147.

Habitat in Europæ pratis aridis, juxta agros. 24

9. **HIERACIUM** *cymosum*. H. foliis lanceolatis integris pilosis, scapo subnudo basi piloso, floribus subumbellatis. *herb. pal. lithu. delph.*

H. caule subnudo, foliis hirsutis, lanceolatis, floribus umbellatis, petiolis ramosis. *Hall. helu. n. 51. **

H. piloselioides majus longifolium, floribus luteis fere umbellatis. *Vaill. act. 706.*

H. murorum angustifolium non sinuatum. *Bauh. pin. 129. prodr. 67. t. 67.*

Pilosella montana hispida, parvo flore. Bauh. pin. 262.

Pilosella, minori flore, hirsutior & elatior non repens. Bauh. hist. 2. p. 1040.

Pilosella altera major. Tabern. p. 197. Hall. R.

Linnaei spec. plant. Europ. Pars II.

Y

206 SYNGENESIA. POLYGAMIA ÆQUALIS.

Habitat in Russia, Dania, Germania, Helvetia. 4.

Radix præmorsa. Folia lanceolata, longiuscula, acuta, integerrima, crassa, utrinque pilis rigidis lupida. Scapus pedalis, adspersus pilis varioribus, basi confertioribus, supra basin instructus foliolo unico. Flores fere umbellati, pedicellis tamen ramosis, quibus notis ab *H. præmorso* differt; ab *H. Auricula* vero differt foliis non supra glabris.

10. *HIERACIUM præmorsum*. *H.* foliis ovatis subdentatis scapo nudo racemoso, floribus superioribus primoribus. *sub. pal. lithu. suec. * delph.*

H. caule nudo, foliis penolatis, ovatis, integerrimis, hirsutis, floribus spicatis. *Hall. helv. n. 51.*

H. foliis ovatis tomentosis, caule nudo floribus spicatis luteolis terminato. *Gmel. sib. 2. p. 32. t. 13. f. 2.*

H. pratense latifolium non sinuatum majus & minus. Bauh. pin. 129.

H. latifolium, præmorsa radice, caule singulari pilosella majori affine. Bauh. hist. 2. p. 1035.

Pilosella major erecta. *Bauh. pin. 262. Rai. hist. 241.*

Intybus aphyllocaulis major. *Thal. harc. 66.*

Habitat in Helvetia, Germania, Uplandia. 24

11. *HIERACIUM aurantiacum*. *H.* foliis integris, caule subnudo simplicissimo piloso corymbifero. *Kniph. cens. 11. n. 56. Jacq. austr. t. 410. delph.*

H. caule subnudo, foliis ovatis, integris, floribus umbellatis aurantiis. *Hall. helv. n. 50.*

H. hortense floribus atro-purpurascens. Bauh. pin. 128. prodr. 65.

H. alpinum non laciniatum, flore fusco. Bauh. pin. 128. prodr. 65.

H. germanicum 1. *Col. ecphr. 2. p. 28. t. 30.*

Pilosella polyclonos repens major synaca, flore amplo aurantiaco. *Morif. hist. 3. p. 78. f. 7. t. 8. f. 7.*

Habitat in Synæ, Helvenæ, Austriæ sylvia. 24

*** *Caule folioso.*

17. *HIERACIUM chondrilloides*. *H.* caule ramoso, foliis

radicalibus lanceolatis dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

dentatis glabris: radicalibus lanceolatis

SYNGENESIA POLYGAMIA ÆQUALIS. 397

Petiolata; integra, glabra. Caulina 3 ad 6, alterna, glabra; longa; dentibus longis, recurvatis; facie Leonrod. autumnalis. Pedunculi ex alis superioribus foliorum, altitudine caulis, foliarii, subnudi, uniflori: Bractea una aut altera; seracea, apice subvillosa. Calyces nigro hispidi, erecti, imbricati foliolis linearibus. Corolla flava.

18. *HIERACIUM porrifolium.* H. caule ramofo foliofo; foliis lanceolato-linearibus subintegerrimis. *Jacq. vind. austr. t. 286. carn. lithu. delph.*

H. caule ramofo, foliis radicalibus linearibus, caulinis minimis. *Ger. prou.*

Hieracium. *Hall. helv. n. 48. excepta varietate. a. R.*

H. trapezontis folio. *Bauh. pin. 129. R.*

H. folio italice, caule foliato. *Segui. ver. 2. p. 270. n. 6: bona descriptio, exceptis synonymis. R.*

H. murorum angustifolium non sinuatum. *Bauh. pin. 189.*

H. phalangii foliis. *Vaill. ad. 1721. p. 184.*

H. montanum, asphodeli foliis acuminatis. *Bocc. mus. 147: t. 106. Rai. suppl. 142.*

Habitat in Alpibus Austriæ, Italiæ, Valis. 24

Caulis pedalis; teres, subnudus folio uno alterove: ramis raris apyillis; Folia radicalia linearia, erecta, dente uno alterove acuta, levia: carina raris villis; Caulina minora, paucissima; Calyx imbricatus, canescens; oblongus. Corolla lutes, non maxima. *Mant. 2. p. 458.*

19. *HIERACIUM murorum.* H. caule ramofo, foliis radicalibus ovatis dentatis; caulino minori: *succ. parif. gallob. päl. fl. herb. lith. burg. lugd.*

H. caule subnudo, paucifloro, foliis ovato-lanceolatis; circa petiolum dentatis. *Hall. helv. n. 46.*

H. scapo multifloro, subnudo; foliis ovatis, dentatis. *Stöp. carn.*

H. caule ramofo, foliis ovatis dentatis. *Fl. lapp.*

20. *Hieracium (pilosissimum) murorum, folio pilosissimo. Bauh. pin. 129.*

Cotchorus. Dalech. hist. 363.

21. *Hieracium (sylvaticum) murorum levinatum minus pilosum. Bauh. pin. 129.*

H. (sylvaticum) caule ramofo foliofo, foliis radicalibus ovatis petiolatis, caulinis sessilibus; pedunculis multifloris.

Gouan. illustr. Hall. helv. n. 46. β. δ. & ε. Distincta species Gouano. R.

Pulmonaria gallosum flore hieracii, Auricula muris major. Tragi. Lob. ic. 387. Gou.

308 SYNGENESIA. POLYGAMIA ÆQUALIS.

Pilosella major f. *Pulmonariae luteae species angustifoliae*.
Bauh. hist. 2. p. 1034. Gou. R.

Pulmonaria gallica semina. Tabern. hist. 504. Blakw. t. 538.

7. *Hieracium macrocaulon hirsutum*, folio longiore. *Rat. angl. 3. p. 169.*

8. *Pulmonaria gallica rotundifolia. Barrel. is. 342. Hall. l. c. var. e. R.*

Habitat in Europæ apricis duris.

20. *HIERACIUM paludosum. H. caule paniculato, foliis amplexicaulibus dentatis glabris, calycibus hirsutis. Fl. suec. ingr. ged. gallob. carn. pal. nass. herb. delph. lithu.*

H. foliis glabris, dentatis, imis petiolaris, superioribus amplexicaulibus, calycibus hirsutis. Hall. helu. n. 45. R.

H. cichorei folio. It. w. goth. 32.

H. montanum latifolium glabrum minus. Bauh. pin. 129. Moris. hist. 3. p. 69. f. 7. t. 5. f. 47.

H. latifolium glabrum, ex valle griessbachiana. Bauh. hist. 2. p. 123. Hall. R.

Habitat in Europæ borealioris nemoribus paludosis.

Calyces pilis nigris exstantibus hirti sunt. Varietas forte sola antecedentis, quamvis ex loco evasit facie diversissima.

22. *HIERACIUM cerinthoides. H. foliis radicalibus obovatis denticulatis; caulinis oblongis semiamplexicaulibus. carn. delph.*

H. foliis radicalibus obovatis obtusis petiolatis denticulatis, caulinis oblongis semiamplexicaulibus acutis. Gouan. illustr. 58. t. 22. f. 4. R.

H. pyrenaicum, foliis cerinthes, latifolium & angustifolium. Tournef. inst. 472.

Habitat in Pyrenæis. 24

Conf. Hieracium caule ramoso folioso, foliis & calyce longo villo barbatis. Hall. helu. I. n. 744.

23. *HIERACIUM amplexicaule. H. foliis amplexicaulibus cordatis subdentatis, pedunculis unifloris hirsutis, caule ramosq. Kniph. cent. 12. n. 56. delph.*

H. foliis ovato-lanceolatis, rariter dentatis, caulinis amplexicaulibus. Hall. helu. n. 36. sec. Gouan. R.

H. (amplexicaule) foliis radicalibus ovato-lanceolatis acutis basi dentatis, caulinis hastato-cordatis amplexicaulibus. Gouan. illustr. 58. quacum denominationis Hier. blattarioides ab ipso combinatur. R.

H. pyrenaicum longifolium amplexicaule. Tournef. inst. 472.

β. *H. pyrenaicum rotundifolium amplexicaule. Tournef. inst.*
472.

Habitat in Pyrenæis.

Affine admodum H. pyrenaico. Planta tota adpersa pilis raris apice glandulosis. Caulis striatus. Folia utrinque viridia dentibus argutis. Rami ex alis superioribus, sæpe biflori. Panicula inter pilos etiam villos habet, unde hinc quasi tomentosa.

24. *HIERACIUM pyrenaicum. H. foliis amplexicaulibus obovato-lanceolatis retrorsum dentatis, caule simplici, calycibus laxis.*

β. *H. (blattarioides) foliis lanceolatis amplexicaulibus dentatis, floribus solitariis, calycibus laxis. Spec. 2.*

H. pyrenaicum, blattariæ folio, minus hirsutum. Tournef. inst. 472.

γ. *Hieracium pilosum.*

Picris (pyrenaica) perianthiis laxis, caule piloso, foliis dentato-sinuatis. Sp. 2. Gouan. Illustr. 53.

Hierac. pyrenaicum, blattariæ folio, minus pilosum. Herm. parad. 184. t. 184.

Pappus plumosus sessilis. Gouan. R.

δ. *Hieracium austriacum.*

Crepis (austriaca) foliis oblongis denticulatis, involucre laxissimo & calycibus hispidis. Jacq. vind. 270. t. 5. austr. t. 441.

ε. *Hieracium helveticum.*

H. foliis lanceolatis serratis hirsutis, calycibus villosis. Hall. helv. n. 40.

H. (conyzæfolium) foliis argute dentatis, radicalibus lanceolato-oblongis acutis, caulinis sagittatis oblongis, pedunculis unifloris, calycibus pilosissimis laxis. Gouan. illustr. 59. R.

H. britannicum quartum. Clus. hist. 2. p. 151. sec. Gouan. R.

H. alpinum asperum, conyzæ facie. Bauh. pin. 128.

Differt a Picride pyrenaica (Hieracio piloso Linn.) pedunculis longis unifloris, apice tumidis, piloso-atris, squamulis vagis destitutis. Calyce pilis atris undique obvoluto, imbricato squamis laxis; Pappo piloso sessili. Gouan. l. c. R.

Habitat in Pyrenæis, Helvetia, Austria. 24

Planta polymorpha ex loco difficilius eximinanda, media inter Hieracia, Crepides & Picrides. Affinis proxime Crepidis sibiricæ. Inquirendum quomodo & quibus ab hac differant β. & problematice propositæ. Syst. veg. 598.

310 SYNGENESIA. POLYGAMIA ÆQUALIS.

25. **HIERACIUM villosum.** H. caule ramoso folioso, foliis hirtis: radicalibus lanceolato-ovatis dentatis: caulinis amplexicaulibus cordatis. *Jacq. austr. t. 87. delph.*

H. foliis villosis integerrimis, radicalibus ligularis, caulinis ovatis, amplexicaulibus, calycibus villosis. *Hall. helu. n. 44.*

H. alpinum latifolium villosum, magno flore. *Bauh. pin. 128. Scheuchz. it. alp. 1. p. 39.*

H. alpinum, latiore folio, pilosum, flore majore. *Pluk. alm. 184. t. 194. f. 2.*

H. alpinum V. villosum. *Clus. hist. 2. p. 141. Hall. opusc. 146.*

β. Pilosella f. Pulmonaria lutea, angustiori folio, valde pilosa, duplex. *Bauh. hist. 2. p. 1034. ic. bona. Hall. R.*

γ. Hieracium caule unifloro, foliis levibus, radicalibus lingulatis, obiter dentatis, caulinis ovato-lanceolatis, amplexicaulibus. *Hall. emed. 3. n. 254. cat. rar. n. 972. helu. n. 44. var. δ. R.*

δ. Variatio glabris foliis. *Seguier. suppl. p. 269. Hall. l. c. R.*

Habitat in Alpibus Bohemicis, Helveticis, Monspeli. 24
Caulis striatus, subpilosus: Ramis altitudine caulis. Folia Radicalia lanceolata, latiuscula, subpilosa, dentibus inermibus. Caulina cordata, amplexicaulia, basi sursum conniventia. Pedunculi uniflori, bractea calycina remotiore. Calyx patulus, minus imbricatus foliolis longitudine subæqualibus adpersus pilis basi nigris: apice (ut tota planta) glandula flava. Semina nigra. Pappo sessili, multiplici, piloso.

26. **HIERACIUM glutinosum.** H. foliis lanceolatis runcinatis scabriusculis, floribus umbellatis.

H. dentis leonis folio, floribus parvis. *Bauh. pin. 127. prodr. 63.*

Habitat in G. Narbonensi. ☉

Habitus *Crepis tectorum*. Folia mollia, glutinosa, summa lanceolata, integerrima. Caulis striatus.

28. **HIERACIUM sprengerianum.** H. caule ramoso folioso, foliis amplexicaulibus oblongis repandis hispidis.

H. pulchrum ramosum sprengerianum. *Bauh. hist. 2. p. 1026. Moris. hist. 3. p. 65. f. 7. t. 5. f. 17.*

Habitat in Lusitania.

Caulis pilis raris rigidulis: ramis alternis brevioribus simplicibus. Folia amplexicaulia, ovata, obtusiuscula, ciliata, rarius pilosa, subtus magis. Calyces terminales, flores pedunculati, sparse pilosi.

SYNGENESIA. POLYGAMIA ÆQUALIS. 311

29. **HIERACIUM sabaudum.** H. caule erecto, multifloro, foliis ovato-lanceolatis dentatis semiamplexicaulibus. *Fl. suec. gallob. pal. sil. lithu. lugd. burg.*

H. foliis hirsutis, dentatis, inferioribus ellipticis, superioribus ovato-lanceolatis. *Hall. helu. n. 35.*

H. caule multifloro, folioso; foliis lanceolatis, dentatis; dentibus glandulosis. *Scop. carn.*

H. virosum. *Pallas it. 1.*

H. fruticosum latifolium hirsutum. *Bauh. pin. 129.*

β. Hieracii sabaudi varietas. 1. *Bauh. hist. 2. p. 1030.*

Hieracium fruticosum subrotundo folio. *Bauh. prodr. Hall. R.*

H. sabaudum altissimum, foliis latis brevibus, crebrius nascentibus. *Morif. hist. 3. p. 71. Hall. R.*

Habitat in Germania. &c. 24

30. **HIERACIUM umbellatum.** H. foliis linearibus subdentatis sparsis, floribus subumbellatis. *Fl. suec. paris. ged. Hall. helu. n. 34. gallob. pal. herb. sil. Flor. dan. t. 680. Kniph. cent. 9. n. 48. lithu. lugd. burg.*

H. fruticosum angustifolium majus. *Bauh. pin. 129. Dill. ephem. Nat. Cur. cent. 5 & 6 app. t. 13. f. 2.*

H. sabaudum. *Dalech. hist. 570.*

β. Var. foliis pene ovatis, vix dentatis, caule humili, pene unifloro.

H. fruticosum angustifolium minus. *Dill. Eph. Nat. Cur. V. VI. append. t. 13. f. 1. Vaill. paris. t. 48. Hall. R.*

γ. Angustissimis, quasi linearibus foliis.

Hieracium fruticosum angustissimo incano folio. *Vaill. Pet. t. 13. Hall. R.*

Habitat in Europæ pascuis siccis. 24

Calyces squarrosi.

993. CREPIS.

Receptaculum nudum. Cal. calyculatus squammis deciduis. Pappus pilosus, stipitatus.

1. **CREPIS pygmaea.** C. foliis ovatis integris villosis petiolatis, caule procumbente. † *delph.*

Hieracium alpinum incanum saxatile, brunellæ foliis integris.

Bocc. mus. 2. p. 33. t. 14. Rai suppl. 142.

Habitat in Alpibus Italiae ad Nursiam prope lacum pilati.

312 SYNGENESIA. POLYGAMIA. EQUALIS.

Cum Hieracio pumilo Linn. conjungis Gouan. Illustr. p. 57. R.

2. CREPIS *bursifolia*. C. foliis pinnatifidis crenatis, scapo nudo paucifloro.

Hieracium siculum, bursæ pastoris folio, Bocc. mus. 2. p. 147. t. 106. & 112. Rai. suppl. 142.

Hieracium minus panormitanum, cichorei folio. Cup. cath. 95 suppl. 3.

Habitat in Sicilia. 2.

3. CREPIS *barbata*. C. involucris calyce longioribus : squamis setaceis sparsis. *Kniph. cent. II. n. 35.*

Hieracium calycibus setis longissimis distantibus patentibus obvallatis. Hort. cliff.

Hieracium boeticum majus. Herm. parad. 185. t. 185.

Hieracium proliferum salcatum. Bauh. pin. 128.

Hieracium calyce barbato. Col. ecphr. 2. p. 28. t. 27. f. 1.

Habitat in Monspelii, Vesuvii, Siciliae, Messanae arenosis maritimis. ☉

Folia lanceolata, glabra, subdentata. Ramuli basi angustiores; quasi unus ex altero prognatus. Flores lutei, ad basin purpurascens, at in disco flosculi atropurpureascentes fere totti. Semina teretia, coronata margine ciliari minutissimo & setis 2 simplicibus longis. Calyx exterior corolla longior.

4. CREPIS *vesicaria*. C. involucris scariosis longitudine calycis, floribus corymbosis, bracteis ovatis.

Hieracium montanum rap. olium. Bauh. pin. 129. prodr. 749. Rai. hist. 237. Moris. hist. 3. p. 63. secundum Burs. VI. 70.

β. Cichorium pratense hirsutum vesicarium. Bauh. pin. 126.

Cichorium sylvestre vesicarium pratense. Col. ecphr. II. p. 238. t. 237.

Habitat in Helvetia, β. in Apulia.

Plantæ α Radix crassa. Folia lyrata, angulata, dentata, scabra. Caulis brevis. Rami longi, nudibifidi s. trifidi, uniflori, pedunculorum Bractee ovatae, foliolosae, concavae, integrae, pallidae. Flores lutei. Calyces singuli cincti. Involucro longitudine calycis, ex foliolis 5. s. 6. patentibus, ovatis, obtusis concavis, membranaceis, corioidis. Planta β foliis a priori manifeste differt, florum vero conjungunt utramque.

Non crescit in Helvetia, nam Hieracium montanum multifolium Bauh. pertinet ad Crepidem biennem. Hall.

SYNGENESIA. POLYGAMIA ÆQUALIS. 313

5. *CREPIS alpina*. C. involucris scariosis longitudine calycis, floribus solitariis.

C. foliis amplexicaulibus oblongis acuminatis: inferioribus superne; summis inferne denticulatis. *Hort. ups.*

Hieracioides annua, endiviæ folio, capite magno. *Vaill. ad. 1721. p. 246.*

Hieracium alpinum, scorsoneræ folio. *Tournef. inst. 472.*

Leontodon calyce toto erecto: inferiore squamis ficcis; foliis amplexicaulibus. *Gmel. sibir. 2. p. 16. t. 5.*

Habitat in Alpibus Italiae. &c. ☉

Calyx pilosus cinctus foliolis calycinis ovatis, glabris, aridis.

6. *CREPIS rubra*. C. foliis amplexicaulibus lyrato-runcinatis. *monsp. Kniph. cens. II. n. 36.*

Hieracium foliis caulinis sinuatis, calycibus ante florescentiam nutantibus *Hort. cliff.*

Hieracium intybaceum, floribus ex purpura rubentibus. *Bauh. pin. 128. Rai. hist. 231.*

Hieracium dentis leonis folio, flore suaverubente. *Bauh. pin. 127.*

Habitat in Apulia. ☉

Pappus stipitatus ut in C. alpina, & vesicaria. Calyx ante florescentiam nutat.

7. *CREPIS foetida*. C. foliis runcinato pinnatis hirtis, petiolis dentatis. *Herb. pal. delph. burg. lugd.*

C. foliis scabris pinnatis, pinnis angulosis, dentatis, retroverfis, extrema triangulari maxima. *Hall. helv. n. 29.*

Hieracium luteum, cichorii sylvestris folio amygdalas amaras olens. *Morif. hist. 3. p. 63. f. 7. t. 4. f. 4.*

Hieracium foliis cichorii sylvestris villosis, odore castorii. *Magn. monsp. 129. t. 128.*

Hieracium castorii odore monspeliensium. *Rai. hist. 232.*

Hieracium orientale altissimum, folio cichorii sylvestris, odore castorii, flore magno. *Tournef. cor. 35.*

Hieracium maximum, erucæ folio. *Tournef. inst. 469. glabrum. Rai suppl. 146.*

Jacobæa sylvatica tomentosa, cichorii sylvestris folio. *Morif. hist. 3. p. 109.*

Senceio hirsutus. *Bauh. pin. 131.*

Erigeron tertium. *Dod. pempt. 641.*

Hieracium apulum, flore suaverubente. *Col. ecphr. I. p. 244.*

Chondrilla purpurascens foetida. *Bauh. pin. 130. prodr. 68. t. 68*

316 SYNGENESIA. POLYGAMIA ÆQUALIS.

Lapsana capillaris. Sp. pl. 1. Gouan. monsp.

Hieracium minus glabrum, foliis eleganter virentibus. Bauh. pin. 127. prodr. 63.

Hieracium minus Dioscoridis. Tabern. 181. Hall. R.

Hieracium dentis leonis folio. hirsutum, flore luteo extus purpurascens. Tournef. paris. 99.

Habitat in Helvetiæ, Italiæ agris.

Folia radicalia lanceolata, obtusa, glabra, subdentata; Caulina lanceolata, hastata, subamplexicaulia, integerrima, acuta, parva. Caules spithamæi, erecti, filiformes, subdichotomi. Flores parvi, lutei. Calyces soli tomentosi. An præcedentis sola varietas.

Cum Crepide sequenti in unam speciem conjungit. Gouan. R.

14. CREPIS Dioscoridis. C. foliis radicalibus runcinatis, caulinis hastatis, calycibus subtomentosis. pal.

Hieracioides vulgaris annua, cichorii folio, flore luteo subtus purpurascens. Vaill. æd. 715. paris. 102.

Hieracium majus erectum angustifolium, caule lævi. Bauh. pin. 127.

Hieracium 1. Tabern. hist. 491.

Habitat in Gallia, Sibiria, Palatinatu. ☉

Caulis pedalis, subangulatus, læviusculus: Ramis rarioribus, teretibus, patentibus. Folia radicalia lyrato-dentata, nuda, glabra, denticulata, vix ciliata. Caulina amplexicaulia, hastata, lineari-lanceolata, disco inferiore litura purpurascens, basi auriculis valde dentatis. Ramea linearia, basi uno dente. Pedunculi longi, nudi, adscendentes, uniflori vix apice incrassati. Calyces farinosi, ante florescentiam non nutantes, calyculo lineari. Corolla flava, ante explicationem s. subtus purpurascens.

15. CREPIS pulchra. C. foliis sagittatis denticulatis, caule paniculato, calycibus pyramidatis glabris. Gouan. monsp. delph.

C. foliis sagittatis dentatis, pedunculis superne turbinato-inflatis. Guett. stamp. Dalib. paris. n. 1.

Chondrilla, hieracii folio, annua. Tournef. inst. 238.

Hieracium pulchrum. Bauh. hist. 2. p. 1024.

Hieracium montanum hirsutum minus. C. B. Rai. hist. 234.

Hieracioides annua glutinosa, floribus parvis. Vaill. æd. 1721.

β. *Lapsana condriilloides calycibus papposis, foliis oblongis subde-*
245. *

...calycibus papposis, foliis oblongis subde-
...calycibus papposis, foliis oblongis subde-

SYNGENESIA. POLYGAMIA ÆQUALIS. 317

Hieracium montanum alterum leptomacrocaulon. Col. ecph.

1. p. 248. t. 249. Rai. hist. 234.

Hieracium annuum montanum, caule canaliculato. Moris. hist. 3. p. 68. f. 7. t. 5. f. 37.

Habitat in Gallia, Italia. ☉

Calyx lævis, angustus, angulatus, auctus squamis minutissimis, viridibus.

16. CARPIS neglecta. *C. foliis amplexicaulibus runcinatis dentatis subpilosis, caule paniculato, calycibus inermi-subspinosis. Ment. 107.*

Habitat in Italia. ☉

Caulis pedalis, ramosus, subpilosus, erectus. Folia radicalia obovato-oblonga, dentata, subpilosa. Caulina amplexicaulia, runcinata, subpilosa dentibus elongatis etiam baseos; summa subhastata. Rami floriferi elongati, nudi, læves, apice biflori s. bifidi. Calyx exterior brevissimus, acutus; interior 8 s. 10 foliolis, in medio una alterave spina inermi notatis. Convenit cum C. pulchra magnitudine & parvis floribus; differt foliis, non integris subdentatis viscosis, sed runcinatis, dentatis, subpilosis, &c. Synonyma inter utramque dispertienda. H. V.

994. ANDRYALA.

Recept. villosum. Cal. multipartitus, subæqualis & rotundatus. Pappus simplex, sessilis.

1. ANDRYALA integrifolia. *A. foliis inferioribus runcinatis, superioribus ovato-oblongis tomentosis. Stamp. Sauv. monsp. Kniph. cent. 4. n. 5. delph. lugd.*

Sonchus villosus luteus major. Bauh. pin. 124.

Sonchus lanatus. Dalech. hist. 1116.

Hieracium villosum. Rai. hist. 231.

β. Andryala (sinuata) *foliis runcinatis. Spec. ed. 2. p. 1137. Mill. dict. n. 3.*

A. foliis inferioribus dentatis, summis integris. Hort. cliff. 387. Sauv. Monsp. 692. Gouan. monsp. 415.

Eriophorus foliis inferioribus ad costam usque laciniatis. Vaill. act. 1721. p. 212.

Sonchus villosus luteus minor. Bauh. pin. 124. prodr. 61.

Habitat Monspeli, inque Sicilia. ☉

2. ANDRYALA ragusina. *A. foliis lanceolatis indivisis denticulatis acutis tomentosis, floribus solitariis.*

218 SYNGENESIA. POLYGAMIA ÆQUALIS.

Hieracium incanum lanuginosum ragulinum, pilosellæ flore.
Herm. Lugdb. 672. t. 673. Mill. ic. 97. t. 146. f. 2.

Habitat in insulis Archipelagi. 24

Folia strictiora, acutiora, acutius dentata quam in reliquis.

3. *ANDRYALA lanata. A. foliis oblongo-ovatis subdentatis lanatis, pedunculis ramosis. Amm. acad. 4. delph.*

Hieracium foliis ovatis, crassis, lanatis. Hall. helv. n. 37.

Hieracium montanum tomentosum. Dill. elik. 181. t. 140. f. 180. Mill. ic. 97. t. 146. f. 1.

Habitat in Europa australi. 24

Planta tota alba est. Receptaculum sæpius non villosum.
Gmel. it. 1. p. 138.

995. HYOSERIS.

Receptaculum nudum. Cal. subæqualis Pappus pilosus calyculatusque.

** Caule nudo.*

1. *HYOSERIS foetida. H. scapis simplicissimis unifloris, foliis pinnatifidis, seminibus nudis. Gouan. monsp. delph. burg.*

Lapsana caule nudo indiviso, foliis semipinnatis, pinnis retroverfis dentatis. Hall. helv. n. 5.

Lamproloma (foetida) scapo unifloro nudo. Scop. carn. 1. p. 397. n. 2. ed. 2. n. 989.

Taraxacomastrum, dentis leonis folio, radice foetida. Vaill. 1721. p. 180.

Leontodontoides alpinus glaber, erysimi folio, radice crassa foetida. Mich. gen. 31. t. 28.

Dens Leonis tenuissimo folio. Bauh. pin. 126. prodr. 62.

Hieracium foetidum III. Col. ecphr. 2. p. 29. t. 31.

Habitat in Alpibus Italiæ, Austriæ superioris. 24

Folia pinnatifida, glabra, lobis latere postico subdentatis. Scapi simplices, nudi. Calyx simplex, basi audus. Semina nuda.

2. *HYOSERIS radiata. H. scapis unifloris nudis, foliis glabris runcinatis: angulis dentatis; apice laciniosis.*

Hyoseris foliis glabris hastato - pinnatis. Sauv. monsp.

Taraxacomastrum dentis leonis folio, pinnatitatem radiato.
Vaill. ad. 703.

SYNGENESIA. POLYGAMIA ÆQUALIS. 319

Dens leonis minor, foliis radiatis. *Bauh. pin.* 129. *Pluk. alm.* 130. t. 37. f. 2.

Habitat in Hispania, G. Narbonensi, circa Tubingam. Gmel. tub. 242. 2.

Pedunculus & rachis foliorum farinacea. Lobi foliorum imbricatim reflexi, utrinque dentati, læves. Semina membranacea, Pappo simplici, sessili.

4. HYOSERIS scabra. H. scapis unifloris: foliis scabris, seminibus papposis.

Taraxaconastrum, erysimi folio, flore minimo. Vaill. ad. 1721. p. 235.

Dens leonis minimus, foliis hirsutis, calycis segmentis a flore delapso erectis & semina complexis. *Rai. suppl.* 147.

Hieracium minimum supinum, tragopogoni capitulis. *Bocc. mus.* 2. p. 146. t. 106.

Habitat in Sicilia.

Folia sublyrata, nuda, vix scabra, opaca, subciliata: Laciniis subalternis, subdentatis: extrema trilobo-angulata. Scapi læves, adpersi polline pellucido, fistulosi, superne incrassati, ad latus inclinati. Calyx oblongus, arcus, sexangularis, subhexaphyllus, foliolis lanceolatis, subcarinatis, æqualibus: foliolis totidem externis basin calycis regentibus. Semina coronata margine scabro pappoque aristato.

6. HYOSERIS minima. H. caule diviso nudo, pedunculis incrassatis. *Fl. suec. monsp. ged. pal. sil. Fl. dan.* t. 201. *delph. lithu. lugd. burg.*

H. angustifolia. Tabern. ic. 180.

Lapsana foliis ovatis, dentatis, caulibus nudis, sub flore tumentibus. *Hall. helu. n.* 4.

Hieracium minus, folio subrotundo. *Bauh. pin.* 127.

Hieracium minimum. *Clus. hist.* 2. p. 142.

Habitat in Europæ arvis apricis.

* * *Caule folioso.*

7. HYOSERIS Hedypnois. H. fructibus ovaris glabris, caule ramoso. *delph.*

Lapsana fructibus subglobosis glabris, seminibus disci calyculato-papposis. *Hort. ups. Sauv. monsp.*

Lapsana calycibus fructus capitatis dehiscentibus, pedunculis incrassatis, floribus nutantibus. *Hort. cliff.*

Rhagadioloides calthæ folio, calyce glabro. *Vaill. ad.* 1721. p. 202.

Hieracium capitulum inclinans, semine adunco. *Bauh. pin.*

320 SYNGENESIA. POLYGAMIA. ÆQUALIS.

Hieracium facie hedypnois. *Lob. ic. 239.*

Hedypnois annua. *Tournef. inst. 478.*

Habitas in Europa australi. ☉

8. HYOSERIS *fragadioides*. H. fructibus ovatis pilosis ;
caule ramofo. *Amph. cens. 5. n. 44. delph.*

Lapiana fructibus globosis pilosis, seminibus disci calyculato-
papposis. *Hort. ups.*

Habitas in Europa australi. ☉

996. S E R I O L A.

Receptaculum paleaceum. *Cal.* simplex. *Pappus*
subplumosus.

2. SERIOLA *astereifolia*. S. hispida, foliis obovatis subden-
tatis. *Jacq. obs. 4. p. 3. t. 79.*

Hypochaeris calycibus aequalibus hispidis. *Hort. ups.*

Achyrophorus hirsutus bellidis obtuso folio. *Vaill. ed.*
739.

Hieracium multicaule, bellidis folio molli. *Bocc. art. 8.*

Habitas in Italia. 2.

4. SERIOLA *urens*. S. urens, caule ramofo, foliis den-
tatis.

Hypochaeris foliis dentatis, caule ramofo folioso, calycibus
hispidis. *Sp. pl.*

Hypochaeris hieracii barbari folio, calyce hispido. *Vaill. ed.*
1721. p. 215.

Hieracium alpinum, dentis leonis folio acuto plosso pro-
cisso, capitulis pungent lanugine obfritis. *Cap. cath. 5.*

Rai. suppl. 146.

Habitas in Sicilia.

Calycis squama a sepe spinali pungentius.

997. H I P O C H A E R I S:

Receptacul. paleaceum. *Cal.* subimbricatus. *Pappus*
plumosus.

HIPPOCHÆRIS *spontanea*. H. caule simplici fol. trifloro
dentatis, cephal.

caule simplici, foliis caulinis ovato-lanceolatis,
capitulis simplicibus. *Hall. herb. 38.*

caule simplici, foliis caulinis ovato-lanceolatis,
capitulis simplicibus. *art. 1. 190.*

Hieracium

SYNGENESIA. POLYGAMIA ÆQUALIS. 221

Hieracium alpinum glabrum, flore singulari, cauli magno, cubitali folioso infidente. *Rai. ext.* 144. *Hall. R.*

Hypochæris non ramosa endiviæ folio prælongo, monanthos. *Vaill. acd.* 1711. p. 215.

Hieracium latifolium pontanum præaltum glabrum, endiviæ folio. *Bocc. mus.* 2. p. 148. t. 113. *Rai. suppl.* 142.

Habitat in Sabaudia montibus.

2. HYPOCHÆRIS maculata. H. caule subnudo: ramo solitario, foliis ovato-oblongis integris dentatis. *Fl. succ. parif. ingr. pal. fil. Flor. dan.* t. 149. *lith. delph.*

H. hirsuta, endiviæ folio, magno flore. *Vaill. acd.* 1721. p. 385.

H. caule unifloro, foliis dentatis, hirsutis. *Hall. helu.* 2. n. 2. t. 1.

H. foliis ovato-oblongis, dentatis; caule folioso, ramoso, hispido. *Scop. carn.*

Achyrophorus maculatus. Scop. carn.

Hieracium alpinum latifolium hirsutis incanum, flore magno. *Bauh. pin.* 128.

Hieracium latifolium 1. *Clus. hist.* 2. p. 139.

β. **Hieracium alpinum**, foliis dentatis, flore magno. *Bauh. pin.* 128. *prodr.* 65. *Hall. helu.* 1. p. 760. t. 24.

Dens leonis foliis integris, caule raris foliis vestito, monanthos fere. *Rai. hist.* 244.

Habitat in Europæ frigidioris pratis asperis. 24

Semina rugosa.

3. HYPOCHÆRIS glabra. H. glabra, calycibus oblongis imbricatis, caule ramoso nudo, foliis dentato-sinuatis. *Hort. upf. * parif. monsp. fil. ged. pal. Flor. dan.* t. 424. *burg. lithu.*

H. foliis glabris, semipinnatis. *Hall. helu.* n. 3. * *gatt.* p. 421.

H. porcellina. *Tabern.* 179. R.

H. chondrillæ folio, parvo flore. *Vaill. acd.* 1721. p. 214.

Hieracium minus, dentis leonis folio oblongo glabro. *Bauh. pin.* 127.

Habitat in Dania, Germania, Belgio, Helvetia. ☉

Corollæ minutæ. Calyx squamis lævibus. Semina disci pappo stipitato; Radix pappo sessili. Hall.

4. HYPOCHÆRIS radicata. H. foliis runcinatis obtusis scabris, caule ramoso nudo lævi, pedunculis squamosis. *succ. monsp. ingr. ged. pom. pal. fil. Flor. dan.* t. 150. *delph. lithu. burg.*

Linnaei spec. plant. Europ. Pars II.

X

322 SYNGENESIA. POLYGAMIA ÆQUALIS.

H. foliis semipinnatis hirsutis, caule nudo, brachiato. Hall. helu. n. 3.

Achyrophorus radicans. Scop. carn.

Hieracium dentis leonis folio obtuso, majus. Bauh. pin. 127. Moris. hist. 3. p. 66. f. 7. t. 4. f. 27.

Hieracium minus, dentis leonis folio subaspero. Bauh. pin. 127. (Vaill.)

Hieracium, dentis leonis folio, bulbosum. Bauh. pin. 127. prod. 63.

Hieracium longius radicans. Lob. ic. 238. Rai. hist. 230.

Hieracium tertium. Dod. pempt. 639.

Hieracium VII. Tabern. hist. 492.

Habitat in Europæ culsioris pascuis. 24

Pedunculi superne incrassati. Corollæ magnæ. Calycis squamæ carina ciliata.

998. L A P S A N A.

Recept. nudum. Cal. calyculatus squamis singulis interioribus canaliculatis.

1. LAPSANA communis. L. calycibus fructus angulatis, pedunculis teoribus ramosissimis. Juec. parif. ged. gall. carn. pal. fil. Fl. dan. t. 500. Kniph. cent. 5. n. 48. Regn. bot. lugd. lith. burg.

L. caule brachiato, foliis ovatis, longe petiolatis, petiolis pinnatis. Hall. helu. n. 6.

Sonchus sylvaticus. 1. 2. 3. Tabern. 192. 193. Hall. R. Lampfana. Dod. pempt. 675.

Soncho affinis Lapsana domestica. Bauh. pin. 124.

Habitat in Europæ cultis.

2. LAPSANA Zacintha. L. calycibus fructus torulosis depressis obtusis sessilibus. monsp.

L. calycibus fructus subrotundis angulatis; pedunculis incrassatis, floribus erectis. Hort. cliff.

Chondrilla verrucaria, foliis cichorei viridibus. Bauh. pin. 130.

Cichorium verrucatum, zacintha. Clus. hist. 2. p. 144.

Habitat in Italia, Oriente. ☉

Folia Radicalia lyrata, acuta; Caulina sagittata, amplius dentata, dentata. Caulis subdichotomus, striatus, rigidus. Flores sessiles, impuberes penduli. Corollæ subu-

SYNGENESIA. POLYGAMIA AEQUALIS. 329

salva, supra flavæ. Calycis exterior ordo erectus, interior basi gibbus, rufescens. Pedunculi incrassati.

3. LAPSANA stellata. L. calycibus fructus undique patentibus: radiis subulatis, foliis caulinis lanceolatis indivisis. Hort. upf. delph.

Hieracium filiqua falcata. Bauh. pin. 128.

Hieracium stellatum. Bauh. hist. 2. p. 1014.

Habitat Monspeli, Bononiæ. ☉

Variat foliis sinuatis, dentatis, indivisis. Gerard.

4. LAPSANA Rhagadiolus. L. calycibus fructus undique patentibus: radiis subulatis, foliis lyratis. Hort. upf. Kniph. cent. 4. n. 36. delph.

Lampfana Rhagadiolus. Scop. carn.

Hieracium falcatum alterum. Rai. hist. 256.

Rhagadiolus lapsanæ foliis. Tournef. cor. 36.

Rhagadiolus edulis hieraciis affinis. Bauh. hist. 2. p. 1014.

Habitat in Oriente, Carniola.

Varietas antecedentis. Gerard, sed forte constans.

999. CATANANCHE.

Recept. paleaceum. Cal. imbricatus. Pappi aristati. pappo s. seto.

1. CATANANCHE cerulea. G. squamis calycinis inferioribus ovatis. Lugdb. monsp. delph.

Catananche cerulea, semisflosculorum ordine simplici. Vahl. ed. 1721. p. 215.

Chondrilla cerulea, cyani capitulo. Bauh. pin. 490.

Chondrilla species tertia. Dod. pempt. 638.

β. Catananche flore pleno ceruleo. Tournef. inf. 478.

Chondrilla cerulea, cyani capitulo, altera. Bauh. pin. 1302.

Habitat in G. Narbonensis collibus sacrofla. 24.

Folia cæna, fragilia.

2. CATANANCHE lutea. G. squamis calycinis inferioribus lanceolatis. monsp.

C. lutea, longo nervoso dematæque folio. Vahl. ed. 1721. p. 216.

Chondrilla cyanoides lutea, coronopæ folio non diviso. Boerh. mus. 2. p. 21. t. 7. Barr. ic. 1135.

Stoebe plantaginis folio. Ald. exot. 287. t. 286.

Habitat in Creta. ☉

322 SYNGENESIA. POLYGAMIA. EQUALIS.

H. foliis semipinnatis hirsutis, caule nudo, brachiatō. Hall. helu. n. 3.

Achyrophorus radicans. Scop. carn.

Hieracium dens leonis folio obtuso, majus. Bauh. pin. 127. Moris. hist. 3. p. 66. f. 7. t. 4. f. 27.

Hieracium minus, dentis leonis folio subaspero. Bauh. pin. 127. (Vaill.)

Hieracium, dentis leonis folio, bulbosum. Bauh. pin. 127. prod. 63.

Hieracium longius radicans. Leb. ic. 238. Rel. hist. 230.

Hieracium tertium. Dod. pempt. 639.

Hieracium VII. Tabern. hist. 492.

Habitat in Europæ cultis pastuis.

Pedunculi superne incrassati. Corollæ magnæ. Calycis squamæ carina ciliata.

998. L A P S A N A.

Recept. nudum. Cal. calyculatus squamis singulis interioribus canaliculatis.

1. *LAPSANA communis. L. calycibus fructus angulatis, pedunculis tenuibus ramulosis. Junc. parif. gallob. carn. pal. fl. Fl. dan. 1. 500. Kniph. cat. 3. n. 48. Rayn. bot. lugd. lith. burg.*

L. caule brachiatō, foliis ovatis, longe petiolatis, petiolo pinnatis. Hall. helu. n. 6.

+ *Sonchus sylvaticus, 1. 2. 3. Tabern. 192. 193. Hall. R. Lampfana. Dod. pempt. 675.*

Sonchus affinis Lapsana domestica. Bauh. pin. 124.

Habitat in Europæ cultis.

2. *LAPSANA Zacintha. L. calycibus fructus torulosis depressis obtusis sessilibus, monsp.*

L. calycibus fructus subrotundis angularis; pedunculis incrassatis, floribus erectis. Hort. cliff.

Chondrilla verrucata, foliis cichorei viridibus. Bauh. pin. 130.

Cichorium verrucatum, zacintha. Clus. hist. 2. p. 144.

Habitat in Italia, Oriente. ☉

Folia Radicalia lyrata, acuta; Caulina sagittata, amplexicausta, dentata. Caulis subdichotomus, striatus, rigidulus. Flores sessiles, impuberes penduli. Corollæ subina-

SYNGENESIA. POLYGAMIA ÆQUALIS. 323

fulvæ, supra flavæ. Calycis exterior ordo erectus, interior basi gibbus, rufescens. Pedunculi incrassati.

3. **LAPSANA stellata** L. calycibus fructus undique patentibus: radiis subulatis, foliis caulinis lanceolatis indivisis; *Hort. upf. delph.*

Hieracium filiqua falcata. Bauh. pin. 128.

Hieracium stellatum. Bauh. hist. 2. p. 1014.

Habitat Monspeli, Bononiæ. ☉

Variat foliis sinuatis, dentatis, indivisis. Gerard.

4. **LAPSANA Rhagadiolus** L. calycibus fructus undique patentibus: radiis subulatis, foliis lyratis. *Hort. upf.* Kniph. cent. 4. n. 36. delph.*

Lampfana Rhagadiolus. Scop. carn.

Hieracium falcatum alterum. Rai. hist. 256.

Rhagadiolus lapsanæ foliis. Tournef. cor. 36.

Rhagadiolus edulis hieraciis affinis. Bauh. hist. 2. p. 1014.

Habitat in Oriente, Carniolia.

Varietas antecedentis. Gerard, sed forte constans.

999. C A T A N A N C H E.

Recept. paleaceum. Cal. imbricatus. Pappus aristatus pappo 5. seto.

1. **CATANANCHE cærulea** C. squamis calycinis inferioribus ovatis. *lugdb. monsp. delph.*

Catananche cærulea, semiflosculorum ordine simplici. Vaill. ad. 1721. p. 215.

Chondrilla cærulea, cyani capitulo. Bauh. pin. 430.

Chondrillæ species tertia. Dod. pempt. 638.

β. Catananche flore pleno cæruleo. Tournef. inst. 478.

Chondrilla cærulea, cyani capitulo, altera. Bauh. pin. 130.

Habitat in G. Narbonensis collibus saxosis. 24

Folia cana, fragilia.

2. **CATANANCHE lutea** C. squamis calycinis inferioribus lanceolatis. *monsp.*

C. lutea, longo nervoso dentatoque folio. Vaill. ad. 2721. p. 216.

Chondrilla cyanoides lutea, coronopi folio non diviso. Bosc. mus. 2. p. 21. t. 7. Barr. ic. 1135.

Stœbe plantaginis folio. Ald. exot. 287. t. 286.

Habitat in Creta. ☉

322 SYNGENESIA. POLYGAMIA. AQUALIS.

H. foliis semipinnatis hirsutis, caule nudo, brachiato. Hall. helv. n. 3.

Achyropterus radicans. Scop. carn.

Hieracium dentis leonis folio obtuso, majus. Bauh. pin. 127. Moris. hist. 3. p. 66. f. 7. t. 4. f. 27.

Hieracium minus, dentis leonis folio subsperio. Bauh. pin. 127. (Vaill.)

Hieracium, dentis leonis folio, bulbosum. Bauh. pin. 127. prod. 63.

Hieracium longius radicans. Lob. ic. 238. Rai. hist. 230.

Hieracium tertium. Dod. pempt. 639.

Hieracium VII. Tabern. hist. 492.

Habitat in Europæ cultis pastuis. 24.

Pedunculi superne incrassati. Corollæ magnæ. Calycis squamis terminis ciliatis.

998. LAPSANA.

Recept. nudum. Cal. calyculatus squamis singulis interioribus canaliculatis.

1. LAPSANA communis. L. calycibus fructus angulatis, pedunculis tenuibus ramulosis, jucc. parif. ged. gallab. carn. pal. fl. Fl. dan. 1. 300. Kniph. cent. 5. n. 48. Regn. bot. lugd. lich. burg.

2. L. caule brachiato, foliis ovatis, longe petiolatis, petiolo pinnatis. Hall. helv. n. 6.

3. Sonchus sylvaticus, 1. 2. 3. Tabern. 192. 193. Hall. R. Lamplana. Dod. pempt. 675.

4. Sonchus affinis Lapsana domestica. Bauh. pin. 124.

Habitat in Europæ cultis.

2. LAPSANA Zacintha. L. calycibus fructus corulosis depressis obtusis sessilibus, monsp.

L. calycibus fructus subrotundis angulatis, pedunculis incrassatis, floribus erectis. Hort. cliff.

Chondrilla verrucata, foliis cichorei viridibus. Bauh. pin. 130.

Cichorium verrucatum, zacintha. Clus. hist. 2. p. 144.

Habitat in Italia, Oriente. ☉

Folia Radicalia lyrata, acuta; Caulina sagittata, amplexicaulia, dentata. Caulis subdichotomus, striatus, rigidus. Flores sessiles, impuberes penduli. Corollæ subnuda.

SYNGENESIA POLYGAMIA EQUALIS. 323

fulva, supra *flava*. Calycis exterior ordo erectus, interior basi gibbus, rufescens. Pedunculi incrassati.

3. LAPSANA stellata, L. calycibus fructus undique patentibus: radiis subulatis, foliis caulinis lanceolatis indivisis. Hort. upf. delph.

Hieracium filiqua falcata. Bauh. pin. 128.

Hieracium stellatum. Bauh. hist. 2. p. 1014.

Habitat Monspeli, Bononiæ. ☉

Variet folius sinuatis, dentatis, indivisis. Gerard.

4. LAPSANA Rhagadiolus. L. calycibus fructus undique patentibus: radiis subulatis, foliis lyratis. Hort. upf. Kmph. cent. 4. n. 36. delph.

Lampfana Rhagadiolus. Scop. carn.

Hieracium falcatum alterum. Rai. hist. 256.

Rhagadiolus lapsanæ foliis. Tournesf. cor. 36.

Rhagadiolus edulis hieraciis affinis. Bauh. hist. 2. p. 1014.

Habitat in Oriente, Carniola.

Varietas antecedentis. Gerard, sed forte constans.

999. CATANANCHE.

Recept. palæstenum. Cal. imbricatus. Pappus aristatus pappo s. seto.

1. CATANANCHE cerulea. C. squamis calycinis inferioribus ovatis. Lugdb. monsp. delph.

Catananche cerulea, semiosculorum ordine simplici. Vaill. ad. 1721. p. 215.

Chondrilla cerulea, cyani capitulo. Bauh. pin. 490.

Chondrilla species tertia. Dod. pempt. 638.

β. Catananche flore pleno ceruleo. Tournesf. inst. 478.

Chondrilla cerulea, cyani capitulo, altera. Bauh. pin. 490.

Habitat in G. Narbonensis collibus saxosis. 24.

Folia cana, fragilia.

2. CATANANCHE lutea. C. squamis calycinis inferioribus lanceolatis. monsp.

C. lutea, longo nervosa dentatæque folio. Vaill. ad. 1721. p. 216.

Chondrilla cyanoides lutea, coronopii folio non diviso. Boreus mus. 2. p. 21. t. 7. Barr. ic. 1135.

Stoebe plantaginis folio. Ald. exot. 287. 2. 286.

Habitat in Creta. ☉

324 SYNGENESIA. POLYGAMIA ÆQUALIS.

Flores inferiores sessiles ad radicem, vix explicandi, magis fertiles. Folia viridia, flexilia, trinervia.

1000. CICHORIUM.

Recept. subpaleaceum. Calyx calyculatus. Pappus sub 5. dentatus, obsolete pilosus.

1. CICHORIUM *Intybus*. C. floribus geminis sessilibus, foliis runcinatis. *Fl. suec. parif. gallob. pal. sil. carn. Knorr. del. 1. t. W. 9. lithu. lugd. burg.*

C. foliis pinnatis, pinnis triangularibus dentatis, floribus sessilibus. *Hall. helu. n. 1.*

C. caule simplici. *Hort. cliff.*

C. sylvestre s. officinarum. *Bauh. pin. 126. Blackw. t. 183.*

Intybus sylvestris. Cam. epit. 285. Fuchsf. hist. 979.

Habitat in Europa ad margines agrorum viarumque. 2.

Variat flore albo; rubro. Hortensem varietatem pingit *Blackw. t. 177. R.*

2. CICHORIUM *Endivia*. C. floribus solitariis pedunculatis, foliis integris crenatis. *Regn. bot.*

Cichorium latifolium s. Endivia vulgaris. Bauh. pin. 125.

Intybum sativum. Dod. pempt. 634.

β. *Endivia crispa. Bauh. pin. 125.*

C. (*crispum*) floribus solitariis pedunculatis, foliis sim-
briatis crispis. *Mill. dict. n. 4. R.*

Habitat - - - 4.

Obs. Cichorium pumilum. Jacq. obs. 4. t. 80. huic affine.

3. CICHORIUM *spinosum*. C. caule dichotomo spinoso, floribus axillaribus sessilibus. *Kniph. cent. 6. n. 29.*

C. spinosum. *Bauh. pin. 126. prodr. 62. t. 62.*

Chondrilla genus elegans, cæruleo flore. *Clus. hist. 2. p. 145.*

Habitat in Cretæ, Siciliæ collibus arenosis maritimis. ♂

1001. SCOLYMUS.

Recept. paleaceum. Calyx imbricatus, spinosus. Pappus nullus.

1. SCOLYMUS *maculatus*. S. floribus solitariis. *delph.*

S. foliis margine attenuatis. *monsp. carn. Kniph. cent. 9. n. 85.*

SYNGENESIA. POLYGAMIA ÆQUALIS. 315

S. annuus. Hort. ups. 244.

S. Chrysanthemus annuus. Dòdart. mem. 4. p. 303.

S. Theophrasti Narbonensis. Clus. hist. 2. p. 153.

Habitat in G. Narbonensi, Italia. ☉

Herba annua radice tenui. Caulis inferne ramosior. Folia glabra, nitida, margine cartilaginea, decurrentia fere ad basin caulis. Flores e divaricaturis solitarii. Corolla minor Antheris fuscis.

2. *SCOLYMUS hispanicus. S. floribus congestis. delph.*

S. foliis margine incrassatis. Mill. dict. n. 2. & & ic. t. 229.

S. vivax. Sauv. monsp.

S. Chrysanthemos. Bauh. pin. 384.

S. Theophrasti hispanicus. Clus. hist. 2. p. 153.

Carduus Chrysanthemos. Dod. pempt. 725.

Habitat in Italia, Sicilia, G. Narbonensi. 24

Herba altero anno florens, radice fusiformi. Caulis inferne minus ramosus. Folia subscabra, cinerascens, margine non cartilaginea, decurrentia, sed desinens ante proximum folium. Flores saepius 4, congesti. Corolla duplo major antheris concoloribus.

1002. A R C T I U M.

Cal. globosus: squamis apice hamis inflexis.

1. *ARCTIUM Lappa. A. foliis cordatis inermibus petiolatis. gallob. carn. pal. sil. Fl. dan. t. 642. Ludw. ed. t. 106.*

Kniph. cent. 3. n. 13. Regn. bot. lugd. lithu. burg. Fl. suec. paris.

Lappa. Hall. helv. n. 161.

Personata s. Lappa major altera. Matth. 1155. Hall. R.

Lappa major s. Arctium dioscoridis. Bauh. pin. 198.

Bardana s. Lappa major. Dod. pempt. 58. Blackw. t. 117.

Personata. Cam. epis. 887.

2. *Lappa major montana, capitulis tomentosis. Bauh. pin. 198. Mill. dict. n. 3.*

Habitat in Europæ cultis ruderatis. ♂

Datur varietas albo flore. R.

3. *ARCTIUM personata. A. foliis decurrentibus ciliato-spinosis: radicalibus pinnatis: caulinis oblongo-ovatis. delph.*

Carduus inermis, foliis imis pinnatis, superioribus ovato-

226 SYNGENESIA. POLYGAMIA ÆQUALIS.

lanceolatis, serratis, squamis calycinis reflexis. Hall. helu. n. 162.

Carduus Personata. Jacq. aust. t. 348.

Carduus mollis, foliis imis laciniatis, superioribus integris, squamis calycinis reflexis. Hall. helu. ed. 1. 678. t. 19. opusc. 232.

Carduus mollis latifolius, lappæ capitulis. Bauh. pin. 377. prodr. 155. Burf. XXI. 13.

Carduus capitulis in vertice congestis lanigeris: squamis reflexis. Gmel. sib. 2. p. 62. t. 24.

Cirsium majus caule elato polycephalon, capitulis majoribus personatæ æmulis. Rai syl. 399. Hall. R.

Habitat in Alpibus Helveticæ, Genevæ, Taureræ, Austriae. ♂

Folia caulina ovato-oblonga, subtus tomentosa, serrata; serraturis spinulosis, nec non spinulis minimis interjectis.

Folia hæc, basi quamvis attenuata, decurrunt per caulem tenuissimis marginibus spinosis. Pedunculi tomentosi & marginibus spinulosis armati. Habitus antecedentis elucet in textura foliorum, panicula, calycis figura, magnitudine.

Cardui species.

Squamæ calycis non desinunt in aculeos hamatos. Hall. R.

3. *ARCTIUM carduelis*, A. foliis pinnatifidis aculeatis.

Cirsium arctioides. Scop. carn. ed. 2. t. 53.

Habitat in Alpibus Carniolæ superioris. D. Scopoli.

Statura *Serratulæ arvensis*. Caulis crispus, aculeatus. Folia amplexicaulia, pinnatifida, laciniata, margine aculeata. Calyces terminales, pedunculati: squamis lineari-setaceis, patentibus incurvis.

1003. SERRATULA.

Cal. subcylindricus, imbricatus muticis.

1. *SERRATULA tinctoria*. S. foliis lyrato-pinnatifidis; pinnæ terminali maxima, flosculis conformibus. Fl. suec. paris. pal. fil. Fl. dan. t. 281. Kniph. cent. 2. n. 83. luth. delph. lugd. burg.

Carduus inermis, foliis glabris serratis, imis ovatis, superioribus semipinnatis. Hall. helu. n. 163.

Carduus tinctorius. Scop. carn.

Serratula. Bauh. pin. 235. Dod. pempt. 24.

Habitat in Europæ borealis pratis. 26

SYNGENESIA. POLYGAMIA ÆQUALIS. 327.

Folia ciliata. Receptaculum *setosum*.

Variat 1) *foliis omnibus integris*. 2) *Foliis infer. integris, superioribus incis.* 3) *Foliis omnibus incis.* 4) *Foliis omnibus laciniatis*. *Hæc varietas est Serratula alpina. Zanon. nov. t. 94. Hall. Datur etiam varietas albo flore. R.*

2. **SERRATULA coronata.** S. foliis lyrato - pinnatifidis : pinna terminali maxima, flosculis radii femineis longioribus.

Serratula præalta centauroides montana. Bocc. mus. 2. p. 45. t. 37.

Carduus inermis, foliis glabris pinnatis : lacinia extima maxima, capitulis squalidis. Gmel. sib. 2. p. 49. t. 20.

Serratula foliis laciniatis. Amm. ruth. n. 181.

Habitat in Sibiria, Italia. 24

Simillima S. tinctoriæ, ut vix distinguatur, nisi Magnitudine tripla; Caule sulcato; Flosculis duplo pluribus; Foliolo declinato sub calyce; sed imprimis Radio flosculis tubulosis, angustioribus, femineis, fertilibus.

Genus ambiguum: Radius femineus indicat Centaureæ genus, sed flosculi radii, æque ac disci, seminiferi, una cum habitu hoc negant. Si plures similes innotescant, novum exposcant genus, quod ab hac unicam vix meretur.

3. **SERRATULA alpina.** S. calycibus subhirsutis ovatis, foliis indivisis. *norv. delph.*

S. foliis petiolatis : radicalibus ovato - oblongis dentatis : caulinis integerrimis. *Fl. suec.*

S. foliis ovato-lanceolatis : radicalibus serratis, caule thyr-sifloro. *Hort. ups.*

Cirsium foliis ovato-lanceolatis, dentatis, subtus tomentosis, calycibus villosis. Hall. helv. n. 179. t. 6.

Cirsium inerme, foliis ex ovato-lanceolatis denticulatis intralanugine candidis. Gmel. sib. 2. p. 67. t. 26.

Carduo-Cirsium minus britannicum, floribus congestis. Pluk. alm. 83. t. 154. f. 3.

β. *Serratula cynoglossifolia.*

Cirsium humile montanum, cynoglossi folio, polyanthum. Moris. hist. 3. p. 148. Dill. elch. 82. t. 70. f. 81. vide Gmel. sib. 2. p. 76. t. 32. Flor. dan. t. 37. Hall. l. c. β.

γ. *Serratula lapathifolia.*

Cirsium polyanthemum, molli hastato folio. Moris. hist. 3. p. 148. f. 7. t. 29. f. 1. t. 22.

Cirsium foliis triangularibus lunate dentatis. Hall. helv. 683.

318 SYNGENESIA. POLYGAMIA EQUALIS.

Carduus mollis, lapathi foliis. *Bauh. pin.* 377.

Carduus mollior. 2. *Clus. hist.* 2. p. 151.

♂. *Serratula angustifolia*.

Cirsium inerme, foliis linearibus utrinque viridibus, calycibus hirsutis. *Gmel. sib.* 2. p. 78. t. 33. *Hall. l. c. var.* 7.

Habitat in Alpibus Lapponiæ, Austriæ, Helvetiæ, Arvonæ, Sibiræ. 24

Planta floribus coruleis distinctissima, foliorum figura diversissima; sed quæ causa produxerit hæc constantes varietates? certe locus non suffecerit.

14. *SERRATULA arvensis.* S. foliis dentatis spinosis. *Spec. paris. herb. pal. sil. Fl. dan.* t. 644. *Kniph. cent.* 6. n. 85. *Regn. bot. lugd. lithu. burg.*

Cirsium caule anguloso, ramoso, foliis semipinnatis, pinnis angulosis, spinosis, calycibus inermibus, cylindricis. *Hall. helv.* n. 171.

Cirsium foliis lanceolatis dentatis spinosis; calycibus squamis tomentosis, inermibus aristatis. *Scop. carn.*

Carduus radice repente, foliis lanceolatis dentatis margine aculeatis. *Fl. lapp.*

Carduus vinearum repens, sonchi folio. *Bauh. pin.* 387. *prodr.* 156.

Carduus in avena proveniens. *Bauh. pin.* 377.

Carduus serpens leuicaulis. *Bauh. hist.* 3. p. 49. R.

Carduus arvensis. *Tabern.* p. 700. R.

Carduus (serratuloides) foliis semiamplexicaulibus decurrentibus sinuato-pinnatifidis, calycibus inermibus globoso-conicis. *Neck. gallob.* p. 338. n. 9.

Ceanothus theophrasti. *Col. ecphr.* I. p. 45. t. 45.

♂. *Cirsium arvense, sonchi folio, radice repente, caule tuberoso.* *Tournef. inst.* 448.

Habitat in Europæ cultis agris. 24

Habitus Cardui est.

1004. C A R D U U S.

Cal. ovatus, imbricatus squamis spinosis. Receptac. pilosum.

* *Foliis decurrentibus.*

1. *CARDUUS leucographus.* C. foliis decurrentibus dentatis spinosis; pedunculis nudis longissimis unifloris, calycibus spinosis inclinatis.

SYNGENESIA. POLYGAMIA ÆQUALIS. 329.

C. foliis sinuatis aculeatis, caule crispo, pedunculis unifloris. *Roy. lugdb.*

Cirsium maculis argenteis notatum. *Tournef. inst. 448.*

Carduus leucographus f. mariae hirsutus, capitulo minore. *Morif. bles. 244.*

Habitat in Campania.

2. CARDUUS lanceolatus. C. foliis decurrentibus pinnatifidis hispidis: laciniis divaricatis, calycibus ovatis spinosis villosis, caule piloso. *gallob. herb. pal. sil. lithua. lugd. delph. burg.*

C. foliis lanceolatis decurrentibus: denticulis superficieque spinosis. *Fl. suec. paris.*

Cirsium foliis semipinnatis, pinnis bilobatis, spinosis capitulis congestis, tomentosis, conicis. *Hall. helu. n. 169.*

Cirsium foliis spinosis, decurrentibus, pinnatifidis; laciniis alternis erectis; calycibus reticulato-ovatis. *Scop. carn.*

Carduus lanceolatus. *Tabern. 699. Hall. R.*

C. lanceolatus latifolius. *Bauh. pin. 385.*

C. lanceolatus f. sylvestris dodonæi. *Bauh. hist. 3. p. 58.*

Habitat in Europæ cultis ruderalis. ♂

Calycis intimæ squamæ inermes, setaceæ, exaridæ.

3. CARDUUS nutans. C. foliis semidecurrentibus spinosis, floribus cernuis squamis calycinis superne patentibus. *Fl. suec. stamp. paris. carn. gallob. pal. sil. Flor. dan. 675. lugd. delph. lithua. lugd.*

C. foliis sinuatis decurrentibus margine spinosis, floribus solitariis nutantibus. *Hort. cliff.*

C. foliis semipinnatis, pinnis angulosis, spinosis, floribus nutantibus, calycinis squamis pungentibus patulis. *Hall. helu. n. 167.*

C. nutans. *Bauh. hist. 3. p. 56. It. gotl. 293. **

C. spinosissimus latifolius sphærocephalus vulgaris. *Bauh. pin. 385.*

Habitat in Europa ad pagos. ♂

Styli reflexi versus latera.

Varietas albo flore reperitur. R.

4. CARDUUS acanthoides. C. foliis decurrentibus sinuatis margine spinosis, calycibus pedunculatis solitariis erectis villosis. *Fl. suec. gallob. pal. sil. Jacq. austr. 1. 249. lugd. delph. burg.*

C. foliis laciniatis: subtus tomentosis ex margine spinosis, capitulis sessilibus congestis aculeatis. *Dalib. paris.*

330 SYNGENESIA. POLYGAMIA ÆQUALIS.

C. foliis ovatis & semipinnatis, subtus tomentosis, spinosis, calycibus molliter spinosis. *Hall. helu. n. 118. sec. Jacq.*

C. polyacanthos. *Schreb. Spicil. p. 15.*

C. acanthoides. *Bauh. hist. 3. p. 59. It. scan. 297. **

C. spinosissimus, capitulis minoribus. *Rai. angl. 3. p. 194.*

C. polyacanthos, capitulis longioribus & tenuioribus foliis albicantibus. *Morif. hist. 3. p. 153.*

Habitat in Europæ ruderatis. ☉

Differt a Carduo nutante calycibus erectis, a C. crispo calycibus solitariis villosis. Calycis squamæ interiores patentæ. Corollulæ semiquinquesidæ, at infima lacinia fere ad basin separata, ut fere bilabiata: labio superiore 4. fido: inferiore simplici, quod non in Sp. 3. 5. Styli flosculis duplo longiores.

5. *CARDUUS crispus. C.* foliis decurrentibus sinuatis margine spinosis, floribus aggregatis terminalibus, squamis calycinis inermibus subaristatis patulis. *suéc. paris. pal. sil. gallob. Flor. dan. t. 621. lith. lugd. burg.*

C. foliis caulinis latissimis, foliis semipinnatis, pinnis angulosis spinosis, floribus fasciculatis oblongis. *Hall. helu. n. 166. R.*

C. spinosissimus angustifolius vulgaris. *Læff. pruss. 34. t. 5.*

C. caule crispo. *Bauh. hist. 3. p. 59. •*

Habitat in Europæ septentrionalioris agris, cultis. ☉

Caulis viridis. Folia subtus ex albido tomentosa, totaliter decurrentia, mitia, in caule subpungentia. Pedicelli albotomentosi. Calyx squarrosus, magnitudine Avellanæ, squamis supra rugosis.

Synonyma Læselii & Bauh. ad præcedentem plantam amendant Schreberus & Jacquinus. R.

6. *CARDUUS polyanthemus. C.* foliis decurrentibus sinuatis ciliatis subtus nudis, floribus pedunculatis congestis.

C. pycnocephalus palustris. *Triumf. obs. 103. t. 103.*

Cirsium palustre lanceolatum, alato caule, polyanthemum. Vaill. ad. 160.

Habitat Romæ. ♂

Simillimus C. crispo, ut ab eo difficile distinguatur, sed neque petioli, neque folia subtus tomentosa. Caulis C. crispæ altior. Folia profunde runcinata, non rigida, obtusiuscula, ciliata spinis inermibus, excavato-venosa, subtus pallidiora, non vero alba. Caulina superiora lanceolata,

SYNGENESIA. POLYGAMIA ÆQUALIS. 338

erosa, decurrentia totaliter, crispa parumque pungentia. Pedunculi crispī, vix arachnoidei. Flores sæpius 3. s. 4. ad apicem subsessiles. Corollæ in Limbo purpureæ, in disco incarnatæ. Antheræ violaceæ. Styli albi Stigmata. purpurea. H. V.

7. **CARDUUS palustris.** C. foliis decurrentibus dentatis : margine spinosis, floribus racemosis erectis, pedunculis inermibus. *Gmel. sib. 2. p. 57. t. 23. f. 2. gallob. pal. herb. Kniph. cent. 6. n. 21. lithu. delph. lugd. burg.*

Cirsium caule recto, subnudo, alato, foliis semipinnatis, spinosis, calycibus molliter spinosis. *Hall. helu. n. 170.*

Cirsium foliis spinosis, decurrentibus, dentatis; floribus erectis, aggregatis; calycibus inermibus. *Scop. carn.*

C. foliis linearibus dentatis spinosissimis, caule crispo spinoso, floribus erectis. *Fl. suec. paris.*

C. palustris. *Bauh. pin. 377. prodr. 156.*

C. spinosissimus erectus angustifolius palustris. *Morif. hist. 3. p. 153. f. 7. t. 32. f. 13.*

C. polycephalus angustifolius. *Dalech. hist. 1743.*

C. lanceolatus angustifolius. *Bauh. prodr. 155.*

Habitat in Europæ pratis subpaludosis. 24

Calyx arde imbricatus, lævis, mucronibus vix conspicuis. Albis floribus variat.

8. **CARDUUS pycnocephalus.** C. foliis decurrentibus pinnatifido-sinuatis pubescentibus spinosis, pedunculis nudis tomentosis, calycibus deciduis. *Jacq. hort. t. 44.*

C. foliis spinosis lanceolatis utrinque tomentosis, radicalibus sinuato-pinnatifidis, caulinis semidecurrentibus sinuatis, pedunculis capitato-multifloris tomentosis inermibus. *Gouan. illustr.*

C. nemorosus italicus. *Barr. obs. 925. t. 417.*

C. pycnocephalus sylvestris. *Triumph. obs. 100. i. 101.*

Habitat in Europa australi. 24

Planta quasi media inter C. crispum & Onopordon. Caulis pedalis s. altior, albido-pilosus, crispus, totaliter foliorum marginibus utrinque decurrentibus, alternatim flexuosis spinosisque. Folia Radicalia sublanceolata, subsinuata; Caulina oblonga, semipinnatifida, dentata, utrinque pubescentia, subtus albidiora, tomentosa s. sublanata. Pedunculi breves, tomentoso-albi. Pedicelli floribus 3 s. 4. Flosculis sæpius 3. s. 4. Calyces magnitudine avellanæ, oblongi,

330 SYNGENESIA. POLYGAMIA AQUALIS.

C. foliis ovatis & semipinnatis, subtus tomentosis, spinosis, calycibus molliter spinosis. Hall. helv. n. 115. sec. Jacq.

C. polyacanthos. Schreb. Spicil. p. 15.

*C. acanthoides. Bauh. hist. 3. p. 59. lt. scan. 297. **

C. spinosissimus, capitulis minoribus. Rai. angl. 3. p. 194.

C. polyacanthos, capitulis longioribus & tenuioribus foliis albicantibus. Moris. hist. 3. p. 153.

Habitat in Europæ ruderatis. ☉

Differt a Carduo nutante calycibus erectis, a C. crispo calycibus solitariis villosis. Calycis squamæ interiores patentiores. Corollulae semiquinquefidæ, at infima lacinia fere ad basin separata, ut fere bilabiata: labio superiore 4. fido: inferiore simplici, quod non in Sp. 3. 5. Styli strobilulis duplo longiores.

5. **CARDUUS crispus.** *C. foliis decurrentibus sinuatis margine spinosis, floribus aggregatis terminalibus, squamis calycinis inermibus subaristatis patulis. succ. parif. pat. gallob. Flor. dan. t. 621. lith. lugd. burg.*

C. foliis caulinis latissimis, foliis semipinnatis, pinnis angustatis spinosis, floribus fasciculatis oblongis. Hall. helv. n. 166. R.

C. spinosissimus angustifolius vulgaris. Less. pruss. 34. t. 1.

C. caule crispo. Bauh. hist. 3. p. 59. •

Habitat in Europæ septentrionalioris agris, cultis. ☉

Caulis viridis. Folia subtus ex albido tomentosa, totaliter decurrentia, mitia, in caule subpungentia. Pedicelli albi tomentosi. Calyx squarrosus, magnitudine Avellanae, squamis supra rugosis.

Synonyma Læselii & Bauh. ad præcedentem plantam amendant Schreberus & Jacquinus. R.

6. **CARDUUS polyanthemus.** *C. foliis decurrentibus sinuatis ciliatis subtus nudis, floribus pedunculatis congestis.*

C. pycnocephalus palustris. Triumph. obs. 103. t. 103.

Cirsium palustre lanceolatum, alato caule, polyanthemum. Vaill. ad. 160.

Habitat Romæ. ♂

Simillimus C. crispo, ut ab eo difficile distinguatur, sed neque petioli, neque folia subtus tomentosa. Caulis C. crispus altior. Folia profunde runcinata, non rigida, obtusiuscula, ciliata spinis inermibus, excavato-venosa, subius pallidiora, non vero alba. Caulina superiora lanceolata,

SYNGENESIA. POLYGAMIA ÆQUALIS. 338

erosa, decurrentia totaliter, crispa parumque pungentia. Pedunculi crispī, vix arachnoidei. Flores sæpius 3. s. 4. ad apicem subsessiles. Corollæ in Limbo purpureæ, in disco incarnatæ. Antheræ violaceæ. Styli albi Stigmata purpurea. H. V.

7. CARDUUS palustris. C. foliis decurrentibus dentatis : margine spinosis, floribus racemosis erectis ; pedunculis inermibus. *Gmel. sib. 2. p. 57. t. 23. f. 2. gallob. pal. herb. Kniph. cent. 6. n. 21. lithu. delph. lugd. burg.*

Cirsium caule recto, subnudo, alato, foliis semipinnatis, spinosis, calycibus molliter spinosis. Hall. helu. n. 170.

Cirsium foliis spinosis, decurrentibus, dentatis ; floribus erectis, aggregatis ; calycibus inermibus. Scop. carn.

C. foliis linearibus dentatis spinosissimis, caule crispo spinoso, floribus erectis. Fl. suec. paris.

C. palustris. Bauh. pin. 377. prodr. 156.

C. spinosissimus erectus angustifolius palustris. Moris. hist. 3. p. 153. f. 7. t. 32. f. 13.

C. polycephalus angustifolius. Dalech. hist. 1743.

C. lanceolatus angustifolius. Bauh. prodr. 155.

Habitat in Europæ pratis subpaludosis. 24

Calyx arcæ imbricatus, lævis, mucronibus vix conspicuis. Albis floribus variat.

8. CARDUUS pycnocephalus. C. foliis decurrentibus pinnatifido - sinuatis pubescentibus spinosis, pedunculis nudis tomentosis, calycibus deciduis. *Jacq. hort. t. 44.*

C. foliis spinosis lanceolatis utrinque tomentosis, radicalibus sinuato-pinnatifidis, caulinis semidecurrentibus sinuatis, pedunculis capitato - multifloris tomentosis inermibus. Gouan. illustr.

C. nemorosus italicus. Barr. obs. 925. t. 417.

C. pycnocephalus sylvestris. Triumph. obs. 100. i. 101.

Habitat in Europa australi. 24

Planta quasi media inter C. crispum & Onopordon. Caulis pedalis s. altior, albido-pilosus, crispus, totaliter foliorum marginibus utrinque decurrentibus, alternatim flexuosis spinosisque. Folia Radicalia sublanceolata, subsinuata ; Caulina oblonga, semipinnatifida, dentata, utrinque pubescentia, subtus albidiora, tomentosa s. sublanata. Pedunculi breves, tomentoso-albi. Pedicelli floribus 3 s. 4. Flosculis sæpius 3. s. 4. Calyces magnitudine avellanæ, oblongi,

332 SYNGENESIA. POLYGAMIA. ÆQUALIS.

imbricati squamis subulatis, erectis, patulis, subspinosis.
 Corolla purpurea limbo erecto, magnitudine Serratula tinctoria. Differt a C. acanthoide pedunculis nudis nec alatis. Calyces maturi toti decidunt.

9. **CARDUUS argentatus.** C. foliis decurrentibus runcinatis spinosis, pedunculis subtomentosis unifloris, calycibus ovatis mucronatis inermibus. Jacq. hort. t. 192.

Cirsium ægyptiacum minus, folius ad limbum argenteis.
 Grange.

Habitat in Ægypto. ☉

Caulis pedalis, flexuosus, lævis, alterne ramosus. Folia totaliter decurrentia margine flexuoso spinoso, dentata spinis pungentibus, adpersa præsertim versus sinus maculis argenteo-niveis. Pedunculi terminales, longitudine totius plantæ uniflori, subaphylli, tomentosi erecti. Calyx ovatus, inermis, magnitudine Avellanae: Squamis arcte imbricatis, mucronatis: apicibus subulatis, inermibus erectis; maturus totus decidit cum fructu. Corolla minima, purpurea mixta pappo, vix dehiscens. Antherae caeruleae. Pappus cinereus.

Affinis C. pycnocephali foliis lacteo-maculatis, pedunculis unifloris. Figura proxima Hamb. Rauw. ii. c. 295.

Cur plantæ Ægyptiacæ plures maculis albis adspargantur ut Carduus marianus, syriacus, peregrinus, Centaurea galactiter, Asperugo Ægyptiaca.

10. **CARDUUS dissectus.** C. foliis decurrentibus lanceolatis, denticulis inermibus, calycibus spinosis. *parif.*

Cirsium anglicum. Lob. ic. 583. Dalech. hist. 584. figura tenuis.

Cirsium majus, singulari capitulo magno, s. incanum variegatum. Bauh. pin. 377.

Habitat in Anglia, Gallia. 24

Folia carent omnino aculeis, diversus itaque a C. heterophyllo & helenioides.

12. **CARDUUS canus.** C. foliis decurrentibus lanceolatis eroso-dentatis ciliato-aculeatis: utrinque arachnoideo-subvillosis. Jacq. austr. I. t. 42. & 43.

Cirsium montanum maximum. Rai. hist. 305.

Cirsium tomentosum radice bulbosa. Bauh. hist. 3. p. 44.

Carduus tuberosus. Jacq. vind. 280. secund. descript.

Habitat in Austria. 24

Radices fusiformes, aggregatae, carnosae, albae (Sisari.)
 Caulis quadripedalis, viridis, angulatus, arachnoidens.

SYNGENESTA. POLYGAMIA ÆQUALIS. 333

Folia lanceolata, subsinuato-dentata, spinulis albis vix pungentibus ciliata, utrinque tomento arachnoideo cana, carinata, in totum decurrentia, alis exstantibus. Pedunculi terminales, elongati, foliis parvis floralibus viridibus. Flores solitarii, purpurei. Calyces inermes squamis extus linea alba. H. V.

14. CARDUUS defloratus. C. foliis decurrentibus lanceolatis serratis subspinoso-ciliatis nudis, pedunculis longissimis lanuginosis unifloris. *Jacq. austr. t. 89. sub Cirsio. delph.*

C. inermis, foliis ciliatis, scapo nudo paucifloro. *Hall. helv. ed. 2. n. 164. t. 4.*

C. foliis rigidis molliter spinosis, scapo longo paucifloro. *Hall. helv. 677. t. 18. f. 2. opusc. 220.*

C. dictus cirsium folio laciniato nigrius. *Bauh. hist. 3. p. 43.*

Cirsium f. Carduus angustifolius. *Bauh. pin. 377. prodr. 155.*

Cirsium singularibus capitulis parvis. *Bauh. pin. 377.*

Cirsium 3. montanum 1. *Clus. hist. p. 149.*

Habitat in Helvetiæ, Austriæ, Monspeli subalpinis. 24

Similis C. serratuloidi, sed Calyces angustiores; Folia decurrentia. Fructus nutans non est.

Flos, quando adolevit, sæpe pendulus. Variat caule toto ramoso & foliis adeo spinosis, ut Carlinam imitentur. Variat etiam albis floribus. Hall. R.

15. CARDUUS monspessulanus. C. foliis decurrentibus lanceolatis subrepandis glabris inæqualiter ciliatis, pedunculis alternis calycibus inermibus. *delph.*

C. foliis lanceolatis glabris integris ciliatis decurrentibus, calycibus inermibus. *Sauv. monsp. 79.*

Carduus Cirsium monspelianum, folio longo glabro. *Bauh. hist. 3. p. 44.*

Cirsium foliis non hirsutis, floribus comp. *Bauh. pin. 377.*

Cirsion. *Dod. pempt. 377.*

Habitat Monspeli.

Habitus C. Serratuloides. Folia lanceolata, utrinque glabra, glauca, integra: radicalia vero repanda, margine spinis setaceis, inæqualibus ciliata. Capitula nonnulla in summitate caulis pedunculis brevibus.

16. CARDUUS tuberosus. C. foliis petiolatis subdecurrentibus subpinnatifidis spinosis, caule inermi, floribus solitariis. *pal. fil. delph. burg.*

Cirsium radicibus bulbosis, foliis semipinnatis, pinnis angu-

334 SYNGENESIA. POLYGAMIA ÆQUALIS.

lofis , ciliatis , floribus nudis. *Hall. helu. n.* 177. R.

Carduus foliis semipinnatis : pinnis semibilobis , capitulis inermibus singularibus. *Sauv. monsp.* 292.

C. pratensis , asphodeli radice , latifolius. *Bauh. pin.* 377. *Morif. hist.* 3. f. 7. t. 29. f. 27. 28.

C. bulbosus monspeliensium. *Lob. ic.* 10.

β. **C. pratensis** , asphodeli radice , foliis tenuiter incis. *Bauh. pin.* 377.

Acanthus sylvestris alter. *Lugdb.* 1444. *Hall.* R.

Jacea aculeata s. tucerosa. *Tabern. ic.* 154.

Habitat Monspeli , Lipsiæ , inque Bohemiæ , Austriæ , Helvetiæ inundatis.

Folia inermia , utrinque viridia , lævia. Radicalia pinnatifida , ceterum caulinis similia. Caulina cordato-lanceolata , dentata , ciliata spinulis inertibus , inæqualibus ; subdecurrentia. Flos *C. heterophylli* , sed minor.

17. **CARDUUS parviflorus**. *C.* foliis basi adnatis lanceolatis nudis eros ciliato-spinulosis inermibus.

Cirsium pratense alpinum polycephalum , asphodeli radice. *Tournef. inst.* 448.

Habitat in subalpinis australibus. 24

Caulis erectus. Folia adnata basi (vix decurrentia ,) lanceolata , repando-dentata , nuda , utrinque viridia , ciliata spinis inertibus , inæqualibus. Flores terminales , pedunculati , magnitudine *Serratulæ tinctoriæ* , albi , stylis longioribus. Calyx ovatus , imbricatus squamis carina nivea , apice spina inermi.

* * Foliis sessilibus.

18. **CARDUUS casabonæ**. *C.* foliis sessilibus lanceolatis integerrimis subtus tomentosis : margine spinis ternatis.

Acarna major , caule non folioso. *Bauh. pin.* 30.

Acarna theophrasti anguillariæ. *Lob. ic.* 486. f. 1. *Dalech. hist.* 1484.

Polyacanthus casabonæ , *Acarnæ* similis. *Bauh. hist.* 3. p. 92.

Habitat in Europa australi. ♂

20. **CARDUUS marianus**. *C.* foliis amplexicaulibus hastatopinnatifidis spinosis , calycibus aphyllis : spinis canaliculatis duplicato-spinosis. *monsp. gallob herb. sil. Ludw. ed.* t. 35. *Blackw.* t. 79. *Kniph. cent.* 1. n. 13. *Knorr. del.* 1. t. M. 3. *Regn. bot. lugd. burg.*

C. squamis calycinis margine apiceque spinosis. *Hort.* paris. monsp.*

SYNGENESIA. POLYGAMIA ÆQUALIS. 335

Silybum nervis foliorum albis. Hall. helu. n. 181.

Cirsium maculatum. Scop. carn.

Carduus albis maculis notatus vulgaris. Bauh. pin. 281.

Carduus mariæ. Dalech. hist. 1475.

Habitat in Angliæ, Galliæ, Italiæ, Germaniæ, &c. aggeribus ruderatis. ☉

21. CARDUUS syriacus. C. foliis amplexicaulibus angulato-spinosis, floribus solitariis subsessilibus obvallatis foliolis subquinis. *Hort. cliff. **

C. albis maculis notatus exoticus. Bauh. pin. 381.

C. latifolius echinos obsoletæ purpuræ ferens. Bauh. pin. 380.

C. luteus syriacus. Cam. hort. 35. t. 10.

Habitat in Syria, Creta, Hispania. ☉

Flores albi & ægyptiacæ varietas floribus purpureis. Forskæl.

22. CARDUUS eriophorus. Carduus foliis sessilibus bifariam pinnatifidis: laciniis alternis erectis, calycibus globosis villosis. *Hort. upf. * monsp. Mill. dict. n. 2. it. ic. t. 293. gallob. pal. Jacq. aust. t. 171. delph. lugd. burg.*

Cirsium foliis pinnatis, pinnis bilobatis, lanceolatis, spinosis, calycibus sphaericis, spinosis, tomentosis. Hall. helu. n. 168.

Cirsium foliis spinosis, semipinnatis; laciniis alternis erectis; calycibus lanigeris, subrotundis. Scop. carn.

Carduus foliis sinuatis decurrentibus: denticulis superficieque spinosis, calycibus lanigeris. Dalib. paris.

C. capite rotundo tomentoso. Bauh. pin. 382.

Carduus tomentosus corona fratrum herbariorum. Lob. ic. 2. p. 9.

C. eriocephalos. Dodon. purg. 445. Clus. pann. 666. Hall. R.

β. Carduus (spurius) foliis semidecurrentibus duplicatopinnatifidis: laciniis alternis erectis, calycibus ovatis. Hort. upf. 249. not.

C. tomentosus pyrenaicus, floribus purpureis glomeratis. Tournef. inst. 441.

Habitat in Anglia, Gallia, Hispania, Lusitania, &c. ♂

25. CARDUUS heterophyllus. C. foliis amplexicaulibus lanceolatis ciliatis integris laciniatisque, caule subunifloro, calyce inermi. *Fl. suec. ingr. Flor. dan. t. 109. sil. Pall. is. 1. p. 62. delph.*

336 SYNGENESIA. POLYGAMIA AQUALIS

Cirsium britannicum, caule multifloro, foliis inferioribus laciniatis. *Scop. ann.*

Carduus calyce inermi, foliis lanceolatis marginibus ciliatis
Fl. lapp.

Cirsium maximum, asphodeli radice. *Bauh. 377.*

Habitat in Europæ frigidioris pratis depressis. 2.

Caulis sesquipedalis s. brevior; Foliis 5 ad 10, non dentatis, ciliatis, subtus niveis; aliis integris, aliis laciniatis. Radix repens.

Cum sequenti conjungit Hallerus helu. n. 180. β. R.

26. *CARDUUS helenioides*. C. foliis amplexicaulibus lanceolatis dentatis: spinulis inæqualibus ciliatis, caule inermi.
*Hort. ups. **

Cirsium foliis ciliatis, subtus tomentosis, radicalibus petiolatis, ovato-lanceolatis, caulinis lanceolatis & semipinnatis. *Hall. helu. n. 180. var. α. t. 7.*

Cirsium britannicum caule unifloro, foliis dentatis. *Scop.*

Cirsium singulari capitulo squamato, vel incanum alterum.
Bauh. pin. 377. Mill. dict. t. 94.

Cirsium anglicum, radice hellebori nigri modo fibroso, folio longo. *Bauh. hist. 3. p. 45.*

Carduus mollis, foliis helenii. *Bauh. pin. 377. videtur.*

Habitat in Anglia, Sibiria. 2.

Antecedenti simillimus, sed duplo altior seu altitudine humana. Radix vix repens, foliis omnibus indivisis, subtus albis, at non ut illi niveis. Caulis sulcatus, totus foliosus. Foliis 40 ad 50, subtus albidis, semiamplexicaulibus (auriculis rotundis, reflexis, adnatis, dentatis, non laciniatis, ciliatis spinis inermibus. Floralibus subulatis.

17. *CARDUUS ferratuloides*. C. foliis subamplexicaulibus, lanceolatis integris: serraturis spinoso-setaceis, pedunculis unifloris. *monsp. Jacq. austr. t. 127. carn. sub Cirsio. fl. lugd.*

C. calycibus inermibus acutis, foliis utrinque viridibus firmis integerrimis inæqualiter ciliatis. *Gmel. fib. 2. p. 52. t. 22. & 23. f. 1.*

Cirsium procerum, folio profunde serrato, utrinque virente, laceæ capitulis minoribus, fibrosa radice. *Amm. ruth. n. 187. R.*

Cirsium angustifolium non laciniatum. *Bauh. pin. 377.*

Cirsium pannonicum r. pratense. *Clus. 2. p. 148.*

Habitat in Sibiria, Helvetia, Monspelii. 2.

Differt a precedenti simillimo: Caule dimidio minore,
vix

SYNGENESIA. POLYGAMIA ÆQUALIS. 337

vix sulcato; Foliis angustioribus; floralibus lanceolatis nec subulatis.

28. CARDEUS tataricus. C. foliis amplexicaulibus lanceolatis: serraturis spinoso-senscatis; floribus triphyllis. *Jacq. austr. t. 90.*

Cirsium foliis ciliatis; semipinnatis; pinnis angulosis; spinosis; superioribus amplexicaulibus. *Hall. helv. n. 176. R.*

Habitat in Sibiria, Helvetia. 2.

Habitus & altitudo C. Serrauloidis, ut facile pro varietate sumeretur, a quo differt Foliis duplo latioribus; magis dentatis: vix ac ne vix subtus tomentosis; Capitulis foliolis 3, rarius duobus obvallatis, nec unico, ut ille. Foliis basi pungentibus. Floribus calyce cylindrico, nec globoso. Colore florum albo nec purpureo, at Pappo plumoso convenit, imprimis vero Foliis radicalibus pinnatifidis. Convenit vero cum Cnico oleraceo his foliis; calycum squamis carina alba; Florum visco fastigiato, nec convexo; sed folia superiora indivisa; flores solitarii, longe pedunculati. Folia floralia parva, nec lata concava s. pallida.

29. CARDUUS flavescens. C. foliis lanceolatis integris inermibus dentatis glabris, floribus aphyllis inermibus spinosis.

Habitat in Hispania. Læsting.

Caulis lævis. Folia lanceolata; basi attenuata; argute dentata, inermia. Calyces cylindrico-ovati, terminati squamis aciformibus, mitibus, patulis. Corolla albo-flavescens.

30. CARDUUS mollis. C. foliis pinnatifidis linearibus subtus tomentosis, caule inermi unifloro. *Jacq. austr. t. 18. carn. sub Cirsio. pal.*

C. foliis decursive interrupteque pinnatis tomentosis, pinnulis integerrimis, caule unifloro; calycibus inermibus squarrosis. *Gouan. illustr.*

C. mollis, laciniato folio. *Bauh. pin. 377.*

C. mollior. 1. *Clus. hist. 2. p. 151.*

Habitat in alpinis Austriae, in Germania, Monspeli.

31. CARDUUS acaulis. C. acaulis; calyce glabro. *Fl. suec. parif. carn. sub Cirsio. gallob. pal. herb. lugd. delph. burg.*

Cirsium floribus acaulis. *Hall. helv. n. 178.*

Carlina minor. *Clus. hist. 156. R.*

Carlina montana minor acaulos. *Barrel. ic. 493. R.*

Linnaei spec. Plant. Europ. Pars II.

338 SYNGENESIA. POLYGAMIA ÆQUALIS.

Carlina acaulis, minore purpureo flore. *Bauh. pin.* 380.

Chamaeleon exiguus. *Bauh. hist.* 3. p. 62. *Trag.* 852.

Habitat in Europæ pascuis apertis depressis. 24

Floribus albis variat. R.

1005. C N I C U S.

Cal. ovatus, imbricatus squamis ramoso-spinosis;
obvallatus bracteis. *Corollulae* æquales.

1. *CNICUS oleraceus*. C. foliis pinnatifidis carinatis, nudis
bracteis concavis integris subcoloratis. *Fl. succ. ged. carn.*
sub Cirsio. gallob. pal. sil. Kniph. cent. 12. n. 29. *delph.*
lith. lugd. burg.

Cirsium foliis ciliatis, amplexicaulibus inermibus, flores
comprehendentibus. *Hall. helu.* n. 173.

Carduus pratensis latifolius. *Bauh. pin.* 376.

Carduus pratensis. *Lob. ic.* 2. p. 11.

β. *Cirsium latissimum*. *Bauh. pin.* 177.

Habitat in Europæ septentrionalioris pratis subnemorosis. 24

Calyces laterales sessiles, sæpius bini, intermedio subpe-
dunculato. Lineola alba secundum carinam squamarum
calycis.

Antherarum appendices minimæ, bifidæ, ut in Carlina
vulgari. *Leers herb.* n. 623. R.

Cnicus oleraceus. L. in sylvis nostris ad rivulos ad huma-
nam altitudinem adsurgit, foliis magis spinosis, ramis in
adultâ stirpe magis elongatis, aphyllis, simillimus tunc
Carduo tatarico *Jacq. austr.* t. 90; an eadem planta? Ce-
terum *Cirsium* *Hall. helu.* n. 176. cujus in *Ad. helu.* tom. 4.
iconem dedi, quodve ætenuis non bene pro *Carduo tatarico*
Linn. habui, a *Jacquini* stirpe abunde discrepat. De la
Chenal in Ad. helu. VIII. p. 141. R.

2. *CNICUS Erisithales* C. foliis amplexicaulibus pinnati-
fidis aristato-ferratis: pedunculis cernuis, calycibus
glutinosi. *Jacq. obs.* 1. p. 28. t. 17. *austr.* t. 310.
delph.

Cirsium foliis ciliatis, imis ovatis, superioribus semipinnatis,
floribus sessilibus obtusis. *Hall. helu.* n. 175. *Comm. gart.*
I. p. 206. t. 8.

Cirsium foliis amplexicaulibus pinnatifidis; calycibus in-
ermibus. *Scop. carn.*

Carduus foliis amplexicaulibus

SYNGENESIA. POLYGAMIA ÆQUALIS. 339

latis spinuloso-ciliatis , calycibus glutinosis nutantibus.

*Jacq. vind. **

Carduus pratensis , foliis tenuibus laciniatis. *Bauh. pin.* 377.

arduo-**Cirsium** maximum profunde laciniatum in foliorum ambitu spinis mollibus hirtum. *Pluk. phyt.* 154. f. 2. (*sed flore purpureo.*)

Cirsium acanthoides montanum , flore flavescente. *Tournef. inst.* 448.

Cirsium alatus polycephalon , foliis imis integris , superioribus latis , profunde laciniatis. *Rai. syllog.* 398. *Hall. R.*

Carduo-**Cirsium** dictum montis piniferi repens. *Volkam. norimb. p.* 85. *Hall. R.*

Erisithales. *Dalech. hist.* 1094.

Habitat in Austriæ , Galliæ pratis subalpinis. 24

Folia pinnato-partita , sessilia , subtus imprimis pubescentia ; Pedunculi aphylli. Flores solitarii , nutantes , lutei : disco convexo longiore. (Flores alias purpurei , alias ochroleuci , etiam albi. *Hall. l. c. R.*)

3. **CNICUS ferox.** C. foliis decurrentibus ligulatis dentato-spinosis , caule ramoso erecto. *Ger. pron. delph.*

C. foliis lanceolatis decursive pinnatis , pinnulis bilobis longissimis , subtus tomentosis , supra strigoso - spinosissimis. *Gouan. illustr.*

Carduus lanceolatus validissimis aculeis. *Tournef. inst.* 450.

Carduus lanceolatus ferocior. *Bauh. hist.* 3. p. 58.

Habitat in Europæ australis montosis , sterilibus. ♂

4. **CNICUS pygmeus.** C. caule unifloro , foliis sublinearibus sessilibus confertissimis , calyce inermi.

Carduus caule unifloro , calyce inermi , foliis sublinearibus , sessilibus. *Jacq. vind. t.* 8.

Serratula pygmæa. *Jacq. austr. t.* 440.

Cirsium pigmæum. *Scop. carn.*

Carduus mollis , folio oblongo , cirsii capitulo. *Bauh. pin.* 377.

Carduus mollior humilis angustifolius. *Clus. hist.* 2. p. 151.

Habitat in Schneeberg Austria. 24

5. **CNICUS Acarna.** C. foliis decurrentibus lanceolatis indivisis , calycibus pinnato - spinosis. *Gouan. monsp. delph.*

Carduus Acarna. *Sp. pl.*

Carduus caule alato spinoso , foliis spinosis spinis duplicibus

Roy. lugdb. 133.

340 SYNGENESIA. POLYGAMIA ÆQUALIS.

Acarna major, caule folioso. *Bauh. pin.* 379.

Chamæleon salmanticensis. *Clus. hist.* 2. p. 155.

β. *Acarna humilis*, caule folioso. *Bauh. pin.* 379.

Picnomon cretæ solonensis. *Dalech. hist.* 1456.

Habitat in Hispaniæ arvis.

Habitus Carlinae. Facies Carthami. Caulis simplex, superne corymbosus. Folia adnata, canescentia, ciliata spinis geminis transversis flavis. Flores terminales, demum axillares, sessiles. Calyces oblongi, imbricati squamis apice spinis innocuis, pinnatis ex spinulis 5 patentibus. Corollæ parvæ, purpurascentes.

6. *CNICUS spinosissimus*. C. foliis amplexicaulibus sinuato-pinnatis spinosis, caule simplici, floribus sessilibus. *Scop. carn. sub Cirsio. delph.*

Cirsium foliis caulinis semipinnatis, pinnis angulosis, spinosissimis, terminantibus mollibus, ciliatis, umbellam floralem comprehendentibus. *Hall. helu. n.* 172. t. 5.

Carlina polycephalos alba. *Bauh. pin.* 380.

Cirsium alpinum spinosissimum, floribus ochroleucis inter flavescencia folia congestis. *Hall. helu. ed.* 1. p. 679. t. 20.

Cirsium spinosum capitulis in vertice inter foliorum valla congestis nudis inermibus. *Gmel. sib.* 2. p. 64 t. 25.

Habitat in Alpibus, Helvetiæ, Austriæ, Tartariæ.

Flores ochroleuci.

7. *CNICUS centauroides*. C. foliis pinnatifidis, calycibus scariosis: squamis acuminatis.

Carduus inermis, foliis pinnatifidis ferratis inermibus, squamis calycinis membranaceis acuminato-lanceolatis. *Hort. cliff.*

Centaureum majus, foliis cinaræ cornuti. *Morif. hist.* 3. p. 131. f. 7. t. 25. f. 2.

Habitat in Pyrenæis: Sibiria. Georgi. it. 1. p. 229. 2/

1006. O N O P O R D U M.

Receptaculum favosum. Cal. squamæ mucronatæ.

1. *ONOPORDUM acanthium*. O. calycibus squarrosis, squamis patentibus, foliis ovato-oblongis sinuatis. *pal. gallob. sil. lugd. pol. delph. burg.*

O. foliis decurrentibus margine spinosis. *Fl. suec. paris.*

C. caule alato, foliis ovatis, dentatis, dentibus angulosis, aristatis. *Hall. helu. n.* 159.

SYNGENESIA. POLYGAMIA ÆQUALIS. 343

Acanos Spina. Scop. carn.

Spina alba tomentosa latifolia sylvestris. Bauh. pin. 382. Læf. pruss. 261. t. 82.

Acanthium. Dod. pempt. 721.

Habitat in Europæ ruderatis, cultis. ♂

Planta canescenti-virens. Caulis alis rectis ad ipsum calycem excurrentibus. Calyce squamis patentibus, subulatis. Floribus albis ludit.

2. *ONOPORDUM illyricum. O. calycibus squarrosis: squamis inferioribus uncinatis, foliis lanceolatis pinnatifidis.*

Jacq. hort. t. 148. delph. burg.

O. foliis decurrentibus ligulatis pinnatifidis. Sauv. monsp.

Spina tomentosa altera spinosior. Bauh. pin. 382.

Carduus, quibusdam dictus Acanthium illyricum. Bauh. hist.

3. p. 55.

Acanthium illyricum. Lob. ic. 1. Barr. ic. 501.

Habitat in Europa australi.

Planta cinereo-cana. Caulis alis undulatis infra calycem spatio desinentibus. Calycis squamæ inferiores reflexo-uncinatae, lanceolatae.

3. *ONOPORDUM arabicum. O. calycibus imbricatis. Jacq. hort. t. 149.*

Carduus tomentosus, acanthi folio, altissimus lusitanicus. Tournef. inst. 441.

Carduus tomentosus, Acanthium dictus, arabicus. Pluk. alm. 85. t. 154. f. 5.

Carduus tomentosus, acanthi folio, altissimus lusitanicus. Barr. ic. 591.

Acanthium altissimum lusitanicum. Moris. hist. 3. p. 153.

Habitat in Lusitania, G. Narbonensi. ♂

1007. CYNARA.

Cal. dilatatus, imbricatus squamis carnofis, emarginatis cum acumine.

1. *CYNARA Scolymus. C. foliis subspinosis pinnatis indivisisque, calycinis squamis ovatis. monsp. Blackw. t. 458.*

C. foliis pinnatis: laciniis ferratis. Hort. ups.

Cinara sylvestris latifolia. Bauh. pin. 384.

β. Cinara hortensis aculeata. Bauh. pin. 383.

Scolymus dioscoridis. Clus. hist. 2. p. 153.

342 SYNGENESIA. POLYGAMA AEQUALIS.

2. *C. hortensis*, foliis non aculeatis. *Bauh. pin.* 383. *Mitt. did. n.* 2.

Habitat in G. Narbonensis, Italiae, Siciliae apricis. 24

2. *CYNARA Cardunculus*. *C.* foliis spinosis : omnibus pinnatifidis, calycinis squamis ovatis.

C. foliis pinnatis : foliolis oblongis indivisis, caule spinoso. *Sauv. meth.*

C. spinosa, cujus pediculi esitantur. *Bauh. pin.* 383.

Scolymus aculeatus. *Tabern. hist.* 1075.

Habitat in Creta. 24

Hybridam s. natam ex praecedentis semine asseverat *J. B.* Flos saturate caeruleus corollis & pistillis, et stamina grisea.

3. *CYNARA humilis*. *C.* foliis spinosis pinnatifidis subtus tomentosis, calycibus squamis subulatis.

Cinara sylvestris baetica. *Clus. car.* 37.

Carduus tingitanus, flore magno caeruleo, folio arachydis divisura subtus incano, spinis durioribus horrido. *Pluk. alm.* 85. t. 81. f. 2.

Carduus humilis tingitanus caeruleus, magno strobilo, tenuius laciniatus. *Moris. hist.* 3. p. 158.

Habitat in agro Tingitano, in Baetica. 24

Conf. Carduus andelsiacus. *Pluk. alm.* 85. t. 19. *Moris. hist.* 3. p. 158.

Semina in hac specie minore, reliquis longe maiora & singularia. Caulis longitudine vix foliorum. Calyx ovato-oblongis squamis patentibus.

1007. CARLINA.

Carl. radiatus squamis marginalibus longis, coloratis.

1. *CARLINA acaulis*. *C.* caule unifloro flore brevior. *mon/p.* *Hall. helv. n.* 183. *carn. fil.* *Blackw. t.* 532. *Knorr. del.* 2. t. *C.* 1. *Regn. bot. lugd. lithu. burg.*

Carlina acaulos, magno flore albo. *Bauh. pin.* 380.

Chamaeleon albus. *Clus. hist.* 2. p. 155. *Cam. epit.* 428.

Habitat in Italiae, Germaniae montibus apricis. 24

Varietas caulifera, pedalis & altior est *Carlina prior*. *Dodon. purg.* 439. *Hall. R.*

2. *CARLINA lanata*. *C.* caule bifido, calycibus sanguineis terminalibus : primo axillari sessili.

SYNGENESIA. POLYGAMIA ÆQUALIS. 343

C. squamis calycinis terminalibus purpureo-rubentibus, floribus inferioribus prolixis. *Ger. prou.*

C. caule trifloro dichotomo: intermedio sessili. *Sauv. monsp.*

C. flore purpureo rubente patulo. *T. Gard. auz. 82. t. 21.*

Acarna atraçtylidis folio, amplo flore. Barr. ic. 483.

Acarna flore purpureo rubente patulo. Bauh. pin. 379.

Acanthoides parva apula. Col. ecphr. 1. p. 29. t. 27. f. 2.

Habitat in Italia, G. Narbonensi. ☉

Calycis squamæ veræ terminantur omnes spina simplicissima; in reliquis autem spinæ compositæ sunt.

Radius purpureus utrinque. Gouan. illustr. R.

3. *CARLINA corymbosa. C.* caule multifloro subdiviso, floribus sessilibus, calycibus radio flavo. *carn. delph.*

Acarna capitulis parvis luteis in umbella. Bauh. pin. 379.

Acarna apula umbellata. Column. ecphr. 1. p. 20. t. 27.

β. *Carlina sylvestris multicaulis, flore flavo, radice perenni. Rai. extr. 86.*

C. sylvestris, flore aureo, perennis. Tournesf. inf. 300.

C. patula, atraçtylis folio & facie. Tournesf. inf. 300.

Habitat in Italia versuris, a mari non remotis. 24

Radius flavus; sed foliola extus & ad basin tantum purpureascentia. Gouan. l. c. R.

Caulis magis lanatus. Flores pauciores quam in C. vulgari. Gerard.

4. *CARLINA vulgaris. C.* caule multifloro corymboso, floribus terminalibus, calycibus radio albo. *Fl. suec. ged. herb. pal. sil. lish. lugd. burg.*

C. foliis radicalibus semipinnatis, pinnis angulosis, spinosis, caulinis ovato lanceolatis, amplexicaulibus. Hall. belu. n. 182.

C. sylvestris vulgaris. Clus. hist. 2. p. 155.

Cnicus sylvestris spinosior. Bauh. pin. 378.

Atractylis mitior. Fuchf. hist. 121.

Habitat in Europæ montosis, aridis, sabulosis. ♂

Corollæ centrales atropurpureæ. Gou. l. c. R.

Vivas observent Botanici australes, qui queant 2, 3, 4. demto colore, rite distingui.

5. *CARLINA racemosa. C.* floribus sessilibus lateralibus paucissimis. *monsp.*

C. squamis calycinis terminalibus longioribus, floribus inferioribus prolixis. Ger. prou.

C. sylvestris minor hispanica. Clus. hist. 2. p. 157.

Acarna flore luteo patulo. Bauh. pin. 379.

844 SYNGENESIA. POLYGAMIA ÆQUALIS.

Habitat in Hispaniæ desertis. ☉

Radii calycina foliola omnino a basi ad apicem setina atropurpurea; cæterum a parte superiore magis flava. *Gou. l. c. R.*

6. *CARLINA pyrenaica*. C. caule multifloro, foliis decurrentibus.

Carduus tomentosus pyrenaicus, spinosissimus, flore albo, Burf. XXI. 43.

Habitat in Pyrenæis versus Hispaniam copiose.

Folia utrinque tomentosa: pinnis sessilibus hastatis dentatis apicibus omnibus spina flavescente. Calyces vix majores *Card. pycnocephalo*: squamis laxis spinosis, erectis, scariosis, angusto-lanceolatis, vix calycis longitudine. Flores purpureascentes.

1008. ATRACTYLIS.

Cor. radiata: corollulis radii 5. dentatis.

1. *ATRACTYLIS gummifera*. A. flore aculi.

Cnicus, carlinæ folio, aculos gummifer aculeatus. Tournef. cor. 33.

Carlina aculos gummifera. Bayh. pin. 380.

Carduus pinea theophrasti. Alp. exot. 126. t. 124. 125.

Chameleon albus dioscoridis. Colum. ecphr. l. p. l. t. 12.

Habitat in Creta, Italia.

2. *ATRACTYLIS humilis*. A. foliis dentato-sinatis, flore radiato: obvallato involucri patente, caule herbaceo.

*A. radiata squamis calycinis truncatis cum spinula setacea. Læf. 11. **

Cnicus aculeatus purpureus humilior. Tournef. inf. 451.

Crocodiloides atractylidis folio, flore purpureo coronato. Vaill. aq. 1718. p. 172.

3. *Carlina minima caulodes hispanica. Barr. rar. 1127. t. 592.*

Habitat Madriti in collibus β circa Narbonam. ♂

Habitus *A. cancellata*, sed folia sinuata. Flos radiatus. Calycis squamæ mucronatæ. Involucrum ex foliis veris consistens.

Varietas β. differt Calycis squamis ovatis non mucronatis, squamis spinula terminatis.

3. *ATRACTYLIS cancellata*. A. involucri

SYNGENESIA. POLYGAMIA ÆQUALIS. 345
tricosis linearibus dentatis , calycibus ovatis , floribus
flosculosis.

A. foliis linearibus dentatis , calycibus conniventibus. *Hort.
cliff.*

A. flore radiato nudo , calyce inermi involucri connivente
obtecto. *Læst. it. 162. **

Acarna capitulis globosis. *Bauh. pin. 379.*

Eryngium parvum palmare , foliis serratis. *Moris. hist. 3. p.
166. f. 7. t. 39. f. 16.*

Carduus parvus. *Bauh. hist. 3. p. 93. Rai. hist. 316.*

Carduus minimus. *Alp. exot. 254.*

Habitat in Hispaniæ , Cretæ agris. ☉

Caulis palmaris simplex , albo-araneosus : Ramis ex alis 2
ulimis. Folia lineari-lanceolata , ciliata , araneosa. Flos
terminalis , cinctus Bracteis 3 , patentibus & inclusus 8 con-
niventibus , in peripheriam globosam inflatam , quarum 3
alternæ apice recurvæ. Bractææ singulæ lineares , pinnato-
dentatæ dentibus bifurcatis. Corolla parva , cærulea , flos-
culosa , rarius radio. Involucrum cancellatum , naturæ stu-
pendum artificium , muscas arcet.

1009. C A R T H A M U S.

Cal. ovatus , imbricatus squamis apice lobovato-
foliaceis.

1. CARTHAMUS tinctorius. C. foliis ovatis integris serrato-
aculeatis. *Ludw. ed. t. 129. Kniph. cent. 7. n. 8. Regn.
bot.*

Cnicus sativus l. Carthamus officinarum. *Bauh. pin. 378.
Rumph. amb. 5. p. t. 79.*

Cnicus vulgaris. *Clus. hist. 2. p. 152.*

Habitat in Ægypto. ☉

2. CARTHAMUS lanatus. C. caule piloso : superne la-
nato , foliis inferioribus pinnatifidis : summis amplexicau-
libus dentatis. *monsp. Mill. dict. n. 2. Blackw. t. 468.
delph. lugd. burg.*

**Atractylis foliis cartilagineis , reticulatis , imis semipinnatis ,
superioribus lanceolatis , amplexicaulibus.** *Hall. helv. n.
192.*

**Atractylis foliis amplexicaulibus , dentato - spinosis ; caule
tomentoso ; floribus flosculosis.** *Scop. carn.*

Carthamus foliis amplexicaulibus acuminato - dentatis, *Hort.
t. Dalib. paris. 250.*

346 SYNGENESIA. POLYGANIA ÆQUALIS.

Attractylis lutea. Bauh. pin. 379.

Attractylis theophrasti & *dioscoridis* fucce sanguinea. Col.
ephr. 1. p. 19. t. 23.

Attractylis. Dod. pempt. 736. Camer. epis. 561.

Habitat in Gallia, Italia, Creta, Helvetia, Carniola. ☉

Flosculi radii neutri. Hall.

5. *CARTHAMUS cæruleus*. C. foliis lanceolatis spinoso-
dentatis, caule subunifloro.

Cnicus cæruleus asperior. Bauh. pin. 378.

Cnicus alter clusii, cæruleo flore. Clus. hist. 2. p. 152.

Habitat in Hispania inter segetes. 24

Caules simplicissimi, sæpe procumbentes, pilosi, uniflori.
Folia supra glabra. Calyx foliosus. Flos cæruleus an-
theris nigris. Semina pappo piloso.

Synonyma ulterius examinanda & a precedenti distin-
guenda.

6. *CARTHAMUS mitissimus*. C. foliis inermibus, radi-
calibus dentatis, caulinis pinnatis. Stamp. parisi. berg.

C. foliis inermibus: radicalibus lanceolatis serratis, caul-
inis pinnatifidis, calycum squamis subvenis, enerviis,
inermibus integerrimis. Gouan. illustr.

Carthamoides cærulea humilis & mitior. Vail. ed. 1718.

772.

Carduncellus montis lupi minor species. Moris. hist. 3.
p. 159.

Habitat circa Parisios, Monspelium.

Semina pappo piloso V. Modo acaulis, modo palmaris.
Gou.

Hortensis vix distinguendus. Caule decumbenti palmari
Foliisque sinuatis longioribus mollibus. Ger. 182.

7. *CARTHAMUS Carduncellus*. C. foliis caulinis linea-
ribus pinnatis longitudine plantæ. Monsp. delph.

C. foliis spinosis: radicalibus obovatis serratis, caulinis
pinnatifidis, calycum squamis trinerviis spinosis. Gouan.
illustr. p. 65. R.

C. foliis caule longioribus ligulatis pinnatifidis. Sacch.
monsp. *

C. foliis radicalibus superne serratis, inferne dentatis;
caulinis dentato-pinnatis. Roy. lugdb.

Cnicus cæruleus humilis montis lupi. Herm. lugdb. 164.

Carduncellus montis lupi. Lob. ic. 2. p. 20.

Eryngium montanum minimum, capitulo magno. Bauh.
pin. 386.

SYNGENESIA. POLYGAMIA ÆQUALIS. 347

Habitat Monspeli.

Semina pappo piloso. V.

8. CARTHAMUS *arborescens. C. foliis ensiformibus sinuato-dentatis.*

Carthamoides lutea altissima & foetidissima. Vaill. ad. 1718. p. 772.

Cnicus hispanicus arborescens foetidissimus. Tournef. inst. 451.

Habitat in Hispania. H

Semina pappo piloso. V.

Caulis ut tota planta, pubescens, firmus, humanæ altitudinis, sempervirens. Folia amplexicaulia, ensiformia s. lanceolata, pedalia, pinnatifido sinuata, dentata, apicibus omnibus subspinosa: costa alba. Flos terminalis, 1. s. 2, sessilis, luteus, suaveolens. Calyx squamis ciliatis, patulis, mucronatis. Stylus corollis duplo longior. Pappus pilosus.

9. CARTHAMUS *corymbosus. C. floribus corymbosis numerosis.*

Echinops floribus fasciculatis, calycibus multifloris.

Chamæleon niger. Dalech. hist. 1454. Dod. pempt. 727.

Chamæleon niger umbellatus, flore cæruleo hyacinthino.

Bauh. pin. 380

Carduus Chamæleon dictus capitulis pluribus minoribus cæruleis corymbatim dispositis. Moris. hist. 8. p. 159. s. 7. 1. 33. f. 17.

Habitat in Apuliæ, Hellesponti, Lemni, Thraciæ campis. 2

Habitus Echinopsis. Qualitas corrosiva. Fructificatio iterum examinanda.

1012. B I D E N S.

Recept. paleaceum. Pappus aristis erectis, scabris.

Cal. imbricatus. Cor. rarius flosculo uno alterove radiante instruitur.

1. BIDENS *tripartita. B. foliis trifidis, calycibus subfoliosis, seminibus erectis. Carn. pal. sil. Blackw. 1. 519. Knorr. del. 2. 1. B. 1. f. 2. delphi. lithua. lugd. burg.*

B. foliis petiolatis trilobatis & quinquelobatis, serratis, floribus circumvallatis. Hall. helu. n. 121.

948 SYNGENESIA. POLYGAMIA ÆQUALIS.

B. corona seminum retrorsum aculeata, seminibus erectis.
Fl. suec.

B. foliis tripartito divisis. *Tournesf. inst.* 462. *Fl. lapp.* 294.
Cannabina aquatica folio tripartitum diviso. *Bauh. pin.* 321.
Hepatorium aquatile. *Dod. pempt.* 595.

β. *Conyza* palustris, foliis tripartito divisis. *Læf. pruss.* 53. *ic.* 10. *Hall. R.*

Habitat in Europæ inundatis. ☉

2. **BIDENS minima.** **B.** foliis lanceolatis sessilibus, floribus seminibusque erectis. *Angl. pat. Flor. dan.* t. 312. *delph. lithua.*

Verbesina minima, *Dill. giss.* 167. *app.* 66. *Rai. angl.* 3. p. 188. t. 7. f. 2.

Habitat in Europæ septentrionalioris paludosis. ☉

B. cernuæ varietatem facit *Hallerus. R.*

5. **BIDENS cernua.** **B.** foliis lanceolatis amplexicaulibus, floribus cernuis, seminibus erectis. *Fl. suec. carn. pom. herb. pal. sil. Knorr. del.* 2. t. B. 1. f. 1. *delphi. lithua. lugd. burg.*

B. foliis sessilibus, floribus nutantibus, circumvallatis. *Hall. helu. n.* 120.

Cannabina aquatica, folio non diviso. *Bauh. pin.* 321.

Eupatorium cannabinum crysanthemum. *Tabern. ic.* 117.

Habitat in Europa ad fontes & fossas. ☉

Conferatur Coreopsis bidens ejusque affinitas.

12. **BIDENS bullata.** **B.** foliis ovatis serratis: inferioribus oppositis, superioribus ternatis: intermedio majore. *Arduin. spec* 2. p. 37. t. 18.

B. palustris annua, foliis subrotundis latioribus plerumque trilobis. *Mich. flor.* 120.

B. foliis ovatis & tripteris, caulibus hirsutis & brachiatis. *Hall. gatt.* 383.

Habitat in America, nunc in Italia. ☉

Caulis erectus, bipedalis. Folia subovata, serrata, crassa, villosa, fere bullata: inferiora simplicia; superiora subternata, subsessilia: foliolis lateralibus minimis. Flores axillares & terminales, solitarii, subsessiles, lutei; Calyces basi cincti foliolis ovato-oblongis, quinis, patentibus. Semina Aristis 2, retro ciliatis.

1013. C A C A L I A.

Recept. nudum. Pappus pilosus. Cal. cylindricus, oblongus, basi tantum subcalyculatus.

11. *CACALIA seracenicæ*. C. caule herbaceo, foliis lanceolatis ferratis decurrentibus.

Solidago foliis lanceolatis decurrentibus, caule angulato. Monsp.

Senecio perennis, solidaginis folio. Vaill. aët. 366.

Virga aurea maxima, radice repente, f. Doria major repens. Moris. hist. 3 p. 123.

Virga aurea f. Solidago seracenicæ latifolia ferrata. Bauh. hist. 2. p. 1063.

Conyza montana, foliis longioribus ferratis, flore e sulphureo albicante. Chom. aët. par. 1705. p. 394.

Habitat in Gallia australi. 24

Simillima senecioni seracenicæ. (Bractææ setuææ.

15. *CACALIA alpina*. C. foliis reniformi-cordatis acutis denticulatis, calycibus subtrifloris. *Delph. lugd.*

C. foliis cordatis ferratis, longissime petiolatis, floribus umbellatis. *Hall. helv. n. 137.*

Tussilago caule ramoso. Sauv. monsp.

α. *Cacalia foliis crassis hirsutis, Bauh. pin. 197. Moris. hist. 3. p. 94. f. 7. t. 12. f. 1. Monnier. obs. 47. Hall. opusc. 207. delphi.*

C. tomentosa. *Jacq. austr. t. 235.*

C. (*alliaris*) calycibus trifloris, foliis cordatis petiolatis dentato-angulatis, caulinis subhastatis, vaginis stipularibus cordato-amplexicaulibus. *Gouan. illustr. 65.*

Cacalia incano folio. Clus. hist. 2. p. 115.

β. *Cacalia foliis cutaneis acutioribus & glabris. Bauh. pin. 198. Moris. hist. 3. p. 94. f. 7 t. 12. f. 6.*

C. glabra *Mill. dict. n. 2.*

C. alpina. *Jacq. austr. t. 234.*

Tussilago Cacalia. Scop. carn.

Cacalia glabro folio. Clus. hist. 2. p. 115.

Cacalia. Lob. ic. 592.

Habitat in Alpibus Helvetiæ, Austriæ. 24

Hanc speciem genere cum antecedentibus convenire docuit autopsia, hinc genere conjungendæ Cacaliæ cum Kleiniis.

Calyx subtriflorus. Calycis foliola conglutinata, Corollulæ. 4-fidæ. Scop.

350 SYNGENESIA. POLYGAMIA. ÆQUALIS.

1015. EUPATORIUM.

Recept. nudum. *Pappus* plumosus. *Calyx* imbricatus, oblongus. *Stylus* semibifidus longus.

* *Calycibus* quadrifloris.

3. EUPATORIUM *scandens*. E. caule volubili, foliis cordatis dentatis acutis.

Conyza scandens, folani folio anguloso. *Plam. ic.* 99. *Clematitis* novum genus, cucumeris folio, virginianum.

Pluk. alm. 109. t. 163. f. 3.

Habitat in aquosis Virginiae. 24

Planta Ind. orient. lævior, foliorum dentibus obsoletioribus, quod Eup. cordatum. *Burm. ind.* 176. t. 58. f. 2.

* * *Calycibus* quinquefloris.

10. EUPATORIUM *altissimum*. E. foliis lanceolatis nervosis : inferioribus extimo subserratis, caule suffruticoso. *Jacq. hort.* t. 164. *Kniph. cent.* 2. n. 23.

E. virginianum, longissimis & angustissimis foliis. *Morif. hist.* 3. p. 97. *Rai. suppl.* 187.

E. folio oblongo, rugoso, ampliori, caule virescente. *Tournef. inst.* 456. R.

Sehr hohes Eupatorium. *Linn. Pflanzenystem* 4. p. 333.

Habitat in Pensylvania. 24

! *Description* Morisoni imperfecta. *Inula* speciem facit *Vaillant*, si vero Flores Morisoni radiati fuissent, vix ad *Eupatoria* eam retulisset Morison.

13. EUPATORIUM *cannabinum*. E. foliis digitatis. *Succ. ingr. ged. carn. gallob pal. sil. Blackw.* t. 110. *Regn. bot. Flor. dan.* t. 745. *lugd. lithua. burg.*

E foliis ovato lanceolatis, tripteris. *Hall. helu.* n. 136.

E. cannabinum. *Bauh. pin.* 320.

E. adulterinum. *Fuchf. hist.* 265.

Habitat in Europa ad aquas. 24

Variat foliis integris. *Rai. syn.* 3. p. 180. *Hall. R.*

1018. STÆHELINA.

Recept. brevissime paleaceum. *Pappus* ramosus. *Antheræ* caudatæ.

2. STÆHELINA *dubia*. S. foliis linearibus denticula-

SYNGENESIA. POLYGAMIA ÆQUALIS. 351

vis, squamis calycinis lanceolatis, pappo calycibus duplo longiore. *Allion. nicæns.* * *Ger. prou.* t. 6.

Gnaphalium caule fruticoso, foliis linearibus denticulatis, capitulis oblongis. *Roy. lugdb.*

Stœchas odorata purpurea. *Lob. illustr. Magn. monsp.* 247. 297.

Stœchadi citrinæ affinis, capitulis longioribus. *Bauh. hist.* 3. p. 156.

Flychrysum sylvestre, flore oblongo. *Bauh. pin.* 265.

Chamæ-Chrysocome prælongis purpureisque jaceæ capitulis. *Barr. ic.* 406.

Jacea capitata, rosmarini folio. *Tournef. inst.* 444. 337.

Habitat in Hispania, G. Narbonensi. \bar{h}

Singularis planta, quasi media inter ferratulas Gnaphalia & Stæhelinas. Pappus pilosus basi connatus. Caulis & Folia subtus tomentosa.

3. *STÆHELINA arborescens.* S. foliis ovalibus. *Schreb. dec.* 1. t. 1.

Centaurea calycibus squamosis, foliis ovatis subtus tomentosis. *Ger prou.*

Raponticoides frutescens, stryacis folio. *Vaill. aed.* 179.

Jacea arborescens, stryacis folio. *Tournef. inst.* 445.

Cyanus arborescens altera, stryacis folio. *Alp. exot.* 33. t. 32.

Frutex rotundo argenteo folio, cyani flore. *Bauh. pin.* 452.

Habitat in Creta, insula Stœchadum; in rupe *Victoriæ* Galloprovinciæ. \bar{h}

Calyx imbricatus squamis muticis. Paleæ multipartitæ.

Pappus ramosus & plumosus. Antheræ ut in Inula, basi dentatæ Flosculi nudi neutri. D. Schreber. Folia petiolata, ovalia, integerrima, subtus sericeo-argentea.

1019. C H R Y S O C O M A.

Recept nudum. *Pappus* simplex. *Cal.* hemisphæricus, imbricatus. *Stylus* vix flosculis longior.

8. *CHRYSOCOMA Linosyris.* C. herbacea, foliis linearibus glabris, calycibus laxis *Fl. suec. monsp. pal.*

Kniph. cent. 2. n. 14. *Hall. helu. n.* 144. *delph. burg.*

C. calycibus laxis. *Hort. cliff.*

Chysocome dioscoridis & plinii. *Col. ecphr.* 1. p. 81. t. 82.

352 SYNGENESIA. POLYGAMIA ÆQUALIS:

Linofyris nuperorum. Lob. hist. 223.

Osyris austriaca. Clus. hist. 1. p. 325.

Linariæ tertium genus. Trag. 358. R.

Linaria folioso capitulo luteo major & minor. Bauh. pin. 213.

Habitat in Europa temperatiore. 24

1022. SANTOLINA.

Recep. paleaceum. Pappus nullus. Calyx imbricatus, hemisphæricus.

1. **SANTOLINA** *Chama-Cyparissus*. S. pedunculis unifloris, foliis quadrifariam dentatis. *Hall. helu. n. 123. Ludw. ed. t. 198. Blackw. t. 346. Kniph. cent. 6. n. 80. Regn. bot.*

Abrotanum femina, foliis teretibus. Bauh. pin. 136.

Abrotanum femina vulgare. Clus. 1. p. 341.

Habitat in Europa australi 7

2. **SANTOLINA** *rosmarinifolia*. S. pedunculis unifloris, foliis linearibus margine tuberculatis.

S. pedunculis unifloris, capitulis globosis, foliis linearibus integerrimis. *Mill. dict.*

Abrotanum femina, foliis rosmarini, majus. Bauh. pin. 137. Moris. hist. 3. p. 12. f. 6. t. 3. f. 22.

Abrotanum femina 4. Clus. hist. 1. p. 342.

β. *Abrotanum femina, foliis rosmarini, minus. Bauh. pin. 137. Kniph. cent. 7. n. 80.*

Santolina (minor.) pedunculis unifloris, foliis linearibus confertis obtusis. Mill. dict. n. 6. R.

γ. *Abrotanum femina viridis. Bauh. pin. 137.*

δ. *Abrotanum femina, flore majore, foliis villosis & incanis. Bauh. pin. 137.*

Santolina (villosa) pedunculis unifloris, calycibus globosis, foliis quadrifariam dentatis tomentosis. Mill. dict.

Habitat in Hispania.

Herba vix suffrutescens. Folia margine utroque bifariam tuberculis crenulata, sed in caule florescente linearia, ad apicem utrinque denticulata. Pedunculi longi terminales, uniflori.

3. **SANTOLINA** *erecta*. S. pedunculis unifloris, foliis bipinnatis, caulibus simplicibus,

Santolinoides

SYNGENESIA. POLYGAMIA ÆQUALIS. 393

Santolineoides alpina saxatilis, foliis glaucis & veluti argenteis, floribus luteis. *Mich. gen.* 31. t. 27.

Pyrethrum alterum minus, cæspitosa radice, anthemidis flore, italicum. *Barr. rar.* 1105. t. 522.

Habitat in monte Moronis Italiae. 24

Flores absque flosculis femineis.

4. *SANTOLINA anthemoides*. S. pedunculis unifloris, foliis bipinnatis, caule ramosissimo villosus.

S. perennis, chamæmeli folio, caule ramoso. *Vaill. act.* 372.

Habitat in Hispania, Italia. Cl. Astræmer, Sibiria. 24

Planta palmaris. Caulis villosus foliosus. Folia Chamomillæ pubescentia. Pedunculi terminales, foliis longiores, fructiferi rigidiusculi.

1023. ATHANASIA.

Recept. paleaceum. Pappus paleaceus, brevissimus; Cal. imbricatus.

4. *ATHANASIA maritima*. A. pedunculis bifloris, foliis lanceolatis crenatis obtusis tomentosis.

Santolina corymbo terminali subdiviso, foliis oblongis integerrimis obtusis. *Hort. cliff.*

Filago maritima. Sp. dict. t. 135.

Gnaphalium maritimum. Bauh. pin. 263.

Chrysanthemum perenne gnaphaloides maritimum. Moris. 3. p. 81. f. 6. t. 4. f. 47.

Habitat in Europæ australis marisque Mediterranei maritimis. 24

Semina ancipitia pappo destituta; an itaque potius *Santolinæ species*? Media inter *Athanasias* & *Santolinas*: Planta tota tomentosa Paleæ longitudine calycis.

POLYGAMIA SUPERFLUA.

1024. TANACETUM.

Recept. nudum. Pappus submarginatus. Cal. imbricatus, hemisphaericus. Cor. radii obsoletæ, 3-fidæ (interdum nullæ omnesque flosculi hermaphroditi.)

5. TANACETUM *annuum*. T. foliis bipinnatifidis linearibus acutis corymbis tomentosis.

T. floribus corymbosis, ramis summis primordialibus, caule longioribus, foliis radicalibus bipinnatis, caulinis rameisque difformibus. *Gouan. illustr. 66.*

T. *annuum* villosum, absinthii austriaci folio. *Vaill. aë. 1716. p. 281.*

Santolina corymbis simplicibus, foliis linearibus confertis. *Mill. dict. 1. 227. f. 1.*

Absinthium corymbiferum *annuum*. *Tournef. inst 458.*

Elichrysum foliis abrotani. *Bauh. pin. 264. Pluk. phys. 160. f. 1.*

Elichryson. *Clus. hist. 1. p. 325. t. 326.*

Habitat in Hispania, Hetruria, Gallia. ☉

Rami longiores quam umbella primaria, divaricati. Folia linearia, pinnato-quinquesida, trifidaque. Differt a congeneribus defectu Radii feminei; ab *Athanasia* defectu palearum.

6. TANACETUM *vulgare*. T. foliis bipinnatis incisisserratis. *Succ. carn. ged. pal. fil. Ludw. ed. 1. 22. Blackw. 1. 464. Knorr. del. 2. 1. T. 2. Regn. bot. lithua. lugd. burg.*

T. foliis pinnatis, pinnis semipinnatis, acute dentatis. *Hall. helv. n. 132.*

T. foliis pinnatis planis: pinnis serratis. *Fl. lapp. 295.*

T. *vulgare* luteum. *Bauh. pin. 132.*

Athanasia f. 1. *Boiss. Hist. 955.*

Artemisia monacha. *Boiss. Hist. 46.*

SYNGENESIA. POLYGAMIA SUPERFLUA. 355

T. Tanacetum (*crispum*) foliis crispis. *Bauh. pin.* 132.
Mattusch. l. c.

T. crispum *Dod. pempt.* 36.

Habitat in Europæ aggeribus. 24

Radius calidiore æstate prodit.

7. TANACETUM Balsamita. T. foliis ovatis integris
ferratis. *Kniph. cent.* 2. n. 92. *Hall. helu. n.* 133. *Regni-*
bot.

Mentha hortensis corymbifera. *Bauh. pin.* 226.

Costus hortensis. *Dalech. hist.* 678.

Balsamita major. *Dodon. com.* 299. *Hall. R.*

Balsamita mas. *Blackw. t.* 98. *R.*

Habitat in Hetruria, Narbona, Helvetia. 24

1025. ARTEMISIA:

Recept. subvillosum, vel nudiusculum. *Pappus* nullus.
Cal. imbricatus squamis rotundatis, conniventibus.
Cor. radii nullæ.

* *Fruticosa, erecta.*

2. ARTEMISIA judaica. A. fruticosa, foliis obovatis
obtusis lobatis parvis floribus paniculatis pedicellatis.
Mat. med. 183. *Stechm. de artem.* p. 20.

A. tota cinerea. *Gron. orient.* 259.

Absinthium halepense tenuifolium, grati odoris, coma
delicatiore. *Pluk. alm.* 4. t. 73. f. 2.

Absinthium Santonicum judaicum & alexandrinum. *Bauh.*
pin. 139.

Habitat in Judæa, Arania, Numidia. 7

Caulis suffruticosus, sesquipedalis, paniculatus, sub-
pubescens, cinereus. Folia obovata, 3. f. 5-loba, obtusa,
plana, subtomentosa, cinerea, minuta, lacinia media
latiore. Panicula racemosa: racemis ultimis pedunculatis
& pedicellatis. Flores subrotundi, depressiusculi, magni-
tudine seminis coriandri.

3. ARTEMISIA æthiopica. A. fruticosa, foliis palmatis
linearibus minutissimis, floribus racemosis pedunculatis.

Abrotanoides africanum, foliis cinereis muscosis, capitulis
florum globosis magnis. *Rai. suppl.* 232.

Habitat in Æthiopia, Hispania. 24

Frutex pedalis, ramosus, albus, obsoletissime tamen-

356 SYNGENESIA. POLYGAMIA SUPERFLUA

tosus. Folia minuta, conferta, subtomentosa: lacinis linearibus angustissimis. Flores racemosi, pedunculati, nutantes, subglobosi. Receptaculum nudum.

4. ARTEMISIA *contra*. A. fruticosa, foliis palmatis linearibus minutis, panicula racemosa, floribus sessilibus.

Habitat in Persia. H. D. Lerche.

Frutex paniculatus, erectus, tomentosus, albus: Ramis magis tomentosis. Folia linearia, saepe conferta, minutissima, obtusiuscula, tomentosa. Panicula e ramis saepius simplicibus: ramis insident. Spicae alternatim sparsae, ovatae, minimae, e minoribus floribus imbricatae minus tomentosae.

5. ARTEMISIA *Abrotanum*. A. fruticosa, foliis setaceis ramosissimis.

A foliis ramosissimis setaceis, caule erecto suffruticosa.

Monsp. carn. Blackw. t. 555.

Abrotanum mas angustifolium majus. Bauh. pin. 136.

Abrotanum mas. Dod. pempt. 21.

Habitat in Syria, Galatia, Cappadocia, Italia, Græcioliæ, Montspeli montibus apricis. H.

Humilis s. pedalis frequenter fruticulus; cultus hominis altitudinem attingit. Rai. extr. 46.

* * Procumbentes, ante florescentiam.

8. ARTEMISIA *campestris*. A. foliis multifidis linearibus, caulibus procumbentibus virgatis. *Fl. spec. carn. pal. Weber. spicill. p. 21. lugd. lithu. burg.*

A. caulibus declinatis, foliis sericeis, caulescentibus, pinnatis, pinnis multifidis acutis. *Hall. helv. n. 131.*

Abrotanum campestre. Bauh. pin. 136.

Ambrosia altera. Cam. epit. 597.

Habitat in Europæ campis apricis, aridis. 24

Receptaculum nudum. Racemi simplices, alterni, floribus solitariis.

9. ARTEMISIA *palustris*. A. foliis linearibus pinnatis integerrimis, floribus glomeratis subsessilibus.

A. herbacea, foliis multifidis corymbis in spica interrupta glomeratis sessilibus. *Gmel. sib. 2. p. 119. t. 55.*

Absinthium capillaceis foliis, flosculis aureis densissime congestis. Amm. ruth. 197.

Habitat in Sibiria.

Caules procumbentes, adscendentes. Folia Coronopi, linearia, pinnatifida: segmentis. 5. s. 7. Flores glomerati, subsessiles, flavi. Receptaculum nudum.

SYNGENESIA: POLYGAMIA SUPERFLUA. 357

10. ARTEMISIA crithmifolia. A. foliis compositis divaricatis linearibus carnosis glabris, caule adscendente paniculato.

Abrotanum hispanicum maritimum, folio crasso splendente & rigido. *Tournef. inst.* 459.

Absinthium inodorum, foliis crithmi lucidis obscure virentibus. *Morif. hist.* 3. p. 11. *Rai. suppl.* 232.

Habitat in Lusitaniæ littoribus arenosis. Læfling.

Habitus *A. campestris*, glaber. **Caules** semipedales, s. pedales, adscendentes, **Panicula** florum terminati. **Folia** pinnatifida trifidaque, linearia, glabra, carnosae. **Flores** *A. campestris*. **Receptaculum nudum.**

11. ARTEMISIA maritima. A foliis multipartitis tomentosis, racemis cernuis, flosculis femineis ternis. *Geß. gallob. vind.*

A. foliis compositis multifidis tomentosis, ramis floriferis nutantibus. *Fl. suec.*

A. foliis compositis multifidis, racemis simplicibus alternis, floribus erectis sessilibus. *Sauv. monsp.*

Absinthium seriphium belgicum. *Bauh. pin.* 139. *Bauh. hist.* 3. p. 178.

A. Absinthium seriphium germanicum. *Bauh. pin.* 139.

β. Absinthium seriphium gallicum. *Bauh. pin.* 139.

Habitat in Europæ septentrionalioris littoribus maris. 24
Receptaculum nudum.

12. ARTEMISIA glacialis. A. foliis palmatis multifidis sericeis, caulibus adscendentibus, floribus glomeratis fastigiatis. *Delph.*

Absinthium foliis sericeis, petiolatis, pinnatis trifidis & quinquefidis. *Hall. helv. n.* 125.

Absinthium alpinum candidum humile. *Bauh. pin.* 139. *prodr.* 71. *Rai. suppl.* 131. *All. pedem.* 5. t. 1. f. 3.

Habitat in Helvetia. Valesia. 24

Statura. *A. rupestris*. **Folia** parva, tripartita, linearia: laciniis trifidis, subfastigiata, albo-sericea, tomentosa, brevissima, longe petiolata, inferiora petiolis caulem vaginantia. **Caules** adscendentes, spithamæi, simplicissimi, subtomentosi, foliosi. **Flores** globosi, lutei, terminales, conglomerati, **Pedunculis** brevissimis, **Calycibus**, subtomentosis. **Receptaculum pilosum.**

13. ARTEMISIA rupestris. A. foliis pinnatis, caulibus adscendentibus hirsutis, floribus globosis cernuis: receptaculo papposo. *Fl. suec.* * *Scop. carn. delph.*

358 SYNGENESIA. POLYGAMIA SUPERFLUA.

A. foliis pinnatis pilosis, ramis adscendentibus, floribus oblongis suberectis. *Gmel. sb. 2. p. 111. t. 49. f. 1.*
(Huic receptaculum nudum. R.)

Abinthium ex alis floridum, foliis angustis sericeis pinnatis, pinnis trifidis & quinquefidis. *Hall. helu. n. 126.*

Abinthium ponticum repens & supinum. *C. B. 139. Pluk. alm. 3. t. 73. f. 1.*

β. Abinthium pumilum palmatum minus, argenteo sericeoque folio. *Booc. mus. 2. p. 81. t. 71. Barr. rar. 106. t. 462.*

Abinthium alpinum incanum. *Bauh. pin. 139. prodr. 71?*

Abinthium orientale tenuifolium argenteum & sericeum, flore magno. *Tournef. cor. 34.*

Abinthium alpinum spicatum, foliis petiolatis bis trifidis, caulibus pinnatis. *Allion. pedem. 1. p. 3. t. 1. f. 2.*

γ. Abinthium viride, foliis multifidis linearibus. *Gmel. sb. 2. p. 131. t. 64. f. 2.*

Habitat in Sibiria, Oelandia rupibus calcareis.

Differt C. B. β. (ex Bursero) planta ramis longioribus, aliisque, forte ex loco natis; Sibirica magis fructuosa est. Planta in rupibus alba sericea; in horto major, streviridis, nuda. Receptaculum pilosum est. Planta ex loco varia.

Ex planta Allionii novam speciem faciunt Hallerus ac Stechmann. quæ est Artemisia Genipi. *Stech. diff. de artemis. p. 17.* seu Artemisia foliis sericeis, radicalibus, petiolatis, pinnatis, incis, caulibus semipinnatis. *Hall. helu. n. 127.* Receptaculum nudum est. R.

* * * Erectæ herbaceæ, foliis compositis.

14. ARTEMISIA pontica. A. foliis, multipartitis subtus tomentosis, floribus subrotundis nutantibus: receptaculo nudo. *Jacq. austr. 1. 99. Ludw. ed. 1. 26. Blackw. 1. 527. Kniph. cent. 6. n. 11.*

A. foliis sericeis, duplicato-pinnatis, pinnulis parallelis. *Hall. helu. n. 129. R.*

Abinthium ponticum tenuifolium incanum. *Bauh. pin. 138.*

Abinthium tenuifolium. *Dod. pempt. 24.*

Habitat in Hungaria interiore, Pannonia, Thracia, Mysia: locis apricis, aridis. 24

15. ARTEMISIA annua. A. foliis triplicato-pinnatis utrinque glabris, floribus subglobosis nutantibus: receptaculo glabro conico. *Mill. dict. n. 14.*

A. radice annua, viridis, foliis pinnatis.

SYNGENESIA. POLYGAMIA SUPERFLUA. 359

calycibus subrotundis fere erectis ex alis copiosissimis.

Gmel. sib. 2. p. 125. n. 108.

Absinthium, tanacetifolio, odoratissimum. *Amm. ruth. 142. t. 193. f. 23. imperfecte.*

β. Absinthium, folio tenui multifido, camphoram spirans. *Bærh. lugdb. 1. p. 126.*

Habitat in Sibiria montosis. ☉ ♂

16. ARTEMISIA tanacetifolia. A. foliis bipinnatis subtus tomentosis nitidis: pinnis transversis, racemis simplicibus. *Mill. dict. n. 20.*

Absinthium foliis radicalibus multoties pinnatis; caulinis pinnatis longe petiolatis, floribus parvis raris sessilibus. *Allion. pedem. 6. t. 2. f. 1.*

Artemisia radice perenni, foliis pinnatis pinnatifidis incisis ferratis, calycibus subrotundis viridibus nutantibus. *Gmel. sib. 2. p. 122. n. 107.*

Habitat in Sibiria, Pedemontio. ℥.

Receptaculum pillosum. Hall.

17. ARTEMISIA Absinthium. A. foliis compositis multifidis, floribus subglobosis pendulis: receptaculo villoso. *Fl. suec. carn. gallob. herb. sil. Ludw. ed. t. 76. Kniph. cent. 4. n. 8. Regn. bot. lithua. delph. lugd. burg.*

Absinthium foliis incanis, pinnatis, pinnis primis bilobatis, pinnulis incisis, lanceolatis, spicis heteromallis, floribus pendulis. *Hall. helu. n. 124.*

Absinthium incanum, foliis compositis latiuscule multifidis, floribus subglobosis pendulis. *Gmel. sib. 2. p. 129. t. 63.*

Absinthium ponticum s. romanum, officinarum s. dioscoroidis. *Bauh. pin. 138.*

Absinthium. *Cam. epis. 452. Blackw. t. 17. Knorr. del. 2. t. A. 17.*

Habitat in Europæ ruderatis, aridis. ℥

Receptaculum pilosum.

18. ARTEMISIA vulgaris. A. foliis pinnatifidis planis incisis subtus tomentosis, racemis simplicibus recurvatis, florum radio quinquefloro. *Carn. gallob. pal. sil. Ludw. ed. t. 153. Blackw. t. 431. Regn. bot. Lugd. lith. burg.*

A. foliis inferne tomentosis, pinnatis, pinnis acute dentatis, spica paniculata erecta. *Hall. helu. n. 130.*

A. foliis pinnatifidis planis laciniatis, floribus erectis. *Fl. suec.*

vulgaris major. *Bauh. pin. 137.*

360 SYNGENESIA, POLYGAMIA SUPERFLUA:

A. vulgaris. *Bauh. hist.* 3. p. 184.

A rubra & alba. *Tatarn.* p. 7 & 8.

Habitas in Europæ cultis, ruderalis. 24

Receptaculum nudum.

*** *Folius simplicibus.*

20. **ARTEMISIA** *carulefcens*. A. foliis caulinis lanceolatis integris : radicalibus multifidis , flosculis femineis ternis.
cap.

Artemisia. Gmel. fib. 2. p. 131. t. 64. f. 1. sec. Pallas. R.

Absinthium maritimum, *lavendulae folio*. *Bauh. pin.* 179.

Marif. hist. 3. p. 7. f. 6. t. 1. f. 5. *Duham. arb.* 1. t. 24. f. 5.

Absinthium angustifolium. *Dod. pempt. 26.*

Abrotanum latifolium rarius, artemisiæ folio. *Col. ecphr.*, 2
p. 75. l. 76.

Habitat in Europæ australis litoribus maris.

Folia utrinque villosa. Flores cylindrici, nutantes : sepalis radii ternis ; hinc nimir affinis *A. maritima*, quoniam Folia suo caelo recedunt.

21. ARTEMISIA *Dracunculus*. A. foliis lanceolatis glabris integerrimis. 126, t. 59 & 60. & f. 1. *cam.*

Abrotanum lini folio acriori & odorato. Tournesf. inf. 459.

Dracunculus hortensis. *Bauh. pin.* 98. *Blackw.* t. 316.

Draco herba. *Dod.* *pempt.* 709.

Habitus in Sibiria, Tartaria. 24

Receptaculum nudum. Hall. flosculi feminei 6, hermaphroditi 12. Toret. app. 106.

22. *ARTEMISIA chinensis*, A. foliis simplicibus tomentosis
obtusis lanceolatis; inferioribus cuneiformibus trilobis.

A. foliis radicalibus a caulinis diversis, *Gmel.fib. 2. p. 61. f. 1. 2.*

A. foliis simplicibus tomentosis obtusis, lanceolatis, inferioribus cuneiformibus bilobis, receptaculo villosa. *Zinn. gatt.* 298. R.

Abſinthium maritimum ſnarum, lavendulæ folio, pulchrioribus Corymbia, inodorum, ſapore aromatico. *Plut. amalth* 3. 4. 153. f. 5.

Habitat in China, Lagerstrom. Sibiria.

Polia tomentosa, pollicem transversum longa : inferiora papiformia : triloba; superiora lanceolata, obtusa. Flores in summis, conferti in Racis in saepius ovatum, simplicibus, subnudi. Calvae las

— sobre Moxa Chubentec

1026. GNAPHALIUM.

Recept. nudum. *Pappus* plumosus. *Cal.* imbricatus, squamis marginalibus rotundatus, scariosis, coloratis.

* *Fruticosa* *Argyrocoma.*

9. GNAPHALIUM *teretifolium*. G. fruticosum, foliis confertis teretiusculis, corymbis ramosis, calycibus extus ferrugineis.

G. fruticosum ramosum, corymbis ramosis, foliis confertis teretiusculis. *Hort. cliff.* 401. *

G. frutescens, foliis tenuissimis teretibus, ramis creberrimis. *Burn. afr.* 217: t. 77. f. 3.

Millefolium æthiopicum, ericæ foliis, incanum, flore specioso. *Pluk. alm.* 251. t. 308. f. 2.

Habitat in Æthiopia. ☿

Calyces glabri toti: extus ferruginei; interius albid. Folia subulata, nuda, subtus tomentosa: revoluta margine.

* * *Fruticosa* *Chrysocoma.*

11. GNAPHALIUM *stæchas*. G. fruticosum, foliis linearibus, corymbo composito, ramis virgatis. *monsp. carn. sil. Blackw.* t. 438. *Kniph. cent.* 6. n. 45. *lugd.*

Filago foliis linearibus, oris contractis, ramis simplicissimis, umbella convexa terminatis. *Hall. helu.* n. 145.

Helichrysum f. *Stœchas* citrina angustifolia, *Bauh. pin.* 264.

Helichrysum f. *Chrysocome* angustifolia vulgaris. *Moris. hist.* 3. p. 401. f. 7. t. 11. f. 7.

Chrysocome f. *Stæchas* citrina minor. *Barr. ic.* 510. 409. 278.

Stœchas citrina. *Dod. pempt.* 268. *Tournef. inst.* 452.

Habitat in Germania, Gallia, Hispania, Oriente; in collibus aridis. ☿

Flosculi omnes androgyni. *Hall. R.*

* * * *Herbacea* *Chrysocoma.*

22. GNAPHALIUM *arenarium*. G. herbaceum, foliis lanceolatis: inferioribus obtusis, corymbo composito, caule simplicissimo. *suec. ged. pal sil. Fl. dan.* 641. *Ludw. ed.* t. 97. *Blackw.* t. 524. *Kniph. cent.* 2. n. 27. *delph. lith.*

Elichrysum f. *Stœchas* citrina latifolia. *Bauh. pin.* 264. *It. el.* 160.

362 SYNGENESIA. POLYGAMIA SUPERFLUA.

Stoechas citrina germanica, latiore folio. *Bauh. hist.* 3. p. 153. *Rai. hist.* 281.

Habitat in Europæ campis arenosis. ☉

27. *GNAPHALIUM luteo-album*. G. herbaceum, foliis semiamplexicaulibus ensiformibus repandis obtusis utrinque pubescentibus, floribus conglomeratis. *Monf. pal. Kniph. cent.* 1. n. 37. *delph. burg.*

Filago foliis mollissimis, tomentosis, obtusis, umbellis convexis, electrinis. *Hall. helv.* 147.

Chrysocome citrina supina, latifolia, italica. *Barrel. ic.* 367. *Hall. R.*

Chrysocome altera. *Clus. hist.* 267. *Hall. R.*

Elichrysum sylvestre latifolium, capitulis conglobatis. *Bauh. pin.* 264.

Gnaphalium majus, lato oblongo folio. *Bauh. pin.* 263.

Pluk. alm. 171. t. 31. f. 16. *Moris. hist.* 3. p. 88. f. 7. t. 11. f. 20.

Gnaphalium ad stoechadem citrinam accedens. *Bauh. hist.* 3. p. 160.

Habitat in Helvetia, G. Narbonensi, Hispania, Lusitania. ☉

Planta valde lanata. Calyces flavo-albicantes, molles, squamis ovato-lanceolatis. Radius femineus multiplex.

Flosculi in ambitu feminei, in centro pauci androgyni. *Hall. R.*

*** *Herbacea Argyrocoma.*

30. *GNAPHALIUM sanguineum*. G. herbaceum foliis decurrentibus lanceolatis tomentosis planis apiculo nudo terminatis. *Amœn. acad.* 4. *

G. ægyptiacum, latiore folio, ramosius, flore ex albicante purpureo. *Pluk. mant.* 91.

G. syriacum. *Barr. ic.* 34.

Gnaphalio montano affinis ægyptiaca. *Bauh. pin.* 264.

Chrysocoma syriaca, flore atro-rubente. *Breyn. cent.* 146.

Baccharis diofcoridis. *Rau. it.* 285. t. 285.

Habitat in Ægypto, Palæstina, Libano, Carmel.

31. *GNAPHALIUM foetidum*. G. herbaceum, foliis amplexicaulibus integerrimis acutis subtus tomentosis, caule ramoso. *Kniph. cent.* 2. n. 28.

G. africanum latifolium foetidum, capitulo argenteo. *Comm. hort.* 2. p. 111. t. 56. *Volkam. norimb.* 194. t. 194.

Conyza africana graveolens, capitulis argenteis. *Pluk. alm.* 117. t. 243. f. 1. *Moris. hist.* 3. p. 115. f. 7. t. 20. f. 32.

Habitat in Æthiopia. ☉

SYNGENESIA. POLYGAMIA SUPERFLUA. 363

Variat Calycibus argenteis & aureis.

38. GNAPHALIUM margaritaceum. G. herbaceum, foliis lineari – lanceolatis acuminatis alternis, caule superne ramoso, corymbis fastigiatis. *Kniph. cent. 12. n. 50.*

Filago foliis lanceolatis, subtus tomentosis, floribus umbellatis. *Hall. helu. n. 146.*

G. latifolium americanum. *Bauh. pin. 263.*

G. americanum. *Clus. hist. 1. p. 327.*

Habitat in America septentrionali, Kamtschatka. 4

Flores omnes androgyni. *Hall. l. c.*

40. GNAPHALIUM dioicum. G. sarmentis procumbentibus, caule simplicissimo, corymbo simplici terminali, floribus dioicis. *suec. ged. carn. gallob. herb. pal. sil. Ludw. ed. 1. 163. Kniph. cent. 3. n. 44. lugd. lith. delph. burg.*

Filago flagellis reptans, sexu distincta, flosculis omnibus androgynis. *Hall. helu. n. 157.*

Gnaphalium caule simplicissimo, floribus coloratis terminato. *Fl. lapp.*

Gnaphalium (mas) montanum, flore rotundiore. *Bauh. pin. 263.*

Pilosella minor. *Dod. pempt. 68. f. exterior.*

Gnaphalium (femina) montanum, longiore & folio, &c. *Bauh. pin. 263.*

Pilosella minor. *Dod. pempt. 68. f. interior.*

Habitat in Europæ apricis aridis. 4

Pappus vix plumosus. *Leers.*

41. GNAPHALIUM alpinum. G. sarmentis procumbentibus; caule simplicissimo, capitulo terminali aphylo, floribus oblongis. *Fl. suec. carn. Flor. dan. 1. 332. delph.*

Filago caule simplici, umbella pauciflora, calycis basi tomentosa. *Hall. helu. n. 150.*

Gnaphalium alpinum minus. *Bauh. pin. 264.*

Filago alpina minor erecta. *Scheuch. alp. 133.*

Habitat in Alpibus. Lapponiæ, Helvetiæ. 4

Caulis simplicissimus, digito brevior, foliis 3 s. 4 lanceolatis. Folia radicalia lanceolato-cuneiformia, supra viridia lævia subtus tomento striato albida, ut margo etiam a latere superiore albus conspiciatur. Ad radicem sarmenta ut G. dioico. Flores terminales, pauci conferti, absque foliis Calyx cylindricus, basi ventricosus, squamis apice membranaceis, lanceolatis, acutis erectis. Flosculi feminei, in planta a me culta, absque staminibus an G. dioici varietas

364 SYNGENESIA. POLYGAMIA SUPERFLUA.

ob sexum, sarmenta, caulem nudum; sed matrem adhuc nondum observavi.

Sarmenta nulla; Flores hermaphroditi. Hall. Scop. R.

***** *Filaginoidea.*

44. GNAPHALIUM sylvaticum. G. caule herbaceo simplicissimo, floribus sparsis. *succ. ingr. ged. gallob. herb. pal. fil. Fl. dan. 254. Eugd. delph. lithu. burg.*

Filago foliis linearibus, alis spiciferis. Hall. kelu. n. 148.

Filago germanica altera secundum folia florida. Lob. advers. p. 201. ic. 482. Hall. R.

Gnaphalium anglicum vel belgicum. Lob. ic. 482. Hall. R.

G. majus, angusto oblongo folio, alterum. Bauh. pin. 263.

G. rectum. Bauh. hist. 3. p. 160.

Habitat in Europæ sylvis arenosis. ♂

Varietas forte? minor calyce simillimo ex alpinis europæis.

Flores in ambitu feminei, in centro hermaphroditi. Hall. R.

45. GNAPHALIUM supinum. G. caule herbaceo simplici procumbente, floribus sparsis. *delph.*

Filago caule simplicissimo, spica pauciflora. Hall. kelu. n. 149.

Gnaphalium supinum, lavandulæ folio. Boer. mus. 107. t. 85.

G. oblongo folio. Scheuchz. alp. 2. p. 134.

Habitat in alpinis Helveticis, Italicis.

Simillimum G. sylvatico, sed humile & caule dum flores decumbente.

46. GNAPHALIUM uliginosum. G. caule herbaceo ramofo diffuso, floribus confertis terminalibus. *succ. carn. norv. gallob. herb. pal. fil. lith. delph. burg.*

Filago caule ramofofissimo, diffuso, umbella sessili terminato. Hall. kelu. n. 151.

Gnaphalium longifolium humile ramofo, capitulis nigris. Rai. hist. 295. angl. 3. p. 18.

G. annuum serotinum, capitulis nigricantibus, in humidis gaudens. Moris. hist. 3. p. 92. ☉

G. medium. Bauh. bas. 78. Hall. R.

Habitat in Europæ paludibus, ubi aqua stagnant. ☉

1027. XERANTHEMUM.

Recept. paleaceum. Pappus setaceus. Cal. imbricatus, radiatus : radio colorato.

• *Receptaculo paleaceo. Pappo quinqueseto.*

1. XERANTHEMUM *annuum*. X. herbaceum, foliis lanceolatis patentibus. *monsp. Jacq. austr. t. 388. carn. Kniph. cent. 4. n. 100. Hall. helu. n. 122. Mill. illustr. delph. burg.*

X. receptaculis paleaceis, seminum pappo quinqueseto. *Hort. cliff.*

X. oleæ folio, capitulis simplicibus incanis non foetens, flore majore violaceo. *Morif. hist. 3. p. 43. f. 6. t. 21. f. 2.*

Jacea oleæ folio, capitulis simplicibus. *Bauh. pin. 272.*

Parmica austriaca. *Clus. hist. 2. p. 11.*

β. Xeranthemum, (*inapertum*.) oleæ folio, capitulis simplicibus, incanum foetens, flore purpurascens minore. *Morif. hist. 3. p. 43. f. 6. t. 12. f. 1.*

X. foliis lineari-lanceolatis, utrinque tomentosis. *Mill. dict. n. 2.*

X. capitulis inapertis. *Hall. helu. ed. 1. p. 709. t. 23.*

Jaceæ oleæ folio, minore flore. *Bauh. pin. 272.*

γ. Xeranthemum (*orientale*) orientale, fructu maximo. *Tournef. cor. 38.*

X. foliis lineari-lanceolatis, capitulis cylindratis, semine maximo. *Mill. dict. n. 3.*

Jaceæ oleæ folio, capitulis compactis. *Bauh. pin. 272.*

δ. Jacea incana, cyani capitulis. *Bauh. pin. 272.*

Habitat α. in Austria, β. in Italia, Helvetia, G. Narbonensi. ☉.

Radii stigma simplex; Semine nudo. Disci stigma subbifidum; Semine 5. aristato.

Planta α robusta, erecta; Seminibus pluribus minoribus; Calyce scarioso; Paleis integris; Radio polypetalis.

Planta β angustior, debilis; Seminibus 6. f. 7. majoribus; Calycis squamis solo margine membranaceo; Paleis laceris; Radio subhexapetalo.

366 SYNGENESIA. POLYGAMIA SUPERFLUA

1028. CARPESIU M.

Recept. nudum. *Pappus*. nullus. *Cal.* imbricatus;
squamis exterioribus reflexis.

1. *CARPESIU M. cernuum*. C. floribus terminalibus. *Hall.*
helv. n. 134. *cern.* *Jacq. austr.* t. 204. *Barrel.* ic. 1142.
delph.

Balsamita conyzæ folio, flore cernuo. *Vaill. ad* 336.

After atticus, foliis circa florem mollibus, *Bauh. pin.* 266.

After cernuus, *Col. esphr.* 1. p. 251. t. 252.

Chrysanthemum conyzoides cernuum, foliis circa florem
mollibus. *Morif. hist.* 3. p. 18. f. 6. t. 5. f. 27.

Habitas in Italia, Helvetia, &c. 26

1029. BACCHARIS.

Recept. nudum, *Pappus* pilosus. *Calyx* imbricatus;
cylindricus. *Flosculi* feminei hermaphroditi
immixti.

4. *BACCHARIS halimifolia*. B. foliis obovatis superne
emarginato-crenatis. *Du Ham. arb.* 1. t. 35.

Senecio virginianus arborescens, atriplicis folio. *Rai. hist.*
1769. *Herm. parad.* 225. t. 225.

Pseudo-Helichrysum virginianum frutescens, halimi
latis foliis glaucis. *Morif. hist.* 3. p. 90. f. 7. t. 10.
f. 4.

Elichryso affinis virginiana frutescens, foliis chenopodii
glaucis. *Pluk. alm.* 134. t. 27. f. 2.

Argyroceme virginiana, atriplicis folio. *Pet. gaz.* t. 7.
f. 4.

Habitas in Virginia. 7

5. *BACCHARIS Dioscoridis*. B. foliis lato-lanceolatis
dentatis sessilibus stipulatis. *Aman. acad.* 4. p. 289*.

Conyza major altera. *Bauh. pin.* 265.

Conyza dioscoridis. *Rauw. it.* 54. t. 54.

Conyza syriæ. *Bauh. hist.* 2. p. 254. *Rai. hist.* 262. *Groc.*
orient. 110.

Habitas in Syria, Ægypto.

Sero flores; in Caldarie viz

1030. C O N Y Z A.

Recept. nudum. Pappus simplex. Cal. imbricatus, subrotundus. Cor. radii trifidæ.

1. *CONYZA squarrosa*. C. foliis lanceolatis acutis, caule herbaceo corymbofo, calycibus squarrosis. *Monf. carn. gallob. fil. Fl. dan. t. 622. pal. lithua. lugd. delph. burg.* C. foliis ovato-lanceolatis, rariter serratis, floribus umbellatis. *Hall. helu. n. 135.*

C. maior vulgaris. *Bauh. pin. 265.*

Conyza. Cam. epit. 612.

Baccharis monspeliensium. Blackw. t. 102.

Habitat in Germaniæ, Belgii, Angliæ, Galliæ siccis. ♂

3. *CONYZA sordida*. C. foliis linearibus integerrimis, pedunculis longis trifloris, caule suffruticoso.

Gnaphalium (sordidum.) fruticosum, foliis linearibus, pedunculis subtrifloris, calycinis squamis lanceolatis.

Spec. ed. monsp.

Elichrysum sylvestre angustifolium, capitulis conglobatis.

Bauh. pin. 264.

Stoechadi citrinæ affinis, capitulis parvis raris. Bauh. hist. 3. p. 156. Rai. hist. 283.

Stoechas citrina spuria, longioribus foliis. Barr. ic. 368. 277.

Ageratum aliud quorundam. Dalech. hist. 778.

Habitat in G. Narbonensi, Italia, ad rupes, muros. ♀

Affinis maxime C. saxatili. Caulis suffruticosus (Lavandulæ), ramosus, albus. Pedunculi recti, longi, apice sæpius triflori. Calyces squamis margine fuscis.

4. *CONYZA saxatilis*. C. foliis linearibus subdentatis, pedunculis longissimis unifloris, caule suffruticoso.

Gnaphalium foliis alternis linearibus acutis planis, pedunculis longissimis unifloris. Hort. cliff.

Helychrysum capitulis singularibus brevibus. Moris. hist. 3. p. 87. f. 7. t. 10. f. 16.

Helichrysum saxatile, singulari capitulo amplo, angusto stoechadis folio. Boc. mus. 142. t. 104.

Elichryso sylvestri, flore oblongo, similis. Bauh. pin. 165. prodr. 123. t. 123.

Habitat in Italia, Istria, Carinthia, Valesia, Palestina, s. b. Spec. ♀

368 SYNGENESIA. POLYGAMIA SUPERFLUA

Caules primum procumbentes, dein erecti. Calyces cylindrici apice petula membranacei. Pappi feminei, plurimi, nudi in ambitu.

1031. ERIGERON.

Recept. audum. Pappus pilosus. Cor. radii lineares, angustissimi.

1. ERIGERON viscosum. E. pedunculis unifloris lateralibus, foliis lanceolatis denticulatis basi reflexis, calycibus squarrosis, corollis radiatis. *Monsp. Jacq. hort. 175. delph. burg.*

E. foliis lanceolatis semiamplexicaulibus dentatis, ramis multifloris, corollis radiatis, caule virgato. *Gouan. illustr.*

Aster foliis serratis, pedunculis simplicibus lateralibus unifloris longitudine folii foliosis. *Hort. cliff. 409.*

Virga aurea major, foliis glutinosis graveolentibus. *Tournef. inst. 580. Vaill. Ad. 579.*

Conyza minor theophrasti, major dioscoridis. *Barr. pin. 167.*

Conyza major. *Dod. pempt. 51. Clus. hist. 2. p. 30.*

Habitat in G. Narbonensi, Hispania, Italia. &

Caules erectus, strictus. Folia amplexicaulis, crebris hirsuta, dispersa pilis, interjectis glandulis viscosa, graveolentia, basi reflexa. Pedunculi diphylli, uniflori, vix foliis longiores. Flores radiati.

2. ERIGERON graveolens. E. foliis sublinearibus integerrimis ramis lateralibus multifloris. *delph. burg.*

E. ramis lateralibus multifloris, foliis lanceolatis integerrimis, calycibus squarrosis. *Spec. fl. 2. monsp.*

E. foliis lanceolato-linearibus integerrimis scissibus decurrentibus, ramis multifloris, corollis radiatis, caule pyramidato. *Gouan. illustr.*

Virga aurea minor, foliis glutinosis & graveolentibus. *Tournef. inst. 484.*

Conyza femina theophrasti, minor dioscoridis. *Barr. pin. 161.*

Conyza minor vera. *Leb. ic. 346. Barr. ic. 370.*

Habitat Monspeli, in Oriente. ☉

Caules erecti, purpurascens. Rami inferne alterni simplices. Folia sessilia, lanceolata, mucronata, scissura, scissura, scissura.

SYNGENESIA. POLYGAMIA ÆQUATIS. 369

la, adspersa pilis minimis apice viscidis, graveolentia. Pedunculi superne laterales, solitarii, adspersi foliolis flore longioribus, uniflori. Calyx oblongus, foliolis patulis, erectis, ante anthesin apice purpurascens. Corolla lutea, parva radio minuto, erecto.

3. ERIGERON *glutinosum*. E. foliis lanceolato-linearibus piloso-viscidis, pedunculis unifloris. *Ger. prou.* †

Conyza montana, foliis glutinosis pilosis. *Bauh. pin.* 265. *Rai. hist.* 265.

Conyza montana saxatilis, hyssopi folio villoso & glutinoso, hispanica. *Barr. ic.* 158.

Conyza montana Myconis. *Dalech. hist.* 1200.

Habitat in Hispaniæ, Galliæ australis montosis maritimis. 24

Pedunculi alterni uniflori pauciores. Flores purpurei.

H. V.

4. ERIGERON *ficulum*. E. calycinis squamis inferioribus laxis florem superantibus, pedunculis foliosis.

Conyza ficula annua, foliis atro-virentibus, caule rubente. *Bocc. sic.* 62. t. 31. f. 4. *Morif. hist.* 3. p. 115. f. 7. t. 20. f. 28. *Pluk. phyt.* 168. f. 2.

Conyza caulibus rubentibus tenuioribus, flore luteo nudo. *Magn. monsp.* 77. t. 76.

Conyza species foliis virgæ aureæ. *Bauh. hist.* 2. p. 1049.

Habitat in Siciliæ, Monspelii paludosis. ☉

Caules rubentes. Flores absque radio, minores: pedunculi secū foliis linearibus, recurvis minutis. Cæterum simillimum E. graveolenti excepto radio omni. Inulæ publicariæ affinis.

6. ERIGERON *canadense*. E. caule floribusque paniculatis, hirtis, foliis lanceolatis ciliatis. *Delph. lugd. lithua. burg.*

E. caule floribusque paniculatis. *Gallob. ged. herb. pal. sil.*

E. floribus paniculatis spicatis. *Hall. helu. n.* 84.

Conyzella. *Dill. cat. app.* 142.

Erigeron racemis lateralibus; foliis imis lanceolatis, dentatis, superioribus linearibus, ciliatis. *Scop. carn.* 1. p. 386. n. 3. ed. 2. n. 1060.

Conyza annua acris alba elatior, linariæ foliis. *Morif. hist.* 3. p. 115. f. 7. t. 20. f. 29. *Bocc. sic.* 85. t. 46.

Virga aurea virginiana hirsuta, flore pallido. *Zanon. hist.* 1. p. 204. t. 78.

Habitat in Canada, Virginia, nunc in Europa australi. ☉

Linnæi spec. plant. Europ. Pars II.

A a

370 SYNGENESIA. POLYGAMIA ÆQUALIS.

10. ERIGERON *ægyptiacum*. E. foliis semiamplexicaulibus . spatulatis dentatis , floribus globosis.

E. serratum. *Forskæhl. ægypt. 148.*

Conyza capitata. f. globosa. *Bocc. sic. 13. t. 7. f. B. Moris. hist. f. 7. t. 20 f. 14. 3. p. 114.*

Jacobæa ægyptiaca , folio glauco coronopi. *Barh. perperam. Habitat in Sicilia , Ægypto. ☉*

Caulis erectus , sesquipedalis , substriatus , virens pubescenti-subviscidus. Folia alterna , sessilia , vix semiamplexicaulia , spatulata , f. obovato-lanceolata , obtusiuscula , subpubescentia , subdentata dentibus fere 5, basi vero integra. Rami ex supremis alis. Flores terminales 4. f. 5. pedunculati , subrotundi , pedunculis purpurascens. Calyx subrotundus squamis subulatis : intimis vix apice sanguinolentis. Corollæ luteæ : flosculis minutis : Radiis nudis , copiosissimis ; pistillis luteis , mox evanescentibus , annulo pappi lato discum quasi cingentibus , *H. V. Mant. 1. p. 112.*

Caulis simplex , cinerascens, Folia lineata. Flores paniculati. Calyx squamis pubescentibus apice patulis. Flosculi feminei brevissimi absque limbo. Stylus bifidus , luteus , capillaris. *Mant. 2. p. 517.*

12. ERIGERON *acre*. E. pedunculis alternis unifloris , succed. gallob. herb. pal. sil. lugd. lituæ. burg.

E. vulgare. *Fl. lapp.*

E. caule alterne ramoso , petiolis unifloris , semiflosculis pappum æquantibus, *Hall. helv. n. 58.*

Conyzoides. *Dill. giff. p. 154.*

Conyza cærulea acris. *Bauh. pin. 265.*

Amellus montanus æquicolorum *Col. ecphr. 2. p. 25. & 26. Habitat in Europæ apricis , siccis. M.*

13 ERIGERON *alpinum*. E. caule subbifloro , calyce subbifuro , foliis obtusis subtus villosis. *carn. delfh.*

E. caule alterne ramoso , petiolis unifloris , semiflosculis pappum superantibus. *Hall. helv. n. 86. R.*

E. uniflorum. *Old fl. dan. t. 292. sec. Linn.*

E. squamis calycinis planis , foliis obtusis subtus villosis *Ger. prou.*

Conyza cærulea alpina major & minor. *Bauh. pin. 265. prodr. 124.*

Asteri montano purpureo similis f. globulariæ. *Bauh. hist. 2. p. 107.*

Habitat in Alpibus Helvetiæ ; Carniolæ. 2

Pappus ruffo-ferrugineus

~~minimè affine sequenti~~ , 21

SYNGENESIA. POLYGAMIA SUPERFLUA. 371
forte sola varietas. Variat extra alpes Caule paniculato ,
Calyce glabro. Varietas floribus albis datur.

14. ERIGERON uniflorum. E. caule unifloro , calyce piloso. *Fl. suet. * norv. delph.*

E. caule unifloro , calyce tomentoso , radiis semiflosculosis.
Hall. helv. n. 87.

E. polymorphum. *Scop. carn.*

Aster caule unifloro , foliis integerrimis , calyce villosa subrotundo. *Fl. lapp. 307. t. 9. f. 3.*

Aster montanus cœruleus omnium minimus , *Rai. hist. 3. p. 120. Hall. R.*

Conyza cœrulea alpina major. *Bauh. pin. 265. prodr. 124.*

Habitat in Alpibus Lapponiæ , Helvetiæ. 24

Vulgo albo flore est , sed datur etiam flore purpureo. Hall. l. c. R.

17. ERIGERON tuberosum. E. foliis linearibus ramis unifloris , caule suffruticoso. *monsp.*

E. foliis lanceolato-oblongis denticulatis , pedunculis longis terminalibus unifloris. *Gouan. illustr.*

E. foliis rigidis , caule sublignoso paucifloro. *Ger. prou.*

E. caule subunifloro , foliis lanceolato-linearibus rigidis , radice tuberosa. *Sauv. monsp.*

α. Chondrilla bulbosa syriaca , foliis angustioribus. *Bauh. pin. 130.*

Chondrilla altera dioscoridis putata. *Clus. hist. 2. p. 145.*

Conyza marina. *Morif. hist. 3. p. 114. f. 7. t. 19. f. 20. bona.*

β. Chondrilla bulbosa syriaca , fol. latioribus. *Bauh. pin. 130.*

Conyza tuberosa lutea , foliis angustis & rigidis. *Magn. monsp. 77. Morif. hist. 3. p. 114. f. 7. t. 20. f. 15.*

γ. Aster gesneri conyzoides. *Morif. hist. 3. p. 118. f. 7. t. 22. f. 7.*

Habitat in Galloprovincia , G. Narbonensi , Syria.

Caulis brevis , lignosus. Rami simplices , uniflori. Folia sparsa , lanceolata , subpetiolata , integerrima. Flores lutei , Asteris facie , terminales , sessiles. Pappus griseus. Media inter Asters & Erigera. Radix præmorsa. Varietates a Gronovio.

1032. TUSSILAGO.

Recept. nudum. Pappus simplex. Cal. squamæ æquales, discum æquantes, submembranaceæ.

1. TUSSILAGO *Anandria*. T. scapo unifloro squamoso erecto, foliis lyrato-ovatis. *Mill. dict. n. 2.*

T. scapo unifloro, calyce clauso. *Hort. ups. 259. t. 3. f. 1. * Gmel. sib. 2. p. 141. t. 68. f. 1.*

*Anandria. Amœn acad. 1. p. 120. t. 11. **

β. Tussilago scapo unifloro, calyce subaperto. *Gmel. sib. 1. p. 143. t. 67. f. 2.*

Habitat in Sibiria. 24

In frigidioribus locis Calyx clausus flore flosculoso: in calidioribus flore radiato patente. Pappus sessilis, pilosus.

4. TUSSILAGO *alpina*. T. scapo subnudo unifloro, foliis cordato-orbiculatis crenatis. *monsp. sil. Jacq. austr. 2. 246. delph.*

Petalites scapo unifloro, foliis reniformibus. Hall. helv. n. 142. Planta adulta; secundum Jacq. R.

Petalites scapo sæpius unifloro, foliofo; foliis orbiculatis, crenato-dentatis. Scop. carn. n. 2.

Tussilago alpina rotundifolia glabra. Bauh. pin. 197.

T. alpina 2. Clus. hist. 2. p. 113.

β. *T. alpina rotundifolia canescens. Bauh. pin. 197.*

T. discolor. Jacq. austr. 1. 247.

T. alpina. Scop. carn. Jacq. austr. R.

Petalites. Hall. helv. n. 142. planta junior.

Tussilago alpina 1, flore evanido. Clus. hist. 2. p. 113.

Habitat in Alpibus Helvetiæ, Austriæ, Bohemiæ, Sibiria. 24

Flos flosculosus: flosculis femineis in ambigua Scapus ut alterave Bractea.

5. TUSSILAGO *Farsura*. T. scapo imbricato unifloro. foliis subcordatis angulatis denticulatis. *Fl. suec. par. sed. tab. herb. carn. pal. sil. n. 27. 11. den. 595. Blackw. 2. 204. Kniph. cent. 6. n. 96. Knorr. del. 1. t. H. 10. Luda. 2. t. 50. Regn. bot.*

Petalites scapo unifloro, flore radiato. Hall. helv. n. 143.

Tussilago corollis radiantibus, scapo unifloro, foliis orbiculato-cordatis angulosis. gallob.

T. vulgaris. Bauh. pin. 197.

SYNGENESIA. POLYGAMIA SUPERFLUA. 373

Tussilago. Clus. hist. 120. Cam. epit. 590. 591.

Habitat in Europæ argillosis, subtus humidis. 24

Flores radiati. Corollulæ radii integerrimæ. Leers. l. c. R.

7. TUSSILAGO frigida. T. thyrsis fastigiato, floribus radiatis. *Flor. lapp. 303. Fl. suec. Oed. dan. t. 61. delph.*

Petasites floribus spicatis erectis, foliis calycinis lanceolatis, flosculis paucissimis androgynis. Hall. helu. n. 141.

T. scapo imbricato, floribus spicatis radiatis, foliis infra incanis acutis. Gmel. sib. 2. p. 150. t. 70.

Cacalia tomentosa. Bauh. pin. 198. prodr. 102. Schuck. alp. 130. t. 18. f. 1.

Habitat in Alpium Lapponiæ, Helvetiæ, Sibiriae convalibus. 24

Variat floribus flosculis & radiatis.

8. TUSSILAGO alba. T. thyrsis fastigiato, flosculis femineis nudis paucis. *Fl. suec. ged. sil. Fl. dan. t. 524. delph. lithu. burg.*

Petasites spicis laxis, umbellatis, flosculis plerisque androgynis. Hall. helu. n. 139.

Petasites minor. Bauh. pin. 197.

Petasites flore albo. Cam. epit. 593.

Habitat in Europa. 24

Flores flosculosi: flosculis femineis in ambitu.

Flores semiflosculos ligulatos vidit La Chenal. diff. p. 8. R.

9. TUSSILAGO hybrida. T. thyrsis oblongo, flosculis femineis nudis plurimis. *gelr. herb. Reich. franc. delph.*

Petasites floribus spicatis pendulis, petiolis multifloris, calycibus æqualibus, flosculis paucissimis androgynis. Hall. helu. n. 140.

Petasites major, floribus pedunculis longis insidentibus. Dill. elth. 309. t. 230. f. 297.

Petasites in medio majoribus flosculis, reliquis minoribus. Buxb. hal. 258.

Habitat in Germania, Hollandia. 24

Flores flosculosi: flosculis hermaphroditis & femineis.

10. TUSSILAGO Petasites. T. thyrsis ovato, flosculis femineis nudis paucis. *lich. lugd. burg.*

T. thyrsis ovato, flosculis omnibus hermaphroditis. Fl. suec. paris. ged. pal. herb. sil. Knorr. del. 2. t. T. Regn. bot.

Petasites floribus dense spicatis, flosculis androgynis. Hall. helu. n. 138.

Tussilago flosculis omnibus tubulosis fertilibus monoclinis; scapo imbricato thyrsifero. Neck. gallob.

374 SYNGENESIA. POLYGAMIA SUPERFLUA

T. major. Cam. epit. 592.

Petalites. Dodon. 597. Blackw. t. 222. major & vulgaris. Bauh. pin. 197.

Habitat in Europa temperiore. 2.

Flosculi feminei (inter hermaphroditos) 2. f. 3. vix petaloidei, ligulati, albi. Linn. fil.

Corollulae femineae in nostrate penitus nullae. Pappus in hac & T. Farfara & T. hybrida sessilis nec stipitatus. Leers l. c. R.

1033. SENECIO.

Recept. nudum. Pappus simplex. Cal. cylindricus, calyculatus: squamis apice sphacelatis.

** Floribus flosculosis.*

7. *SENECIO vulgaris. S. corollis nudis, foliis pinnato sinuatis amplexicaulibus, floribus sparsis. Fl. suec. paris. ged. Hall. helu. n. 58. pal. fil. Fl. dan. t. 513. Blackw. t. 132. Knorr. del. 2. t. S. 5. Kniph. cent. 6. n. 84. lithu. lugd. burg.*

S. floribus hermaphroditis, foliis amplexicaulibus, pinnato sinuatis. Scop. carn. 1. p. 380 n. 1. ed. 2. n. 1063.

S. fol. amplexicaulibus subbipinnatifidis sinuatis: flosculis omnibus tubulosis monoclinis. Neck. gallob. 353.

S. foliis pinnatifidis denticulatis: laciniis æqualibus patentissimis: rachi lineari. Hort. cliff. 406.

S. minor vulgaris. Bauh. pin. 131. Fl. lapp. 296.

S. Erigeron. Tabern. 168.

Erigeron minus. Dod. pempt. 641.

Habitat in Europæ cultis, ruderatis, succulentis. ☉

** * Floribus radiatis: radio revoluto.*

11. *SENECIO lividus. S. corollis revolutis, foliis amplexicaulibus lanceolatis dentatis, squamis calycinis brevissimis intactis. Hort. ups. * Kniph. cent. 10. n. 79.*

S. foliis sinuatis: basi latioribus amplexicaulibus, floribus corymbosis, radiis disco brevioribus. Roy. lugdb. 165.

Habitat in Hispania. ☉

12. *SENECIO trilobus. S. corollis revolutis, foliis amplexicaulibus, squamis calycinis sphacelatis laceris. Hort. ups. * Jacobæa hispanica minus laciniata, petalis brevissimis. Vaill. paris. 21.*

SYNGENESIA. POLYGAMIA SUPERFLUA. 375

Habitat in Hispania. ☉

13. SENECIO viscosus. S. corollis revolutis, foliis pinnatifidis viscidis, squamis calycinis laxis longitudine perianthii. *Fl. suec. pal. sil. delph. lith. lugd. burg.*

S. foliis viscidis, pinnatis, pinnis semipinnatis, angulosis, calycibus hirsutis. *Hall. helu. n. 60.*

S. corollis revolutis – radiantibus, foliolis calycinis perianthium subæquantibus, foliis pinnatifidis. *Neck. gallob.*

S. foliis pinnatifidis sessilibus: laciniis angulosis: superioribus latioribus, florum radiis disco brevioribus. *Roy. lugdb. 142.*

S. hirsutus viscidus graveolens. *Dill. elth. 347. t. 258. f. 336.*

S. incanus pinguis. *Bauh. pin. 131.*

Habitat in Europæ pagis, urbibus. ☉

Variat corollis non radiatis. *Hall. l. c. R.*

14. SENECIO sylvaticus. S. corollis revolutis, foliis pinnatifidis denticulatis, caule corymbofo erecto. *ged. pal. sil. delph. burg. lithu.*

S. foliis pinnatifidis denticulatis, florum radiis revolutis linearibus. *Fl. suec. parif.*

S. foliis pinnatis, pinnis alterne majoribus, semipinnatis, angulosis, calycibus levibus, radiis brevissimis revolutis. *Hall. helu. n. 59.*

S. corollis revolutis – radiantibus, foliis subpinnatifidis floribus corymbosis. *Neck. gallob.*

S. minor, latiore folio, f. montanus. *Bauh. pin. 131. Dill. elth. 258. t. 258. f. 337.*

Jacobæa, senecionis folio incano, perennis. *Rai. hist. 285. Hall. ien. 177. t. 4.*

Senecio montana mollior, pallido parvo flore. *Barrel. ic. 401. Hall. R.*

Senecium montanum. *Tabern. 169. R.*

β. Jacobæa sicula chrysanthemi cratici folio. *Tournef. Barrel. ic. 262. n. 2. rudis. Gouan. R.*

Jacobæa pumila gallica. *Bocc. sic. t. 41. f. 1. optime. Gouan.*

Habitat in Europæ borealis sylvis cæduis. ☉

Var. β. crescit in Luco Gramuntio inter lapides; glabra est & pusilla. *Gouan. l. c. R.*

15. SENECIO nebrodensis. S. corollis revolutis, foliis lyratis sinuatis obtusis petiolatis, caule hirsuto.

Jacobæa montana, senecionis pallido parvo flore. *Barr. rar. 1081. t. 401.*

376 SYNGENESIA. POLYGANIA SUPERFLUA.

Jacobæa nebrodensis altera, oblongo folio latiori minus incano lacerato. *Rai. suppl.* 179.

Habitat in Sicilia, Hispania, Pyrenæis. Alstræmer. ☉

Statura *S. vulgaris*, sed Folia majora, obtusiora, repando-sinuata, integerrima. Caulis minus divisus, valde hirsutus. Pedunculi solitarii, viscid. Flores solitarii, magnitudinis *S. Jacobææ*, radio minus revoluta. Calyx non sphacelatus.

*** Floribus radiatis: radio patente. Foliis pinnatifidis.

20. *SENECIO elegans*. *S.* corollis radiantibus, foliis pinnatifidis æqualibus patentissimis: margine incrassato receptato.

S. corollis radiantibus, foliis pinnatifidis æqualibus patentissimis: rachi inferne angustata. *Hort. ups. Kniph. com.* 7. n. 85.

Alter africanus annuus, senecionis foliis. *Comm. hort.* 2. p. 59. t. 30.

Jacobæa capensis, flore purpureo, senecionis folio dentato. *Seb. mus.* 1. p. 33. t. 22. f. 1.

β. *Jacobæa africana* frutescens, flore amplo purpureo elegantissimo, senecionis folio. *Volk. norib.* 225. t. 225.

Habitat in Æthiopia.

Calycis squamæ subciliatæ. Varietas β. major Caule frutescente ab α. videtur orta.

22. *SENECIO crucifolius*. *S.* corollis radiantibus, foliis pinnatifidis dentatis subhirsutis, caule erecto. *Fl. jacc. pol. delph. lith. burg.*

S. scanorensis. *It. scan.*

S. Jacobæa selandica incana. *Lob. illustr.* 76.

Jacobæa incana altera. *Bauh. pin.* 131.

Jacobæa artemisiæ folio, radice repente. *Vaill. ed.* 566.

Jacobæa altissima, foliis crucæ artemisiæve similibus æmulis. *Rupp. ien.* 164.

Jacobæa incana repens. *Barr. rar.* 1075. t. 153.

Habitat in aggeribus Europæ temperatæ.

Culta villositatem exuit. *Villich.* Calycis squamæ in apicem rubellum, minime autem nigrum terminantur. *Pollich. l. c.*

Cum *S. Jacobæa* jungit. *Haller R.*

23. *SENECIO incanus*. *S.* corollis radiantibus, foliis utrinque tomentosis subpinnatis obtusis, corymbo subrotundo. *monsp. carn. fil. delph.*

S. tomentosus, foliis pinnatis, pinnis subrotundis, umbellæ globosæ. *Hall. helv.* n. 61.

SYNGENESIA. POLYGAMIA SUPERFLUA. 377

Jacobæa pumila, absinthii melliferi foliis incanis floribus velut in umbellam positis. *Pluk.* 194. t. 39. f. 6.

Chrysanthemum alpinum incanum, foliis laciniatis. *Bauh.* pin. 134.

Chrysanthemum alpinum I. *Clus. hist.* I. p. 333.

Jacobæa alpina incana minor. *Barr. ic.* 262. *Bocc. mus.* 2. p. 20. t. 8.

Absinthium alpinum umbelliferum tenuifolium & minus. *Bauh.* pin. 140.

Habitat in Alpibus Helvetiæ, Austriæ, Pyrenæorum, Monspeli.

Caules *digitales*, *foliosi*. Folia *Achilleæ clavennæ* s. *Othonnæ maritimæ* licet parva, *pinnatifida*, *obtusæ*, *alba*, *lineata*.

24. **SENECIO** *abrotanifolius*. S. corollis radiantibus, foliis pinnato multifidis linearibus nudis acutis, pedunculis subbifloris. *lugd. burg.*

S. corollis radiantibus, foliis pinnato multifidis linearibus nudis acutis, floribus corymbosis. *Spec. p. 2. vind. monsp. carn. Jacq. fl. austr.* t. 79.

S. foliis decompositis. *Guett. stamp.*

Solidago foliis vaginosis, pinnatis, pinnis perangustis, acutis, simplicibus & bifidis. *Hall. helu. n.* 71.

Achillea montana Artemisiæ tenuifoliæ facie. *Lob. adv.* 333. *ic.* 746. *Hall. R.*

Ageratum ferulaceum. *Munt. ic.* 87. *Bauh. hist.* 3. P. I. p. 152. *Hall. R.*

Jacobæa alpina, foliis multifidis *Etscheriana*, flore luteo plerumque gemello. *Pluk. alm.* 194.

Chrysanthemum alpinum, foliis abrotani multifidis. *Bauh.* pin. 134.

Chrysanthemum alpinum II. *Clus. hist.* I. p. 333. t. 334.

Habitat in Alpibus Stiriacis, Pyrenaicis, Carinthiæ, Horto Dei. 24

Folia *radicalia bipinnatifida*; *Caulina simpliciter pinnatifida*. Pedunculi *pubescentes*, 3. f. 4. sæpius *biflori*.

26. **SENECIO** *Jacobæa*. S. corollis radiantibus, foliis pinnato-lyratis: laciniis lacinulatis, caule erecto. *Fl. suec. lugdb. paris. carn. gallob. tub. pal. sil. Kniph. cent.* 6. n. 83. *Knorr. del.* I. t. I. *Regn. bot. lugd. lish. burg.*

S. foliis lineatis, pinnis dentatis, extrema maxima. *Hall. helu. n.* 62.

Jacobæa vulgaris laciniata. *Bauh. pin.* 131.

378 SYNGENESIA. POLYGAMIA SUPERFLUA.

Herba s. Jacobi. *Fuchf. hist.* 742.

β. *Jacobaea vulgaris*, flore nudo. *Rai. angl.* 2. p. 82. *Morif. hist.* 3. p. 108. *Pollich. l. c.*

Jacobaea vulgaris laciniata, flore discoide. *Vaill. act.* 1720. p. 383.

Habitat in Europæ pascuis humentibus. 24

Varietas discoidea β. vix differt specie; cum sit simillima; frequentissima in dunt Belgicis.

Variat foliis inferioribus ovatis serratis, reliquis pinnatis; & radii ligulis crassis, brevibus, striatis ac punctatis, profunde trilobatis. *Pollich. l. c. R.*

* * * *Floribus radiatis: radiopatente. Foliis indivisis.*

29. *SENECIO linifolius*. S. corollis radiantibus, foliis linearibus integerrimis, corymbo subsquamato, caule herbaceo.

Solidago foliis linearibus integerrimis, corymbo simplici. *Hort. cliff.*

Linariæ aureæ affinis. *Bauh. pin.* 213. *prodr.* 107.

Jacobaea linifolia hispanica & italica. *Bocc. mss.* 2. p. 604. 49. *Barr. rar.* 97. t. 802.

Habitat in Hispania, Italia, Russia. 24

Pedunculi squamis minutis acutis adspersi. Calyx sectionis, licet non sphacelatus.

30. *SENECIO paludosus*. S. corollis radiantibus, foliis ensiformibus acute serratis subtus subvillosis, caule striato. *Fl. suec. ged. fl. manufr. pal.* *Fl. dan.* t. 385. *delph. lichen. lugd. burg.*

S. foliis linearibus serratis, floribus umbellatis. *Hall. helv.* n. 66. R.

S. foliis subensiformibus serratis subtus incanis; florum radio subrevoluto. *Neck. gallob.*

S. foliis integris serratis. *Guet. stamp.*

Solidago caule simplici, corymbo terminali, foliis subtus tomentosis, flosculis femineis sedecim. *Fl. suec.*

Jacobaea foliis longis integris & mucronatis. *Morif. hist.* 3. p. 110. f. 7. t. 19. f. 22.

Conyza palustris serratifolia. *Bauh. pin.* 266.

Consolida palustris. *Tabern.* 555. *Hall.*

Lingua major. *Dalech. lugdb.* 1037. R.

Conyza aquatica maxima serrato folio. *Thal. hanc.* 31. ic. 3. R.

β. *Solidago palustris altissima incana & tomentosa, foliis longioribus serratis.* *Ann. ruth.* 220. *Gmel. sib.* 2. p. 161.

SYNGENESIA. POLYGAMIA SUPERFLUA. 379

Habitat in Europæ paludibus maritimis. 24

Variat foliis utrinque tomentosis.

31. SENECIO nemorensis. S. corollis radiantibus octonis , foliis lanceolatis biserratis subtus villosis , caule ramoso. *monsp. Jacq. obs. 3. p. 15. t. 65. 66. & austr. t. 184. pal. delph.*

S. foliis ovato lanceolatis , serratis , subtus hirsutis , floribus umbellatis. *Hall. helu. n. 64.*

Virgaurea f. Solidago saracenica latifolia serrata. *Bauh. hist. 2. p. 1063. Plukn. 390. t. 235. f. 1. Hall. R.*

Jacobæa nemorensis , latiore rigidiore & hirsuto folio. *Rupp. ien. 142.*

Solidago foliis lanceolatis serratis , floralibus ad pedunculum minimis. *Gmel. sib. 2 p. 165.*

Habitat in Germaniæ , Sibiricæ nemoribus. 24

32. SENECIO saracenicus. S. corollis radiantibus , floribus corymbosis , foliis lanceolatis serratis glabriusculis. *monsp. ged. Jacq. austr. t. 186. pal. sil. Kniph. cent. 4. n. 78. lith. lugd.*

S. foliis lanceolatis serratis glabris , floribus umbellatis. *Hall. helu. n. 65.*

S. floribus radiatis , corymbosis ; foliolis sessilibus , lanceolatis , serratis. *Scop. carn.*

S. foliis lanceolatis serratis , floralibus pedunculos æquantibus. *Roy. lugdb.*

Solidago caule simplici , corymbo terminali : pedunculis partialibus alternis nudis longitudine folii. *Hort. cliff. 410.*

Solidago saracenica. *Fuchsf. hist. 728. Bauh. hist. 2. p. 1062.*

Virga aurea angustifolia serrata. *Bauh. pin. 268.*

Jacobæa , subrotundo minus laciniato folio. *Bauh. pin. 131.*

Habitat in Helvetiæ , Monspeli , &c. montanis , nemorosis. 24

Dignoscitur facie , latis foliis , magnitudine , radice maxime reptante.

33. SENECIO Doria. S. corollis radiantibus , floribus corymbosis , foliis subdecurrentibus nudis lanceolatis denticulatis , superioribus sensim minoribus. *Jacq. austr. t. 185. Kniph. cent. 10. n. 78. delph.*

Solidago caule simplici , corymbo terminali , pedunculis terminalibus nudis foliolo subjecto brevissimo. *Hort. cliff.*

Virga aurea major f. Doria. *Bauh. pin. 268.*

Alisma monspeliensium. f. Dori. *Bauh. hist. 2. p. 1064.*

Habitat in Oriente , Austriæ , Germaniæ sylvis , Monspeli ad Ladi ripas. 24

380 SYNGENESIA. POLYGAMIA SUPERFLUA.

Conf. Doria orientalis, limonii folio. *Dill. elth.* 125. t. 105. f. 125.

Folia utrinque lævia. Bractææ setaceæ sub calyce. Folia caulis superiora evadunt minuta, ut corymbus quasi longius pedunculatus appareat; in præcedenti folia majuscula usque ad corymbum continuant. Similis Cacaliæ saracenicæ.

34. SENECIO Doronicum. S. corollis radiantibus, caule indiviso subunifloro, foliis indivisis ferratis: radicalibus ovatis subtus villosis, *monsp. prou. carn. Jacq. austr. V. app.* t. 45. *delph.*

S. foliis crassis, ferratis, imis petiolatis subrotundis, caulinis lanceolatis. *Hall. helv. n.* 67.

Solidago Doronicum. Sp. pl.

Doronicum foliis lanceolatis alternis dentatis; caulinis amplexicaulibus, caule ramoso. Roy. lugdb.

Doronicum longifolium hirsutæ asperum. Bauh. pin. 185.

Doronicum II. austriacum I. Clus. hist. 2. p. 17.

β. Doronicum foliis oblongo - lanceolatis ferratis. Monnier. obs. 152.

Doronicum integro & crasso hieracii folio. Magn. monsp. 295. *Seg. ver. suppl.*

γ. Doronicum foliis lanceolatis denticulatis subtus tomentosis, caule unifloro. Roy. lugdb.

Doronicum helveticum humile crassis foliis & incanum. Bauh. pin. 185. *prodr.* 97.

δ. Tussilago alpina, folio oblongo. Bauh. pin. 197. *prodr.* 101.

ε. Virga aurea montana ferrata, magno flore. Bauh. pin. 268.

Habitat in Alpibus Pyrenæis, Monspeliacis, Helveticis, Austriacis, Italicis. ℥

Caulis simplicissimus, villosus. Folia Radicalia ovato-oblonga, petiolata, serrata, obtusiuscula, crassiuscula, villosa subtus, supra glauca. Caulina parva, lanceolata, & fere subulata. Flos terminalis, sæpe unicus, crassiusculus. Calyx calyculatus.

Varietates progressivas α - ε vidit D. Jacquin, at omnes negat convenire. D. Haller.

1034. A S T E R.

Recept. nudum. Pappus simplex. Cor. radii plabra.

10. *Cal. imbricati squamæ inferiores patulæ.* 2,

* * *Herbacei integrifolii ; Pedunculis nudis.*

6. **ASTER alpinus.** A. foliis spatulatis hirtis : radicalibus obtusis , caule simplicissimo unifloro. *delph.*

A. foliis lanceolatis hirtis : radicalibus obtusis , caule simplicissimo unifloro. *Spec. pl. 2. monsp. Jacq. austr. t. 88.*

A. foliis radicalibus petiolatis , ovatis , scabris , caule unifloro. *Hall. helu. n. 82.*

A. montanus cæruleus , magno flore , foliis oblongis. *Bauh. pin. 267. ?*

Aster VII. austriacus V. *Clus. hist. 2. p. 15.*

β. A. atticus alpinus alter. *Bauh. pin. 267.*

A. atticus cæruleus , alpinus. *Bauh. pin. 125. **

γ. A. hirsutus austriacus cæruleus , magno flore , foliis subrotundis. *Bauh. pin. 267. prodr. 124.*

Habitat in Austria , Vallesia , Helvetia , Pyrenæis , Montpellier. 24

Folia caulina 2 , rarius 3 , Gouan. Calyx æqualis.

Variat floribus albis. Hall.

8. **ASTER Tripolium.** A. foliis lanceolatis integerrimis carnosissimis glabris , ramis inæquatis , floribus corymbosis. *Fl. succ. carn. Fl. dan. t. 615.*

A. maritimus Tripolium dictus. *Rai. hist. 270. Hort cliff. 408.*

Tripolium majus cæruleum. *Bauh. pin. 267.*

Tripolium. *Dod. pempt. 379.*

Habitat in Europæ littoribus maritimis & ad Sibiria , Germaniæ lacus salsos. 24

9. **ASTER Amellus.** A. foliis lanceolatis obtusis scabris triner-
viis integris , pedunculis nudiusculis corymbosis , squamis
calycinis obtusis. *monsp. pal. Kniph. cent. 8. n. 12. Regn.
bot. Jacq. austr. t. 435. delph. lugd. lithua.*

A. foliis lanceolatis scabris semiamplexicaulibus subserratis ,
calycibus laxis : squamis lanceolatis. *Roy. lugdb.*

A. caule folioso , foliis scabris , ciliatis , lanceolatis , obtusis ,
calycinis squamis ovatis , ciliatis. *Hall. helu. n. 83.*

A. foliis semiamplexicaulibus , lanceolatis , scabris ; squamis
calycinis obtusis. *Scop. carn.*

A. atticus cæruleus vulgaris. *Bauh. pin. 267.*

380 SYNGENESIA. POLYGAMIA SUPERFLUA.

Conf. Doria orientalis, limonii folio. *Dill. elth.* 125. t. 105. f. 125.

Folia utrinque lævia. Bractææ setaceæ sub calyce. Folia caulis superiora evadunt minuta, ut corymbus quasi longius pedunculatus appareat; in præcedenti folia majuscula usque ad corymbum continuant. Similis Cacaliæ saracenicæ.

34. SENECIO *Doronicum*. S. corollis radiantibus, caule indiviso subunifloro, foliis indivisis ferratis: radicalibus ovatis subtus villosis, *monsp. prou. carn. Jacq. austr. V. app.* t. 45. delph.

S. foliis crassis, ferratis, imis petiolatis subrotundis, caulinis lanceolatis. *Hall. helv. n.* 67.

Solidago Doronicum. Sp. pl.

Doronicum foliis lanceolatis alternis dentatis; caulinis amplexicaulibus, caule ramoso. *Roy. lugdb.*

Doronicum longifolium hirsutie asperum. Bauh. pin. 185.

Doronicum II. austriacum I. *Clus. hist.* 2. p. 17.

β. *Doronicum* foliis oblongo - lanceolatis ferratis. *Monnier. obs.* 152.

Doronicum integro & crasso hieracii folio. *Magn. monsp.* 295. *Seg. ver. suppl.*

γ. *Doronicum* foliis lanceolatis denticulatis subtus tomentosis, caule unifloro. *Roy. lugdb.*

Doronicum helveticum humile crassis foliis & incanum. Bauh. pin. 185. *prodr.* 97.

δ. *Tussilago alpina*, folio oblongo. *Bauh. pin.* 197. *prodr.* 101.

ε. *Virga aurea montana ferrata*, magno flore. *Bauh. pin.* 268.

Habitat in Alpibus Pyrenæis, Monspeliacis, Helveticis, Austriacis, Italicis. 24

Caulis simplicissimus, villosus. Folia Radicalia ovato-oblonga, petiolata, serrata, obtusiuscula, crassiuscula, villosa subtus, supra glauca. Caulina parva, lanceolata, & fere subulata. Flos terminalis, sæpe unicus, crassiusculus. Calyx calyculatus.

Varietates progressivas α - ε vidit. D. Jacquin, at omnes negat convenire. D. Hailer.

Recept. nudum. Pappus simplex. Cor. radii pinnati.
 10. *Cal. imbricati squamæ inferiores patulæ.*

* * *Herbacei integrifolii ; Pedunculis nudis.*

6. **ASTER alpinus.** A. foliis spatulatis hirtis : radicalibus obtusis , caule simplicissimo unifloro. *delph.*

A. foliis lanceolatis hirtis : radicalibus obtusis , caule simplicissimo unifloro. *Spec. pl. 2. monsp. Jacq. austr. t. 88.*

A. foliis radicalibus petiolatis , ovatis , scabris , caule unifloro. *Hall. helu. n. 82.*

A. montanus cæruleus , magno flore , foliis oblongis. *Bauh. pin. 267. ?*

Aster VII. austriacus V. *Clus. hist. 2. p. 15.*

β. A. atticus alpinus alter. *Bauh. pin. 267.*

A. atticus cæruleus , alpinus. *Bauh. pin. 125. **

γ. A. hirsutus austriacus cæruleus , magno flore , foliis subrotundis. *Bauh. pin. 267. prodr. 124.*

Habitat in Austria , Vallesia , Helvetia , Pyrenæis , Montpellier. 24

Folia caulina 2 , rarius 3 , Gouan. Calyx æqualis.

Variat floribus albis. Hall.

8. **ASTER Tripolium.** A. foliis lanceolatis integerrimis carnosissimis glabris , ramis inæquatis , floribus corymbosis. *Fl. succ. carn. Fl. dan. t. 615.*

A. maritimus Tripolium dictus. *Rai. hist. 270. Hort cliff. 408.*

Tripolium majus cæruleum. *Bauh. pin. 267.*

Tripolium. *Dod. pempt. 379.*

Habitat in Europæ littoribus maritimis & ad Sibiria , Germaniæ lacus salsos. 24

9. **ASTER Amellus.** A. foliis lanceolatis obtusis scabris triner-
 viis integris , pedunculis nudiusculis corymbosis , squamis
 calycinis obtusis. *monsp. pal. Kniph. cent. 8. n. 12. Regn.
 bot. Jacq. austr. t. 435. delph. lugd. lithua.*

A. foliis lanceolatis scabris semiamplexicaulibus subserratis ,
 calycibus laxis : squamis lanceolatis. *Roy. lugdb.*

A. caule folioso , foliis scabris , ciliatis , lanceolatis , obtusis ,
 calycinis squamis ovatis , ciliatis. *Hall. helu. n. 83.*

A. foliis semiamplexicaulibus , lanceolatis , scabris ; squamis
 calycinis obtusis. *Scop. carn.*

β. A. atticus cæruleus vulgaris. *Bauh. pin. 267.*

380 SYNGE

Carf. Dora

105. f.

Folia

caule

fructu

fructu

FLUVA. 381

MYGAMIA SUPERFLUA:

i. *Blackw.* t. 109.

ii. *Cluj. hist.* p. 16.

tron. 8.

is. *Asperis collibus.* 24

canonem Anello. *Virg.*

his lanceolatis-linearibus stratis
bus corymbois fastigiatis, pedun-

rib. 2492.

ro. rarius simplicibus, foliis lig-

. *Griff. cir.* 47. t. 11.

ria. 257.

flore. *Barr. ic.* 606. bene.

A. minor narbonentium, tripolii flore, linearis folio. *Lab.*

ic. 347. bene. *Pluck. alm.* 56. t. 271. f. 3.

Habitat in Hungaria interamni, Hispania, Monspeli.

Varietate n. luteo flore habet *Gmel. fib. it.* 1. p. 172. R.

* * * * Herbacei Serratifolii; Pedunculis lœvibus.

25. ASTER annuus. A. foliis ovatis: inferioribus crenatis,
caule corymbofo, pedunculis nudis, calycibus hemispha-

ricis. *Hart. ufs.* * *Fl. dan.* t. 486.

A. annuus ramosus albus latifolius canadensis. *Murif. hist.*

3. p. 122.

Bellis ramosa umbellifera. *Cor. canad.* 193. t. 194.

Habitat in Canada: Dania. ☉

* * * * Herbacei Serratifolii, pedunculis squamosis.

35. ASTER chinensis. A. foliis ovatis angulatis dentatis pe-
tiolaris, calycibus terminalibus patentibus foliosis. *Korr.*

del 1. t. S. 3. 4. 5.

A. chenopodii folio, annuus, flore ingenti specioso. *Dist.*

elth. 38. t. 34. f. 38.

Habitat in China. ☉

Fructificatio nonnihil differt a reliquis: plena est eximis.

1035. SOLIDAGO.

Recept. nudum. Pappus simplex. Cor. radium circiter

5. Cal squam. imbricatz, clausz.

2. SOLIDAGO

ternis recur.

tubiformi

Vit. p. inter

alm. 389. t.

nonculstro-corymbosa, re-

curvatus, foliis trinerviis

canadensis. *Plac.*

125.

SYNGENESIA. POLYGAMIA SUPERFLUA. 283
Virga aurea americana, foliis serratis angustis subtus nervosis.
Morif. blaf. 322.

Habitat in Virginia, Canada. 24

Folia angusto-lanceolata, vix manifeste serrata, fere glabra.

11. SOLIDAGO Virgaurea. S. caule subflexuoso angulato, racemis paniculatis erectis confertis. *ged. gallob. herb. pal. fil. Fl. dan. t. 663. Ludw. ed. t. 150. Blackw. t. 169. Kniph. cens. 7. n. 89. Knorr. del. 1. t. W. 11. Regn. bot. delphi. lugd. burg.*

S. caule erecto, racemis alternis erectis. *Fl. suec. paris.*

S. caule anguloso, hirsuto, foliis ovato-lanceolatis, dentatis, racemis multifloris spicatis. *Hall. helv. 69.*

S. caule paniculato; foliis ovato-lanceolatis, serratis. *Scop. carn.*

S. floribus per caulem simplicem undique sparsis. *Fl. lapp.*

Virga aurea vulgaris latifolia. Bauh. hist. 2. p. 162.

Virga aurea latifolia serrata. Bauh. pin. 268.

Habitat in Europæ pascuis siccis. 24

12. SOLIDAGO minuta. S. caule simplicissimo, foliis caulinis integerrimis, pedunculis exillaribus unifloris. *delph.*

Virga aurea humilis alpina. Morif. blaf. 329.

Virga aurea omnium minima, floribus maximis. Herm. par. 245. t. 245. Bærh. lugdb. 1. p. 97.

Virga aurea montana biuncialis pumila. Pluck. alm. t. 390. 235. f. 7. 8.

Virga aurea montana minor. Barrel. ic. 783. Bocc. mus. 169. t. 118. Scheuchz. it 2. p. 144. it 4. p. 463. Hall. R.

Habitat in Pyrenæis. 24

1036. C I N E R A R I A.

Recept. nudum. Pappus simplex. Cal. simplex, polyphyllus, æqualis.

3. CINERARIA sibirica. C. racemo simplici, foliis cordatis obtusis denticulatis lævibus caule simplicissimo monophyllo.

C. racemo composito, foliis crenatis lævibus hastato-cordatis petiolaris. *Gouan. illustr. 69. R.*

Othoana foliis cordatis, caule subnudo simplicissimo. Hort. up. 173. Hall. plant. 588. t. 14.

Jacobæastrum cacaliæ folio. Aum. ruth. 221. t. 24.

Virga orientalis, cacaliæ folio. Tournef. cor. 37.

SYNGENESIA: POLYGAMIA SUPERFLUA. 385

*Senecio foliis imis petiolatis cordatis, obiter dentatis; caulibus lingulatis sessilibus. Hall. helu. n. 68. **

Jacobaea montana lanuginosa angustifolia non laciniata: Bauh. pin. 131. Morf. hist. 3. p. 112. f. 7. t. 12. f. 28.

Jacobaea montana non laciniata nostras. Rai. hist. 272.

Jacobaea montana polyanthos, flore aureo, foliis longis & integris. Barr. ic. 260.

Solidago foliis lanceolatis sessilibus integris denticulatis, floribus umbellatis, involuero setaceo. Gmel. sib. 2. p. 134. t. 71.

J. Cineraria helenitis.

Othonna foliis lanceolatis integris subdentatis villosis, floribus umbellatis. Sp. pl. Gouan. monsp. Reyg. ged.

Conyza helenitis mellita incana: Lob. ic. 347. ex figura.

Conyza incana. Bauh. pin. 265.

Habitas in Alpibus Pyrenæis, Helveticis, Austriacis, Monspelii, Sibiricis. 2.

Polymorpha, in solo natali extricata a Jacquinò, cui addo & helenitis J. facie nimis affinem, inquirent de hac ejus populares alterius.

Vas. B. in mare alpinis; γ. in montosis herbidis; Foliis latioribus, magis dentatis; Radio luteo distinctas Gerard.

Cinerariæ Jacq. austr. 1. 176. 177. 179. 180. & 181. comia modo mihi videntur in duas species conjungi posse; unam nempe foliis ex cordato-oblongis, petiolatis, petiolis nudis; alteram foliis inferioribus ovato-oblongis, petiolis quidem instructis, sed membranaceis latiusculis, superioribus lanceolato ellipticis vel ligulatis saltem, amplexicaulibus. Prior species contineret Cinerariam t. 176 & 177. Senecionem; Hall. helu. n. 63. Altera vero sub se comprehenderet Cinerarias tab. 179. 180. 181. Jacq. & Seneciones. Hall. helu. n. 67 & 68. Linnaeus omnes sub Ciner. alpina conjungit. De C. crispa: Jacq. t. 178. non visa nil certi pronuntio, videtur tamen ad primam speciem pertinere. Hartenut De la Chenal in Act. Helu. VIII. p. 137. R.

9. CINERARIA maritima. C. floribus paniculatis, foliis pinnatifidis tomentosis; laciniis sinuatis, caule frutescente. Kniph. cent. 6. n. 68. delph.

Othonna 2. Hort. upf. 273.

Solidago foliis pinnatifidis; laciniis sinuatis, corymbis fasciculatis. Hort. cliff. 140. Gron. orient. 277.

Senecio alpina. Bauh. pin. 131.

in plant. Europ. Pars II.

Bb

386 SYNGENESIA. POLYGAMIA SUPERFLUA

Cineraria. *Dod. pempt.* 642.

Habitat ad maris inferi littora. 7

1037. I N U L A.

*Recept. nudum. Pappus simplex. Antherae basi in
fetas duas desinentes.*

1. *INULA Helenium*. I. foliis amplexicaulibus ovatis rugosis : subtus tomentosis , calycum squamis ovatis. *gallob. herb. ged. fil. Ludw. t. 59. Kniph. cent. 7. n. 36. Rega bot. Flor. dan. t. 728. lithu. delph.*

After foliis ovatis rugosis amplexicaulibus : subtus tomentosis , calycum squamis ovatis patulis. *Fl. suec. paris.*

After foliis ovato-lanceolatis , ferratis , subtus tomentosis , calycinis ovato lanceolatis maximis. *Hall. helu. n. 72.*

After *Helenium*. *Scop. carn.*

Helenium vulgare. *Bauh. pin.* 267.

Helenium. *Cam. epit.* 35. *Fuchsf. hist.* 242. *Blackw. t. 473.*

Habitat in Anglia , Belgio , &c. 24

2. *INULA odora*. I. foliis amplexicaulibus dentatis hirsutissimis : radicalibus ovatis , caulinis lanceolatis , caule paucifloro.

After luteus , radice odora. *Bauh. pin.* 266:

Afteris altera species apula. *Col. ecphr.* I. p. 251. t. 253.

Conyza altera apula. *Morif. hist.* 3. p. 113. f. 7. t. 21. f. 6.

Habitat in Italia , Galloprovincia , Narbona , Tubingæ.
Gmel. sub. 257.

Folia mollia : caulina amplexicaulia. Flos 1. f. 2. pendunculatus.

3. *INULA Oculus Christi*. I. foliis amplexicaulibus oblongis integris hirsutis , caule piloso corymboso. *Jacq. austr. t. 223. delph.*

After caule superne ramoso ampliato , foliis lanceolatis amplexicaulibus integris , calycibus laxis terminalibus. *Hort. cliff.*

Conyza pannonica lanuginosa. *Bauh. pin.* 265. *Morif. hist.* 3. p. 113. f. 7. t. 19. f. 1.

Conyza 3 austriaca. *Ciusf. hist.* 2. p. 20.

Habitat in Austria , Tubingæ. *Gmel. sub.* 257. 24

Planta tota hirsuta est.

4. *INULA britannica*. I. foliis amplexicaulibus lanceolatis

SYNGENESIA. POLYGAMIA SUPERFLUA. 387

distinctis serratis subtus villosis, caule ramoso erecto villoso. *Fl. suec. Fl. dan. t. 413. delph. lugd. burg.*

α. Conyzeis affinis. Bauh. pin. 265.

Conyza palustris repens britannica dicta. Meris. hist. 3. p. 113. f. 7. t. 19. f. 8.

β. Conyza aquatica, asteris flore aureo. Bauh. pin. 266. prodr. 124. Burs. XV. 41.

Habitat in Lusatia, Bavaria, Scania, Sibiria.

Habitus I. dysentericæ, sed caulis altior, erectior. Folia angustiora, argutius serrata. Varietas β. differt solum Caule & Foliis subtus magis villosis. Radii petala angustissima.

5. INULA dysenterica. I. foliis amplexicaulibus cordato-oblongis subtomentosis, caule villoso paniculato, squamis calycinis setaceis. monsp. carn. sub Astere. gallob. pal. sil. Fl. dan. t. 410. Kniph. cent. 12. n. 62. delph. lugd. burg.

Aster foliis amplexicaulibus, undulatis, subtus tomentosis. Hall. helv. n. 79.

Aster foliis semiamplexicaulibus oblongis, caule paniculato hirsuto, floribus latis, squamis calycinis setaceis. Fl. suec. stamp. paris.

Aster pratensis autumnalis, conyzæ folio. It. gotl.

Conyza media, asteris flore luteo, s. tertia dioscoridis. Bauh. pin. 265.

Conyza I. Tabern. hist. 1243.

Calaminthæ tertium genus. Fuchsf. hist. 436.

Habitat in Europæ fossis subhumidis. 24

Calycis squamæ subulatae, molles recurvatae, marginē coloratae.

8. INULA Pulicaria. I. foliis amplexicaulibus undulatis, caule prostrato, floribus subglobosis: radio brevissimo. ged. carn. sub Astere. gallob. herb. pal. sil. Flor. dan. t. 613. Blackw. t. 103. lith. delph. lugd. burg.

Aster foliis amplexicaulibus undulatis, hirsutis, radiis brevissimis. Hall. helv. n. 80.

Erigeron foliis undulatis amplexicaulibus. Fl. suec. 693. 758.

Conyza ramis secundum flores gradatim enascentibus, foliis lanceolatis obtusis. Hort. cliff.

Conyza major, flore globoso. Bauh. pin. 266.

Conyza minima. Dod. pempt. 52.

β. Helenium palustre annuum, hyssopi foliis crispis. Vaill. ed. 575.

388 SYNGENESIA. POLYGAMIA SUPERFLUA?

Aster palustris fruticosus minimus. Barr. ic. 369.

Conyza minor exotica. Bauh. pin. 265.

Conyza minor hispanica ex oris gaditanis. Pluk. ment. 56. t. 384. f. 2.

Conyza media crispa lacustris, flore nullatenus radiato. Rai. suppl. 152.

Habitat ad vias & plateas Europæ temperata. ☉

Varietas β. major, rigidior, vix meretur distingui.

10. *INULA spiræifolia. I. foliis subsessilibus ovato-oblongis nudis reticulatis confertis serrulatis, floribus terminalibus subsessilibus.*

Conyza media monspeliensi affinis multiflora. Bauh. hist. 2. p. 1049.

Habitat in Italia.

Flores vix pedunculati sunt.

11. *INULA squarrosa. I. foliis sessilibus ovalibus laevibus reticulato-venosis subcrenatis, calyce squarroso. delph.*

Aster conyzoides odoratus luteus. Tournef. inst. 483.

Aster luteus latifolius glaber, foliis rigidis & minutissime crenatis. Pluk. alm. 37. t. 16. f. 1.

Habitat in Italia, Monspeli. 24

Caulis pubescens, striatus, sæpe uniflorus. Folia sessilis, scabriuscula, nuda, subserrata, margine magis scabra. Flos magnus, terminalis, sessilis. Calyx foliolis lanceolatis, exterioribus majoribus.

12. *INULA salicina. I. foliis lanceolatis recurvis serrato-scabris, ramis angulatis, floribus inferioribus altioribus. delph. lithu. lugd. burg.*

I. foliis sessilibus lanceolatis recurvis serrato-scabris, floribus inferioribus altioribus, ramis subangulatis glabris. Sp. pl. 2. monsp. carn. sub Astere. pat. fil. Kniph. cent. 6. n. 49.

Aster foliis lanceolatis amplexicaulibus serrato-ciliatis glabris acuminatis recurvis, floribus solitariis, caule striato. Fl. suec. parif.

Aster foliis glabris, ciliatis, venosis. Hall. helv. n. 76.

Aster luteus latifolius glaber, foliis rigidis & minutissime crenatis. Pluk. 57. t. 16. f. 1. Rai. syn. 3. p. 163. Hall. R.

Aster montanus luteus, salicis glabro folio. Bauh. pin. 266.

Subgenum luteum I. Tabern. hist. 716.

Habitat in Europæ borealis pratis uliginosis; asperis. 24

Diff. 1. In Italia. Caule glabro, superno sulcato, angulato; foliis, margine tantum scabris,

SYNGENESIA. POLYGAMIA SUPERFLUA. 389

13. INULA hirta. l. foliis sessilibus lanceolatis recurvis subserrato-scabris, caule teretiusculo subpiloso, floribus inferioribus altioribus. *ged pal. Jacq. austr. t. 358. Kniph. cent. 12. n. 63. lithu.*

Aster foliis ovatis, venosis, scabris, hirsutis. *Hall. helu. n. 75.*

Aster foliis lanceolatis sessilibus integerrimis, caulibus simplicibus unifloris, squamis calycinis subulatis. *Dalib. parif.*

Aster luteus, hirsuto salicis folio. *Bauh. pin. 266.*

Aster tertius pannonicus clusii luteus, folio hirsuto salicis. *Bauh. hist. 2. p. 1047.*

Habitat in Gallia, Sibiria, Germania, Genevæ. *24*

Habitus Inulæ salicinæ, sed Folia latiora, obtusiora, vix manifeste serrata, venosa, scabra præsertim margine & carina: Caulis foliis pilisque rigidiusculis adpersus, ubique teres, nec sulcato-angulatus. Calyx quasi foliosus.

Folia utrinque hirta lanugine subrufa. *Syst. Veg. 638.*

Confer. Aster conyzoides odoratus luteus. *Segu. ver. 2. p. 211.*

Aster luteus, cisti hirsutis & crenatis foliis, summo caule ramoso, floribus amplis. *Pluk. alm. 57. t. 78. f. 4.*

Aster luteo flore, aliis Conyzæ species. *Cam. epit. 907.*

Veronæ lecta differt Foliis ovalibus, nec lanceolatis aut serratis; Calyx videtur indicare eandem speciem.

15. INULA germanica. l. foliis sessilibus lanceolatis recurvis scabris floribus subs fasciculatis. *pal. Jacq. austr. t. 134. Kniph. cent. 4. n. 32. delph. lith.*

Aster foliis lanceolatis amplexicaulibus oris reflexis, ramis multifloris, calycibus oblongis laxis. *Gmel. fib. 2. p. 181. t. 78. f. 1.*

Conyzæ affinis germanica. *Bauh. pin. 266. Morif. hist. 3. f. 7. t. 19. f. 26.*

Aster thuringiacus altissimus latifolius montanus, flore luteo parvo. *Hall. ien. 181.*

Habitat in Misnia, Pannonia, Sibiria, &c. *24*

Habitus I. Salicinæ. Folia eadem; at Flores cylindrici conferti ad apicem caulis in corymbum.

16. INULA ensifolia. l. foliis sessilibus linearibus acuminatis nervosis glabris sparsis, caule subunifloro. *Scop. carn. Jacq. austr. t. 162.*

Aster montanus saxatilis luteus, angusto nervoso acuto & molli plantaginis folio, multicaulis. *Bocc. mus. 1. p. 18.*

390 SYNGENESIA. POLYGAMIA SUPERFLUA:

Aster luteus, linearis rigido glabro folio. *Bauh. pin.* 267.

Habitat in Austria inferiore; *Tubingæ Gmel. tub.* 24.

17. *INULA crithmifolia*. l. foliis linearibus carnosis tricusp-
pidatis. *monfr. Scop. carn. sub Senecione.*

Aster flore terminali, foliis linearibus tricusp-
pidatis. *Guett. stamp*

Crithmum maritimum, colore asteris atticis, *Bauh. pin.* 288.

Crithmum crysanthemum. Dod. pempt. 706.

*Habitat in Angliæ, Galliæ, Lusitanæ, Hispaniæ mari-
timis.* 24

18. *INULA provincialis*. l. foliis subserratis subtus tomen-
tosis: radicalibus petiolatis ovatis, caule erecto un-
floro

l. caule folioso tomentoso unifloro, foliis radicalibus omni-
pinnatifidis integrisque, caulinis linearibus integerrimis
Gouan. illustr. R.

Solidago foliis tomentosis, ovatis & semipinnatis. Hall. helv.
n. 70. sec. Gouan. R.

Jacobæa rotundifolia incana. Bauh. pin. 131. *prodr.* 69.
Burf. VI. 127.

Habitat in Collendaneis Galloprovinciæ. 24

*Radix fibris validis. Caulis pedalis, simplicissimus,
erectus, villosus. Folia Radicalia ovata, petiolata, obuse
serrata, subtus villosa, obtusiuscula; Caulina pauca, lan-
ceolata, sessilia, minus serrata, subtus villosa: summa li-
nearia. Flos sat magnus, flavus.*

*Raubini Synonymon ad Senecionem Doronicum refert
Haller.*

19. *INULA montana*. l. foliis lanceolatis hirsutis integerr-
imis, caule unifloro, calyce brevi imbricato. *pat. lugh.
lithu. delph. burg.*

*Aster foliis ellipticis, integerrimis, tomentosis, caule uni-
floro. Hall. helv. n. 81.*

Aster montanus luteus mas & femina. 3. 4. Tabernæm.

monfr.

Aster montanus hirsutus. Lob. ic. 350. *Hall. R.*

Aster atticus luteus montanus villosus, magno flore. Bauh.

monfr.

Aster folius luteus. Bauh. hist. 3. p. 1046.

Monspeliis, in Hispania, &c. 24

*Aster villosa est. Calyx & structura plantæ
Gris.*

l. foliis lanceolatis oblongis den-

SYNGENESIA. POLYGAMIA SUPERFLUA. 391
riculatis, floribus congestis terminalibus subsessilibus.
delph.

Conyza pyrenaica, foliis primulae veris. *Herm. parad.* 127.

Conyza latifolia viscosa suaveolens, flore aureo e galloprovincia. *T. Garid. aix.* 125. t. 23.

Conyza praelata, caule alato, odorata. *Becc. mus.* 1. p. 168. t. 121. *Rai. suppl.* 153.

Habitat in Italia, Galloprovincia, Pyrenæis. ♂

Conf. *Conyza bifrons*, quam eandem absque radio statuunt Botanici: nostra est Caule pedali, rigidulo, corymbosa, fortiter odora ut *Tanacetum*. Flores lutei radio brevi.

25. *INULA foetida*. I. foliis lanceolato-linearibus integerrimis, corymbis ramosis, radiis florum brevissimis.

Solidago foliis lanceolato-linearibus, &c. *Hort. cliff.*

Conyza melitenensis, retusis foliis. *Bocc. succ.* 26. t. 13.

Pseudo-Helichrysum melitense, retusis foliis viridibus flore luteo radiato. *Moris. hist.* 3. p. 90.

Habitat in Melita. ☉

Affinis *Senecioni foetido*. *Habitus Cinerariae.*

1038. A R N I C A.

Recept. nudum. Pappus simplex. Corollulae radiis filamentis 5 absque antheris.

1. *ARNICA montana*. A. foliis ovatis integris: caulinis geminis oppositis. *carn. herb. pal. sil. Fl. dan.* t. 63. *Blackw.* t. 595. *Knorr. del.* 2. t. A. 16. *lugd. lithu. delph.*

A. foliis conjugatis ovatis integerrimis. *Hall. helv.* n. 90.

Doronicum foliis caulinis oppositis. *Fl. suec. Hall. opusc.* 103.

Doronicum foliis oblongis-ovalibus. *Fl. lapp.* 304.

Doronicum, plantaginis folio, alterum. *Bauh. pin.* 185.

Alistna. Maush. diosc. 934.

Diuretica. *Renealm. spec.* 118.

2. *Arnica alpina.*

Doronicum foliis lanceolatis. *Fl. lapp.* 305. *Roy. lugd.* 160.

Caltha alpina. Tabern. ic. 336.

Habitat in Alpibus & prasis Europae frigidioris. 24

392 SYNGENESIA. POLYGAMIA SUPERFLUA.

Alpina foliis angustissimis, & Pratenſi ovatis maxime differt.

Filamenta quinque radii caſtrata non adſunt. Hall. R.

3. ARNICA ſcorpionides. A. foliis alternis ſerratis. auſtr. 2. 349. monſp. Scop. ſub Aſtere, delph.

A. foliis alternis, argute ſerratis, radicalibus petiolatis. Hall. helv. n. 89.

Doronicum auſtriacum, 3. Cluſ. pann. 520. 521. Blackw. 4. 503. Hall. R.

Doronicum radice ſcorpion brachiata. Bauh. pin. 184.

Doronicum brachiata radice. Dalech. hiſt. 1203. Lob. 2. 649.

β. Doronicum radice dulci. Bauh. pin. 184.

Habitat in Helvetia, Auſtriæ ſubalpinis humidis. 26

Semina omnia pappoſa.

1039. DORONICUM.

Recept. nudum. Pappus ſimplex. Cal. ſquamæ duplicis ordinis æquales, diſco longiores. Sem. radii nuda pappoque deſtituta.

1. DORONICUM Pardalianches. D. foliis cordatis obtuſis denticulatis: radicalibus petiolatis; caulinis amplexicaulibus. Mill. diſc. t. 128. monſp. Jacq. auſtr. 2. 350. Lud. eſc. t. 57. Kniph. cent. 2. n. 21. Blackw. 1. 239. Reg. bot. delph. burg.

D. foliis cordatis, obtuſis, imis longe petiolatis, ſuperioribus amplexicaulibus. Hall. helv. n. 88.

D. foliis auriculatis, ſubdentatis; dentibus glanduloſis; calyce ſubito. Scop. carn.

D. foliis cordatis denticulatis, caule ramoſo. Hort. cliſſ.

D. maximum; foliis caulem amplexantibus. Bauh. pin. 185. Cam. epit. 823.

Doronicum VII, auſtriacum 3. Cluſ. hiſt. 2. p. 19.

β. D. radice ſcorpion. Bauh. pin. 184.

D. latifolium. Cluſ. hiſt. 2. p. 16.

Aconitum Pardalianches. Dod. purg. 305.

Habitat in Alpibus Helvetiæ, Pannoniæ, Valleſiæ, Germaniæ. 26

Semina radii nuda. Caulis ramoſus.

2. DORONICUM plantagineum. D. foliis ovatis acutis

SYNGENESIA. POLYGAMIA SUPERFLUA. 393

subdentatis, ramis alternis. *paris. monsp. sub. delph. burg. lugd.*

D. plantaginis folio. *Bauh. pin. 184.*

D. minus officinarum. *Dalech. hist. 1202.*

β. **D.** plantaginis folio, lusitanicum. *Tournef. instr. 488.*

Habitat in Lusitania, Hispania, Gallia, Germania. 24

Caulis ramosus.

3. DORONICUM Bellidiastrum. **D.** scapo nudo simplicissimo unifloro. *Jacq. austr. t. 400. Scop. carn. sub Astere. delph. burg.*

Arnica caule nudo, unifloro, foliis ovato-lanceolatis, serratis. *Hall. helv. n. 92.*

Bellidiastrum alpinum, foliis brevioribus hirsutis, caule palmari, flore albo. *Mich. gen. 32. t. 29.*

Bellis sylvestris media, caule carens. *Bauh. pin. 261.*

Bellis media. *Clus. hist. 2. p. 44. Cam. epit. 654. Hall. opusc. 174. **

β. **Bellis** caule pedali s. bipedali nudo, foliis magnis latis, floribus rubris & albis. *Mentz. pug. t. 8.*

Habitat in Alpibus Helveticis, Italicis, Tyrolensibus, locis umbrosis; etiam Tübingæ. Gmel. tub. p. 259. 24

Semina radii in hac etiam papposa sunt. Ruberrimo flore reperit Haller.

1041. HELENIUM.

Recept. nudum, Radii paleaceum. Pappus 5. aristatus. Cal. simplex, multipartitus, Corollulæ radii semitrifidæ.

1. HELENIUM autumnale. **H.** foliis serratis.

H. foliis decurrentibus. *Hort. ups.*

Chrysanthemum americanum perenne, caule alato, foliis angustato glabro. *Morif. hist. 3. p. 24. f. 6. t. 6. f. 74.*

Aster floridanus aureus, caule alato. *Pluk. amalth. 43. t. 372. f. 4.*

Aster luteus alatus. *Corn. canad. 52. t. 63.*

Habitat in Americæ septentrionalis subhumidis. 24

Varietas hirsuta datur; an satis distincta?

Alteram speciem foliis integerrimis Mill. (dict.) non vidit

394 SYNGENESIA. POLYGAMIA SUPERFLUA

1042. BELLIS.

Recept. nudum, conicum. *Pappus* nullus. *Cal.* hemisphaericus : squamis aequalibus. *Sem.* obovata.

1. *BELLIS perennis* B. scapo nudo. *pal. carn. fil. gallob.*
Flor. dan. t. 503 *Ludw. ed. t. 184.* *Blackw. t. 200.*
Knorr. del t. M. Regn. bot. lugd. burg.

B. scapo nudo unifloro. *Fl. suec. parif.*

B. sylvestris minor. *Bauh. pin. 267.*

B. sylvestris. *Dod. pempt. 265.*

B. *Bellis hortensis.*

B. hortensis, flore pleno. *Bauh. pin. 261.* *Mill. dict. n. 3.*
Kriph. cent. 1. t. 10.

7. B. hortensis, flore albo bullato. *Tournef. inst. 491.*

8. B. hortensis rubra, flore multiplici fistuloso. *Tournef.*
inst. 491. *Blackw. t. 530.*

9. B. hortensis prolifera.

Habitat in Europæ apricis pascuis. 2.

2. *BELLIS annua.* B. caule subfolioso.

B. caule ramoso multifloro ; pedunculis longissimis unifloris.
Gouan. illustr. 1.

B. minima pratensis, caule folioso. *Bac. mus. 2. p. 96. t. 35.*

B. maritima, foliis agerati. *Bauh. pin. 261. prodr. 121.*
Burt. XIV. 70.

B. hortensis pediculo folioso. *Bauh. pin. 262. prodr. 121.*

B. minima, caule folioso pubescens. *Cup. cath.*

B. *Leucanthemum annuum italicum.* *Mich. gen. 34.*

Habitat in Sicilia, Hispania, Monspeli. ☉

1043. BELLUM.

Recept. nudum. *Semina* conica corona paleacea
8. phylla, pappoque aristata. *Cal.* foliolis aequalibus.

1. *BELLIUM bellidivides.* B. scapis nudis filiformibus.

Bellis (droseraefolia) stolonibus reptantibus, scapis unifloris,
foliis spathulatis. *Gouan. illustr. p. 69.*

Bellis annua minima. *Triumf. obs. 82. t. 82.*

Bellis maritima minima, rosis folis folio, pyrenaica. *Boca*
mus. 149. Rai. suppl. 222.

SYNGENESIA. POLYGAMIA SUPERFLUA. 395

Bellis foliis parvis, floribus albis, alpina. *Mentz. pug.* 1. 8.

Habitat in Italiæ, Romæ aridis, olim cultis; primo vere. In herbidis Insulæ baleariæ maritimis copiose. *Gouan.* ☉

Facies *Bellidis* omnino, sed pappo diversissima. Folia radicalia, obovata, integerrima, erectiuscula, supra scabra, petiolo breviora. Propagines filiformes, apice radicales, foliiferi & floriferi. Scapi filiformes, teretes, foliis longiores, uniflori, flore ante anthesin cernuo. Calyx foliolis 10, lanceolatis, scabriusculis, parallelis. Corolla radiis albis, ovalibus, emarginatis disco luteo: corollulis 4. fidis. Semina subpilosa. Pappo paleis 8. subrotundis, & setis 8. longioribus, alternantibus. H. V.

2. *BELLIUM minutum*. B. caule folioso capillari. *Mont.* 286.

Pectis (*minuta*) foliis ovatis petiolatis. *Spec.* 2. p. 1250.

Schreb. Nov. Act. upf. vol. 1. p. 81. t. 5. f. 2.

Bellis cretica fontana omnium minima. *Tournef. cor.* 37.

Vaill. act. 547.

Habitat ad Orientis fontes. ☉

Planta inter omnium minutissimas. Caulis capillaris, pollicaris; tota planta lævis, ascendens. Folia obovata, integerrima, lævia, petiolis breviora. Pedunculus terminalis. 1. f. 2. longitudine totius, capillaris, uniflorus. Calyx magnitudine seminis *Brassicæ*. Planta oculo armato pilis raris armata videtur.

1044. TAGETES.

Recept. nudum. Pappus aristis 5, erectis. Cal. 1. phyllus, 5. dentatus, tubulosus. Flosculi radii 5, persistentes.

1. *TAGETES patula*. T. caule subdiviso patulo. *Ups. Kniph. cent.* 10. n. 90. *Knorr. del.* 1. t. S. 7.

Tanacetum africanum s. *Flos africanus minor*. *Bauh. pin.* 132.

Flos africanus. *Dod. pempt.* 255.

β. *Tagetes indicus minor*, multiplicato flore. *Tournef. inst.* 488. *Knorr. del.* 1. t. S. 8.

T. minor, flore fulvo maculato. *Dill. elth.* 273. t. 279. f. 361.

Habitat in Mexico. ☉

596 SYNGENESIA. POLYGAMIA SUPERFLUA.

2. *TAGETES erecta*. T. caule simplici erecto, pedunculis nudis unifloris. *Knorr. del. 1. t. S. 6. a.*

Tanacetum africanum majus, simplici flore. *Bauh. pin. 133.*

Caryophyllus indicus. *Cam. epit. 406.*

β. *Tagetes maximus rectus*, flore maximo multiplicato. *Bauh. h. fl. 3. p. 100. Knorr. del. 1. t. S. 6.*

γ. *Tagetes mexicanus*, flore fistuloso, simplex. *Col. ecphr. 2. p. 47. t. 46. f. 1.*

δ. *Tagetes mexicanus alter polyanthos*, fistuloso flore. *Col. ecphr. 2. p. 48. t. 46.*

Habitat in Mexico. ☉

1046. Z I N N I A.

Recept. paleaceum. Pappus aristis 2, erectis. Cal. ovato-cylindricus, imbricatus. Flosculi radii 5, persistentes, integri.

1. *ZINNIA pauciflora*. Z. floribus sessilibus. *Kniph. cen. 7. n. 100.*

Z. caule villoso. *Reich. in Nov. Act. A. N. C. vol. 6. p. 173. R.*

Verbesina foliis oppositis ovatis acutis integerrimis scabris. *Ad. petrop. 1763. p. 325. t. 13.*

Chrysogonum foliis sessilibus amplexicaulibus, calycibus sessilibus squamosis. *Sp. pl. 1.*

B. dens calyce oblongo, seminibus radii corolla non decidua coronatis. *Mill. dict. 1. 64.*

Rudbekia foliis oppositis. *Zinn. gest. 409. t. 1.*

Ienica. *Hill. exot. n. 29.*

Crassina. *Sceptn. diff. acad.*

Habitat in Peru. ☉

Caulis teres, erectus, compositus. Folia opposita, sessilia, amplexicaulia, cordato-lanceolata, glabra, subtrinervia, integerrima, margine scabra. Calyx terminalis, sessilis, solitarius, glaber, cylindricus. Corolla lutea, persistens. Rami ex alis foliorum, non flori proximis, cauli similes, altiores.

2. *ZINNIA multiflora*. Z. floribus pedunculatis. *Linn. det. 23. t. 12. Kniph. cent. 12. n. 100.*

Z. ramis villosis, numerosis: lateralibus primarium superantibus. *Jacq. obs. 2. p. 19. t. 40.*

SYNGENESIA. POLYGAMIA SUPERFLUA. 392
Z. caule piloso. Nov. Act. A. N. C. VI. p. 173. R.

Habitat in Louisiana. ☉

Caulis hirtus. Folia petiolata. Radius ruber.

Variat radio luteo. R.

1048. CHRYSANTHEMUM.

Receptaculum nudum. Pappus nullus. Cal. hemisphaericus, imbricatus : squamis marginalibus membranaceis.

*** Leucanthema.**

3. CHRYSANTHEMUM atratum. C. foliis omnibus cuneiformi-oblongis incisicarnosis. vind. delph.

Bellis alpina major, folio rigido. Bauh. pin. 261. prodr. 120.

Bellis alpina major, foliis angustis rigidis. Bauh. hist. 3 p. 115.

β. Bellis alpina minor, folio non rigido. Bauh. hist. 3. p. 115.

Bellis montana major, folio acuto. Bauh. pin. 261. prodr. 121.

Habitat in Alpium Helvetiae, Austriae pratensibus.

Folia Radicalia cuneiformia, apice lobata; Caulina lanceolata, serrata. Caulis uniflorus. Calyx margine atro. Cum Ch. Leucanthemo conjungit. Haller.

4. CHRYSANTHEMUM alpinum. C. foliis cuneiformibus pinnatifidis: laciniis integris, caulibus unifloris. prou. delph.

Pyrethrum caule unifloro, foliis glaucis, petiolatis, palmatis. Hall. helv. n. 96.

Chrysanthemum pallidum minimis imisque foliis incisic. Barrel. ic. 421. Gouan. R.

Chamæmelum montanum incanum absinthioides italicum. Barr. ic. 457. Gouan. R.

Chamæmelum alpinum. Bauh. pin. 136.

Leucanthemum alpinum tenuifolium. Clus. hist. 1. p. 335. Barr. ic. 458. f. 3.

Habitat in Alpibus Helvetiae, ad Thermas Piperinas; etiam circa Tubingam. Gmel. tub. p. 261.

Caules stoloniferi. Foliis petiolatis, pinnatifidis: laciniis saepe 8, integris, acutis, distantibus. Caules terminales simplicissimi, vix spithamei, uniflori. Foliis 3 s. 4. linearibus,

398 SYNGENESIA. POLYGAMIA SUPERFLUA.
plerumque indivisis, imprimis superioribus. Calycis squamæ
nigræ scariosæ.

5. CHRYSANTHEMUM Leucanthemum. C. floribus amplexicaulibus oblongis: superne ferratis, inferne dentatis. *fuec. parif. ged. pal. tub. fil. Ludw. ed. t. 183. Kniph. cent. 6. n. 28. Blackw. 42. lugd. lithu. burg.*

Matricaria foliis radicalibus petiolaris, ovatis, crenatis; caulinis amplexicaulibus dentatis. Hall. helm. n. 98. multas habet varietates.

Matricaria foliis simplicibus, amplexicaulibus; receptaculis convexis. Scop. carn.

Chrysanthemum foliis oblongis ferratis. Fl. lapp.

Bellis sylvestris, caule folioso, major. Bauh. pin. 261.

Bellis major. Fuchf. hist. 148. Cam. epit. 635.

β. Bellis montana, folio obtuso crenato. Bauh. pin. 261. prodr. 121. Barr. ic. 458. f. 2. Ger. prov. 206.

γ. Leucanthemum vulgare, caule villis canescente. Tournef. inst. 492.

Habitat in pratis Europæ. 24

Semina nigra striis albis capitulo luteo cylindrico cavo.

6. CHRYSANTHEMUM montanum. C. foliis imis spatulato-lanceolatis ferratis, summis linearibus. *Jacq. obs. 4. p. 9. t. 91. tub. fil. delph. lith.*

Leucanthemum montanum minus. Tournef. inst. 492.

Bellis montana minor. Bauh. hist. 3. p. 115. Magn. monsp. 36.

Habitat Monspeli, Tubingæ, in Silesia.

Varietas præcedentis secundum Gerardum.

7. CHRYSANTHEMUM graminifolium. C. foliis linearibus subintegerrimis, caule simplicissimo. *Jacq. obs. 4. t. 92.*

C. foliis radicalibus oblongis dentatis, caulinis linearibus integerrimis. Gouan. illustr.

Leucanthemum gramineo folio. Tournef. inst. 493.

*Bellis montana, gramineis foliis. Magn. monsp. 291. * Hort. 31. t. 31.*

Habitat Monspeli. 24

8. CHRYSANTHEMUM monspeliense. C. foliis imis palmatis; foliolis linearibus pinnatifidis. *Jacq. obs. 4. p. 10. t. 93.*

Bellis montana major, foliis chrysanthemi cretici angustioribus. Magn. monsp.

Leucanthemum montanum, foliis chrysanthemi. Tournef. inst. 492.

SYNGENESIA. POLYGAMIA SUPERFLUA. 399

Habitat Monspelii. 24

9. **CHRYSANTHEMUM** *Balsamita*. C. foliis ovatis oblongis serratis auriculatis. *Jacq. obs. 4. p. 8. t. 89.*

Leucanthemum orientale, costi hortensis folio. Tournef. cor. 37.

Bellidoides balsamitæ majoris folio facie & odore. Vaill. ad 550.

Habitat in Oriente. 24

Simillimum Tanaceto Balsamitæ, sed radiatum. Caulis subangulatus, ramosus. Folia lanceolato-ovata, obtusiuscula, serrata, subtomentosa, petiolata, auricula utrinque parva. Flos Asteris annui. Calyx fere Achilleæ. Corolla lutea radio albo. Pappus nullus.

10. **CHRYSANTHEMUM** *inodorum*. C. foliis pinnatis multifidis, caule ramoso diffuso. *ged. herb. pal. fil. Flor. dan. t. 696. lithu.*

Matricaria inodora, receptaculis hemisphæricis, radiis patentibus, seminibus coronato-marginatis, squamis calycinis margine obsoletis. Fl. suec. 2.

Chamæmelum inodorum annuum humilius, foliis obscure virentibus. Moris. hist. 3. p. 36. Rai angl. 3. p. 186.

Chamæmelum inodorum s. Cotula non fœtida. Bauh. bist. 3. p. 122.

β. *Chamæmelum maritimum. It. w. goth. 148.*

Habitat in ruderatis Sueciæ & præstantioris Europæ. ☉ Calyx margine scarioso hanc Chrysanthemis adsociat.

11. **CHRYSANTHEMUM** *Achilleæ*. C. foliis bipinnatis: pinnis imbricatis, caule stricto multifloro.

Parthenium foliis tenuissimis, achilleæ cæsuris. Mich. nen. 34. t. 29.

Habitat in Italia. 24.

Caulis erectus, subangulatus, pedalis, multiflorus. Folia Millefolii, sed octuplo majora: punctis eminentibus vix conspicuis adspersa; subtus pilis albis raris; apicibus mucrone terminali albido.

12. **CHRYSANTHEMUM** *corymbosum*. C. foliis pinnatis inciso-serratis, caule multifloro. *monsp. Jacq. austr. t. 379. herb. pal. cub. Kniph. cent. 6. n. 27. delph. burg.*

Pyrethrum foliis pinnatis, pinnis semipinnatis, acute dentatis, floribus umbellatis. Hall. kelu. n. 95.

Matricaria foliis pinnatis; pinnis semipinnatis: laciniis

400 SYNGENESIA. POLYGAMIA SUPERFLUA:

dentatis; semine coronato, denticulato. *Strep. carni*
sub Pyrethro.

Tanacetum montanum inodorum, minore flore. Bauh.
pin. 132.

Tanacetum leucanthemum. Tabern. hist. 379.

Tanacetum inodorum 2. Clus. pann. 551. Hall. R.

Bellis montana minor Tanaceti folio. Pluk. 66. t. 84.
f. 1. Hall. R.

β. Tanacetum inodorum, flore majore. Bauh. pin. 132.

Tanacetum inodorum 1. Clus. pann. 550. Hall. R.

Tanacetum latifolium inodorum, magno bellidis flore.
Barr. ic. 781. Hall. R.

Habitat in Thuringiæ, Bohemiæ, Helvetiæ, Sibiriciæ sy-
vaticis montosis. 24

* * *Chrysanthema.*

15. *CHRYSANTHEMUM pedinatum. C. foliis pinnatis*
linearibus parallelis acutis integerrimis, pedunculis soli-
tariis unifloris.

Chamæmelum montanum incanum absinthioides italicum.
Barr. ic. 457? repræsentat florem & habitum.

Leucanthemum fruticans pinnatis superioribus quomodocumque
foliis, imis vermiculatis & torosis. Barr. ic. 422. foliis
refert.

Habitat in Hispania, Italia. 24

Caules brevissimi, crassiusculi, prostrati, repentes. Fol-
pinnata, parva, (Tanaceti frutescentis) pubescentia.
Pedunculus altior, basi folio uno alterove subulato. In-
flavus.

Nomen Barrel. ad Ch. alpinum refert Hallerus. R.

16. *CHRYSANTHEMUM segetum. C. foliis amplex-*
caulibus: superne laciniatis; inferne dentato-serratis.
Fl. suec. parif. pal. lugd. burg.

C. segetum. Clus. hist. 1. p. 334.

C. segetum vulgare glaucum. Moris. hist. 3. p. 15. f. 6.
t. 4. f. 1.

Bellis lutea, foliis profunde incis, major. Bauh. pin. 262.

Habitat in Scania, Germaniæ, Belgii, Angliæ, Gallia
agris. ☉

17. *CHRYSANTHEMUM Myconis. C. foliis lingulatis*
obtusis serratis, calycinis squamis æqualibus. Jacq. obs.
4. p. 10. t. 94.

C. myconis. Dalech. hist. 873. t. 873.

C. latifolium. Bauh. hist. 3. p. 105.

Bell.

SYNGENESIA. POLYGAMIA SUPERFLUA. 407

Bellis lutea; foliis subrotundis. *Bauh. pin.* 262.

Habitas in Lusitaniæ, Hispaniæ, Italiæ, agris. ☉

Simillimum antecedenti, sed *Caulis erectus, lavis, teretiusculus. Folia extrorsum latiora, obtusa, obtuse serrata, laevia, semiamplexicaulia. Pedunculus terminalis, striatus, obscurior, apice non incrassatus. Calyx squamis apice scariosis. Corolla saturate lutea. Radius brevis, tridentatus. Seminum corona membranacea.*

18. **CHRYSANTHEMUM italicum.** C. foliis bipinnatis ferratis, florum radiis longitudine disci, caule procumbente.

Parthenium foliis tenuissimis, achilleæ cæsuris. Mich. gen. 34. s. 29.

Habitas in Italia. Arduini.

Simillimum *millefoliato*, sed *Caulis magis ramosus, multiflorus, erectior. Florum radius albidus, longitudine disci.*

Obs. Synonymon Michellii ad. Ch. achilleæ jam adducitur; forsan Ch. achilleæ & italicum una eademque planta bis posita. R.

19. **CHRYSANTHEMUM millefoliatum.** C. foliis bipinnatis dentatis, caule decumbente, corollæ radiis disco brevioribus. *Syst. Veg.* 643.

Anthemis (millefolia) foliis bipinnatis: lacinii linearibus integris, pedunculis nudis longissimis. Spec. ed. 2. p. 1263. Roy. lugdb. 271.

Pyrethrum foliis incanis, triplicato-pinnatis, tomentosis. Hall. gatt. 393. R.

A. foliis pinnatis; foliolis lineari-lanceolatis basi sursum auctis. Mill. ic. 6. s. 9.

Leucanthemum, folio absinthii, alpinum class. Bærh. lugdb. 1. p. 107.

Pyrethrum foliis duplicato-pinnatis: pinnulis incisis, pedunculis unifloris, caule procumbente. Gm. sib. 2. p. 207. s. 86. f. 1. 2.

Habitas in Sibiria.

Discus omnino absque paleis. Differt a Ch. bipinnato disci corollarum diametro non superante longitudinem calycis, a Chr. bipinnatum disco duplo latiore gaudet. Affinis C. italico.

21. **CHRYSANTHEMUM coronarium.** C. foliis pinnatis incisis extrorsum latioribus. *Kniph. cent.* 1. s. 14.

Matricaria foliis amplexicaulibus, pinnatis, pinnis semipin-
Linnaei spec. plant. Europ. Pars II. C c

102 SYNGENESIA. POLYGAMIA SUPERFLUA
natis, acute dentatis, semiflosculis latissimis. *Hall. helv.*
n. 99.

Chrysanthemum foliis matricariae. Bauh. pin. 134.

C. majus, folio profundius laciniato, magno flore. Bauh.
pin. 134.

C. creticum. Clus. hist. 1. p. 334. Moris. hist. 3. p. 16.
f. 6. t. 4. f. 2. 3.

Habitas in Creta, Sicilia, Helvetia. ☉

22. CHRYSANTHEMUM *flosculosum*. *C. flosculis omni-*
bus uniformibus hermaphroditis. monsp. Kniph. cent. 4. n. 18.

Tanacetum foliis integris rigidis dentatis, scapo uniflora.
Hall. gart. 370.

Balsamita foliis agerati. Vaill. ac. 336.

Bellis major spinosa, petalis carens s. nuda. Moris. hist. 3.
p. 29. f. 6. t. 9. f. 16.

Bellis spinosa, foliis agerati. Bauh. pin. 262.

Bellis spinosa. Alp. exot. 327. t. 326.

B. Cotula (grandis) caule simplici erecto, foliis lanceolatis
dentatis. Spec. ed. 2. p. 1257. Jacq. obs. 4. p. 4. t. 81.

Habitas in Africa, Creta. ☿

Flosculi 5. fidi. Caulis simplex, erectus, sesquipedalis,
angulatus, purpurascens. Rami superiores elongati, uni-
flori; inferiores steriles, foliosi. Folia amplexicaulia, sin-
gulata, argute dentato-ferrata. Mant. 2. p. 473.

Sempervirens. Folia obovata, sinuato-dentata, rigida.
Flos terminalis, solitarius. G. Syst. Veg. 643.

1049. MATRICARIA.

Recept. nudum. Pappus nullus. Cal. hemispha-
ricus, imbricatus: marginalibus solidis, ac-
tiusculis.

1. MATRICARIA *Parthenium*. *M. foliis compositis pl-*
nis: foliolis ovatis incisis, pedunculis ramosis. Fl. Linn.
ed. t. 192. Fl. dan. t. 674. Kniph. cent. 5. n. 57. Rep-
bot. delph. lugd. burg.

M. foliis pinnatis, pinnis semipinnatis, laciniis obtusis, flo-
ribus umbellatis. Hall. helv. n. 100.

M. vulgaris s. sativa. Bauh. pin. 133.

Matricaria. Dod. pempt. 35. Blackw. t. 192.

Habitas in Europæ cultis, ruderalis. 24 ♂

SYNGENESIA. POLYGAMIA SUPERFLUA. 103

Calyx scariosus, ut Chrysanthemi. Variat foliis cristatis; floribus plenis; disco nudo absque semiflosculis. Hall. R.

2. **MATRICARIA** *maritima*. M. receptaculis hemisphaericis, foliis bipinnatis subcarnosis: supra convexis, subius carinatis.

Chamaemelum maritimum perenne humilius, foliis brevioribus crassius obscure virentibus. Rai. angl. 3. p. 186. t. 7. f. 1. It. w. goth. 148.

Habitat in Europæ septentrionalis litoribus maris.

3. **MATRICARIA** *suaveolens*. M. receptaculis conicis, radiis deflexis, seminibus nudis, squamis calycinis marginæ æqualibus. Fl. suec. sub. herb. delph.

M. leucanthemos annua suaveolens, chamaemeli folio, ovariiis albicantibus. Vailt. parisi. 1720.

Chamaemelum tenue incisum, foliis floris deorsum protensis; umbone extuberante, suavius oleas. Cup. cathol. suppl. 17.

Habitat in Europa. ☉

4. **MATRICARIA** *Chamomilla*. M. receptaculis conicis, radiis patentibus, seminibus nudis, squamis calycinis marginæ æqualibus. Fl. suec. parisi. ged. pal. sil. Ludw. ed. t. 131. Kniph. cont. 9. n. 64. Blackw. t. 298. lugh. lithu. burg.

M. foliis capillaribus duplicato-pinnatis, pinnis lanceolatis, bifidis trifidisque. Hall. helv. n. 101.

M. foliis supradecompositis, setaceis, receptaculis oblongis. Scop. carn.

M. foliis supradecompositis setaceis, pedunculis solitariis. Hort. cliff.

Chamaemelum vulgare s. Leucanthemum, dioscoridis. Bauh. pin. 135.

Anthemis vulgaris. Lob. ic. 770. Fl. tapp. 309.

Habitat in Europæ agris, cultis. ☉

5. **MATRICARIA** *argentea*. M. foliis bipinnatis, pedunculis solitariis. Hort. cliff. 419.

M. monoleucanthemos, foliis argenteis plerumque conjugatis. Vailt. ad. 1720. p. 369.

Chamaemelum orientale incanum millefolia folio. Tournef. cor. 37.

Habitat in Oriente.

1050. COTULA.

Recept. subnudum. *Pappus* marginatus. *Corollula* disci 4. fide; radii fere nullæ.

1. *COTULA anthemoides*. C. foliis pinnato-multifidis dilatatis, floribus flosculosis.

C. foliis pinnato-multifidis dilatatis, corollis radio destitutis *Spec. 2.*

Ananthocyclus chamæmeli folio V. *Dill. elth.* 26. t. 23. f. 23.

Chrysanthemum exoticum perpusillum nudum, foliis coronopi. *Pluk. alm.* 101. t. 274. f. ult.

Habitat in Insula Hælenæ, Hispania.

Caulis pilosus. Folia pinnatifida, inciso. Pedunculi longi, solitarii, uniflori. Flos globosus, cernuus, luteus. Calyx non imbricatus. Receptaculum nudum.

2. *COTULA aurea*. C. foliis pinnato-setaceis multifidis, floribus flosculosis cernuis.

C. foliis pinnatis setaceis, caulibus procumbentibus *Læst. it. **

Chamæmelum aureum peregrinum, capite sine foliis. *Bank. hist.* 3. p. 119.

Habitat in Hispania, Europa australi. ☉

Flores absque paleis, flosculis quadrifidis.

3. *COTULA coronopifolia*. C. foliis lanceolato-linearibus amplexicaulibus dentatis, floribus flosculosis. *Oed. in.* t. 341.

C. foliis lanceolato-linearibus amplexicaulibus pinnatifida *Spec. pl. Kniph. cens.* 10. n. 27.

Ananthocyclus coronopi folio V. *Dill. elth.* 27. t. 1. f. 26.

Chrysanthemum exoticum minus, capite aphyllis, chamæmeli nudæ facie. *Breyn. lent.* 156. t. 76.

Bellis annua, capite aphyllis luteo coronopi folio, caulibus procumbentibus. *Herm. lugdb.* 86. *Morif. hist.* 3. p. 30. f. 6. t. 6. f. ult.

Habitat in Æthiopia, nunc in Prissæ inundatis prope Emden, Mæhringio teste. ☉

Recept. paleaceum. Pappus emarginatus. Sem. lateribus membranaceis.

3. *ANACYCLUS aureus*. A. foliis bipinnatis teretiusculis. incanis excavato-punctatis.

Chamæmelum luteum, capite aphyllō. *Bauh. pin.* 135.

Chamæmelum aureum peregrinum, capite sine foliis. *Bauh. hist.* 3. p. 119.

Anthemis Chrysanthemum herbariorum. *Lob. ic.* 771.

Habitat in Europa australi & Oriente. ☉ 24. H. V.

Caules plures, spithamæi, erectiusculi, substriati, obsolete canescentes villis. Folia bipinnata, teretiuscula, linearia, subcanescentia villis decumbentibus: pinnulae patentes, mucronatae, excavato-punctatae: infimæ caulem amplexantur. Supremum folium simpliciter pinnatum. Pedunculi terminales, solitarii, longitudine caulis, uniflori, aphylli, similiter subpubescentes. Calyx hemisphaericus, pubescens imbricatus squamis apice scariosis obscurioribusque. Corollæ Discus semiovatus. Flosculi disci 5. fidi. Radii feminei plurimi, parvi absque limbo. (In *Santolina anthemioide* simili sunt nulli flosculi feminei) Receptaculum concavum, paleis lanceolatis, concavis, luteis.

Synonyma. *Colutea aurea* forte ad hanc plantam potius spectant.

4. *ANACYCLUS valentinus*. A. foliis decompositis linearibus: laciniis divisis teretiusculis acutis, floribus flosculosis.

Chrysanthemum valentinum. *Clus. hist.* 1. p. 332.

Buphthalmum lanuginosum, foliis millefolii. *Bauh. pin.* 135.

Buphthalmo tenuifolio simile. *Bauh. hist.* 3. p. 125.

Chamæmelum tenuifolium, flore bullato aureo. *Barr. rar.* 1095. t. 450.

Habitat ad Reg. Valentini agros & vias. ☉

Conf. *Anthemis valentina*, cui corollæ similiter aristatae; sed Calyx minus scariosus; forte sola varietas est.

Recept. paleaceum. *Pappus* nullus. *Calyx* hemisphaericus, subaequalis. *Flosculi* radii plures quam. 5.

* *Radius* discolore seu alba.

1. **ANTHEMIS** *Cato*. A. florum paleis rigidis pungentibus.

A. italica *orvenlis* annua major vulgarissima, flore maximo disco pungente. *Mich. gen.* 32.

Chamæmelum annuum ramosum, corulae foetidae floribus amplioribus, capitulis spinosis. *Morif. hist.* 3. p. 36. f. 6. t. 8. f. 11. *prou.* 12. *Till. pis.* 38. t. 19.

Bellis montana tanacetii foliis, caule singulari, annua. *Pluk. alm.* 65. t. 17. f. 5.

Habuat in Italiae arvis.

Pedunculi vix *superne* incrassati. *Calyx* arctus squamis vix manifeste pice nigricantibus. Floris fructiferi discus hemisphaericus calyce plano, magnitudine *Pruni*, adeoque vix penes maximus.

2. **ANTHEMIS** *altissima*. A. erecta, foliis pinnatis: pinnarum basibus denticulo reflexo asperis. *delph.*

A. erecta, foliorum apicibus subspinosis. *Spec. pl.* 2. *Keph. cent.* 11. n. 4.

A. foliorum serraturis setaceis, radiis florum albis. *Sam. monsp.*

Chamæmelo affine. *buphrthalmum* italicum segetum altissimum. *Bauh. hist.* 3. p. 120.

Chamæmelum *Leucanthemum* hispanicum, magno flore. *Bauh. pin.* 135. *prodr.* 70.

Habuat in Italiae, Hispaniae, G. Narbonensis, agris, A. Sauvages. ☉

Caulis multiflorus, segete fere altior, striatus, erectus. *Pedunculi* *superne* parum incrassati. *Calyces* globri. *Pinnatae*, *subtruncatae*, *terminatae* seta rigidiuscula longioribus calycis.

Folia pinnata: pinnis dentato-pinnatifidis; ad basin pinnarum subius denticulus reflexus, unde subius foliorum asperitas manifestissima. *Mant.*

3. **ANTHEMIS** *maritima*. A. foliis pinnatis dentatis or

SYNGENESIA. POLYGAMIA SUPERFLUA. 407

foliis nudis punctatis, caule prostrato, calycibus subtomentosis. *Kniph. cens. 10. n. 8.*

A. maritima annua odorata præcox, flore albo, caule purpurescente. *Mich. gen. 33. Till. pis. t. 19. f. 3.*

Matricaria maritima. *Bauh. pin. 134.*

Chamæmelum maritimum. *Bauh. hist. 3. p. 122.*

Habitat Monspeli, inque Italia. 24

Caules late prostrati, læves, purpurescentes. Folia pinnata, incisa, nuda, punctis excavatis adspersa, versus basin confertius dentata, sub basi linea transversa elevata purpurescente. Pedunculi terminales, solitarii, substriati, pubescentes, superne crassiores. Flores odore *Matricariae Parthenii*.

4. *ANTHEMIS tomentosa.* *A.* foliis pinnatifidis obtusis planis, pedunculis hirsutis foliosis, calycibus tomentosis.

A. foliis bipinnatis, foliolis trifidis, corollulis biaristatis. *Gouan. illustr. p. 70. R.*

Chamæmelum majus, tenuifolium, caule rubente. *Magnol. monsp. sec. Gouan. R.*

Chamæmelum, coronopi folio, tomentosum. *Vaill. ad. 1720. p. 412.*

Chamæmelum maritimum inaequum, folio absinthii crasso. *Boerh. lugdb. 1. p. 110.*

Habitat in Græciæ maritimis, circa Monspelium. 24

Corollæ laciniis 2 exterioribus majoribus. Caules pedales uniflori. Folia - tomentoso - sericea.

5. *ANTHEMIS mixta.* *A.* foliis simplicibus dentato-lacinatis.

A. foliis singularibus pinnato-dentatis. *Guett. stamp.*

Anthemis maritima lanuginosa annua, foliis crassis pyrethri sapore, semiflosculis albis juxta discum luteis. *Mich. gen. 32. t. 32. f. 1.*

Chamæmelum annuum ramosum, coronopi folio, flore mixto. *Morif. hist. 3. p. 36. f. 6. t. 18. f. 15.*

Chamæmelum lusitanicum latifolium, f. coronopi folio. *Breyn. cent. 149. t. 74.*

Bellis pumila crenata agerati æmula crenis bicornibus asperiusculis. *Pluk. alm. 65. t. 17. f. 4.*

Bellis maritima foliis Agerati. *Bauh. prodr. pin. Gouan.*

Habitat in Italia, Gallia. ☉

Folia non composita, sed laciniata; Radius florum albus, basi luteus.

408 SYNGENESIA. POLYGAMIA SUPERFLUA

6. *ANTHEMIS alpina*. A. foliis dentato-pinnatis linearibus integerrimis, caule villosulo-unifloro, petalis ovatis, paleis sphacelatis. *Jacq. austr. VI. app. 1. 30.*

Chamæmelum alpinum saxatile perenne, flore albo singulari, calyce nigricante. *Till. pis. 39. 1. 19. f. 1. Segu. ver. 3. p. 282.*

Habitat in Alpibus Tyrolis, Baldi, Tübingæ. *Gmel. sub p. 262. 2.*

Confer. *Anthemis fruticosa*, foliis linearibus carnosiss. *Ger. prout. 209. 1. 8.*

8. *ANTHEMIS nobilis*. A. foliis pinnato-compositis linearibus acutis subvillosis. *Lud w. ed. 1. 135. Blackw. 1. 526. Regn. bot. burg.*

A. foliis pinnato-decompositis: laciniiis setaceis. *Hort. cliff*
Chamæmelum foliis subhirsutis, nervo duro, pinnis pinnatis pinnulis lanceolatis incis. *Hall. helv. n. 102.*

Matricaria receptaculis conicis, radiis deflexis, seminibus nudis, squamis calycinis margine æqualibus. *Fl. succ. parif.*

Chamæmelum nobile f. *Leucanthemum odoratum*. *Bauh. pin. 135.*

Chamæmelum odoratum. *Dod. pempt. 270.*

β. Chamæmelum nobile, flore, multiplici. *Bauh. pin. 135.*

Chamæmelum romanum, flore multiplici. *Cam. epit. 64. Kniph. cent. 5. n. 13.*

Habitat in Europæ pascuis apricis. 2.

Variat floribus semiplenis: floribus plenis; & fructu nudo. R.

9. *ANTHEMIS arvensis*. A. receptaculis conicis, paleis setaceis, seminibus coronato-marginatis. *Fl. succ. parif. ged. carn. herb. pal. fil. Kniph. cent. 11. n. 5. lugd. lish. burg.*

Chamæmelum foliis glabris, duplicato-pinnatis, nervo foliaceo, pinnulis lanceolatis, semine levi. *Hall. helv. n. 103.*

Corula non foetida. *Bauh. hist. 3. p. 120.*

Chamæmelum inodorum. *Bauh. pin. 135. Rai. angl. 1. 185. Russ gen. 173.*

Habitat in Europæ, præsertim Sueciæ, agris. 3.

Planta canescens. Caulis diffusus Paleæ lanceolatae.

10. *ANTHEMIS Corula*. A. receptaculis conicis: paleis setaceis, seminibus nudis. *Fl. succ. parif. carn. herb. pal. fil. Blackw. 1. 67. Regn. bot. delph. lish. lugd. burg.*

SYNGENESIA. POLYGAMIA SUPERFLUA. 409

A. foliis pinnato-decompositis : laciniis linearibus caulibus diffusis , radice annua. *Roy. lugdb.*

Chamæmelum foliis glabris , duplicato-pinnatis , nervo foliaceo , pinnulis lanceolatis , semine exasperato. *Hall. helu. n. 104.*

Chamæmelum foetidum. *Bauh. pin. 135.*

Chamæmelum foetidum f. *Cotula foetida. Bauh. hist. 3. p. 36.*

Habitat in Europæ ruderatis , præcipue in Ucraina. ☉

Paleæ in hac minutissimæ sunt.

Variat 1) floribus plenis 2) floribus absque radio 3) floribus semiflosculis fistulosis. R.

11. ANTHEMIS montana. **A.** foliis pinnato-multifidis planis : laciniis linearibus acutis trifidis , pedunculo longissimo.

A. foliis pinnatis , pinnis integerrimis pinnatifidisque , pedunculis longis erectis unifloris. *Gouan. illustr. p. 71. R.*

A. fruticosa foliolis linearibus carnosis , pedunculis longis erectis nudis unifloris. *Ger. gallopr. 209. t. 8. bene. Gou. R.*

Abinthium montanum , chamæmeli flore magno. *Bauh. pin. 140. Bauh. hist. 1. p. 184.*

Abinthium montanum , abrotani feminae flore. *Col. phyt. 1. p. 217. t. 33.*

Chamæmelum alpinum , abrotani folio. *Vaill. aët. 591.*

Bellis incana , chrysanthemi cretici folio. *Bocc. mus. 2. p. 136. t. 98.*

Bellis pumila monanthos , foliis nasturtii petræi. *Rai. suppl. 222.*

Habitat in Italia , Helvetia , in Pyrenæis. Gouan.

In Syst. Veg. omissa est hæc species R.

12. ANTHEMIS Pyrethrum. **A.** caulibus simplicibus unifloris decumbentibus , foliis pinnato-multifidis. *monsp. Blackw. t. 390. Kniph. cent. 11. n. 6.*

Chamæmelum specioso flore , radice longa fervida. *Shaw. afr. 138.*

Pyrethrum flore bellidis. *Bauh. pin. 148.*

Habitat in Arabia , Syria , Creta , Apulia , Germania , Bohemia , Monspelii , Apennino. 24

Caules plures , prostrati , sæpius uniflori , rarius ramosi. Radius floris albus , subtus purpureus. Radix crassa , sapore Polygalæ Senegæ.

*** Radio concolore f. luteo.*

13. ANTHEMIS valentina. **A.** caule ramoso , foliis pubes-

§10 SYNGENESIA. POLYGAMIA SUPERFLUA.

centibus tripinnatis setaceis, calycibus villosis pedunculatis
monsp. Mill. dict. n. 10. Kniph. cent. 9. n. 8.

Buphthalmum cotulae folio. Bank. pin. 134.

Buphthalmum tenuifolium, folio fere millefolio. Bank. hist. 3. p. 124.

Buphthalmum creticum, cotulae facie. Breyn. cent. 150. & 55.

Buphthalmum alterum penz. Dalech. hist. 853.

β. Buphthalmum flore luteo subtus purpurascens Bank. pin. 134.

Buphthalmum flore purpurascens. Cam. epis. 652.

Buphthalmum narbonense. Clas. hist. 2. p. 332.

γ. Chamæmelum foetidum maritimum. Vaill. ad. 589. Bank. hist. 3. p. 121.

Habitas in G. Narbonensi. ☉

Affinis admodum Anacyclo valentino.

*Caulis diffusus, purpurascens. Pedunculi superius in-
 crassati. Calycis squama villosa, sed intima scarina,
 crenata, glabra.*

*Varietas β ex Aquis mortuis Monspelti, radice sub
 sanguinea.*

*Corollulas aristatas observavit in hac specie Fåhrus; &
 similes in aliis?*

14. ANTHEMIS repanda. A. foliis simplicibus ovato-
 lanceolatis repando-crenatis.

*Chrysanthemum lusitanicum, agerati folio. Tournef. in
 492.*

*Cry anthemum parvum f. Bellis lutea parva latifolia. Bank.
 hist. 3. p. 105.*

Habitas in Hispania. Lusitania.

*Caules simpliciusculi, erecti. Folia ovato-oblonga, re-
 pando-crenata, petiolata, alterna, obtusa, vix villosa.
 Flores terminales, subsolitarii, globosi. Calyx minus im-
 bricatus. Radius angustus; Discus convexus flavus. Se-
 mina tetragona, coronata squamis quatuor, quarum duae
 in setas desinunt.*

16. ANTHEMIS cindoria. A foliis bipinnatis serratis
 subtus tomentosis, caule corymboso. *ged. herb. pol. &
 Blackw. t. 439. Kniph. cent. 12. n. 10. Regem. in
 Flor. dan. t. 741. lithua.*

*Chamæmelum foliis hirsutis, pinnatis, pinnulis simpli-
 bus lanceolatis. Hall. helv. n. 105.*

*Buphthalmum caule ramoso, foliis pinnatis, linearibus
 dentatis serratis, floribus*
succ.

SYNGENESIA. POLYGAMIA SUPERFLUA. 408
Buphtbalmum tanacetii minoris foliis. Bauh. pin. 134.
Chrysanthemum foliis tanacetii. Læc. pous. 47. t. 9. flore
aureo italicum. Barr. ic. 465.

β. Anthēmis Triumfetti. Leers. L. c. var. β.

Buphtbalmum alpinum, flore candido. Triumf. obs. 79.
t. 80.

Bellis alpina, parthenii foliis, elatior. Pluk. alm. 66.
t. 17. f. 6.

Chamæmelum tanacetii minoris folio, flore albo majore.
Vaill. ad. 591.

Habitas in Sueciæ, Germaniæ apricis pratis siccis. 2f

Planta β. differt radio perfecte explicato albo, basi
samen luteo; reliqua conveniunt ut vix species diversa
eruat. Calycis squamæ interiores utrinque apice ciliatæ.

1053. ACHILLEA.

Recept. paleaceum. Pappus nullus. Cal. ovatus,
imbricatus. Flosculi radii circiter 4.

** Corollis flavis.*

2. ACHILLEA Ageratum. A foliis lanceolatis obtusis
acute serratis. dict. n. 8. ic. t. 10. monsp. Ludw. ed.
t. 27. Blackw. t. 300. Kniph. cens. 6. n. 1. Regn. bot.

Ageratum foliis serratis. Bauh. pin. 221.

Balsamita minor. Dod. pempt. 295. Moris. hist. 3. p.
38. f. 6. t. 5. f. 2.

Habitas in G. Narbonensi, Hetruria. 2f.

4. ACHILLEA tomentosa. A. foliis pinnatis hirsutis :
pinnis linearibus dentatis. Gmel. it. 1. p. 137. t. 25.
f. 2. Kniph. cens. 5. n. 3. delph.

A. foliis linearibus pinnatifidis pubescentibus, foliolis tri-
partitis transversalibus : media productiore. Hort. ups.

A. foliis hirsutis, pinnatis, pinnis æqualibus, pinnatis,
pinnulis integris & trifidis, floribus luteis umbellatis.
Hall, helu. n. 106.

Millefolium tomentosum luteum. Bauh. pin. 140.

Stratiotes millefolia, flavo flore. Clus. hist. 1. p. 330.

Habitas in G. Narbonensi, Vallesia, Tataria. 2f.

5. ACHILLEA pubescens. A. foliis pinnatis : foliolis
lanceolatis incisis serratis subtus lanigeris.

Millefolium orientale, foliis tanacetii incanis, radiis pal-

412 SYNGENESIA. POLYGAMIA SUPERFLUA

lute luteis. *Barrh. lugdb. 1. p. 112.*

Parmica orientalis, foliis tanacetii incanis, semisfoculis
florum pallide luteis. *Tournef. cor. 37.*

Matricaria tomentosa & incana, achilleæ folio, flore aureo.
Vaill. ad. 1722. p. 286.

Habuat in Oriente. 24.

Receptaculum paleaceum adesse negat Vaillant: nec ob-
servavi ipse.

* * *Corollis radio albis.*

9. *ACHILLEA macrophylla*. A. foliis pinnatis planis in-
ciso-ferratis: extremis majoribus coadunatis. *delph.*

A. foliis pinnatis planis inciso-ferratis, extremis confluen-
tibus. *Hall. helv. n. 115.*

A. foliis bipinnatis: pinnulis linearibus indivisis. *San-*
meth.

Dracunculus alpinus, foliis scabiosæ. *Bauh. pin. 9.*
prodr. 39.

Parmica alpina, matricariæ foliis. *Triumph. obs. 83. t.*
23. Boer. mus. 2. p. 150. t. 110. Barr. rar. 119.

t. 991.

Corymbifera millefolii umbella, folio alato & laciniato.
Rai. hist. 345.

Habuat in Alpibus Helvetiæ Italiæ.

Folia magnitudine Tanacetii. Caulis subpubescens, 1
pedalis. Flores Parmicæ, calycis squamæ nigro margi-

11. *ACHILLEA Clovenna*. A. foliis laciniatis planis ob-
tusis tomentosis. *car. Jacq. austr. 1. 76.*

A. foliis sinuato-laciniatis planis villosis nitidis obtus.
Hort. cliff.

A. foliis tomentosis, longe petiolatis, palmatis. *Hel-*
helv. n. 114.

Abinthium alpinum umbelliferum latifolium. *Bank. pin.*
139.

Abinthium alpinum umbelliferum. *Chus. hist. 1. p. 34.*

Dracunculus argenteus. *Moris. hist. 3. p. 40. f. 6. a. 10.*
f. 5.

Habitat in Alpibus Austriæ, Pannoniæ, Carinthiæ
ris. 24.

12. *ACHILLEA Parmica*. A. foliis lanceolatis acumi-
natis argute ferratis. *gallob. pal. sit. Flor dan. a. 64.*

Ludw. ed. 1. 118. lugd. litua. burg.

A. foliis integris minutissime ferratis. *V. FL. succ. pot.*

A. foliis linearibus, lanceolatis, acutissime ferratis. *Hel-*
helv. n. 117.

SYNGENESIA. POLYGANIA SUPERFLUA. 413

Dracunculus pratensis, serrato folio. *Bauh. pin.* 98.

Parmica vulgaris. *Clus. hist.* 2. p. 12. *Blackw.* 1. 276.

Parmica vulgaris, flore pleno. *Clus. hist.* 2. p. 12.
Kniph. cens. 5. n. 2.

Habitat in Europa temperata. 24.

13. *ACHILLEA alpina*. A. foliis lanceolatis dentato-serratis : denticulis tenuissime serratis.

A. foliis semipinnatis, pinnullis minutissime serratis. *Gmel. sib.* 2. p. 196. *Hall. helu.* n. 116.

A. foliis pinnatifidis, laciniis dentatis, umbella conferta. *Gouan. illustr.* p. 71. R.

Parmica alpina, foliis profunde incis. *Tournef. inst.* 497.

Parmica alpina, incanis serratis foliis. *Herm. lugdb.* 694.

Habitat in Alpibus Sibiriae, Helvetiae. 24.

An locus potuerat ex praecedenti formasse hanc?

14. *ACHILLEA atrata*. A. foliis pinnullis pectinatis integriusculis, pedunculis villosis. *Jacq. aust.* 1. 77. *Scop. carn. delph.*

A. foliis glabris, pinnatis, pinnis simplicibus & multifidis. *Hall. helu.* n. 111.

Chamaemelum alpinum saxatile perenne, flore albo singulari, calyce nigricante. *Till. pis.* 1. 19. *Hall. R.*

Matricaria alpina, chamaemeli foliis. *Bauh. pin.* 134.

Parthenium alpinum. *Clus. hist.* 1. p. 336.

Habitat in Alpibus Helvetiae, Valesiae, Austriae : locis asperis humidis. 24

Margines calycis atri & sphacelati. Folia interdum bipinnata.

15. *ACHILLEA nana*. A. foliis pinnatis dentatis hirsutissimis, floribus glomerato-umbellatis. *Delph.*

A. foliis tomentosis, pinnatis, pinnis simplicibus & incis. *Hall. helu.* 2. n. 113.

A. foliis pinnatis lanugine-obductis, floribus albis umbellatis. *Hall. helu.* 713. *Allion. pedem.* 3. 1. 2. f. 3.

Millefolium alpinum, incanum, flore specioso. *Bauh. hist.* 3. p. 138. *Moris. hist.* 3. p. 39. f. 6. 1. 11. f. 11. *Rai. hist.* 347.

Millefolium alpinum tomentosum odoratum nanum. *Bocc. mus.* 2. p. 166. 1. 120.

Habitat in Alpibus Helvetiae, Valesiae.

Planta parva. Paleae inter flosculos luteos disci sunt apice atrae.

16. *ACHILLEA magna*. A. foliis bipinnatis subpilosis :

414 SYNGENESIA. POLYGAMIA SUPERFLUA

lacinis linearibus dentatis : auriculis decussatis. *Delph.*
Millefolium maximum, umbella alba, *Bauh. pin. 140.*
prodr. 72.

Habitat in Europa australi. 24.

Simillima A. *millefolio*, sed duplo major. Caulis ad-
 spersus pilis raris. Folia similia *Millefolio*, sed majora,
 basi quasi auriculata auriculis duobus : altera inferiore sub
 folio ; altera superiore supra costam prominente, methodo
 plane eadem, quæ in *Athamanta libanotide*.

17. *ACHILLEA Millefolium*. A foliis bipinnatis nudis :
 lacinis linearibus dentatis, caulibus superne sulcatis. *Fl.*
suec. parif. ged. carn. gallob. pal. fil. Ludw. ed. 1.
67. Kniph. cent. 7. n. 1. Regn. bot. Flor. dan. 1.
737. Lugd. litua. burg.

A. foliis pinnatis, pinnis semipinnatis, incisiss & integris.
Hall. helu. n. 108.

A. foliis pinnato pinnatis. *Fl. lapp. 311.*

Millefolium vulgare album. Bauh. pin. 140. Blackw.
1. 18.

β. *Achillea tanacetifolia*, flore purpureo. *Wail. ad*
593. Kniph. cent. 5. n. 1.

Millefolium purpureum majus. Bauh. pin. 140.

Millefolium purpureum. Tabern. hist. 130.

γ. *Millefolium alpinum. Clus. pann. 562. Hall. R.*

Habitat in Europæ pascuis pratibusque. 24

58. *ACHILLEA nobilis*. A foliis bipinnatis : inferioribus
 nudis planis, superioribus obtusis tomentosis, corym-
 bis convexis conterritissimis. *Scholl. barb. pal. lina-*
delph.

Millefolium nobile. Trag. hist. 476.

Tanacetum minus album, odore champhoræ. Bauh. pin.
132.

Habitat in Helvetia? Germania, Bohemia, G. Narbo-
nenfi, Tataria. 44

Folia bipinnata, obtusa, tomentosa, at vero Folia in
 ramis summis floralibus simplicia, semipinnata, inae-
 rima, tomentosa, obtusa. Corymbi dense conferti. Caues
 teretes, non sulcati. Florum radius demum reflectitur,
 ut non appareat.

19. *ACHILLEA odorata*. A. foliis bipinnatis ovalibus ne
 diuiculis, corymbis fastigiatis confertis:

A. pinnis æqualibus confertis tomentosis : pinnaulis cili-
 lariter divisis. *Hall. helu. ed. pr. p. 716. Savv. mon.*

SYNGENESIA. POLYGAMIA SUPERFLUA. 418

Millefolium minimum crispum, flore albo, hispanicum.

Barr. rar. 1116. t. 991.

Millefolium odoratum minus monspeliense, Moris. blas., Magn. monsp. 177.

Habitat in Helvetia, Narbona. Hispania. 22.

Folia aequaliter bipinnata, pinnulis distinctis, integerrimis, acutis. Caulis vix palmo altior.

Varietatem A. Nobilis credit Gerard.

Cum A. Millefolio nunc conjungit Hallerus. R.

20. *ACHILLEA cretica. A. foliis linearibus: pinnis subrotundis retrorsum imbricatis, caule tomentoso.*

Millefolium incanum creticum. Bauh. pin. 130. pr. 72.

Millefolium creticum. Bauh. hist. 3. p. 139.

Habitat in Creta. 24.

Habitus A. Millefolii. Caulis tomentosus. Folia linearia, subtomentosa, retrorsum imbricata: pinnis reniformibus, supra quadrifariam imbricatis. Flores albi.

1057. SIEGESBECKIA.

Recept. paleaceum. Pappus nullus. Cal. exterior, 5. phyllus, proprius, patens. Radius dimidiatus.

1. *SIEGESBECKIA orientalis. S. petiolis sessilibus, calycibus exterioribus linearibus majoribus patentibus. Kniph. cent. 2. n. 87.*

*Siegesbeckia. Hort. cliff. 412. t. 23. * Hort. ups. 267. **

Bidenti similis foliis latissimis serratis Buxb. cent. 3. p. 29. t. 52.

Cichorio affinis Lapsana sinica, menthastri foliis, calyce fimbriato hispido. Pluk. amalth. 58. t. 380. f. 2.

Habitat in China, Media ad pagus. 24.

1058. VERBESINA.

Recept. paleaceum. Pappus aristatus. Cal. 2. plicul ordine. Flosculi radii circiter 5.

1. *VERBESINA alata. V. foliis alternis decurrentibus undulatis obtusis. Kniph. cent. 4. n. 93.*

Cannabia indica, foliis integris, alato caule. Magn. hort. 40. t. 40.

116 SYNGENESIA. POLYGAMIA ÆQUALIS.

Chrysanthemum curassavicum, alato caule, floribus aurantiis. *Herm. par.* 125. t. 225.

Chrysanthemum americanum, caule alato, flore sphylo globofo aurantio, foliis baccharidis. *Comm. hort.* 1. p. 5. t. 3.

Chrysanthemum americanum bidens alatum, flore parvo aurantiaco. *Pluk. alm.* 100. t. 84. f. 3.

Chrysanthemum conycoides curassavicum, abrotani feminæ flore aurantio. *Volk. norib.* 106. t. 106.

Habitat in Curassao, Surinamo. 24

A reliquis maxime differt habitu & structura, ut fuit proprii generis.

1059. BUPHTHALMUM

Receptaculum paleaceum. *Pappus* margo obdolan-

Sem. latera, præsertim radii marginata. *Sig.*

floscul. hermaphroditorum indivisum.

* *Asterisci.* T.

3. *BUPHTHALMUM spinosum*. B. calycibus acutifoliis, foliis alternis lanceolatis amplexicaulibus integerrimis, caule herbaceo. *Monsp. Blackw.* t. 272.

Aster luteus, foliis ad florem rigidis. *Bauh. pin.* 26.

Aster legitimus clusii alter s. spinosus luteus. *Barr. d. W.*

Habitat in G. Narbonensi, Hispania, Italia, Creta ad margines agrorum. ☉

Radius multiplex, tenuissimus. Stigma hermaphroditum bifidum.

4. *BUPHTHALMUM aquaticum*. B. calycibus ellipticifoliis sessilibus axillaribus, foliis alternis oblongis serratis, caule herbaceo. *Monsp.*

Asteriscus annuus lusitanicus odoratus. B. *Seb. def.* p. 47. t. 29. f. 7.

Chrysanthemum conyzoides lusitanicum. *Breyn. cent.* 1. t. 77.

Habitat in Creta, Lusitania, Massiliæ. ☉

Caulis dichotomus : *Flores* sessiles in dichotomis.

5. *BUPHTHALMUM maritimum*. B. calycibus oblongifoliis pedunculatis, foliis alternis spatulatis, caule herbaceo. *Kniph. cent.* 3 n. 23.

SYNGENESIA. POLYGAMIA SUPERFLUA. 417

Aster supinus lignosus ficulus, conyzæ odore. *Bocc. mus.*

2. p. 161. t. 129.

Aster supinus luteus massilioticus. *Barr. ic.* 1151.

Aster 2 supinus. *Clus. hist.* 2. p. 13.

Aster luteus supinus. *Bauh. pin.* 267.

Asteriscus maritimus perennis (& annuus) patulus. *Tournef. inst.* 498.

Habitat ad maris Mediterranei littora. 2

Stigma hermaphroditicum bifidum. Willich. Fors-kæhl.

* * *Asteroida. T.*

7. **BUPHTHALMUM** *salicifolium*. B. foliis alternis lanceolatis subserratis villosis, calycibus nudis, caule herbaceo. *Monsp. carn. Jacq. austr.* t. 370. *deph.*

B. foliis hirsutis, imis periolatis, ovatis, superioribus amplexicaulibus, lanceolatis; rariter dentatis. *Hall. helv.* n. 118.

Aster luteus major, foliis succisæ. *Bauh. pin.* 266.

Aster 3. austriacus I. *Clus. hist.* 2. p. 13.

Asteroides hirsuta. *Mich. flor.* 12. t. 3. 4.

Conyza major altera. *Thal. herc.* 21. t. 2.

Habitat ad radices Alpium Austriæ, Stiriae, Helvetiae, Lusatiæ, Monspelii. 2

Affine nimium sequenti.

8. **BUPHTHALMUM** *grandiflorum*. B. foliis alternis lanceolatis subdenticulatis glabris, calycibus nudis, caule herbaceo. *Delph.*

Asteroides alpina, salicisfolio. *Tournef. cor.* 30. *Mich. flor.* 12. t. 5.

Aster luteus angustifolius. *Bauh. pin.* 266.

Chrysanthemum perenne minus, salicis glabro folio, ramosum. *Morif. hist.* 3. p. 21. f. 6. t. 7. f. 52.

Habitat in Alpibus Austriæ, Italiæ, Monspelii. 2

Simillimum Inulæ salicinæ, sed Calyx tantum duplici foliorum ordine, ut in B. salicifolio.

Pappus est margo ciliaris & Calyx æqualis; an itaque potius Inulæ seminibus subnudis?

Cum præcedenti jungit Gouan illustr. p. 71.

9. **BUPHTHALMUM** *speciosissimum*. B. foliis alternis amplexicaulibus ovatis nudis serratis subciliatis, caule unifloro.

B. caule erecto unifloro, foliis amplexicaulibus dentatis: inferioribus ovatis obtusis; superioribus lanceolatis acutis.

Ard. spec. 1. p. 26. t. 12.

Linnaei spec. plant. Europ. Pars II.

D d

118 SYNGENESIA. POLYGAMIA SUPERFLUA:

Asteroides orientale, petalis folio, flore maximo. Tournef. cor. 51.

Habitas in montibus Brixienfibus. 24

Planta in Schreb. cent. 1. 6. videtur maxime affinis; sed folia huic magis profunde serrata.

Radix fusiformis, tuberosæ similis. Caulis bipedalis, leviter lineis albis tenuissimis; inferne petiolis vaginatus, superne parum ramosus in pedunculis. Folia alterna, subcordata, acuta, serrato-dentata, compacta, venis reticulata, subciliata: Petiolis basi dilatatis, submembranaceis, amplexicaulibus; superiora obsoletiora. Pedunculi subtriquetri. Calyx communis imbricatus foliolis ovatis, obtusis, magnis. Corolla radiata: Radii femineis, æque hermaphroditis masculis. Radii ligulata, biplicata, tridentata, lutea. Disci tubulosa, 4. fida, erecta, petiolata. Stamina disci 4, fusca antheris connatis. Filicolum Radii Germen ovatum. Styli 2 filiformes. Disci Germen nullum. Stylus cylindricus, obtusus, luteus, pubescens. Sermen Radii ovatum, membranaceum, subcavum. Semen disci nullum. Receptaculum paleaceum, hemispharicum, Paleis fuscis. D. Zœga. An Euph. species?

POLYGAMIA FRUSTRANEA.

1060. HELIANTHUS.

Recept. paleaceum, planum. Pappus diphyllus. Calyx imbricatus, subsquarrosus.

1. *HELIANTHUS annuus*. H. foliis omnibus trinerviatis, pedunculis incrassatis, floribus compositis. Kniph. cent. 12. n. 54. Knort. del. 1. t. S. 1. Mill. 24.

H. radice annua. Hort. ups.

Helenium indicum maximum. Bauh. pin. 276.

Herba maxima. Dod. pempt. 264.

Chrysis. Ronsal. spec. 84. t. 83.

Habitas in Peru, Mexico. ☉

3. *HELIANTHUS cultiflorus*. H. foliis inferioribus ovatis, superioribus ovatis. Kniph. cent. 12. n. 55.

SYNGENESIA. POLYGANIA FICIFOLIA. 409

H. radice tereti inflexa perenne. *Herb. Am. 288.*
Helenium indicum ramatum. *Bark. pin. 277.*
Corona Solis minor femina. *Tibet. 2. 74.*
Chrysanthemum americanum majus perenne, hirsutum
 foliis & floribus. *Moriz. hist. 3. p. 22. f. 6. 2.*
 159. f. 2.

Habitas in Virginia. 24

Calyx confert squamis 40 ad 50.

Caulis & Pedunculi scabri. Folia ~~ovata-ovatis~~ *Calyces*
 late imbricati; nec squarrosi aut cerni.

4. *HELIANTHUS tuberosus*. *H.* foliis ovatis-cuneatis
 triplinerviis. *Jacq. hort. 2. 261.*

H. radice tuberosa. *Hort. nyp. 263.*

Helenium indicum tuberosum. *Bark. pin. 277.*

Chrysanthemum latifolium brasilianum. *Bark. pin. 277.*

Flos Solis farnesiacus. *Col. cap. 2. p. 11. 2. 13.*

Habitas in Brasilia. 24

1061. R U D B E C K I A

Recept. paleaceum, conicum. *Pappas* margine 4-
 dentato. *Cal. 2. plici ordine squamarum.*

1. *RÜDBECKIA laciniata*. *R.* foliis compositis laciniatis. *Kniph. cens. 4. n. 69.*

Chrysanthemum americanum perenne, foliis divisis distinctius virentibus, majus. *Moriz. hist. 3. p. 22. f. 6. 2. 6. f. 53 & 54.*

Doronicum americanum, laciniato folio. *Bark. pin. 316.*

Aconitum helianthemum canadense. *Corn. canad. 178. 1. 179.*

Habitas in Virginia, Canada. 24

3. *RÜDBECKIA hirta*. *R.* foliis indivisis spatulato-ovatis triplinerviis, radii petalis emarginatis. *Kniph. cens. 2. n. 77. Knorr. del. 1. t. F. 1.*

R. ramis indivisis unifloris, foliis ovato-lanceolatis. *hirta*
*Buttn. canon. 227. **

R. foliis lanceolato ovatis alternis indivisis, pedalis radii integris. *Gron. virg. 1. p. 181.*

Obeliscotheca integrifolia, radio aureo, umbone atro-rubente. *Dill. elth. 295. t. 218. f. 285.*

Chrysanthemum helenii folio, umbone floris grandiusculo
 D d a

720 SYNGENESIA. POLYGAMIA FRUSTRANEA.
prominente. *Pluk. alm.* 99. t. 242. f. 2. *Morif. hist.* 3. p.
23. *Rai. suppl.* 210.

Habitat in Virginia, Canada. ☉. 24.

Radius luteus. Caulis simplex alterove florè.

1062. COREOPSIS.

Receptaculum paleaceum. *Pappus* bicornis. *Cal*
erectus, polyphyllus, basi radiis patentibus
cinctus.

10. COREOPSIS *Bidenx*. C. foliis lanceolatis ferratis op-
positis amplexicaulibus. *Fl. succ. ged. herb. fil. Kniph*
cent. 12. n. 34. *delph. lithua.*

Hepatorium aquatile, folio non diviso, flore aureis petalis
undique radiato. *Triumf. obs.* 65.

Chrysanthemum aquaticum *bidenx*, folio non diviso, flore
aureis petalis undique radiato. *Morif. hist.* 3. p. 17. f. 6.
t. 5. f. 25.

Eupatorium cannabinum *chrysanthemum*. *Barr. ic.* 1209.

Habitat ad fossas Europæ, *Pensylvaniæ.* ☉

Si observationibus evincatur hanc *BIDENTIS* cernuæ
solam esse varietatem, ad *Bidentes* reducatur; sin confus-
sit, permaneat.

Est varietas Bidentis cernuæ secundum Hallerum Pelt-
chium aliosque. R.

1066. CENTAUREA.

Receptaculum setosum. *Pappus* simplex. Cor.
radii infundibuliformes, longiores, irreg-
lares.

* *Jaceæ: calycibus lævibus inermibus.*

1. CENTAUREA *Crupina*. C. calycibus inermibus: se-
mis lanceolatis, foliis pinnatis ferratis. *Kniph. cent.* 7. n.
9. *burg.*

C. calycibus inermibus: squamis lineari-subulatis, foliis pin-
natis ferratis. *Hort. upf. monsp. delph.*

Serratula foliis radicalibus petiolatis ovatis, caulinis pin-
natis, linearibus, denticulatis, ciliatis. *Hall. bot.* 1
197.*

SYNGENESIA. POLYGAMIA FRUSTRANEA. 421

Chondrilla foliis laciniatis ferratis, purpurascente flore. *Bauh. pin.* 130.

Senecio-Carduus apulus. *Col. ecphr.* 1. p. 32. t. 34.

Habitat in *Hetruriæ*, *G. Narbonensis*, *Orientis collibus*; in *Helvetia*. ☉

Foliola scabra: infima recurvata. Corollæ disci 3; radii 5, quadrifidæ. Semen sæpius 1. f. 2. in flore.

2. *CENTAUREA moschata*. *C. calycibus inermibus subrotundis glabris: squamis ovatis, foliis lyrato-dentatis. Kniph. cent. n. 24. 12. Knorr. del. 2. t. C. 4. f. 2.*

Cyanus orientalis major moschatus, flore purpureo & albo. *Moris. hist. 3. p. 135. f. 7. t. 25. f. 5.*

β. Centaurea Amberboi.

C. calycibus inermibus subrotundis glabris, squamis ovatis obtusis, foliis laciniatis ferratis. Mill. dict. n. 9. Kniph. cent. 6. n. 25. Knorr. del. 2. t. C. 4. f. 1.

Cyanus orientalis major, foliis magis dissectis, flore luteo. *Moris. hist. 3. p. 135. f. 7. t. 25. f. 9.*

Cyanus orientalis, flore luteo fistuloso. *Dodart. mem. 267.*

Habitat in *Persia*, *Russia*. ☉

Varietas β Corollulis radii magnis differt, qui autem velit hanc specie distinguere, forte poterit, quamvis habitus plantæ unam ex altera prognatam satis declares.

3. *CENTAUREA erucifolia*. *C. calycibus inermibus: squamis lanceolatis, foliis lanceolatis subdentatis lanuginosis.*

Jacea foliis erucæ lanuginosis. Tournef. inst. 444.

Amberboi, erucæ folio, majus. Vail. act. 1728. p. 181.

Stœbe major, foliis erucæ mollibus lanuginosis. Bauh. pin. 273.

Jacea major exotica. Pluk. phyt. 39. f. 3. mala.

Habitat 24

5. *CENTAUREA alpina*. *C. calycibus inermibus: squamis ovatis obtusis, foliis pinnatis glabris integerrimis: impari ferrato.*

Centaureum alpinum luteum. Bauh. pin. 117. prodr. 56. Moris. hist. 3. p. 132. f. 7. t. 25. f. 5.

Centaureum majus luteum. Corn. canad. 69. t. 70.

Habitat in *Baldo*.

6. *CENTAUREA Centaurium*. *C. calycibus inermibus: squamis ovatis, foliis pinnatis: foliolis decurrentibus, ferratis.*

Centaureum majus, foliolis lacinias plures diviso. Bauh. pin. 117.

412 SYNGENESIA. POLYGAMIA FRUSTRANEA.
Centaureum majus L. vulgare. *Clas. hist. 2. p. 10. Mart.*
1. 43.

Habitat in Alpibus Gargano, Baldo, Tarunia. 24

* *Cyanus* : calycinis squamis serrato-ciliatis.

7. *CENTAUREA phrygia* C. calycibus recurvato-plumosis, foliis indivisis oblongis scabris. *Fl. succ. mansp. carth. gallob. fl. Fl. dan. 1. 520. Kniph. cent. 11. n. xy. delph. burg. lugd.*

C. calycibus ciliatis : ciliis setacea recurvatis. *Hort. ups. parif.*

Cyanus foliis subhirsutis, ovaro-lanceolatis, glandulosis, dentatis, plumulis calycinis reflexis. *Hall. helv. n. 188.*

Jacea latifolia & angustifolia, capite hirsuto. *Bauh. pin. 271.*

Jacea & austriaca, capite villoso. *Clas. hist. 2. p. 7.*

β. *Jacea alba*, hirsuto capite. *Bauh. pin. 271.*

Habitat in Helvetia, Austria, Finlandia, Germania. 24
Calycis plura a pluvia eriguntur.

8. *CENTAUREA capillata* C. calycibus recurvato-plumosis, foliis imis pinnatis dentatis : summis lanceolatis.

Cyanus foliis calycinis plumosis reflexis, foliis pinnatis. *Act. gen. 1. p. 202. t. 6.*

Habitat in Sibiria Gmelinus. Hispania.

9. *CENTAUREA uniflora* C. calyce recurvato - plumosis, foliis lanceolatis subdentatis tomentosis. *Ger. prae. ch. delph.*

C. calycibus recurvato-plumosis, foliis lanceolatis mucronatis, radicibus dentatis, summis integerrimis. *Gouan. illustr. 172. R.*

Cyanus foliis lanceolatis tomentosis, calycinis plumosis flexis. *Hall. helv. n. 189. R.*

Cyanus alpinus, capitulo reticulato. *Bocc. mus. 2. p. 11. t. 2.*

Habitat in Alpibus Europae australis. 24

Affinis C. phrygiae, ejusque progenies aliena. Folia lanceolata, subdentata. Flos unicus, sessilis. Mart. 17.

10. *CENTAUREA linifolia* C. calycibus recurvato-plumosis, foliis linearibus sparsis.

C. calycibus plumosis, foliis linearibus integerrimis scabro-ciliatis mucronato-spinosis. *Gouan. illustr. bot. 73. R.*

SYNGENESIA. POLYGAMIA FRUSTRANEA. 423

Jacea hispanica pumila, *linariae* folio. *Tournef. inst.* 445.
Barr. ic. 162.

Jacea minor purpurea parvis & integris foliis, *Barr. ic.* 139.
Gouan.

Habitat in Hispania, Italia. 24

Caulis brevis, saepe decumbens. Folia linearia, integra, scabra, in ramis valde conferta. Calyx omnino C. phrygiae.

11. CENTAUREA pectinata. *C. calycibus recurvato-plumosis: foliis lyratis denticulatis; rameis lanceolatis integerrimis.*

C. calycibus erecto-plumosis, foliis lyratis denticulatis: rameis lanceolatis integerrimis. Spec. ed. Gouan. illustr. 72.

Jacea montana incana aspera, capitulis hispidis. Bauh. pin. 272. *prodr.* 128.

Jacea montana incana laciniata, capitulo hispido. Bauh. prodr. 128.

Jacea cum squamis pinnatis s. capite villoso, elatior & humilior. Bauh. hist. 3. p. 29.

Habitat in Gallia Narbonensi, Galloprovincia.

Calyces C. phrygiae, terminales, sessiles. Rami magis divaricati. Folia oblonga, amplexicaulia, lyrata, dentibus versus basin majoribus. Folia floralia lanceolata, integerrima, pubescentia. Spec. plant. 1287.

12. CENTAUREA nigra. *C. calycibus ciliatis: squamula ovata: ciliis capillaribus erectis, foliis lyrato-angulatis, floribus flosculosis. ged. pal. Reich. fl. manufr. cum icone. delph. burg.*

C. squamis calycinis ciliatis inermibus, foliis pinnatis: foliolis lineari lanceolatis scabris. Sauv. monsp.

Jacea foliis radicalibus semipinnatis, caulinis ovato-lanceolatis. Hall. helv. n. 184. R.

Jacea nigra laciniata. Bauh. pin. 271.

Jacea austriaca. VI. Clus. hist. 2. p. 7.

Jacea nigra. Rai. hist. 325. *angl.* 3. p. 199.

β. Centaurea calycibus ferratis ciliatisque, foliis lanceolatis dentatis, floribus sessilibus. Hort. cliff.

Habitat in Anglia, Helvetia, Austria, Germania, Montpelii. ♂

Differt a C. phrygia, quod squamula ciliaris sit ovata, erecto patens, nec recurva & subulata.

Similis C. laceae, sed absque radio neutro.

424 SYNGENESIA. POLYGAMIA FRUSTRANEA:

13. CENTAUREA *pullata*. C. calycibus ciliatis verticillato-foliosis, foliis lyratis dentatis obtusis. *Mill. ic. 152. f. 2. delph.*

C. *inermis*, calycum ciliis pinnatis, foliis oblongis villosis humifusis, floralibus verticillatis ligulatis. *Sauv. monsp. 288.*

Jacea *humilis alba*, hieracii folio. *Bauh. pin. 271. Morif. hist. 3. p. 140. f. 7. t. 28. f. 18.*

Jacea *monspeffulana*, cui in squamis fibræ nigræ, interdum acaulis. *Bauh. hist. 3. p. 29.*

Jacea *pumila serpens* acaulis ferme, flore albo. *Lob. ic. 54. Habitat in G. Narbonensi, Hispania, Oriente. ☉*

Calyx foliis cingitur sessilibus, intra hæc folia Squæ calycinae linearilanceolatae margine atro, apice setis filiciliatae. Folia in caule sæpe lyrata sunt. Planta humilis & colla purpurascens radio magno.

14. CENTAUREA *montana*. C. calycibus ferratis, foliis lanceolatis decurrentibus caule simplicissimo. *monsp. carn. Jacq. austr. t. 371. pal. Kniph. cent. 3. n. 24. delph. lith. burg.*

Cyanus caule unifloro, alato, foliis ellipticis lanceolatis. *Hall. helv. n. 170.*

Cyanus *montanus latifolius* f. *Verbasculum cyanoides*. *Bauh. pin. 273.*

Cyanus *montanus*, caule folioso, capitulo oblongo. *Burm. mus. 2. p. 20. t. 2.*

Cyanus *major*. *Lob. ic. 548. Blackw. t. 66.*

β. Jacea *integrifolia humilis*. *Bauh. pin. 271. prodr. Burf. XV. 29.*

Habitat in Alpibus Helveticis, Austriacis, Germanicis.

Planta β. foliis minoribus, parum basi decurrentibus, ad proximum folium, a vulgari differt.

Variat floribus albis; foliis incis. *Hall. R.*

15. CENTAUREA *Cyanus*. C. calycibus ferratis: foliis linearibus integerrimis; infimis dentatis. *succ. parif. gallob. fil. Ludw. ed. t. 55. Kniph. cent. 6. n. 24. del. 1. t. K. 4. 5. Regn. bot. lugd. lith. burg.*

Cyanus foliis imis ellipticis, dentatis, superioribus linearibus semiflosculis latissimis. *Hall. helv. n. 191.*

Centaurea foliis imis simplicibus, dentatis; superioribus bipinnatis; supremis linearibus. *Scop. carn.*

Cyanus *segerum*. *Bauh. pin. 273.*

Cyanus *vulgaris*. *Lob. ic. 546. Blackw. t. 270.*

SYNGENESIA. POLYGAMIA FRUSTRANEA. 425

β. *Cyanus hortensis*, flore simplici. *Bauh. pin.* 273.

γ. *Cyanus hortensis*, flore pleno. *Bauh. pin.* 273.

Habitat inter Europæ segetes biennes. ☉

Variat floribus albis; roseis, purpureis.

16. CENTAUREA paniculata. C. calycibus ciliatis: squamis planis, foliis bipinnatifidis rameis pinnatifidis linearibus, caule paniculato. *Leysf. hall. monsp. ged. Jacq. austr. t.* 320. *pal. fil. lithu. lugd. delph. burg.*

C. calycibus ciliatis, oblongis, foliis pinnatifidis. *Gmel. fib.* 2.

• *Cyanus pinnulis foliorum pinnatis, linearibus.* *Hall. helu.* 187.

Centaurea foliis bipinnatis linearibus; caule paniculato. *Scop. carn.*

C. micranthus. *Gmel. it. i. p.* 135. *t.* 23. *f.* 1.

C. calycibus ciliatis, foliis filiformibus bipinnatis, caule folioso ramoso. *Sauv. monsp.* 264

C. calycibus ciliatis oblongis, foliis scabris, caulium pinnatis, ramorum simplicissimis linearibus. *Gmel. fib.* 2. *p.* 98. *t.* 43.

Jacea non spinosa, foliis magis divisis, elatior, capitulis minoribus non splendentibus. *Morif. hist. 3. p.* 140. *f.* 7. *t.* 28. *f.* 15.

Stœbe major, cauliculis non splendentibus. *Bauh. pin.* 273.

Centaurei majoris species tenuifolia. *Bauh. hist. 3. p.* 31.

Habitat in G. Narbonensi, Austria, Hispania, Verona, Sibiria, Germania. ☉

Flores purpurei pistillis albis. Folia ramea pinnatifida.

Varietatem hujus etiam habemus Floribus majoribus, magnitudine C. Jaceæ, Foliis canescentibus, facieque non-nihil diversam.

18. CENTAUREA ragusina. C. calycibus ciliatis, foliis tomentosis pinnatifidis: foliolis obtusis ovatis integerimis: exterioribus majoribus. *Kniph. cent. 3. n.* 25.

Jacea cretica lutea, foliolis cinaræ. *Morif. hist. 3. p.* 141. *f.* 7. *t.* 27. *f.* 22.

Jacea arborea argentea ragusina. *Zan. hist. 107. t.* 43. *Mill. dict. t.* 152. *f.* 1.

Stœbe montana nivea, capite cardui, subrotundis foliorum lobis. *Barr. ic.* 309.

Habitat in Creta. Epidauro. 24

Folia alba quasi papposa, pinnata lobis rotundatis.

426 SYNGENESIA. POLYGAMIA FRUSTRANEA.

Flos solitarius, pedunculatus, luteus. Radio parvo
 stylis longis. Calyx magnus squamis acutis, ciliatis,
 flavescens.

19. CENTAUREA Cineraria. C. calycibus ciliatis ter-
 minati-filibus, foliis tomentosis bipinnatifidis : lobis
 acutis.

Jacea montana candidissima, flores foliis. Bauh. pin. 272.
 prod. 126. Moris. hist. 3. p. 141. f. 7. t. 26. f. 20.

β. Jacea cineraria laciniata, flore purpureo. Triumph. alt.
 72. Moris. hist. 3. p. 141. Barr. ic. 348. 347. Jacq.
 hort. 1. 92. est distincta species. R.

Habitat in Italia. 24

Folia alba quasi papposa : caulina bipinnata lobis inf.
 acutis : extremo palmato ; Ramea pinnatifida. Cili
 cilia fusca. Flos purpureus, Stylis albi ; radius corollae
 longior.

20. CENTAUREA argentea. C. calycibus ferratis, foliis
 tomentosis : radicalibus pinnatis : foliolis uniauratis.
 Scop. carn. 2. delph.

C. calycibus ciliatis villosis, foliis cuneiformibus super
 ferratis inferne dentatis. Hort. cliff.

Jacea cretica laciniata argentea, flore parvo flavescens
 Tournef. cor. 31. Barr. ic. 218.

Habitat in Creta.

Caulis ad radicem valde tomentosus. Folia Radica
 pinnata dense tomentosa : foliolis subaequalibus, ovalibus,
 basi inferiore tantum lobulo acutis. Caulina folia simplici,
 cuneiformia, ad basin altero margine instructi dentibus
 obtuso. Flores lutei. Folia alba, ut C. cineraria.

22. CENTAUREA sempervirens. C. calycibus ciliatis,
 foliis lanceolatis ferratis : infimo dente substipulari,
 inferioribus hastatis. Bocc. sic. 73. t. 39. f. 3. Dodon.
 mem. 85.

C. calycibus ciliatis, foliis lanceolatis ferratis : inferioribus
 hastatis. Spec. plant 2.

Jacea lusitana sempervirens. Moris. hist. 3. p. 139. f. 7.
 t. 28. f. 9.

Habitat in Lusitania. 24

caulis, pubescens ; inferne griseus. Folia lanceo-
 lata, pubescentia. Capitula : inferiora hastata
 superiora, versus caulem saepius falcata. Pediculi
 calycis, pubescentes.

SYNGENESIA. POLYGAMIA FRUSTRANEA. 417

Flores *flosculosi*. Folia *basi dente elongato*, unde quasi *stipulis instructus caulis*. *Syst. Veg. p. 654.*

23. CENTAUREA Scabiosa. C. calycibus ciliatis, foliis pinnatifidis: pinnis lanceolatis. *Fl. suec. parif. ingr. ged. pal sil. lugd. lithu. delph. burg.*

C. foliis pinnatifidis; pinnis lanceolatis. *Scop. carn.*

Cyanus foliis pinnatis, pinnis dentatis obtusis, squamis calycinis triangularibus. *Hall. helu. n. 186.*

Scabiosa major, squamatis capitulis. *Bauh. pin. 269.*

Scabiosa major altera, squamatis capitulis. *Bauh. pin. 269. ex Hall.*

Scabiosa major, *Dalech. hist. 1066.*

Habitat in Europæ septentrionalis pratis. 24

24. CENTAUREA stæbe. C. calycibus ciliatis oblongis, foliis pinnatifidis linearibus integerrimis. *delph.*

Stœbe incana, cyano similis, tenuifolia. *Bauh. pin. 273.*

Stæbe austriaca humilis. *Clus. hist. 2. p. 10.*

Habitat in Austria.

* * * *Rhapontica: calycinis squamis aridis scariosis.*

26. CENTAUREA orientalis. C. calycibus squama ciliatis, foliis pinnatifidis: pinnis lanceolatis. *Mill. dict. n. 17.*

Kniph. cent. 4. n. 16. Gmel. sib. 2. p. 98.

C. calycibus ciliatis, foliis pinnatis glabris; foliolis lanceolatis integerrimis. *Hort. ups.*

Cyanus foliis radicalibus: partim integris, partim pinnatis, bractea calycis ovali, flore sulphureo. *Hall. act. angl. 1745. vol. 43. n. 472. p. 94.*

Habitat in Sibiria. 24

Differs a C. Scabiosa, cui simillima, non tantum Corolla

sulphurea, sed præcipue Calyce, cujus foliola terminata squama ovata, ferruginea, ciliata, ciliis longitudine ipsius squamæ, pectinatis; cum in C. Scabiosa foliorum calycis margo niger ciliato-lacerus sit, nec in squamam eductus.

27. CENTAUREA Behen. C. calycibus scariosis, foliis radicalibus lyratis: lobis oppositis, caulinis amplexi-

caulibus. *Mat. med. 192. Gron. orient. 273. Lepech. 24. p. 264.*

Rhaponticoides lutea, foliis inferioribus dissectis, cæteris antharmi. Vail. act. 1718. p. 229.

Maratula affinis, capitulo squamoso luteo ut & flore. Bauh. pin. 235.

Behen album. Rauw. itin. 388. t. 288. Bauh. hist. 3. p. 36.

24. 37. Bauh. pin. 234.

426 SYNGENESIA. POLYGAMIA FRUSTRANEA.

Flos solitarius, pedunculatus, luteus. Radio parvo
stylis longis. Calyx magnus squamis acutis, ciliatis,
flavescentibus.

19. **CENTAUREA Cineraria.** C. calycibus ciliatis ter-
minali-sessilibus, foliis tomentosis bipinnatifidis : lobis
acutis.

Jacea montana candidissima, floebs foliis. *Bauh. pin.* 272.
prodr. 126. *Morif. hist.* 3. p. 141. f. 7. t. 26. f. 20.

β. *Jacea cineraria laciniata*, flore purpureo. *Triumph. obs.*
72. *Morif. hist.* 3. p. 141. *Barr. ic.* 348. 347. *Jacq.*
hort. t. 92. cui *distincta species.* R.

Habitat in Italia. 24

Folia alba quasi papposa : caulina bipinnata lobis incisis
acutis : extimo palmato ; Ramea pinnatifida. Calycis
cilia fusca. Flos purpureus ; Styli albi ; radius corolla via
longior.

20. **CENTAUREA argentea.** C. calycibus ferratis, foliis
tomentosis : radicalibus pinnatis : foliolis uniauritis.
Scop. carn. 2. *delph.*

C. calycibus ciliatis villosis, foliis cuneiformibus superne
ferratis inferne dentatis. *Hort. cliff.*

Jacea cretica laciniata argentea, flore parvo flavescente.
Tournef. cor. 31. *Barr. ic.* 218.

Habitat in Creta.

Caulis ad radicem valde tomentosus. Folia Radicalia
pinnata dense tomentosa : foliolis subæqualibus, ovalibus,
basi inferiore tantum lobulo auratis. Caulina folia simplicia,
cuneiformia, ad basin altero margine instructi denticulo
obtusos. Flores lutei. Folia alba, ut *C. cineraria*.

22. **CENTAUREA sempervirens.** C. calycibus ciliatis,
foliis lanceolatis ferratis : infimo dente substipulatis ;
inferioribus hastatis. *Bocc. sic.* 73. t. 39. f. 3. *Dodara*
mem. 85.

C. calycibus ciliatis, foliis lanceolatis ferratis : inferioribus
hastatis. *Spec. plant.* 2.

Jacea lusitanica sempervirens. *Morif. hist.* 3. p. 139. f. 7.
t. 28. f. 9.

Habitat in Lusitania. 24

Caulis teres, pubescens ; inferne griseus. Folia lanceo-
lata, ferrata, pubescentia, mollia : inferiora hastata.
Stipulae lanceolatae, versus caulem sæpius falcatae. Pedun-
culi superne incrassati. Calycis cilia grisea.

SYNGENESIA. POLYGAMIA FRUSTRANEA: 417

Flores *stosculosi*. Folia *basit dente elongato*, unde *quasi stipulis instructus caulis*. *Syst. Veg. p. 654.*

23. CENTAUREA Scabiosa. C. calycibus ciliatis, foliis pinnatifidis: pinnis lanceolatis. *Fl. Sueo. parisi. Ingr. ged. pal sil. lugd. lithu. delph. burg.*

C. foliis pinnatifidis; pinnis lanceolatis. *Scop. carn.*

Cyanus foliis pinnatis, pinnis dentatis obtusis, squamis calycinis triangularibus. *Hall. helu. n. 186.*

Scabiosa major, squamatis capitulis. *Bauh. pin. 269.*

Scabiosa major altera, squamatis capitulis. *Bauh. pin. 269. ex Hall.*

Scabiosa major, *Dalech. hist. 1066.*

Habitat in Europæ septentrionalis pratis. 24

24. CENTAUREA Stæbe. C. calycibus ciliatis oblongis, foliis pinnatifidis linearibus integerrimis. *delph.*

Stæbe incana, cyano similis, tenuifolia. *Bauh. pin. 273.*

Stæbe austriaca humilis. *Clus. hist. 2. p. 10.*

Habitat in Austria.

* * * *Rhapontica: calycinis squamis aridis scariofis.*

26. CENTAUREA orientalis. C. calycibus squama ciliatis, foliis pinnatifidis: pinnis lanceolatis. *Mill. dict. n. 17.*

Kniph. cent. 4. n. 16. Gmel. sib. 2. p. 98.

C. calycibus ciliatis, foliis pinnatis glabris; foliolis lanceolatis integerrimis. *Hort. upf.*

Cyanus foliis radicalibus: partim integris, partim pinnatis, bractea calycis ovali, flore sulphureo. *Hall. act. angl. 1745. vol. 43. n. 472. p. 94.*

Habitat in Sibiria. 25

Differt a C. Scabiosa, cui simillima, non tantum Corolla sulphurea, sed præcipue Calyce, cujus foliola terminata squama ovata, ferruginea, ciliata, ciliis longitudine ipsius squamæ, pectinatis; cum in C. Scabiosa foliorum calycis margo niger ciliato-lacerus sit, nec in squamam educus.

27. CENTAUREA Behen. C. calycibus scariofis, foliis radicalibus lyratis: lobis oppositis, caulinis amplexicaulibus. *Mat. med. 192. Gron. orient. 273. Lepech. I. p. 264.*

Rhaponticoides lutea, foliis inferioribus dissectis, cæteris catharmi. Vaill. act. 1718. p. 229.

Serratulæ affinis, capitulo squamoso luteo ut & flore. Bauh. pin. 235.

Behen album. Rauw. itin. 388. t. 288. Bauh. hist. 3. p. 36. t. 37. Bauh. pin. 234.

430 SYMPLICESIA. PORTULACA PORTULACA

Centaurium majus orientale caeruleum, pluri folio, l.

Jacea. Comm. rar. 39. & 39.

Habitus in Occidente, Sibiria. 27

Flos luteus: Radius 5 folio nunc.

36. *CENTAUREA conferta. C. calycibus brevibus, 5-*

sepalatis: radicalibus lanceolatis, caulis pinnatis, caule simpliciter. monsp. Belgii.

Jacea montana incana, capite pini. Bauh. pin. 272. M. 462. & 153.

Chamaeaea non aculeata. Loh. it. 7.

Habitus Montipellii et sanctis glauca. 28

Papposorum plumosus. Dill.

**** *Saxat. calycibus spinis pedunculatis.*

37. *CENTAUREA fructifera. C. calycibus peduncu-*

latis, foliis decurrentibus spinulosis angustis-dentatis.

Jacea lacustris, fructu folio, l. parva luteolae purpureae capite spinoso. Bauh. pin. 272. monsp. 272. M. 462. & 673. Pluk. p. 39. f. 1.

Habitus in maribus maris Medicamentum. 29

Folia interiore pedunculata, inferiore, luteolae-auris

superiore sessile, versus basem dentata: superiora folia

curvata, lanceolata. Caulis simplex, angustatus

decurrentibus, repando-spinulosis. Ramis caule longis

Flos solitarius, major, saepe depressus caulem aut

Calyx ovatus, viridis specis flavescentibus, maxime

fere recurva. Floris Radius 5 folia, major, purpurea

Descus albidus, apice purpureus.

38. *CENTAUREA serotina. C. calycibus peduncu-*

latis, foliis decurrentibus rotundatis oblongis: inferius

dentatis. monsp.

Jacea, foliis serotinis candidis, purpurea. Bauh. pin. 272.

Jacea purpurea hispanica americana. Bauh. hist. 3. p. 33. M.

Jacea bifolia, caule 2-ato, capite magno nuchitum. Pl.

phyl. 38. f. 1.

Jacea bifolia, spinosis capitis pelchis, flore purpureo

Bauh. hist. 3. p. 33. Rai. lig. 320.

Habitus in Hispania. 24

Planta tomentosa, vix pedalis: Caulibus simplicibus

Folia lato-lanceolata, serrata, decurrentia, juniora

spinosis. Calyces terminales, ovati: Squamis apice

Spinis setaceis 9. pluribusque. Corollae purpureae

duco albedo pauciflora.

39. *CENTAUREA romana. C. calycibus peduncu-*

SYNGENESIA. POLYGAMIA FRUSTRANEA. 429

Stoebe calyculis argenteis. Bauh. pin. 273.

Stoebe salmantica 3. Clus. hist. 3. p. 10.

Habitat in Helvetia, Hispania, Sibiria. ♂

33. *CENTAUREA Rhapontica. C. calycibus scariosis, foliis ovato-oblongis denticulatis integris petiolatis: subtus tomentosis. delph.*

Centaureum foliis petiolatis, lanceolatis, ad pediculum emarginatis. Hall. helv. n. 160.

Rha capitatum Inulæ s. Centauræ majoris facie. Lob. ic. 288. R.

Rhaponticum alterum angustiori folio. Lob. ic. 288. Hall. R.

Centaureum majus germanicum. Cord. hist. 4. c. 59. Hall. R.

Rhaponticum folio helenii incano. Bauh. pin. 117.

Rhaponticum angustifolium incanum. Bauh. pin. 117. Hall.

Rha s. Rhei ut existimatur. Dod. pempt. 389.

Habitat in Alpibus Helveticis, Veronæ.

Folia subcordato-oblonga, serrata, subtus albo-tomentosa. Caulina pauca; aliquando pinnatifida. Flos solitarius, magnus. Flosculi neutri nulli. Hall.

34. *CENTAUREA babylonica, C. calycibus scariosis, foliis subtomentosis decurrentibus indivisis: radicalibus lyratis.*

*Serratula (babyloniæ) foliis tomentosis decurrentibus subintegerrimis concavis canaliculatis: radicalibus lyratis, floribus thyrsoides - spicatis. Spec. ed. Gouan. monsp. illustr. **

Raponticoides lutea altissima, helenii folio, alato caule. Vaill. act. 182. n. 27.

Jacea latissima, laciniato folio. Bauh. pin. 235.

Jacea maxima. Alp. exot. t. 282.

Jacea babylonica. Bauh. prodr. 235.

Habitat in Oriente. 24

Videtur Cent. glastifoliæ proxima. Caulis ramosus & ramis angulatis. Folia radicalia lanceolato-ovata, subdenticulata, basi sæpe bidentata, scabriuscula, petiolata, magnitudine Belladonnæ. Caulina lanceolata, decurrentia (ut in Centaur. glastifolia) Calyx leviter scariosus. Corollæ lutea. Flores in thyrsum subsessiles pedicellis subquadrifloris.

35. *CENTAUREA glastifolia. C. calycibus scariosis, foliis indivisis integerrimis decurrentibus. Gmel. sib. 2. p. 83. Gouan. illustr. 75. Kniph. cent. 4. n. 14.*

430 SYNGENESIA. POLYGAMIA FRUSTRANEA.

Centaureum majus orientale erectum, gladi folio, flore luteo. Comm. rar. 39. t. 39.

Habitat in Oriente, Sibiria. 24

Flos luteus: Radio disco minore.

36. *CENTAUREA conifera. C. calycibus scariolis, foliis tomentosis: radicalibus lanceolatis, caulinis pinnatifidis caule simplici. monsp. delph.*

Jacea montana incana, capite pini. Bauh. pin. 272. Mill. dict. t. 153.

Chamæleon non aculeatus. Lob. ic. 7.

Habitat Monspelii in saxosis glareosis. 24

Pappus seminum plumosus. Dill.

* * * * *Stæbæ: calycinis spinis palmatis.*

37. *CENTAUREA sonchifolia. C. calycibus palmato-spinosis, foliis decurrentibus spinulosis repando-dentatis.*

Jacea laciniata, sonchi folio, s. jacea latifolia purpurea, capite spinoso. Bauh. pin. 272. prodr. 128. Herm. lugdb. 331. t. 675. Pluk. phyt. 39. f. 1.

Habitat in maritimis maris Mediterranei. ☉ 24.

Folia inferiora petiolata, indivisa, lanceolato-ovata; superiora sessilia, versus basin dentata; suprema subdecurrentia, lanceolata. Caulis simplex, angulatus foliis decurrentibus, repando-spinulosis. Ramus caule longior. Flos solitarius, magnus, saepe deprimis caulem ad ætrem. Calyx ovatus, viridis spinis flavescentibus, media longiore, fere recurva. Floris Radius 5, fidus, major, purpureus; Discus albidus, apice purpurascens.

38. *CENTAUREA seridis. C. calycibus palmato-spinosis, foliis decurrentibus tomentosis oblongis: infimis sinuato-dentatis. monsp.*

Jacea, foliis seridis candicantibus, purpurea. Bauh. pin. 272.

Jacea purpurea hispanica muricata. Bauh. hist. 3. p. 33. descr.

Jacea latifolia, caule alato, capite magno turbinato. Pluk. phyt. 38. f. 1.

Jacea latifolia, spinosis capitulis pulchris, flore purpureo.

Bauh. hist. 3. p. 33. Rai. hist. 320.

Habitat in Hispania. 24

Planta tomentosa, vix pedalis: Caulibus subramosis.

Folia lato-lanceolata, serrata, decurrentia, serraturis subspinosis. Calyces terminales, ovati: Squamis apice palmatis Spinis seraceis 9. pluribusque. Corollæ purpureo radio, disco albido pauciflora.

39. *CENTAUREA romana. C. calycibus palmato-spinosis,*

SYNGENESIA. POLYGAMIA FRUSTRANEA. 531

foliis decurrentibus inermibus & radicalibus pinnatifidis : impari maximo.

Jacea spinosa cretica. *Zern. hist. 141. & 142.*

Cyanus crucæ folio, flore rubro. *Barr. rar. 87. t. 504.*

Habitat in Campania Romana. ☉

40. *CENTAUREA sphærocephala.* C. calycibus palmato-spinosis, foliis ovato-lanceolatis petiolatis dentatis.

Jacea sphærocephala spinosa tingitana. *Herm. lugdb. 332. t. 333. Moris. hist. 3. p. 143. f. 7. t. 27. f. 9.*

Habitat in Mauritania, Hispania.

41. *CENTAUREA Isnardi.* C. calycibus palmato-spinosis, foliis lyrato dentatis hispidis subamplexicaulibus, floribus sessilibus terminalibus.

Calcitrapoides procumbens, cichorii folio, flore purpureascente. *Isn. paris. 1719. p. 164. t. 9.*

Habitat in Europa australi ? 24

Varietas forte sola C. pullatæ vel ejus hybrida proles.

42. *CENTAUREA napifolia* C. calycibus palmato-spinosis foliis decurrentibus sinuatis spinulosis : radicalibus lyratis. *Hort. ups. 272. **

Jacea cyanoides altera, alato caule. *Her. par. 189. t. 189. bon.*

Jacea peregrina napifolia, echinatis capitulis, caule lato. *Pluk. alm. 192. t. 94. f. 2. Moris. hist. 3. p. 154. f. 7. t. 26. f. 20.*

Habitat in Creta. ☉

Folia radicalia lyrata. Calycis spinæ palmatæ, flavæ. Corolla radiata purpurea : Neutræ 5. fidæ : laciniis duabus interioribus minoribus.

43. *CENTAUREA aspera.* C. calycibus palmato-trispinosis, foliis lanceolatis dentatis. *delph.*

C. calycibus spinosis, foliis lanceolatis dentatis. *Roy. lugdb.*

Stoebe squamis asperis. *Bauh. pin. 273. Rai. hist. 319.*

Jacea, cum spinosis capitulis, purpurea tenuifolia. *Bauh. hist. 3. p. 33.*

Jacea supina maritima laciniata, capite spinoso, flore rubello. *Bocc. mus. 2. p. 35. t. 26.*

β. *Carduus pumilus supinus, Jaceæ folio incano & tenui.* *Bocc. rar. 919.*

Habitat Monspelii, inque Hetruria, Lusitania.

Calycis spinæ. 3. 5. plices.

***** *Calcitrapæ, calycinis spinis compositis.*

44. *CENTAUREA benedicta.* C. calycibus duplicato-spinosis lanatis involucreatis, foliis semidecurrentibus denti-

432 SYNGENESIA. POLYGAMIA FRUSTRANEA.
culato-spinosis. *Kniph. cent. I. t. 16. Ludw. ed. t. 6.*
Blackw. t. 476. Regn. bot. delph.

Cnicus caule diffuso, foliis dentato-sinuatis. Hort. ups.

Carduus sylvestris hirsutus f. Carduus benedictus. Bauh. pin.
378.

Carduus benedictus. Cam. epit. 562.

Habitat in Chio, Lemno, Hispania ad versuras. ☉

Radix flosculi parvi, trifidi.

45. CENTAUREA eriophora. C. calycibus duplicato-spinosis lanatis, foliis semidecurrentibus integris sinuatisque caule prolifero. *Kniph. cent. n. 10.*

C. calycibus duplicato-spinosis, foliis decurrentibus integris. *Hort. cliff.*

Calcitrapa lutea, alato caule, capite eriophoro. Vaill. ad.
1718. p. 212.

Carduus lusitanicus canescens, alato caule, capite lanuginoso. Tournef. inst. 441.

Habitat in Lusitania. ☉

Caulis prolifer. Folia inermia. Radicalia lanceolata. Caulina laciniata; Ramea linearia. Calyx magnus, valde lano intertextus. Corolla parva, lutea. Flosculi neutri quadrifidi hermaphroditi breviores.

47. CENTAUREA Calcitrapa. C. calycibus subduplicato-spinosis sessilibus, foliis pinnatifidis linearibus dentatis, caule piloso. *pal. Kniph. cent. II. n. 25. Regn. bot. lugd. burg. delph.*

C. calycibus subduplicato-spinosis sessilibus, foliis linearibus pinnatifidis. *Hort. cliff. gelr. paris. monsp.*

Calcitrapa foliis pinnatis, pinnis dentatis, floribus in alis sessilibus. Hall. helv. n. 194.

Centaurea foliis pinnatifidis; pinnis linearibus, acutis, dentatis; calycibus sessilibus. Scop. carn. sub Rhapon-tico.

Carduus stellatus f. Calcitrapa. Bauh. hist. 3. p. 89.

Carduus stellatus, foliis papaveris erratici. Bauh. pin. 387.
Rai. hist. 317.

Carduus muricatus vulgo Calcitrapa dictus. Clus. hist. 2. p. 7.

Hippophæstum. Col. phytob. 85. t. 24.

Habitat in Helvetia, Anglia & Europa australiori secus vias.

Flos huic purpureus, nec ut in Solstitiali, melitenfi aus-sicula luteus. Caulis prolifer. Spinæ calycis albæ. Variat albo flore.

SYNGENESIA, POLYGAMIA FRUSTRANEA. 433

48. *CENTAUREA calcitrapoides*. C. calycibus subduplicato spinosis; foliis amplexicaulibus lanceolatis indivisis serratis. *Carduus stellatus*, foliis integris serratis. *Magn. monsp.* 292.

Habitat Monspelii & in Palæstina.

Folia C. *Calcitrapæ*, sed foliis indivisis.

49. *CENTAUREA solstitialis*. C. calycibus duplicato spinosis solitariis, foliis ramulis decurrentibus inermibus lanceolatis, radicalibus lyrato-pinnatifidis. *paris. monsp. carn. Kniph. cent. 3. n. 23. delph. burg.*

Calcitrapa caule alato, foliis tomentosis, radicalibus semipinnatis, caulinis lanceolatis, dentatis. *Hal. helv. n. 193.*

Carduus stellatus luteus, foliis cyathi. *Bauh. pin. 387. Rat. hist. 317.*

Carduus stellatus mitior apulus. Col. ecphr. i. p. 30. l. 31.

Habitat in Gallia, Anglia, Italia, Germania, &c. ☉

Folia radicalia lyrata, acuminata. Flores terminales solitarii. Pedunculi longissimi, foliosi; ad calycem nullum foliolum. Spinæ calycis albæ, longiores sub calyce, basi tantum dentatæ.

50. *CENTAUREA melitensis*. C. calycibus duplicato spinosis terminalibus confertis; foliis decurrentibus lanceolatis sinuosis inermibus. *Kniph. cent. 4. n. 15.*

Jacea melitensis, capitulis conglobatis. *Bocc. sec. 61. t. 356. Moris. hist. 3. p. 145. Rat. hist. 323.*

Habitat in Melita, Monspelii. ☉

Folia radicalia lyrata, obtusa. Flores terminales; conferti, 2. f. 3. Pedunculi brevissimi. Spinæ suo calyce non longiores, supra basin etiam dentatæ; ad Calycem foliolum unum alterumve, calycis ipsius longitudine.

51. *CENTAUREA sicula*. C. calycibus ciliato spinosis; foliis decurrentibus lyratis inermibus incanis; capitulis terminalibus. *monsp.*

Jacea sicula, cichorii folio, flore luteo, capite spinoso. *Magn. monsp. 138. Moris. hist. 3. p. 144. f. 7. t. 28. f. 16.*

Habitat in Sicilia. 24

Folia Radicalia lyrata, obtusa denticulata; scabræ; caulina lanceolata, decurrentia. Calyces magni, ovati; Squamis spina majore fusca.

52. *CENTAUREA centauroides*. C. calycibus ciliatis spinosis; foliis lyrato-pinnatis integerrimis; lacinia impari majore. *Kniph. cent. 6. n. 23. Regn. bot.*

Jacea lutea spinosa centauroides. Bauh. pin. 272.

Linnaei spec. plant. Europ. Pars II.

E 6

334 SYNGENESIA. POLYGAMIA FRUSTRANEA.

Jacea laevis maxima centauroides apula. Col. ecphr. 1. p. 33.
t. 35.

Habitat in Italia, Hispania, Monspeli.

Calycis squamæ ciliatæ spina terminali, sed interiores squamæ scariosæ, muticæ, ciliatæ.

53. *CENTAUREA collina. C. calycibus ciliatis inermi-spinosis, foliis radicalibus bipinnatifidis, caule acutangulo.*

Jacea lutea, capite spinoso. Bauh. pin. 272.

Jacea luteo flore. Clus. hist. 2. p. 8.

Habitat in Italia, Hispania, Monspeli, Carniolia. 2.

54. *CENTAUREA rupestris. C. calycibus ciliatis spinosis, foliis bipinnatis linearibus.*

Jacea laciniata lutea. Bauh. pin. 272.

Jacea montana minima tenuifolia. Col. ecphr. 1. p. 36.
t. 35. f. 2.

Habitat in Italia. 2.

Affinis C. collinæ, calycinis squamis fusco-ciliatis terminalis spina debili, simplici, squama ipsa brevior; Squamis vero intimis scariosis. Caulis parum ramosus, vix angulatus. Folia fere omnia bipinnata, linearia, excepto margine lævia; minime vero lanceolata pinnis cornu cervi modo ramosis.

* * * * * *Crocodiloidea, spinis simplicibus.*

56. *CENTAUREA salmantica. C. calycibus setula subspinosa exstante glabris, foliis lyrato-runcinatis ferratis. Jacq. hort. t. 64. burg.*

C. calycibus setula reflexa, spinosis glabris, foliis lyrato-runcinatis, ferratis. Sp. pl. monsp.

Jacea major, foliis cichoraceis mollibus, flore stamineo.

Morif. hist. 3. p. 140. f. 7. t. 26. f. 14.

Jacea major, foliis cichoraceis mollibus lanuginosis. Bauh. pin. 273.

Stoebe salmantica 1. Clus. hist. 2. p. 9. Rai. hist. 324.

Habitat in Europa australi. 2.

Caulis paniculatus: ramis simpliciusculis. Floribus terminalibus. Folia Radicalia lanceolata, serrata, hirta, serraturis mucronatis. Caulina runcinata, hispido-scabra, serraturis subspinosis. Calyces glaberrimi, ovati. Spinæ calycinæ parvæ, ut vix manifestæ. Corollæ albæ aut purpureæ: masculis 4. partitis, non majoribus.

57. *CENTAUREA cichoracea. C. calycibus setaceo-spinosis, foliis decurrentibus indivisis ferrato-spinosis. †*

SYNGENESIA. POLYGAMIA FRUSTRANEA. 435

Jacea foliis cichoreis , caule alato , flore purpureo. *Rei. suppl.* 203. *Till. pis.* 84. t. 27.

Habitat in monte argentario Italiae.

58. *CENTAUREA muricata*. C. calycibus simplicissime spinosis , foliis inferioribus pinnatifidis , superioribus lanceolatis , pedunculis longissimis.

Jacea cyanoides , echinato capite. *Bauh. pin.* 272. *Morif. hist.* 3. p. 195.

Cyanoides flos. *Dod. pempt.* 251.

Habitat in Hispania. ☉

59. *CENTAUREA peregrina*. C. calycibus setaceo-spinosis ; foliis lanceolatis petiolatis : inferne dēntatis.

Centaureum majus , folio molli acuto lanceolato , flore aureo magno , calyce spinoso. *Bark. lugdb.* 1. p. 144.

Habitat in Europa australi. ♂

61. *CENTAUREA nudicaulis*. C. calycibus setaceo-spinosis , foliis indivisis : superioribus subdentatis , caule simplici nudiusculo unifloro. † *Ger. proa.* 187. t. 5.

Jacea folio cerinthes , purpurascēte flore , e rupe victoriæ. *Herm. parad.* 190. t. 190.

Jacea inrybacea , capite rubro spinoso. *Barr. ic.* 1218. *Bocc. mus.* 2. p. 60. t. 48.

Habitat in Italia , rupe victoriæ Galloprovinciæ. 24

Folia prima obovata , integerrima ; reliqua ovato-lanceolata , basi dentata. Caulis subnudus.

62. *CENTAUREA Crocodilium*. C. calycibus scariosis simplicissime spinosis , foliis pinnatifidis integerrimis : lacinia extima majore dentata.

Crocodilium vulnerariæ foliis. *Vaill. ac.* 183.

Cyanus purpureus , capite spinoso , syriacus. *Barr. rar.* 927. t. 503. *bon.*

Habitat in Syria. ☉

Caulis Ramis striatis , scabris. Folia lyrata , pinnata aut indivisa. Ped. longi , uniflori. Calycis squamæ ovatae , gibbae marginibus membranaceis , albis , apice spina simplici terminatis. Flores Neutri maximi , 5. fidi , subæquales , purpurascētes ; Hermaphrodisi albidī apicibus purpurascētibz , longitudine radii.

65. *CENTAUREA galactites*. C. calycibus setaceo spinosis ; foliis decurrentibus sinuatis spinosis subtus tomentosis. *monsp.*

Carduus tomentosus , capitulo minore. *Bauh. pin.* 382.

Carduus galactites. *Bauh. hist.* 3. p. 54.

436 SYNGENESIA. POLYGAMIA NECESSARIA:
β. Carduus creticus non maculatus, caule alato. *Tournef.*
cor. 31.
Habitas in Europa australi. 26

POLYGAMIA NECESSARIA.

1073. CALENDULA.

Recept. nudum. Pappus nullus. Cal. polyphyllus
æqualis. Sem. disci membranacea.

1. CALENDULA *arvensis*. C. seminibus cymbiformibus
muricatis incurvatis: extimis erectis pratensis. *Fl. suet.*
pal. lugd. delph.

C. foliis lanceolatis, amplexicaulibus, dentatis, floribus
concoloribus. *Hall. helv. n.* 94.

Caltha officinalis. *Scop. carn.*

Caltha arvensis. *Bauh. pin.* 275.

Caltha minima. *Bauh. hist.* 3. p. 103.

Habitas in Europæ arvis. ☉

Affinis nimium C. officinali Foliis subdentatis, sed ex-
cordato-lanceolatis, nec spatulatis. Floribus minoribus.
Seminibus angustioribus, linearibus: exterioribus erectis,
longioribus.

3. CALENDULA *officinalis*. C. seminibus cymbiformibus
muricatis incurvatis omnibus. *fil. Ludw. et.* 1. 79. *Blackw.*
1. 106. *Kniph. cent.* 6. n. 18. *cent.* 11. n. 19. *Knorr. del.*
1. 1. R. 21. *Regn. bot. burg. lugd.*

C. seminibus radii cymbiformibus echinatis: bicornibus.
Hort. upf. parif.

Caltha vulgaris. *Bauh. pin.* 275.

β. Caltha polyanthos major. *Bauh. pin.* 275.

γ. Caltha floribus reflexis. *Bauh. pin.* 275.

δ. Caltha prolifera, majoribus floribus. *Bauh. pin.* 275.

Habitas in Europæ australis arvis. ☉

3. CALENDULA *pluvialis*. C. foliis lanceolatis sinuato-
denticulatis, caule folioso, pedunculis filiformibus. *Hort.*
upf. 274.

C. scabra. *Berg. cap.* 310.

C. foliis dentatis. *Roy. lugdb.* 177. *Mill. dict. n.* 4. *in*
1. 75. f. 2.

SYNGENESIA. POLYGAMIA NECESSARIA.

C. seminibus radii obsolete serratis, disci cordatis. *Hort. cliff.* 425.

C. humilis africana, flore intus albo, foris violaceo simplici. *Herm. lugdb.* 104. t. 105. *Breyn. ic.* 26. t. 14. f. 1. *Stiff. bot.* 59. t. 59.

Caltha africana, flore intus albo extus ferrugineo. *Morif. hist.* 3. p. 14. f. 6. t. 3. f. 8.

Habitat in Æthiopia. ☉

6. CALENDULA nudicaulis. C. foliis lanceolatis sinuato-dentatis, caule subnudo. *Kniph. cent.* 2. n. 10.

C. africana, foliis scabiosæ argenteæ viridibus. *Herm. afr.* 5.

Bellis florum pediculis pene aphyllis, foliis incis. *Comm. hort.* 2. p. 66. t. 33.

Caltha africana, flore intus albo extus leviter violaceo, femine plano cordato. *Boerh. lugdb.* 1. p. 113.

Habitat in Æthiopia.

Caulis simplicissimus, erectus, inferne foliosus. Folia spatulata, integerrima, scabra rarius dente angulato. Semina orbiculata.

1074. A R C T O T I S.

Recept. villosum f. paleaceum. Pappus corona pentaphylla. Cal. imbricatus: squamis apice scariosis.

* Receptaculo villoso.

1. ARCTOTIS calendulacea. A. flosculis radiantibus sterilibus, foliis runcinatis subtomentosis. *Syst. Veg.* 659.

2. Arctotis (Calendula) flosculis radiantibus sterilibus duodenis subintegris, foliis lyratis nigro-denticulatis. *Spec. ed.* 2. p. 1306. *Mill. dict. n.* 4. *Kniph. cent.* 10. n. 10.

A. foliis ovatis dentatis: petiolis longissimis superne dentatis, caule ramoso. *Hort. cliff.* 412.

Anemone affinis æthiopica, fibrosa radice, flore asteris, taraxaci foliis subincanis. *Herm. lugdb.* 43. t. 42.

Anemonospermos africana, jacobææ maritimæ foliis, flore sulphureo. *Comm. rar.* 36. t. 36.

Habitat in Æthiopia. ☉

Pedunculi villosi, adpersi setis rubris, ante & post florescentiam nutantes, Radius flavus, subtus purpureascens.

438 SYNGENESIA. POLYGAMIA NECESSARIA.

noctu contortus. Corollulæ disci extus nigrae. Antheris nigris. Calycis squamæ exteriores patentes subulatae, brevissimæ, hispidae. Ex spec. plant. 2. p. 1306.

β. *Arctotis (hypochondriaca)* flosculis radiantibus sterilibus trifidis media inflexa.

A. tristis flosculis radiantibus vicenis tripartitis. *Spec. pl. 2. p. 1306. Mill. dict. n. 1.*

A. foliis pinnatifidis, caule herbaceo, petalis radii profunde trifidis. Roy. lugdb. 179.

Anemonospermos afra, foliis & facie taraxaci incanis. Brey. ic. 27. t. 15.

γ. *Arctotis (tristis)* flosculis radiantibus sterilibus quadripartitis. *Mant. 479. Berg. cap. 320.*

Anemonospermos africana, Jacobææ maritimæ foliis, flore sulphureo. Comm. rar. 36. t. 36.

Caules patuli, hirsuti. Folia runcinata, crassiuscula, fragilia, serraturis spinulosis vagis, pubescentia, subtus villoso-alba. Pedunculi striati, longi hirti floccosi. Calyces ante anthesin cernui, squamis apice villoso-ciliatis. Discus niger corollulis luteis apice nigris. Pistilla clava lutea. Radius flosculis radiantibus 4. f. 5. partitis, luteis subtus plumbeis; lacinia intermedia divaricata obvallante discum. Mant. 479.

δ. *Arctotis (coruscans)* flosculis radiantibus sterilibus tripartitis; exteriore trifida.

ε. *Arctotis (superba)* flosculis radiantibus sterilibus quinquepartitis, foliis pinnatis dentatis: laciniis bipartitis. *Spec. ed. 2. p. 1305. Amæn. acad. 4. p. 331. **

Jacobæa africana, stœbes facie, flore ampliore, petalis radialibus trifidis. Volk. norib. 224. t. 224.

Habitas in Æthiopia.

Pedunculi villosi, adspersi setis hyalinis, noctu decumbentes, nunquam vero cernui. Radius subtus virescens, noctu connivens. Corollulæ disci luteæ, Antheris luteis.

1075. OSTEOSPERMUM.

Recept. nudum. Pappus nullus. Cal. polyphyllus.

Sem. globosa, colorata, ossea.

1. *OSTEOSPERMUM spinosum. O. spinis ramosis. Hort. cliff. 424.*

SYNGENESIA. POLYGAMIA NECESSARIA 439

Chrysanthemoides osteospermum africanum odoratum spinosum & viscosum. Comm. hort. 2. p. 85. t. 43.

Chrysanthemum africanum frutescens spinosum. Volk. nerib. 105. t. 105. Moris. hist. 3. l. 6. t. 3. f. 56.

Habitat in Æthiopia. ☿

Maxime affinis O. pififero. Fructus oblongo pedunculo pendulus Baccis in orbem digestis, fusco - purpurascens, asseis, globosis.

1076. O T H O N N A.

Recept. nudum. Pappus subnullus. Cat. 1. phyllus, multifidus, subcylindricus.

3. *OTHONNA pedinata. O. foliis pinnatifidis: laciniis linearibus parallelis. Mill. dict. n. 3. ic. t. 194. f. 2.*

Jacobæa africana frutescens, foliis absinthii umbelliferi incanis. Comm. hort. 2. p. 137. t. 69. Rai suppl.

β. *Jacobæa absinthites, tomentosis cinerariæ foliis, æthiopica, calyce integro. summis. pris. dentato. Pluk. alm. 106.*

Habitat in Æthiopia. ☿

Planta sæpe tota tomentosa est. Caulis cicatrifatus.

1079. F I L A G O.

Recept. nudum. Pappus nullus. Cat. imbricatus. Flosculi feminei inter squamas calycis locati.

1. *FILAGO acaulis. Filago (pygmæa) floribus acaulibus sessilibus, foliis floralibus majoribus. monsp.*

Santolina caule simplicissimo, foliis superioribus majoribus. Roy. lugdb. 147.

Gnaphalium (species omnes). Vaill. art. 1719. p. 314.

Gnaphalium roseum hortense. Bauh. pin. 263. prodr. 122. Barr. ic. 127.

Habitat in Europæ australis & Orientis stagnis exsiccatis. ☉

Flores sessiles radicales intra orbem foliaceum. Raro caulescit.

2. *FILAGO germanica. F. panicula dichotoma, floribus rotundatis axillaribus hirsutis, foliis acutis. gallob. carn. sub. Gnaphalio. herb. pal. sil. delph. lith. burg. lugd.*

E c 4.

440 SYNGENESIA. POLYGAMIA NECESSARIA.

F. caule dichotomo, corymbo sphaerico, in angulo sessilibus.
Hall. helv. n. 153.

Gnaphalium caule erecto dichotomo, floribus in alis sessilibus. *Fl. suec.*

Gnaphalium vulgare majus. *Bauh. pin. 263.*

Gnaphalium germanicum. *Bauh. hist. 3. p. 158.*

Gnaphalium. *Fuchs. hist. 222.*

Habitat in Europa. ☉

Semina pappum gerunt. R.

3. FILAGO pyramidata. **F.** caule dichotomo, floribus pyramidatis pentagonis axillaribus, flosculis femineis seriatis.
Sp. pl. delph.

Gnaphalium medium. *Bauh. pin. 263.*

Habitat in Hispania. ☉

Caulis bipollicaris, erectus. Ramis radicalibus. Folia lanceolata, obtusiuscula. Flores e dichotomia caulis & terminales, sessiles, pentagono-pyramidati, tomentosi ut tota planta, aggregati.

4. FILAGO montana. **F.** caule subdichotomo erecto, floribus conicis terminalibus axillaribusque. *Flor. suec. pal. herb. delph. lugd.*

F. caule dichotomo, floribus glomeratis spicatis. *Hall. helv. n. 115.*

Gnaphalium caule erecto diviso, floribus pyramidatis axillaribusque. *Fl. suec.*

Gnaphalium minus repens. *Bauh. pin. 263.*

Gnaphalium minimum. *Lob. ic. 481. Bauh. hist. 3. p. 129. Rai. hist. 296.*

Habitat in Europæ sabulosis, montosis. ☉

Flosculi utriusque sexus. Hall. Leers. R.

5. FILAGO gallica. **F.** caule dichotomo erecto, floribus subulatis axillaribus, foliis filiformibus. *Polich. pal. delph. burg. lugd.*

Gnaphalium vulgare medium. *Moris. hist. 3. f. 7. t. 11. f. 14. Pet. herb. t. 18. f. 12.*

Gnaphalium minimum alterum nostras, stœchadis citrinæ foliis tenuissimis. *Pluk. alm. 172. t. 298. f. 2.*

Gnaphalium vulgare tenuifolium. *Bauh. hist. 3, p. 159. Hall. R.*

Habitat in Gallia, Helvetia, Anglia, Germania.

Folia subtomentosa, tamen glabra, nec hirsuta. (Flosculi utriusque sexus. Hall. R.)

6. FILAGO arvensis. **F.** caule paniculato, foliis conicis la-

SYNGENESIA. POLYGAMIA NECESSARIA. 441

teralibus. *Flor. suec. carn. sub Gnaphalio. herb. pal. delph. lugd. lithu. burg.*

F. caule erecto, ramosissimo, foliis mollissimis, tomentosis, florum acervis sessilibus, in spicas continuatis, *Hall. helu. n. 154.*

Gnaphalium paniculatum hirsutum. Fl. suec. i. n. 679.

Gnaphalium majus, angusto oblongo folio. Bauh. pin. 263.

Filago incana tomentosa erecta. Vaill. ad. 1729. p. 391.

Habitat in Europæ campis sabulosis. ☉

Semen hermaphroditum tantum pappo destitutum. Leers.

7. *FILAGO Leontopodium. F. caule simplicissimo, capitulo terminali bracteis hirsutissimis radiato. delph.*

F. floribus sessilibus, umbellatis, foliis umbellam excedentibus, tomentosis. *Hall. helu. n. 152.*

Gnaphalium Leontopodium. Scop. carn. Jacq. austr. t. 86.

Gnaphalium alpinum, magno flore, folio oblongo. Bauh. pin. 264.

Gnaphalium alpinum pulchrum. Bauh. hist. 3. p. 161.

Gnaphalium alpinum. Clus. hist. 1. p. 328.

Leontopodium. Dod. pempt. 68.

8. *Gnaphalium magno flore, brevi folio. Bauh. pin. 264.*

Habitat in Alpibus Helvetiæ, Valesiæ, Corinthi, Austriæ, Sibiriae, Germaniæ. Gmel. sub. p. 267. 2.

Planta canescens, terminata stella foliacea lanuginosa, e cujus centro Flores hermaphroditi quinquesidi. Bracteis hirs radiantibus insident Flores partim feminei 4. fidi, partim incompleti, destituti staminibus, stylo, germine. Scop. ann. 2. p. 63.

1080. M I C R O P U S.

Recept. paleaceum. Pappus nullus. Cal. calyculatus.

Radius corollæ nullus. Flosculi feminei a squamis calycinis involuti.

1. *MICROPUS supinus. M. caule procumbente, foliis geminis. delph.*

M. foliis floralibus oppositis, fructibus muricatis. Gouan. illustr. p. 74.

Gnaphalium supinum, echinato semine. Pluk. alm. 171. 4. 187. f. 6. Rai. suppl. 191.

INESIA. POLYGAMIA SEGREGATA.

halum supinum, semine echinato. *Morif. hisp.*

franzæ, Italiæ, Orientis maritimis. 24

1080. *US erectus*. M. caule erecto, calycibus
coarctatis, foliis solitariis. *delph.*

M. feminibus compressis lanatis inermibus. *Læfl. hisp.* 166.
t. 1. f. 3. *

Gnaphalium caule ramoso diffuso, floribus confertis lan-
rectis. *Hort. cliff.*

Gnaphalium minus repens. *Magn. monsp.* *

Gnaphalium minus, latioribus foliis. *Bauh. pin.* 263.

Gnaphalium plateau. 3. *Clus. hisp.* 1. p. 229.

Filago f. *Impia* capitulis compressis. *Vaill. parif.* 52.

Leontopodium verius *dioscoridis*. *Barr. ic.* 296.

Habitat in Orientis, Galliæ, Hispaniæ collibus. ☉

POLYGAMIA SEGREGATA.

1081. ELEPHANTOPUS.

Cal 4. florus. *Cor.* ligulatz, hermaphroditæ.

Recept. nudum. *Pappus* fetaceus.

1. *ELEPHANTOPUS scaber*. E. foliis oblongis scabris.
Hort ups.

E. erectus, foliis oblongo-ovatis rugosis serratis: floralibus
cordiformibus ternatis, capitulis remotis terminalibus.
Brown. jam. 312.

E. conyzæ folio. *Dill. elth.* 126. t. 106. f. 126. *Vaill. ad.*
1719.

Bidens frutescens, foliis oblongis utrinque acuminatis ve-
nosis & lanuginosis. *Breyn. ic.* 31. t. 34.

Echinophoræ indicæ affinis, semine & floribus in capi-
tulis lævibus, in caulium cymis. *Pluk. alm.* 132. A
388. f. 6.

Anaschovadi. *Rheed. mal.* 10. p. 13. t. 7. *Burm. ind.* 189.

Habitat in Indiis. 24

Cal. 1. florus. *Cor.* tubulosa, hermaphrodita.

Recept. fetosum. *Pappus* obsoletus.

1. **ECHINOPS** *sphærocephalus*. E. capitulis globosis, foliis pubescentibus. *carn. Scholl. barb. delph.*

E. calyculis unifloris, caule multifloro, foliis spinosis supra nudis. *Sp. pl.* 1.

E. floribus capitatis, calycibus unifloris. *Hort. upf. paris.*

E. caule ramoso, foliis semipinnatis, pinnis dentatis, dentibus aristatis. *Hall. helv. n.* 158.

E. foliis supra villosis hispidis, subtus tomentosis, caule multifloro. *Gouan. illustr.*

Echinopus major. *Bauh. hist.* 3. p. 69.

Carduus *sphærocephalus* latifolius vulgaris. *Bauh. pin.* 381.

Chalcepos. *Dalech. hist.* 1480.

Habitat in Italia, Austria, Germania. 24

Caulis ramosus, multiflorus, pubescens villis patentibus.

Folia supra pilosa. *Calycis squamæ* omnes ad apicem usque pubescentes.

3. **ECHINOPS** *Ritro*. E. capitulo globoso; foliis pinnatifidis supra glabris. *n. 2. ic. t.* 130. *Scop. carn. delph.*

E. caule subunifloro. *Hort. upf.*

E. foliis supra glaberrimis, subtus tomentosis, caule multifloro corymbofo. *Gouan. illustr.* 64.

E. caule subunifloro, foliis duplicato pinnatifidis, foliolis latiusculis vicinis. *Gmel. sib.* 2. p. 100.

Carduus *sphærocephalus* cæruleus minor. *Bauh. pin.* 381.

Ritro floribus cæruleis. *Lob. ic.* 2. p. 8.

Crocodylium monspeliensium. *Dalech. hist.* 1476.

β. *Echinops* caule subunifloro, foliis duplicato-pinnatifidis, foliolis linearibus remotis. *Gmel. sib.* 2. p. 102. t. 46.

E. minor. *Bauh. hist.* 3. p. 72.

Habitat in Sibiria, Gallia, Italia collibus aridis. 24

Caulis albus, tomentosus. *Folia* supra glabra, subtus albo-tomentosa. *Calycis squamæ* ad apicem usque ciliatæ.

Annua pronuntiat *Gouan.* R.

4. **ECHINOPS** *strigosus*. E. capitulis fasciculatis, calycibus lateralibus sterilibus, foliis supra strigosis. *Læst hisp.* 159. *

Carduus tomentosus, capitulo majore. *Bauh.*

GENESIA. MONOGAMIA.

luisolia annua. *Herm. parr.* 224. s. 224.

Lob. ic. 2. p. 9.

nat. in hispania. ☉

Folia pinnatifida, latiuscula: laciniis lanceolatis, spina minima terminatis, lateribus inermibus, supra pilis rigidis strigosa, subtus tomentosa. Calyces squamis carinatis, superne, laeves, basi ciliati, non in capitulum globosum congesti, sed fasciculati: calycibus lateralibus s. inferioribus minoribus, sterilibus s. floscula destitutis.

MONOGAMIA.

1090. JASIONE.

Cal. communis 10. phyllus. *Cor.* 5. petala, regularia.

Caps. infera bilocularis.

1. *JASIONE montana*. Jasiona, *ged. gallob. herb. pal. sil.* *Fl. dan.* 1. 319. *Kniph. cent.* 12. n. 58. *Sabb. hort.* 1. 1. 76. *luthu. lugd. delph. burg.*

2. foliis lineari-lanceolaris obsolete serratis. *Fl. succ. paris.* *Rapunculus* foliis linearibus suba speris, spica planiuscula, petalis liberis. *Hall. helv. n.* 678.

Rapunculus scabiosae capitulo caeruleo. *Bauh. pin.* 92.

Rapuntium montanum capitatum leptophyllum. *Col. cepha.* 1. p. 226. s. 227.

Aphyllanthes. *Dalech. hist.* 804.

Habitas in Europæ collibus siccissimis. ☉

Floribus albis variat.

1091. LOBELIA.

Cal. 5. fidus. *Cor.* 1. petala, irregularis. *Caps.*

infera, 2. s. 3. locularis.

* *Foliis integerrimis.*

3. *LOBELIA Dortmanna*. L. foliis linearibus bilocularibus integerrimis, caule subaudo. *Fl. succ. Fl. dan.* 1. 39.

L. foliis bilocularibus subulatis. *Fl. lapp.*

SYNGENESIA. MONOGAMIA. 44

Dortmannia lacustris, floribus sparsis pendulis. *Rudb. ed. ups.* 1720. p. 97. t. 2.

Gladiolus stagnalis f. *lacustris*. *Rudb. elys.* 2. p. 202. f. 13.

Gladiolus stagnalis dortmanni. *Clus. cur.* 40. *Rai hist.* 1911.

Habitat in Europæ frigidissimæ lacubus & ripis. 24

13. *LOBELIA Cardinalis*. L. caule erecto, foliis lanceolatis serratis, spicâ terminali secunda. *Kniph. cens.* 4. n. 42. *Knorr. del.* 2. t. L. 2. *Mill. illustr.*

Rapuntium galeatum virginianum, coccineo flore majore. *Moris. hist.* 2. p. 466. f. 5. t. 5. f. 54.

Rapuntium maximum, coccineo spicato flore. *Hern. mex.* 879. t. 880.

Habitat in Virginia. 2L

14. *LOBELIA siphilitica*. L. caule erecto, foliis ovato-lanceolatis subserratis, calycum sinibus reflexis. *Mai. men.* 194. *Amœn acad.* 4. p. 527. *Gron. virg.* 134. *Kniph. cent.* 8. n. 60.

Rapunculus galeatus virginianus, flore violaceo majore. *Moris. hist.* 2. p. 466. f. 5. t. 5. f. 55.

Rapunculus americanus, flore dilute cæruleo. *Dolet. mem.* 297.

Trachelium americanum, flore cæruleo. *Rob. ic.*

Habitat in Virginie sylvis aridis, lutosis. 24

Caulis simplex, erectus, pedalis, angulis pilis rigidulis & foliorum marginibus decurrentibus. Folia alterna, sessilia, latius lanceolata, serrata, scabriuscula. Flores axillares, solitarii, brevissime pedunculati, cærulei. Calyx serrato-denticulatus: laciniis lanceolatis, sinibus reflexis, germeni regentibus (ut in *Campanulis Medio* &c.) Corolla cærulea, angulata: laciniis subæqualibus carina ciliatis; palato; 2 gibbositatibus. *Mant.* 482.

18. *LOBELIA urens*. L. caule erectiusculo, foliis inferioribus subrotundis crenatis, superioribus lanceolatis serratis floribus racemosis.

L. caule erecto, foliis lanceolatis subdentatis, spica laxa longa terminali. *Læfl. it.* 167. *

L. foliis oblongo-ovatis, floribus laxè spicatis. *Gæst. stamp. Dalib. parif.*

Rapunculus galeatus blesensis f. *soloniensis*, flore violaceo minore. *Moris. hist.* 2. p. 407. f. 5. t. 5. f. 56.

Rapuntium urens soloniense. *Moris. bles.* 300. *Boec. sic.* 20. t. 11. *Rai. hist.* 746. *Monnier. obs.* 131.

Draba flore cæruleo galeato. *Bauh. prodr.* 33. *pin.* 110.

446 SYNGENESIA. MONOGAMIA.

Habitat in Gallia, Hispania. ☉

Caulis strictus, simplex, pedalis, nudiusculus, angulis scabris. Folia radicalia elliptica, glabra, obtusiuscula, subdenticulata; Caulina remotissima, 8. lanceolata, sessilia, basi quasi agglutinata, argute serrata, lævia, remota. Racemus terminalis, strictus: floribus erectis remotis. Calyx 5. fidus, æqualis, scaber laciniis subulatis, erectis. Corolla cærulea, scabra, palato maculis 2 pallidis: Labii inferioris laciniis lanceolatis dependentibus. Mant. 488.

* * * *Caule prostrato; foliis incis.*

20. *LOBELIA Laurentia. L. caule prostrato, foliis lanceolato-ovalibus crenatis, caule ramoso, pedunculis solitariis unifloris longissimis.*

Laurentia annua minima, flore cæruleo. Mich. gen. 18. & 14.

Rapunculus aquaticus repens, flore cæruleo inaperto.

Rai. suppl. 383. Bocc. mus. 35. t. 27.

β. *Rapuntium creticum minimum, bellidis folio, flore maculato. Tournef. cor. 9.*

Habitat in Italia, Creta. ☉

Bractea setacea in medio pedunculi.

1092. V I O L A.

Cal. 5. phyllus. Cor. 5. petala, irregularis, postice cornuta. Caps. supera, 3. valvis, 1. locularis.

* *Acaules.*

3. *VIOLA pinnata. V. acaulis, foliis pinnatifidis. Gmel. sib. 4. p. 101. t. 49. f. 4.*

V. acaulis, foliis multifidis obtusis. Hall. helu. n. 561.

V. martia inodora, folio instar Petroselini eleganter insecto.

Act. Nat. C. vol. 3. n. 3. R.

V. montana laciniato folio. Clus. hisp. 309. Hall. R.

V. acaulis, foliis palmato-multifidis & laciniatis. Misc. taur. 3. p. 181. t. 2. f. 2. Hall. R.

V. alpina, folio in plures partes dissecto. Bauh. pin. 199.

V. montana, folio multifido. Bauh. hist. 3. p. 544.

Habitat in Sibiria & Alpibus Europæ.

6. *VIOLA hirta. V. acaulis, foliis cordatis piloso-hispidis.*

Fl. suec. parif. Ingr. Hall. helu. n. 559. herb. Fl. dan. t. 618. delph. burg.

V. martia hirsuta inodora. Moris. hist. 2. p. 475. f. 5. t. 35. f. 4.

V. trachelii folio. Rai. hist. 1051.

Viola. Brunsf. herb. 1. p. 137. f. 3.

Habitat in Europæ frigidioribus nemoribus. 2f

Radix crassiuscula, erecta, eminens, cicatricibus nudis. Folia cordata, oblongiuscula, ut Petioli, subtus pilosa. Pedunculi radicales; dorso canaliculati supra bracteas. Petalum cornutum emarginatum.

7. *VIOLA palustris. V. acaulis, foliis reniformibus. Hall. helu. n. 560. Fl. suec. paris. ged. Flor. dan. t. 83. lithu. delph.*

V. foliis subrotundo-cordatis: pedunculis radicatis. Fl. lapp.

V. alpina minima Nummulariæ folio. Bocc. mus. 163. t. 127. Allion. t. 2 f. 2. Hall. R.

V. palustris rotundifolia glabra. Moris. hist. 2. p. 475. f. 5. t. 35. f. 5.

Habitat in Europæ frigidioribus paludibus. 2f

Cum V. canina male combinat Gmel. sib. 4. p. 98. R.

8. *VIOLA odorata. V. acaulis, foliis cordatis, stolonibus reptantibus. Hall. helu. n. 558. gallob. herb. pal. sil. Fl. dan. t. 309. Ludw. ed. t. 175. Blackw t. 55. Koenig. del. 1. t. V. Kniph. cent. 3. n. 97. Regn. bot. lugd. lithu. burg.*

V. acaulis, stolonibus teretibus reptantibus, pedunculis radicalibus. Fl. suec.

V. pedunculis radicalibus; foliis cordatis, stolonibus reptantibus. Scop. carn.

V. martia purpurea, flore simplici odoro. Bauh. pin. 119.

V. odorata. Renealm. spec. 141. t. 140.

V. nigra f. purpurea. Dod. pempt. 156. t. 1. 2.

β. *V. martia alba. Bauh. pin. 199.*

γ. *V. martia, multiplici flore. Bauh. pin. 199.*

Habitat in Europæ nemoribus. 2f

Radix repens, filiformis. Folia cordata, rotundata, nudiuscula, subtus pubescentia. Pedunculi radicales, minime canaliculati supra bracteas. Petalum cornutum emarginatum.

Huius prolem hybridam: Floribus majoribus, campanulatis; Petalis fere æqualibus conniventibus, omnibus basi calcaratis, Aquilegiæ flores, dentis petalis, referentibus, semel legis b. Leers l. c. Vera Violæ Peloria R.

* * *Caulescentes.*

9. *VIOLA canina. V. caule adultiore adscendente, foliis oblongo-cordatis. Gmel. sib. 4. p. 98. t. 49. f. 1. ingr. gallob. herb. pal. sil. lugd. lithu. burg.*

V. caulibus adscendentibus floriferis, foliis cordatis. Fl. suec. paris.

148 SYNGENESIA. MONOGAMIA

V. caule procumbente ramoso, foliis petiolatis cordatis
Hall. helu. n. 563.

V. pedunculis caulinis; stipulis ciliatis; foliis cordatis:
Scop. carn.

V. foliis cordatis oblongis, pedunculis subradicatis:
Fl. lapp.

V. martia inodora sylvestris. Bauh. pin. 199.

V. cærulea maritima inodora sylvatica, in acumine semini
ferens. Bauh. hist. 3. p. 543.

Habitat in Europæ apricis. 4

Caules teretes, hinc plani, laeves. Folia cordata glabra.
Stipulae dentato-ciliatae. Corolla cærulea cornu albo.

Floribus albis ludis. Varietatem singularem Violæ mirabili
accedentem describit Leers. l. c. Aliam varietatem foliis
mucronatis oblongis & strictioribus habet Thalius. hanc
p. 130. R.

10. VIOLA montana. V. caulibus erectis, foliis cordatis
oblongis. Fl. suec. Kniph. cent. 4. n. 99. lugd. delph.
lithu. burg.

V. caule lato, erecto, foliis ellipticis, crenatis, stipulis
semipinnatis. Hall. helu. n. 567.

V. sylvestris longifolia. Tabern. 305. R.

V. flore cæruleo, longifolia. Riv. t. 119.

V. foliis ovato-lanceolatis, caule erecto, stipulis dentis:
Roy. lugdb. 430. Dalib. paris. 269.

V. martia arborescens purpurea. Bauh. pin. 199.

V. erecta, flore cæruleo & albo. Moris. hist. 2. p. 475:
f. 5. t. 7. f. 7.

V. arborescens. Cam. epit. 911.

Habitat in Alpibus Lapponiæ, Austriæ, Germaniæ;
Baldo. 24

11. VIOLA cenista. V. caulibus filiformibus indivisis
prorepentibus, foliis ovatis petiolatis integerrimis glabris;
stipulis indivisis pedunculo solitario.

V. multicaulis, foliis ovatis, petiolatis, integerrimis. Hall.
helu. n. 565.

V. foliis ovalibus integerrimis uniformibus, pedunculis
caulinis All. pedem. 14. t. 3. f. 4.

Habitat in Cenesii alpinis Ronche dictis. 24

Caules filiformes, perennantes, indivisi, sub terra prore-
pentes, summitate adscendentes, simplicissimi. Folia ovata,
minima, petiolata, glabra, ad apicem caulis conferta. Pe-
dunculus solitarius, longus. Flos magnus, cæruleus: calcar
petalis brevior.

13. *VIOLA mirabilis*. V. caule triquetro, foliis reniformi-
cordatis, floribus caulinis apetalis. *Jacq. austr.* t. 19.

V. floribus radicalibus corollatis abortientibus, caulinis
apetalis seminiferis. *Fl. suec.*

V. montana latifolia, flores e radice, semina in cacumine
ferens. *Dill. elth.* 408. t. 303. f. 390.

Habitat in Germaniæ, Sueciæ, nemoribus. 27

Corollæ albidæ petalo impari venis violaceis.

14. *VIOLA biflora*. V. caule bifloro, foliis reniformibus
serratis. *fil. Fl. dan.* t. 46. *delph.*

V. foliis reniformibus, pedunculis caulinis. *Fl. suec.*

V. caule debili, paucifloro, foliis petiolatis reniformibus,
obtusè dentatis. *Hall. helu. n.* 564.

V. pedunculis caulinis; stipulis lanceolatis; foliis renifor-
mibus. *Scop. carn.*

V. flore luteo. *Riu.* t. 119. *Hall. R.*

V. alpina rotundifolia lutea. *Bauh. pin.* 199.

V. rotundifolia montana major. *Pluk. alm.* 388. t. 234. f. 1.

V. alpina rotundifolia minor lutea. *Pluk. alm.* 388.
t. 233. f. 7.

*Habitat in Alpibus Lapponiæ, Austriæ, Helvetiæ, An-
gliæ.* 28

* * * *Stipulis pinnatifidis; Stigmate urceolato.*

16. *VIOLA tricolor*. V. caule triquetro diffuso, foliis ob-
longis incisis, stipulis pinnatifidis. *lithu. lugd. delph. burg.*

V. caule triquetro diffuso, foliis oblongis incisis, stipulis
dentatis. *Fl. suec. paris. gallob. herb. pal. fil.*

V. pedunculis caulinis angulatis, stipulis oblongis pinnato-
dentatis: foliis ovato-oblongis crenatis. *Hort.* 527.

V. pedunculis caulinis triquetris: stipulis laciniatis; foliis
oblongis crenatis. *Scop. carn.*

V. arvensis. *Murray. prodr.* 73.

V. caule diffuso, ramoso, foliis ovatis, dentatis, flore
calyce paulo majori. *Hall. helu. n.* 569.

V. bicolor arvensis. *Bauh. pin.* 200.

Trinitatis herba. *Fuchf. hist.* 803.

Jacea altera. *Cam. epit.* 913.

β. *Viola tricolor hortensis repens*, *Bauh. pin.* 199. *Gmel.*
l. c. Pollich. Leers.

V. (*tricolor*) caule anguloso, diffuso, foliis ovatis, dentatis,
flore calyce duplo longiore. *Hall. helu. n.* 568. *Murray.*
prodr. 73. *Flor. dan.* t. 623. *Kniph. cent.* 7. n. 99. *Knorr,*
del. 1. t. D. *Blackw.* t. 44.

Linnaei spec. plant. Europ. Pars II.

Gg

450 SYNGENESIA. MONOGAMIA.

V. tricolor. Renealm. spec. 144. t. 140. Riv. t. 120.

Jacea f. Flos trinitatis.. Cam. epis. 912.

Habitat in Europæ cultis. ☉

17. *VIOLA grandiflora. V. caule triquetro simplici, foliis oblongiusculis, stipulis pinnatifidis delph.*

V. caule erecto, paucifloro, foliis imis subrotundis, caulinis ellipticis, stipulis semipinnatis maximis. Hall. helv. n. 566. t. 17. an? videtur cum sequenti confundere. R.

V. caule triquetro erecto, foliis cordatis crenatis, stipulis dentatis, radice repente. Hudj. angl.

V. montana tricolor odoratissima. Bauh. pin. 109.

V. montana lutea grandiflora. Bauh. pin. 200. Barr. ic. 691. 692.

Habitat in Alpibus helveticis, Pyrenaicis. 24

Caulis altus, ramosus, foliosus, stipulatus omnino V. tricoloris, sed Flos maximus, flavus: Petalis ovatis, longitudine foliorum; Calcar calyce duplo longiore, sed petalis brevior.

Separanda a V. calcarata, cui tamen nimis affinis; ambo enim sunt progenies V. tricoloris.

18. *VIOLA calcarata. V. caule abbreviato, foliis subrotundis, stipulis pinnatifidis, nectariis calyce longioribus delph.*

V. alpina purpurea, exiguis foliis. Bauh. pin. 199.

V. montana cærulea grandiflora. Tournef. inst. 420.

Melanium montanum. Dalech. hist. 1204.

Habitat in Pyrenæis, Helveticis alpibus.

Quæ demum natura ex herba V. tricolore, adjectis floribus in hac planta.

19. *VIOLA cornuta. V. caule elongato, foliis oblongo-ovatis, stipulis pinnatifidis, nectariis subulatis corollæ longioribus.*

V. foliis cordato-lanceolatis, serratis, stipulis lanceolatis, acute dentatis. Hall. helv. n. 570.

V. pyrenaica longius caudata, reucræ folio. Tournef. inst. 421.

V. pyrenaica, folio reucræ, serotina. Rai suppl. 310.

Habitat in Pyrenæis.

Differt a V. calcarata quod huic Caulis longus. erectus, illi vero subcaulis; huic Petala oblonga foliis minora; illi rotunda foliis majora; huic folia oblongo-ovata; illi rotundato-ovata; cæterum nimis affinis.

20. *VIOLA arborescens. V. caule fruticoso, foliis lanceolatis integerrimis.*

V. hispanica fruticosa longifolia. *Tournesf. inst.* 421.

V. hispanica fruticans. *Barr. ic.* 368.

V. montana pumila angustifolia, flore inodoro: *Bauh. pin.* 199.

Habitat in Hispania. 7

1093. IMPATIENS.

Cal. 2. phyllus. *Cor.* 5. petala; irregularis, nectaris cucullato. *Caps.* supera. 3 valvis.

* *Pedunculis unifloris.*

5. IMPATIENS *Balsamina*. I. pedunculis unifloris aggregatis, foliis lanceolatis: superioribus alternis, nectaris flore brevioribus. *Blackw. t.* 583. *Kniph. cent.* 12. n. 61. *Knorr. del. hort.* 2. t. B. 3. flore pleno. *Regni bot.*

I. pedunculis confertis unifloris: *Hort. cliff.*

Balsamina femina. *Bauh. pin.* 306.

Balsamina. *Dod. pempt.* 671.

Habitat in India. ☉

* * *Pedunculis multifloris.*

7. IMPATIENS *Nolitangere*. I. pedunculis multifloris solitariis, foliis ovatis, geniculis caulinis tumentibus. *Fl. suec. parif. ged. carn. sub.* *Balsamina herb. pal. sil. dan.* t. 588. *délph. lithu. lugd. burg.*

I. pedunculis solitariis multifloris. *Hort. cliff. Roy. lugdb.* caule angulato. *Gort. gelr.*

I. caule geniculato, pedunculis ramosis, foliis petiolatis, ovatis, crenatis. *Hall. helu. n.* 557.

Balsamita lutea polonica. *Barr. ic.* 1297. R.

Balsamina lutea f. *Noli me tangere.* *Bauh. pin.* 306.

Noli me tangere. *Col. ecphr.* I. p. 149. t. 150.

Habitat in Europæ, Canadae nemoribus. 24

CLASSIS XX.

GYNANDRIA.
DIANDRIA.

1094. ORCHIS. *Nectarium corniculatum.*
 1095. SATYRIUM. *Nectarium scotiforme.*
 1096. OPHRYS. *Nectarium subcarinatum.*
 1097. SERAPIAS. *Nectarium ovatum, subtus gibbum.*
 1100. CYPRIPEDIUM. *Nectarium inflato-ventricosum.*

TRIANDRIA.

1103. SISYRINCHIUM. 1-gyna. Cal. o. Cor. 6. petala
 plana. Stigmata 3. Caps. 3.
 locularis, infera.

PENTANDRIA.

1110. PASSIFLORA. 3. gyna. Cal. 5. partitis. Cor.
 5. petala. Bacca peltata.

HEXANDRIA.

1111. ARISTOLOCHIA. 6. gyna. Cal. o. Cor. 1. petala
 Caps. 6. locularis.

POLYANDRIA.

1121. CALLA. Spatha. Cal. o. Cor. 6. Stami
 mixta pistillis.
 1119. ARUM. Spatha. Cal. o. Cor. o. Stami
 supra pistilla.
 1123. ZOSTERA. Folium. Cal. o. Cor. o. Sem.
 alterna, nuda.

GYNANDRIA.

DIANDRIA.

1094. ORCHIS.

Nectarium corniforme pone florem.

* * *Bulbis indivisis.*

8. ORCHIS *bifolia*. O. bulbis indivisis, nectarii labio lanceolato integerrimo: cornu longissimo, petalis patentibus. *Fl. suec. paris. Flor. dan. t. 235. austr. noru. herb. pal. sil. blackw. t. 588. Ludw. eccl. t. 95. Kniph. cent. 6. n. 65. delph. lithu. burg. lugd.*

O. radicibus oblongis, labello lineari. *Hall. helu. n. 1285. t. 35.*

O. labio lanceolato, simplici; cornu setaceo, longissimo, subangulato. *Scop. carn.*

Satyrium flore albo. *Riu. t. 12.*

O. alba bifolia minor, calcar oblongo. *Bauh. pin. 83. Vaill. paris. 151. t. 30. f. 7. Segu. ver. t. 15. f. 10.*

Testiculi species V. *Cam. epit. 625.*

9. Orchis trifolia major & minor. *Bauh. pin. 83. Leers. l. c.*

γ. Orchis bifolia altera. *Bauh. pin. 82.*

Orchis bifolia latissima. *Bauh. pin. 82.*

Habitat in Europæ pascuis asperis. 2f.

Flos albus. Petala exteriora patentia. Labium, lanceolatum, integerrimum. Calcar subulatum longissimum.

10. ORCHIS *cucullata*. O. bulbis indivisis, nectarii labio trifido, petalis confluentibus, caule nudo.

O. radice rotunda, cucullo tridentato. *Gmel. sib. 1. p. 16. t. 3. f. 2.*

Habitat in Sibiria. 2f.

Folia radicalia ovata bina.

11. ORCHIS *globosa*. O. bulbis indivisis, nectarii labio resupinato trifido: media emarginata; labio brevi, petalis apice subulatis. *Jacq. t. 265. ca. barb. delph.*

434 GYNANDRIA. DIANDRIA.

O. radicibus subrotundis, spica densissima, petalis exterioribus aristatis. *Hall. helu. n. 1272. t. 27. epust. 226.*

O. Halleri. *Crantz. austr.*

O. flore globoso. *Bauh. pin. 81.*

Habitat in Helvetia, Austria, Carniola, Germania.

Corolla resupinata. Labium trifidum, media emarginata. Bulbi fusiformes. Spica densissima brevis. Flores resupinati, purpurascens. Calcar germinis brevius.

12. *ORCHIS pyramidalis.* *O.* bulbis indivisis, nectarii labio bicorni trifido æquali integerrimo: cornu longo, petalis sublanceolatis. *Fl. suec. *, Jacq. austr. t. 266. delph. burg. lugd.*

O. radicibus subrotundis, spica densa, calcare longissima, labello bicorni, æqualiter tripartito. *Hall. helu. n. 1286. t. 35.*

O. labio trifido, auriculato; cornu setaceo longissimo sered. *Scop. carn.*

O. bulbis indivisis, nectarii labio trifido antice bidentato: cornu longo, petalis acuminatis. *Ad. upf. 1740.*

O. radicibus subrotundis, bracteis flore brevioribus, nectarii labio trifido, seta longissima. *Hort. cliff.*

O. purpurea, spica congesta pyramidalis. *Raf. 3. p. 377. t. 18. Segu. ver. t. 15. f. 11.*

O. flore conglomerato. *Riz. t. 14.*

Cynosorchis militaris montana, spica rubente conglomerata. *Bauh. pin. 81. prodr. 28.*

β. *Cynosorchis latifolia*, hians cuculo, altera. *Bauh. pin. 81. Crantz. l. c.*

γ. *Cynosorchis latifolia*, spica compacta. *Bauh. pin. 81.*

Habitat in Helvetiæ, Belgii, Angliæ, Galliarum arenis, cretaceis.

Flos purpureus. Petala ovato-lanceolata; lateralia reflexa. Labium bicornis, trifidum, æquale, integerrimum. Calcar longissimum. Varietatem albo flore habet *Mappus.*

13. *ORCHIS coriophora.* *O.* bulbis indivisis, nectarii labio trifido reflexo crenato: cornu brevi, petalis conniventibus. *paris. Jacq. austr. t. 1. 2. carn. pal. fl. ind. Fl. dan. t. 224. delph. lugd. burg.*

O. bulbis indivisis, labello trifido punctato: lacinula media longiore, cornu germinibus brevior. *Sauv. monsp.*

O. radicibus subrotundis, galea connivente, labello trifido reflexo. *Hall. helu. n. 1284. t. 34.*

O. cimicifera. *Crantz. austr.*

O. flore parvo viridi purpureo. *Riu. hex. t. 20.*

Orchis odore hirci minor. *Bauh. pin. 82. Vaill. paris. 149. t. 31. f. 30. 31. 32. Hall. it. herc. 7.*

Tragorchis minor, flore fuliginoso. *Rai. hist. 1213.*

Tragorchis minor & verior gemmæ. *Lob. ic. 177.*

β. Orchis odore hirci minor, spica purpurascente. *Bauh. pin. 82.*

Habitat in Europæ australioris, Orientis pascuis. 24

Flores viridi-purpurascens, foetidi. Petala conniventia. Labium reflexum, trifidum, æquale, serratum, acutum: media integerrima. Calcar germine brevius.

15. ORCHIS Morio. O. bulbis indivisis, nectarii labio quadrifido crenulato: cornu obtuso adscendente, petalis obtusis conniventibus. *delph. lugd. burg. lithu.*

O. bulbis indivisis, nectarii labio quadrifido crenulato: cornu obtuso, petalis omnibus conniventibus. *Fl. suec. paris. austr. pal. tub. sil. Fl. dan. t. 253. Riu. hex. t. 19.*

O. radicibus subrotundis, galeæ petalis lineatis, labello trifido, crenato, medio segmento emarginato. *Hall. helu. n. 1282. t. 33.*

O. labio quadrifido, crenato, cornu obtuso, rugoso, germinibus brevior. *Scop. carn.*

O. radicibus subrotundis, nectarii labio quadrifido æquali crenulato: cornu obtuso. *Roy. lugdb. 25.*

O. morio femina. *Bauh. pin. 82. Vaill. paris. t. 31. f. 13. 14. Segu. ver. t. 15. f. 7.*

Triorchis serapias mas. *Fuchf. hist. 559.*

Habitat in Europæ sylvis asperis. 24

Flores purpurei, pauciores. Petala omnia obtusa, lineata, conniventia. Labium reflexum, lateribus crenulatum, 3-fidum media emarginata. Calcar obtusum, adscendens.

16. ORCHIS mascula. O. bulbis indivisis, nectarii labio quadrilobo crenulato: cornu obtuso, petalis dorsalibus reflexis. *Fl. suec. austr. pal. Flor. dan. t. 457. delph. lugd. burg. lithu.*

O. radicibus subrotundis, petalis lateralibus reflexis, labello trifido, segmento medio longiori bifido. *Hall. helu. n. 1283. t. 33.*

O. labio quadrifido crenato; cornu obtuso lævi, germinum longitudine. *Scop. carn.*

Palmata major. *Riu. hex. t. 10. Segu. ver. t. 15. f. 6.*

Orchis morio mas, foliis (non) maculatis. *Bauh. pin. 81. Vaill. paris. 151. t. 31. f. 12. It. gottl. 171. **

416 GYNANDRIA. DIANDRIA.

Tessiculus IV. Cam. epit. 624.

Orchis foliis sessilibus non maculatis. *Bauh. pin.* 82.

Satyrium mas. *Blackw. t.* 53.

Habitas in Europæ lucis. 24

Flores purpurei, plurimi. Petala exteriora acutiuscula, reflexa. Labium crenulatum, 3. fidum: media bifida, lateralibus longiore. Calcar obtusum, submarginatum.

17. ORCHIS *astulata*. O. bulbis indivisis, nectarii labio quadrifido punctis scabro: cornu obtuso, petalis distinctis. *Fl. succ. parif. tab. pal. Flor. dan. t.* 103. *delph.*

O. radicibus subrotundis, labello quadrifido, calcare brevissimo. *Hall. Helv. n.* 1273. *t.* 28.

O. labio quadrifido, maculato; cornu geminibus & bractea brevior. *Scop. carn.*

O. amœna. *Crantz. austr.*

O. militaris pratensis humilior. *T. Vail. parif. t.* 31. *f.* 35. 36. *Segu. ver. t.* 15. *f.* 4.

O. militaris minor, minima. *Rin. hex. t.* 17.

O. pannonica IV. *Clus. hist. 1. p.* 268.

Cynorchis militaris pratensis humilior. *Bauh. pin.* 81.

Habitas in Europæ temperatae pratis. 24

Flores Spica albido-rubra apice purpurascens. Petala distincta, conniventia. Labium trifidum: media biloba productione; punctatum punctis scabris. Calcar breve, obtusiusculum.

18. ORCHIS *militaris*. O. bulbis indivisis, nectarii labio quinquefido punctis scabro: cornu obtuso, petalis conniventibus. *Fl. succ. parif. tab. norv. pal. sil. austr. delph. lith. lugd. burg.*

O. radicibus subrotundis, labello quadrifido, laciniis anterioribus latiusculis, posterioribus remotis, fere convergentibus. *Gmel. fib. 1. p.* 14. *R.*

O. labio quinquefido, maculato; bractea colorata, cornu & geminibus brevior. *Scop. carn.*

O. radicibus subrotundis, spica conica, labello quadrifido, brachioli & pectore perangustis. *Hall. Helv. n.* 1277. *t.* 28.

O. (*Rivini*) bulbis subrotundis, spica conica, petalis conniventibus, nectarii labio trifido, lacina media productione bifida. *Gouan. illustr.*

Orchis galea & alis cinereis. *Bauh. hist. 2. p.* 755.

O. mascula. *Rin. hex. t.* 15.

O. galea & alis cinereis. *Bauh. hist. 2. p.* 755.

Orchis mas latifolia. Fuchs. hist. 554.

Cynosorchis latifolia, hiantē cucullo, major. Bauh. pin. 80.

β. Orchis magna, latis foliis, galea fusca vel nigricante. I. B. Rai. angl. 3. p. 378. t. 19. f. 2 Pollich. l. c.

O. radicibus subrotundis, spica longa, labello quadrifido, brachiolis angustis, crurculis latis, serratis. Hall. helu. n. 1276. t. 31.

O. fusca. Jacq. austr. t. 176.

γ. Orchis latifolia, hiantē cucullo minor. F. Vaill. paris. t. 31. f. 22. 23. 24.

Cynosorchis latifolia, hiantē cucullo, minor. Bauh. pin. 81.

δ. Orchis militaris major. Tournef. inst. 438. t. 247. Vaill. paris. 148. t. 31. f. 21. 27. 28. Segu. ver. 2. p. 122. t. 11. f. 2. Riu. hex. t. 16. Pollich. l. c.

O. purpurea. Hudf. angl.

Cynorchis militaris major. Bauh. pin. 81.

ε. Orchis flore simiam referens. Bauh. pin. 82. Pollich. l. c.

O. Zoophora cercopithecum exprimens oreades. Col. ecphr. 1. p. 320.

Habitat in Europæ temperatæ pratis. 2.

Flores spicati, purpureo-canescentes. Petala connata Labium trifidum: media biloba productiore punctis scabris. Calcar breve obtusiusculum.

19. ORCHIS papilionacea. *O. bulbis indivisis, nectarii labio indiviso crenato emarginato ampliato: cornu subulato, petalis conniventibus. Scop. carn.*

O. papilionem expansum referens. Bauh. pin. 83.

Habitat in Hispania Alstræmer, in Carniolia calidiore. Scop.

Statura O. militaris & flos fere ejusdem, sed Labium maximum, magnitudine totius unguis pollicis, latius quam longum, retusum s. emarginatum, indivisum, denticulatum, uti flos purpurascens.

21. ORCHIS pallens. *O. bulbis indivisis, nectarii labio trifido integerrimo, cornu obtuso mediocri, petalis patentibus. Mant. 292. Jacq. austr. 1. t. 45.*

O. radicibus subrotundis, petalis galeæ linearis, labello trifido, integerrimo. Hall. hellu. n. 1281. t. 30.

O. panonica VII. Clus. hist. 1 p. 267.

O. præcox alba barba luteola. Rupp. gen. 1. p.

*O. foetida sylvatica præcox, flore albo barba luteola. Rupp. ien. 2. p. 297. **

O. bulbosa, floribus flavescentibus. Segu. suppl. 247. t. 8. f. 3.

458 GYNANDRIA. DIANDRIA.

Habitat in Europæ sylvestris. 24

Statura *O. morionis*, facies *O. sambucinae*, sed Bulbi non palmati, Flos flavus. Labium saturatius flavum, immaculatum, trifidum, integerrimum: media brevior, emarginata: cornu mediocre, subemarginatum, adscendens.

* * *Bulbis palmatis.*

22. *ORCHIS latifolia.* *O.* bulbis subpalmatis rectis, nectararii cornu conico: labio trilobo lateribus reflexo, bracteis flore longioribus. *succ. par. austr. ged. tub. pal. fil. Fl. dan. t. 266. Kniph. cent. 7. n. 65. Knorr. del. hort. 2. t. O. 4. f. 1. lugd. lith. delph. burg.*

O. radicibus palmatis, caule fistuloso, bracteis maximis, labello trifido serrato, medio segmento obtuso. *Hall. helv. n. 1279. t. 32.*

O. radicibus palmatis, bracteis flore longioribus, nectararii labio trifido, cornu germinibus brevior. *Hort. cliff.*

O. palmata pratensis latifolia, longis calcaribus. *Bauh. pin. 85. Vaill. paris. t. 31. f. 1. 2. 3. 4. 5.*

O. palmata pratensis maculata. *Bauh. pin. 85.*

O. palmata palustris tota rubra. *Ei. ib. 86.*

O. palmata palustris altera & tertia. *Rin. herap. t. 19.*

Satyrium femina, Blackw. t. 405.

β. *Orchis palmata palustris latifolia. Bauh. pin. 85.*

γ. *O. palmata montana altera. Ei. ib.*

δ. *O. palmata palustris maculata. Ei. ib.*

Habitat in Europæ pratis. 24

Flores spicati. Petala lateralia reflexa. Labium lateribus reflexum, serratum, trilobum. Calcar conicum. Bracteis flore longioribus; Foliis immaculatis; & Caule fistuloso ab *O. maculata* evidenter differt.

23. *ORCHIS incarnata.* *O.* bulbis palmatis, nectararii cornu conico: labio obsolete trilobo serrato, petalis dorsalibus reflexis. *Fl. succ. 2. delph.*

O. radicibus palmatis, bracteis coloratis, labello circumferrato, trilobato, lobo medio emarginato. *Hall. helv. n. 1280.*

O. palmata lutea, floris labio maculato. *Segu. ver. 3. p. 349. t. 8. f. 5.*

Habitat in Europa. 24

Varietas forte sequentis; cum qua conjungit *Hallerus.*

24. *ORCHIS sambucina.* *O.* bulbis subpalmatis rectis, nectararii cornu conico, labio ovato subtrilobo, bracteis longitudine florum. *Fl. succ. * sub. noru. Jacq. austr. t. 108. pal. fil. delph.*

GYNANDRIA. DIANDRIA. 459

Orchis, Hall. n. 1280. var. β . cum præcedente conjungit.

O. palmata lutea; labio floris maculato. Segu. ver. 249. t. 8. f. 5.

O. palmata, sambuci odore. Bauh. pin. 86. Rudb. clys. 2. p. 213. f. 9.

O. pannonica VIII. Clus. hist. 1. p. 269.

Habitat in Europa. 24

Flores spicati, rariores, lutei, Petala lateraliter reflexa, Labium lateribus reflexum, serratum, trilobum: media emarginata punctis purpureis. Calcar emarginatum, mediocre.

25. *ORCHIS maculata*. *O. bulbis palmatis patentibus, nectarium cornu germinibus brevioribus: labio plano: petalis dorsalibus patulis.* suec. parisi. austr. tub. sil. Regn. bot. delph. lithu. burg.

O. radicibus palmatis, caule solido, labello trifido, serrato, medio segmento acuminato. Hall. helv. n. 1278. t. 32.

O. labio trifido maculato; cornu conico; germinibus brevioribus. Scop. carn.

Palmata maculata, non maculata & angustifolia maculata, Riu. hexap. t. 8. & 11.

Orchis palmata montana maculata. Bauh. pin. 85.

O. palmata pratensis maculata. Bauh. pin. 86. Vaill. parisi. t. 31. f. 9, 10.

Satyrium basilicum femina. Dod. pempt. 240.

Habitat in Europæ pratis succulentis. 24

Petala 3 exteriora erecta; 2 interiora conniventia. Labium planum, trilobum, crenatum: intermedia angustissima integerrima. Bractæe non flore longiores. Caulis solidus. Folia maculata. Flore albo variat.

26. *ORCHIS odoratissima*. *O. bulbis palmatis, nectarium cornu recurvo brevioribus: labio trilobo, foliis linearibus.* norv. herb. Jacq. austr. t. 264. delph.

O. radicibus palmatis, labello obtuse trifido, concolore, calcare germine breviori. Hall. helv. n. 1274. t. 29.

O. conopsea varietas. Jacq. vind. 293. huc pertinet. Jacq. R.

O. bulbis palmatis, nectarium cornu recurvo; labio ovato acuto, foliis linearibus. Spec. pl. ed. 2. monsp.

Orchis palmata angustifolia minor odoratissima. Bauh. pin. 86. prodr. 32. t. 30. Rai. hist. 1225. Segu. ver. 3. p. 250. t. 8. f. 6.

Habitat in Italia, Gallia, Germania, rarissima in Suecia.

Flores purpurei, fragrantiores. Petala dorsalia expansa. Labium æquale, trifidum: media latiore acuta. Calcar breve.

460 GYNANDRIA. DIANDRIA.

27. ORCHIS *conopsea*. O. bulbis palmatis, nectarii cornu setaceo germinibus longiore: labio trifido; petalis duobus patentissimis. *Juec. * paris. * austr. carn. ged. tub. Fl. dan. t. 224. lithu. lugd. burg. delph.*

O. radicibus palmatis, calcare longissimo, labello unicolore, obtuse trifido. *Hall. helu. n. 1287. t. 29.*

O. angustifolia non maculata. *Riu. hex. t. 11.*

O. radicibus palmatis, bracteis longitudine floris, nectarii labio trifido, seta germinibus longiore. *Hort. cliff. 429.*

O. palmata minor, calcaribus oblongis. *Bauh. pin. 85. Vail. paris. t. 30. f. 88. Rudb. elys. 2. p. 212. f. 5.*

Satyrium basilicum mas. *Fuch. hist. 712.*

β. Orchis palmata angustifolia minor. *Bauh. pin. 85.*

γ. Orchis palmata pratensis maxima. *Bauh. pin. 85. Pollich. L. c. β.*

Habitat in Europæ pratis montosis. 24

Petala exteriora patentia. Labium crenulatum, æquale, trifidum: intermedia integerrima. Calcar setaceum, longissimum, (us in O. pyramidalis.) Similis O. odoratissima & pyramidalis.

* * * * Bulbis fasciculatis.

31. ORCHIS *hyperborea*. O. bulbis fasciculatis, nectarii cornu longitudine germinis: labio lineari integerrimo truncato.

O. bulbis fibrosis, calcare germine brevior, labio tripartito: laciniis æqualibus integerrimis. *Oed. dan. 333.*

Habitat in Islandia. Koenig.

Caulis palmaris cum spica. Folia alterna, lanceolata, longitudine caulis: Spica ovata: Bracteis lineari-lanceolatis, longitudine florum. Corollæ viridi-lutescentes: Petalum supremum latius, ovatum; lateralia 2 superiora lanceolata, ad primum conniventia; lateralia 2 inferiora oblonga. Labium vix petalo longius.

32. ORCHIS *abortiva*. O. bulbis fasciculatis filiformibus, nectarii labio ovato integerrimo, caule aphylo, *paris. noru. carn. sub Serapiade. herb. Jacq. austr. t. 193. delph. burg.*

O. radicibus cylindricis, bracteis flore brevioribus, labello trifido obtuso, seta germinibus brevior. *Sauv. monsp.*

Epipactis aphylla calcare longo, labello ovato lanceolato. *Hall. helu. n. 1288. t. 36.*

Pseudo-Limodorum austriacum. *Clus. hist. 1. p. 270.*

Orchis abortiva violacea. *Bauh. pin. 86.*

Limodorum. Hall. opusc. 212.

*Habitat in Galliæ, Helvetiæ, Angliæ, Italiæ, Germaniæ
sylvis umbrosis. 24*

Caulis aphyllus, squamis vaginatus. Flores violacei.

1095. SATYRIUM.

*Nectarium scrotiforme f. inflato - didymum pone
florem.*

1. SATYRIUM hircinum. S. bulbis indivisis, foliis lanceo-
latis, nectarii labio trifido : intermedia lineari elongata
obliquapræmorsa. *ups. 1740. t. 18. paris. monsp. Jacq.
austr. t. 367. pal. lugd. delph. burg.*

*Orchis radicibus subrotundis, labello longissimo, tripartito,
plicato. Hall. helu. n. 1368. t. 25.*

Orchis hircina. Scop. carn. Crantz. austr. p. 484.

*Orchis barbata foetida. Bauh. hist. 2. p. 756. Vaill. paris.
t. 30. f. 6. Segu. ver. t. 15. f. 1. Riv. hex. t. 18.*

*Orchis barbata, odore hirci, brevior latioreque folio.
Bauh. pin. 20. Moris. hist. 3. p. 491. f. 12. t. 12. f. 9.*

Habitat in Galliæ, Cantii, Germaniæ &c. campestribus. 24

*Planta alta. Flores albi, hircosi. Labium trifidum :
intermedia longissima lineari obliqua præmorsa ; lateralibus
brevissimis subulatis integerrimis.*

2. SATYRIUM viride. S. bulbis palmatis, foliis oblongis
obtusis : nectarii labio lineari trifido ; intermedia obsoleta.
suéc. paris. pal. Flor. dan. t. 73. delph. burg.

*Orchis radicibus palmatis, galea connivente, labello tri-
sulco, calcare brevissimo. Hall. helu. n. 1269. t. 26.
Segu. ver. 2. t. 16. f. 18.*

Orchis viridis. Crantz. austr. Scop. carn.

Satyrium foliis oblongis caulinis. Fl. lapp.

Orchis palmata, flore viridi. Bauh. pin. 86. prodr. 30.

*Orchis palmata, flore galericulato dilute viridi. Læf. pruss.
192. t. 59.*

3. Orchis palmata batrachites. *Bauh. pin. 86. t. Vaill. paris.
153. t. 31. f. 6. 7. 8. Pollich. l. c.*

Habitat in Europæ frigidioris asperis. 24

*Flores virescenti - pallidi. Petala conniventia. Labium
trifidum : lateralibus angustioribus ; media minore.*

3. SATYRIUM nigrum. S. bulbis palmatis, foliis linea-
ribus, nectarii labio resupinnato indiviso. *delph.*

S: bulbis palmatis ; foliis linearibus, floribus resupinatis,

462 GYNANDRIA. DIANDRIA.

nectarii labio indiviso ovato acuminato. *Fl. suec. Jacq. austr. t. 368. Scop. carn. sub Orchide.*

Saryrium foliis linearibus. *Roy. prodr. 14.*

Orchis radicibus palmatis, spica densissima, flore resupinato, calcare brevissimo. *Hall. helv. n. 1271. t. 27.*

Orchis miniata. *Crantz. austr. n. 7.*

Orchis palmata angustifolia alpina; nigro flore. *Bauh. pin. Segner. ver. 133. t. 15. f. 17.*

Palma christi minor. *Cam. epit. 627.*

Palmata angustifolia, flore supinato, calcare brevissimo. *Hall. opusc. 228.*

Habitat in Alpibus Helveticis, Lapponicis. 24

Flores atropurpurei, odoratissimi, supinati. Petala patensia. Labium indivisum, ovato-lanceolatum, crenatum. Calcar brevissimum. Variat flore roseo.

5. SATYRIUM albidum. S. bulbis fasciculatis, foliis lanceolatis, nectarii labio trifido acuto: lacinia intermedia obtusa. *Fl. suec. Jacq. vind. Fl. dan. t. 115. Scop. carn. sub Orchide. delph. lugd.*

Saryrium scanense. *It. scan.*

Orchis radicibus confertis, teretibus, calcare brevissimo, labello trifido. *Hall. helv. n. 1270. t. 26.*

Orchis alpina. *Crantz. austr.*

Orchis palmata, palmis inversis, flore albo. *It. prodr. 182. R.*

Helleborine Broccenbergense. *Riu. hex. t. 3.*

Pseudo Orchis alpina, flore herbaceo. *Mich. gen. 301. 26.*

Limodorum montanum, flore albo dilute virescente. *chom. Ad. parif. 1705. p. 517.*

Habitat in Scania, Germaniae; Helvetiae, Averniae praeis sylvaticis. 24

Labium trifidum, acutum: media obtusiore.

5. SATYRIUM. *Epipogium*, S. bulbis compressis dentatis, caule vaginato, nectarii labio resupinato, indiviso. *Jacq. austr. t. 84. delph.*

6. bulbis compressis dentatis, caule vaginato, spiculis resupinatis, nectarii labio albo indiviso. *Spec. pl. t. 1. p. 1358.*

Epipogium. *Gmel. sib. 1. p. 12. t. 2. f. 2.*

Epipactis caule aphylo, flore supinato: labello ovato-lanceolato, calcare ovato turgido. *Hall. helv. n. 1289. Ad. bernens. 5. p. 309.*

Habitat in Sibiriae, Austriae, Helvetiae, Germaniae, Arvensibus umbratis. 24

GYNANDRIA. DIANDRIA. 463

Flores *resupinati*. Caulis *aphyllus*, *vaginatus*. Labium *indivisum*, *ovato-lanceolatum*.

7. SATYRIUM *repens*. S. bulbis fibrosis, foliis ovatis radicalibus, floribus secundis *suec. parif. norv. n. 321. t. 6. f. 1. Jacq. austr. t. 369. delph. burg.*

Epipactis foliis petiolatis, ovato-lanceolatis, floribus tetrapetalis hirsutis. *Hall. helu. n. 1295. t. 22. Ad. helu. 4. p. 114.*

Orchioides. *Trew. comm. noric. 1736. t. 6. f. 7.*

Satyrium foliis ovatis radicalibus, *Fl. lapp.*

Orchis minor, sticulis albis, s. radice repente. *Cam. hort. III. t. 35.*

Pseudo Orchis. *Bauh. pin. 84.*

Pyrola angustifolia polyanthos, radice geniculata. *Laf. pruss. 210. t. 63.*

Epipactis foliis ovatis radicalibus. *Gmel. sib. 1. p. 13.*

β. Orchis radice repente, foliis maculis nigris & albis adspersis. *Mentz. pug. t. 3. f. 4. 5.*

Habitat in Sueciæ, Angliæ, Sibiria, Helvetiæ sylvis, in pinetis Germaniæ. *Reich. fl. franc. n. 651. 2.*

Flores secundi. Petala tria, pubescentia, conglutinata. Labium lanceolatum s. ovato-oblongum. Calcaris loco carina cymbiformis. Radix repens. Folia ovata, petiolata, fusco tessellata.

1096. OPHRYS.

Nectarium subtus, subcarinatum.

1. OPHRYS *Nidus avis*. O. bulbis fibroso-fasciculatis, caule vaginato aphylo, nectarii labio bifido. *austr. sub Epipactide. Scop. carn. pal. sil. Fl. dan. t. 181. lithu. delph. burg. lugd.*

Neottia bulbis fasciculatis, nectarii labio bifido. *Fl. suec. parif.*

Epipactis aphylla, flore inermi, labello bicorni. *Hall. helu. n. 1290. t. 37.*

Orchis abortiva fusca. *Bauh. pin. 86.*

Nidus avis. *Lob. ic. 195. Riv. hex. t. 7.*

Habitat in Sueciæ, Germaniæ, Galliæ nemoribus.

2. OPHRYS *Corallorhiza*. O. bulbis ramosis flexuosis, caule vaginato aphylo, nectarii labio trifido. *Crantz. austr. sub Epipactide. Fl. dan. t. 451. lithu. delph.*

- 264 **GYNANDRIA. DIANDRIA.**
Neottia bulbis reticulatis, nectario labio bifido. *Fl. alp.*
Neottia radice reticulata. *Fl. lapp.*
Corallorhiza Hall. *helu.* n. 1301. t. 44.
Corallorhiza *Neottia*. *Scop. carn.*
Epactis aphylla radice coralloide. *Crantz. austr.* 464.
Orobanche radice coralloide. *Bauh. pin.* 88.
Orobanche sueonum radice coralloide, flore albo. *Roth. elys.* 2. p. 234. f. 16.
Orobanche spuria s. *Corallorhiza*. *Rupp. gen.* 284. t. 2.
Orobanche radice coralloide ruberrima. *Menz. pug.* t. 9. f. 11.
Habitas in Europæ desertis. 2.
 Stamina 4. f. in singulis loculis 2. *Hall.*
 Conf. *Orobanche virginiana*. *Pluk. phyt.* III. f. 13.
 3. **OPHRYS** *spiralis* O. bulbis aggregatis oblongis, caule subfolioso, floribus spirali-secundis, nectarii labio indiviso crenulato. *Paris. Oed. dan. Riv. hex.* t. 14. *Se-guier. ver.* t. 8. f. 9. *Vaill. paris. pal. carn. sub Serapiade delph. lugd. burg.*
Epipactis bulbis cylindricis, spica spirali, labello crenulato. *Hall. helu.* n. 1294. t. 38.
Triorchis alba odorata minor. *Bauh. pin.* 84.
Satyrium odoriferum. *Bruff. herb.* 1. p. 105.
Testiculus odoratus. *Lob. ic.* 186.
 : *Triorchis* s. *Tetrorchis* alba odorata major. *Bauh. pin.* 84. *Pollich. l. c.*
 7. *Orchistrum æstivum* palustre album odoratum. *Mich. gen.* 30. t. 26.
 8. *Epipactis* foliis plerisque ex lineari-lanceolatis. *Gmel. fib.* 2. p. 13. t. 3. f. 1.
Habitas in Sibiria, Germania, Gallia, Austria, Helvetia, Italia graminosis.
 Flores spica secunda spirali, intus albi. Petala exteriora tria conglutinata. Labium subrotundum, ciliato-crenulatum.
 9. **OPHRYS** *ovata*. O. bulbo fibroso, caule bifolio, foliis ovatis, nectarii labio bifido. *Scop. carn. Crantz. austr. sub Epipactide. rub. gallob. Gagneb. in Ad. helu.* 2. f. 56. t. 6. *pal. Fl. dan.* t. 137. *delph. burg. lithu. lugd.*
 O. foliis ovatis. *Fl. suec. fib. paris.*
Epipactis foliis binis ovatis, labello bifido. *Hall. helu.* t. 1291. t. 27.
Ophrys bifolia. *Bauh. pin.* 87.
Ophrys *Fuchs. hist.* 565.

Bifolium. Riv. hex. t. 7.

β. Ophrys trifolia. Bauh. pin. 87.

Habitat in Europæ subhumidis præis: 22

Folia caulina, duo, opposita, ovata. Labium lineare, semibifidum.

6. OPHRYS cordata. O. bulbo fibroso, caule bifolio, foliis cordatis. carn. norv. delph.

O. foliis cordatis. Fl. suec.

Epipactis foliis binis, cordatis, labello bifido, polluce bidentato. Hall. helv. n. 1292. t. 22. f. 4.

Ophrys minima. Bauh. pin. 87. prodr. 31.

Bifolium minimum. Bauh. hist. 3. p. 534.

β. Ophrys minima, floribus purpureo-croceis. Menz. pag. t. 9. f. 2.

Habitat in Europæ frigida sylvis humentibus.

Folia caulina, 2, opposita, cordata. Labellum bifidum basi utrinque dente notatum, quod non in O. ovata.

* * Bulbis rotundis.

7. OPHRYS lilifolia. O. bulbo subrotundo, scapo nudo, foliis lanceolatis, nectarii labio integro, petalis dorsalibus linearibus. Fl. suec. Ehret. ad. angl. 1764. n. 53. p. 81. t. 4.

O. scapo nudo, foliis radicalibus ovato-oblongis dimidii scapi longitudine. Gron. virg. I. p. 185.

O. radicibus ovatis tunicatis, scapo nudo. Hort. cliff. 429. (vix memini)

β. Equipactis foliis binis ovatis, bulbis pyriformibus foliosa vagina cinctis. Hall. ad. 4. p. 120.

Bifolium bulbosum. Dod. pempt. 292.

Habitat in Virginiae, Canadae, Sueciae, &c. paludibus. 24

Scapus nudus, triqueter. Folia radicalia, nervosa. Flores rubri, sed petala 3 angusta, linearia, viridia.

8. OPHRYS Læselii. O. bulbo subrotundo: scapo nudo trigono, nectario labello ovato lithu.

O. diphyllos bulbosa. Læs. pruss. 180. t. 58.

Habitat in Sueciae, Borussiae paludibus.

Radix bulbus adscendens ramo. Folia lanceolata, bina; longitudine fere scapi. Scapus nudus, superne triqueter. Petala angusta, reflexa. Nectarii labium ovatum; magnum respectu reliquorum. Flores in scapo 3 ad 8, nec ultra.

9. OPHRYS paludosa. O. bulbo subrotundo; scapo subnudo pentagono, foliis apice scabris, nectario labio integro. Flor. suec. pal. delph.

Linnaei spec. plant. Europ. Pars II.

4 Q 8

466 GYNANDRIA DIANDRIA

Orchis minima bulbosa. Rai. suppl. 587.

Orchis bifolia minor palustris. Pluk. alm. 270 t. 247. f. 21

Bifolium palustre. Rai. angl. 3. p. 385.

Habitat in Suecia, Germania, Russia paludibus arsis.

Bulbus ovals, incurvus, infra radicans. Scapus pterogonus, maxima parte nudus. Folia radicalia 3. f. 4 spatulata, alterna, apice praesertim in pagina interiori scabra, in exteriori breviora. Flores in racemo plurimi, virides, flavescens. Petala 2. lateralibus ovato-oblonga, reflexo-erecta; 2. interiora lateralibus linearia, recurva; 1 supremum rectum, fornicem flaminum formans. Labium nectarii lanceolato-ovatum, reflexum, integrum. Bulbi coherens deorsum quasi cernatim ramato interstincti. Differt ab *O. Læselii* floribus pluribus & labio floris angusto; ab *O. monophyllo* foliis non ovatis apice scabris, pedicellis non capillaceis.

10. *OPHRYS monophyllos.* *O.* bulbo rotundo, scapo nudo, folio ovato, nectarii labio integro. *delph.*

Epipactis folio unico, amplexicauli, spica proluxa multiflora. *Hall. helv. a. 1293. t. 36.*

Ophrys monophyllos bulbosa. *Læss. pruss. 180. t. 57.*

Monochrys ophioglossoides. *Mentz. pag. 6. 5. f. 1. 2.*

Pseudo Orchis monophyllos. *Clus. hist. 1. p. 269.*

Habitat in Borussia, Moldavia, Helvetiae paludibus sylvaticis. 24

Flores minui, virides ut precedentis.

11. *OPHRYS Monorchis.* *O.* bulbo globoso, scapo nudo, nectarii labio trifido cruciato. *pal Fl. dan. t. 102. lith.*

Herminium bulbo supra radiato, nect. labio trifido. *Fl. suec.*

Herminium radice globosa. *Fl. lapp.*

Orchis radice subrotunda, labello hastato. *Gmel. sib. 1. p. 11. t. 4. f. 1.*

Orchis bulbo unico, subrotundo, labello cruciformi. *Hall. helv. u. 1262. t. 22.*

Orchis Monorchis. *Crantz. austr. Scop. carn.*

Orchis odorata moschata s. *monorchis.* *Bauh. pin. 84.*

Orchis trifolia, floribus spicatis herbaceis. *Segur. var. 2. p. 131. t. 16.*

Orchis coleo unico s. *Monorchis* flosculis pallide viridibus. *Læss. pruss. 184. t. 61.*

Monorchis. *Mich. gen. 30. t. 26. Rupp. gen. 421. t. 2.*

g. *Orchis lutea*, hirsuto folio. *Bauh. pin. 84.*

γ. *Triorchis lutea*, folio glabra. *Bauh. pin. 84.*

δ. *Triorchis lutea altera.* *Bauh. pin. 84.*

Monorchis bifolia, flore pallide virente, praeulica. *Menz.* pug. t. 5. f. 5.

Habitat in Europæ pratis uliginosis.

Flores lutea-virentes, suaveolentes. Labium trifidum: lateribus transversis. Radix globosa, solitaria.

12. *OPHRYS alpina*. O. bulbis ovatis, scapo nudo, foliis subulatis, nectarii labio indiviso obtuso utriusque unidentato. *Fl. suec. Jacq. vind. t. 9. * carn. sub Orchide el. dan. t. 452. delph.*

Orchis radicibus subrotundis, labello ovato, utrinque denticulo notato. *Hall. helu. n. 1263. t. 22. f. 1.*

Orchys graminea. Crantz. austr.

Chamae Orchis alpina, folio gramineo. *Bauh. pin. 81. prodr. 29.*

Habitat in Alpibus Lapponiæ, Helvetiæ, &c.

Radix Bulbis duobus, subrotundis (nec uno infra radicato, ut præcedentes tres), supra radicatis. Folia radicalia sæpius 4, filiformia, subulata, pollicem longa. Caulis nudus, longitudine foliorum. Spica floribus 4. f. 5. alternis. Petala 5. conniventia: barba ovato-oblonga, integra, vix petalis longiore. Vaginae 2 obvestiunt basin plantæ & includunt folia cum scapo inferne.

14. *OPHRYS antropophora*. O. bulbis subrotundis, scapo folioso, nectarii labio lineari tripartito: medio elongato bifido. *Oed. fl. dan. t. 103. lugd. delph. burg.*

Neottia bulbis subrotundis, nectarii labio quadrifido. *paris.*

Orchis radicibus subrotundis, spica longa, flore inermi, labello perangusto quadrifido. *Hall. helu. n. 1264. t. 23.*

Orchis, flore nudi hominis effigiem repræsentans, femina. *Bauh. pin. 82. paris. 147. t. 31. f. 19. 20.*

Orchis antropophora oreades. Col. ecphr. I. p. 320.

Habitat in Italia, Lusitania, Gallia, Helvetia.

Flores spica oblonga. Petala conniventia. Labium lineare, tripartitum: media longiore bifida.

15. *OPHRYS insectifera*. O. bulbis subrotundis, scapo folioso nectarii labio subquinelobo. *monsp. pal. lugd. delph. burg. lithu.*

Cypripedium bulbis subrotundis, foliis oblongis caulinis. *Fl. suec.*

α. Ophrys insectifera, myodes. Gunn. norv. t. m. 2. t. 5. f. 1.

Orchis muscaria. Scop. carn.

468 GYNANDRIA. DIANDRIA

- Orchis radicibus subrotundis, labello sericeo bisulco, brachiolis brevibus subulatis. *Hall. helv. n. 1265. t. 24.*
- Orchis vesperam referens. *Riu. her. t. 13. 19.*
- Orchis muscæ corpus referens minor, galea & alis herbidis. *Bauh. pin. 83. Vaill. parif. 147. t. 31. f. 17. 18. Kniph. cent. 3. n. 68.*
- Orchis myodes prima, floribus muscam exprimens. *Lob. ic. β. Orchis muscam referens major. Bauh. pin. 83.*
- γ. Orchis muscam referens lutea. *Bauh. pin. 83.*
- δ. Orchis muscam cæruleam majorem representans. *Breyn. Morif. hist. 3. p. 494. f. 12. t. 13. f. 11.*
- ε. Orchis myodes lutea lusitanica. *Breyn. 175. Morif. hist. 3. p. 495. f. 12. t. 13. f. 15.*
- ζ. Orchis myodes fusca lusitanica. *Breyn. cent. 41.*
- η. Ophrys insectifera, arachnites. *Mill. dict. n. 7.*
- Orchis arachnites. *Scop. carn.*
- Orchis radicibus subrotundis, labello holosericeo emarginato, appendiculato. *Hall. helv. n. 1266. t. 24.*
- Orchis araneam referens. *Bauh. pin. 84. Tournef. inst. 434. t. 247. f. C. Vaill. parif. t. 30. f. 10. 11. 12. 13.*
- Orchis arachnitis. *Lob. ic. 135.*
- θ. Orchis fucum referens: colore rubiginoso. *Bauh. pin. 83. Vaill. parif. 146. t. 31. f. 15. 16.*
- ι. Orchis fucum referens major, foliis superioribus candidis & purpurascens. *Bauh. pin. 83. Vaill. parif. 146. t. 30. f. 10. 11. 12. 13.*
- Orchis fucum referens, flore subvirente. *Bauh. pin. 83.*
- κ. Orchis scarabæum referens. *Breyn. Morif. hist. 3. p. 494. f. 12. t. 13. f. 10.*
- λ. Orchis cercopithecum exprimens lusitanica. *Breyn. Morif. hist. 3. p. 494. f. 12. t. 13. f. 4.*
- Habitat in Europa temperiori. 2c*

Varietas Myodes α. & Arachnites β. fugitivis oculis inspicienti diversissima apparent, qui vero has confert cum congeneribus, omnesque varietates simul inspicit, primam originem ex una facile perspiciet, nec limites distinguendas obtinebit, quamvis constantes persistant.

1097. SERAPIAS

Nectarium ovatum, gibbum, labio ovato.

1. SERAPIAS *latifolia*. S. bulbis fibrosis, foliis ovatis amplexicaulibus, floribus pendulis. *pal. norv.* t. 5. f. 3. 6. *delph. burg. lithu.*

S. bulbis fibrosis, nectarii labio obtuso crenato petalis brevioribus. *Fl. suec. parif.*

S. bulbis fibrosis, nectarii labio obtuso crenato, petalis æquali, fructibus pendulis. *Huds. angl.* 341.

S. bulbis fibrosis, floribus secundis, labello planiusculo, petalis ovatis. *Scop. carn.*

S. bulbis fibrosis, caule multifloro, floribus spicatis, nectario labio petalis brevioribus. *Ger. prou.*

Epipactis foliis amplexicaulibus, ovato-lanceolatis, labello lanceolato. *Hall. helu. n.* 1297. t. 40. *Act. helu.* 4. p. 108.

Epipactis (*Helleborine*) cymba cartilaginea depressa, vernice splendente inuncta; flore minore. *Crantz. austr. fasc.* 6. p. 467. t. 1. f. 6. c.

Damasonium flore mixto. *Riv. hex.* t. 6.

Helleborines recentiorum genus II. *Clus. pann.* p. 275. & III. p. 276. *Hall. R.*

Epipactis s. Helleborine. *Cam. epit.* 187.

Helleborine latifolia montana. *Bauh. pin.* 186.

Helleborine altera, flore atro-rubente. *Bauh. pin.* 187.

Habitat in Europæ sylvis. 24

Folia amplexicaulia. Spica multiflora, longa. Petioli longi. Flores Fructusque penduli. Nectarium lineis nullis.

2. SERAPIAS *longifolia*. S. bulbis fibrosis, foliis ensiformibus sessilibus, floribus pendulis. *Oed. dan.* t. 267. *pal. sil. Kniph. cent.* 2. n. 82. *delph. burg.*

S. bulbis fibrosis, nectariis labio obtuso longitudine petalorum. *Ger. prou.*

S. (*palustris*) bulbis fibrosis, floribus sparsis, patulis: labello concavo, petalis longiore. *Scop. carn.* 1.

Epipactis foliis ensiformibus, caulinis, floribus pendulis, labello obtuso, oris plicatis. *Hall. helu. n.* 1296. t. 39. *Act. helu.* 4. p. III.

Epipactis (*palustris*) caule folioso multifloro, labello articulado vomeriformi, floribus pendulis. *Crantz. austr. fasc.* 6. p. 462. t. 1. f. 5.

170 GYNANDRIA. DIANDRIA.
Helleborine angustifolia palustris f. pratensis. *Bauh. pin.* 87.
Riu. hex. t. 3.

Habitat in Europæ asperis, sylvis. 24. In pratis udis. *Poll. Scop. Hall.*

Caulis præcedente altior. Spica longa multiflora. Flores fructusque penduli. Flores duplo majores petalis latioribus. Nectarium reliquis petalis cohærens: cymba lineis purpureis. *Hall.*

3. SERAPIAS grandiflora. S. bulbis fibrosis, foliis ensiformibus, floribus erectis, nectarii labio obtuso petalis brevioribus. *Huds. angl. Oed. dan.* 506. *pal. Scop. carn. delph. burg.*

S. bulbis fibrosis, caule paucifloro, floribus distantibus, nectarii labio petalis brevioribus. *Ger. prax.*

S. Damasonium. *Mill. dict.*

Epipactis caule paucifloro, lineis obtusi labelli levibus. *Hall. helu. n.* 1298. t. 41.

Epipactis (alba) caule folioso paucifloro, labello articulo obtuso labro reflexo. *Crantz. austr. p.* 460. *fasc.* 6. t. 1. f. 42.

Helleborine alba barba luteola. *Riu. hex.* t. 4.

Helleborine flore albo, Damasonium montanum latifolium. *Bauh. pin.* 187.

6. Helleborine montana angustifolia spicata. *Bauh. pin.* 187. *Pollich. l. c.*

Damasonium flore albo. *Riu. hex.* t. 5.

Habitat in Europæ sylvis. 24

Flores albi magni. Fructusque erectus. Labello brevius acutum lineis eminentibus. *Hall.*

4. SERAPIAS rubra. S. bulbis fibrosis, foliis ensiformibus, floribus erectis: nectarii labio acuto. *Oed. dan.* t. 345. *pal. fl. herb. * delph. burg.*

Epipactis caule paucifloro, lineis acuti labelli undulatis. *Hall. helu. n.* 1299. t. 42.

Epipactis (purpurea) caule folioso paucifloro, labello articulo-rostriformi acuto. *Crantz. austr.*

Damasonium flore roseo. *Riu. hex.* t. 6.

Helleborine montana angustifolia purpurea. *Bauh. pin.* 187.

Helleborine flore carneo. *Bauh. pin.* 187.

Helleborines genus 5. *Clus. pann.* 575.

Habitat in sylvis Europæ. 24

Flores purpurei magni Fructusque erecti. Labello articulo-rostriformi acuto. Fructusque erecti. Labello articulo-rostriformi acuto. *Hall.*

GYNANDRIA. DIANDRIA. 471

5. **SERAPIAS** *Lingua*. S. bulbis subrotundis, nectarii labio trifido acuminato glabro, petalis longiore. *car. sub* Orchide.

Orchis radicibus subrotundis, petalis galeæ conglutinatis, longissime rostratis, labello trilobo, ovato lanceolato. *Hall. helu n. 1267.*

Satyrium radicibus subrotundis, cucullo clauso monopetalo: labello ovato - lanceolato. *Sauv. monsp. **

Orchis montana italica, flore ferrugineo; lingua oblonga. *Bauh. pin. 84. prodr. 29.*

β. Orchis montana italica, lingua oblonga altera. *Bauh. pin. 84.*

Orchis macrophylla. *Col. ecphr. 2. p. 321. t. 322.*

Habitas in Italia, Lusitania, Gallia. 24

6. **SERAPIAS** *cordigera*. S. bulbis subrotundis, nectarii labio trifido acuminato maximo basi barbato.

Orchis montana italica lingua trifida. *Rudb. elys. 2. p. 204. f. 20.*

Habitas in Hispania, Italia, Oriente.

Similis S. Linguae, sed major. Nectarii labium subcordatum, basi media barbatum, lateribus utrinque auricula erecta obtusa.

1100. CYPRIPEDIUM.

Nectarium ventricosum, inflatum, cavum.

1. **CYPRIPEDIUM** *Calceolus*. C. radicibus fibrosis, foliis ovato lanceolatis caulinis. *Fl. suec. Hall. helu. n. 1300. sub Calceolo. ged. Kniph. cent. 10. n. 35. Knorr. del. hort. 1. t. M. 2. lithu. delph.*

C. foliis ovato lanceolatis. *Fl. lapp.*

Calceolus (*marianus*) foliis ovato-lanceolatis, petalis interioribus ligulatis. *Crantz. austr.*

Calceolus foliis ovato-lanceolatis. *Gmel. sib. 1. p. 2. t. 2.*

Calceolus marianus. *Dod. pempt. 180. f. 1. 2.*

Helleborine flore rotundo s. Calceolus. *Bauh. pin. 187.*

β. Helleborine Calceolus dicta mariana, caule folioso flore luteo minore. *Pluk. mant. 101. t. 418. f. 2.*

Helleborine virginiana s. Calceolus flore luteo majore. *Morif. hist. 3. p. 488. f. 12. t. 11. f. 15.*

γ. Calceolus marianus canadensis. *Corn. canad. 204.*

Helleborine flore majore purpureo. *Morif. hist. 3. p. 488. f. 12. t. 11. f. 17.*

δ. Calceolus minor, flore vario. *Amm. ruth. 133. t. 22.*

472 GYNANDRIA. TRIANDRIA.

Calceolus foliis ovatis binis. *Gmel. fib. 1. p. 5.*

Habitat in Europæ, Asiæ, Americæ septentrionalibus. 24
Flos tetrapetalus.

2. CYPRIPIEDUM *bulbosum*, C. bulbo subrotundo, folio subrotundo radicali.

C. folio subrotundo. *Fl. lapp. 319. t. 12. f. 5.*

Serapias scapo uniflora. Gmel. fib. 1. p. 7. t. 2. f. 1.

Orchis lapponensis monofolia. Rudb. elys. 2. p. 209. f. 10.

Habitat in Lapponia, Russia, Sibiria. 24

Flos pentapetalus.

TRIANDRIA.

1103. SISYRINCHIUM.

Monogyna. *Spatha* diphylla. *Petala* 6, *plum*
Capf. 3. *locularis*, *infera*.

1. SISYRINCHIUM *Bermudiana*. 8. foliis ~~angustifolius~~
enerviis. Syst. Veg. 682.

8. caule foliisque ancipitibus. *Spec. pl. 2.*

8. (*angustifolium*) foliis lineari-gladiolatis, pedunculis longioribus. *Mill. dict. n. 2.*

8. cæruleum parvum, gladiato caule, virginianum, *Pluk. alm. 368. t. 61. f. 1.*

Bermudiana graminea, flore minore cæruleo. *Dill. dict. 49. t. 41. f. 49.*

8. *Sisyrrinchium bermudense*, floribus parvis ex cæruleo & aureo mixtis. *Pluk. alm. 348. t. 61. f. 2.*

8. (*bermudianum*) foliis gladiolatis amplexicaulibus, pedunculis brevioribus. *Mill. dict. n. 1.*

Bermudiana iridis folia, radice fibrosa. *T. Dill. elb. 48. t. 41. f. 48.*

Habitat α in Virginia, β in Bermudis. 24

Planta α, in loca natali, Folia saepe & Scapum filiformiter, at si probe examinetur, samentum ancipitiū; hæc per gradus ad cultæ staturam adscendere observatur.

In var, α Caulis alatus; in β teres.

PENTANDRIA.

1108. AYENIA.

Monogyna. *Cal.* 5. phyllus. *Petala* connata in stellam unguibus longis. *Antheræ* 5. sub stella. *Caps.* 5. locularis.

1. AYENA *pusilla*. A. foliis cordatis glabris. L. *ed. stockh.* 1756. p. 23. t. 2. *

A. foliis ovatis acutis serratis, germine pedicellato, nectario plano stellato. *Læfl. is.* 200. *

Dayenia inermis, foliis oblongo cordatis marginibus dentatis, floribus axillaribus. *Mill. dict.* t. 118.

Urticæ folio anomala, flore pentaphyllo purpureo, fructu pentacocco muricato, *Sloan. jam.* 90. *hist.* 1. p. 209. t. 132. f. 2.

Habitat in Jamaica, Cumana, Peru, ☉
Flores varii abortiunt, an sexu ?

1110. PASSIFLORA.

Trigyna. *Cal.* 5. phyllus. *Nectarium* corona. *Bacca* pedicellata,

* *Foliis indivisis.*

1. PASSIFLORA *serratifolia*. P. foliis indivisis ovatis serratis. *Amæn. acad.* 1. p. 217. f. 1. * *Jacq. hort.* t. 10.

P. foliis ovato lanceolatis integris serratis. *Hort. cliff.*

Granadilla americana folio oblongo leviter serrato, petalis ex viridi rubescentibus. *Mart. cent.* 36. t. 36.

Habitat Surinami.

7. PASSIFLORA *laurifolia*. P. foliis indivisis ovatis integerrimis, petiolis biglandulosis, involucris dentatis. *Amæn. acad.* 1. p. 220, f. 6. * *Jacq. obs.* 1. p. 35. *hort.* t. 162.

P. foliis solitariis oblongis integerrimis, floribus solis involucre tripartito dentato. *Roy. lugdb.* 132.

474 GYNANDRIA. PENTANDRIA.

Parborea, laurinis foliis, americana. Pluk. alm. 282. t. 211. f. 3.
Clematis indica, fructu citriformi, foliis oblongis. Plum.
amer. 64. t. 80. Rai suppl. 341.

Marquias. Mer. surin. 21. t. 21.

Habitat in Surinamo. L

Bacca ovata sapidissima.

* * *Foliis bilobis.*

13. PASSIFLORA Vespertilio. *P. foliis bilobis basi rotundatis glandulosisque: lobis acutis divaricatis subtus punctatis. Amæn. acad. 1. p. 223. f. 11. **

P. foliis obverse lunatis: punctis duobus meliiferis sub basi. Hort. cliff.

Granadilla bicornis, flore candido: filamentis intortis. Dill. elth. 164. t. 137. f. 164.

Granadilla folio lunato, flore parvo albo, fructu succulento, ovato. Mart. cent. 52. t. 52.

Habitat in America. H

Folia subtus punctata, basi punctis 2 majoribus excavatis, setula nulla inter lobos. Petioli eglandulosi.

* * * *Foliis trilobis.*

15. PASSIFLORA rotundifolia. *P. foliis subtrilobis obtusis subrotundis subtus punctatis. Jacq. obs. 2. t. 26. f. 1.*

Granadilla folio hederaceo, flore albo, fructu globoso villoso. Plum. spec. 6. ic. 138. f. 1. Barr. obs. pref. 4. titul. f. 2.

Habitat in America australiori.

Folia subtus inter nervos longitudinali ordine punctata. Petioli eglandulosi.

16. PASSIFLORA punctata. *P. foliis subtrilobis oblongis subtus punctatis intermedio minore. Amæn. acad. 1. p. 224. f. 12. * Fabric. helmst. 349.*

P. foliis bilobis obtusis, nectario multifido. Roy. lugdb. 260.

Granadilla folio tricuspidi obruso & oculato. Few. peruv. 1. p. 718. t. 11. Hort. angl. 25.

Habitat in Peru. L

Flos albidus. Petala calyce breviora. Nectarium ambitus exterior radiis 30. luteis, longitudine fere petalorum, clavatis; Intermedius ordo capillaris brevior; Intimus plicatus, truncatus, etiam viridis. Receptaculum rectum operculo concavo striato albido. Folia fere biloba sunt. Petioli eglandulosi.

18. PASSIFLORA minima. *P. foliis trifidis integerrimis: lobis sublanceolatis; intermedio longiore. Jacq. hort. 2. 20. Amæn. acad. 1. p. 229. f. 18. Læsl. ii. 247. **

GYNANDRIA. PENTANDRIA. 491

P. f. Flos passionis curassavicus, folio glabro trilobato & angusto, flore flavescente omnium minimo. *Pluk. alm.* 282. t. 210. f. 3.

Granadilla folio glabro & angusto, flore virescente minimo. *Tournef. inst.* 240. R.

Habitat in Curassao.

Folia subpubescentia, ultra medium trifida. Petioli eglandulosi. Flores minuti. Caulis senior suberosus evadit.

19. PASSIFLORA suberosa. P. foliis trilobis subpeltatis, cortice suberoso. *Amæn. acad.* 1. p. 226. f. 14. *Mill. dict. n.* 5. *Jacq. hort.* t. 163.

P. foliis trilobis peltatis integris: glandulis duabus sub dorso basi convexis. *Roy. lugdb.* 261.

Clematis indica, folio hederaceo, major, fructu olivæformi. *Plum. amer.* 70. t. 84. *Rai suppl.* 339.

Passifloræ affinis, hederæ folio, americana. *Pluk. alm.* 202. t. 210. f. 4.

8. Clematis indica, folio angusto trifido; fructu olivæformi. *Plum. amer.* 70. t. 85.

Flos passionis minor, foliis in tres lacinias non serratas profundius divisis, flore luteo. *Sloan. Jam.* 104. *hist.* 1. p. 320.

Habitat in Dominica, Antillis. *H.*

Caulis perennis: Cortice suberoso, fisso, albo. Flores axillares, gemini, albi, apetal. Bacca violacea. Folia in tenellis adhuc plantis peltata; in adultis vero vix ac ne vix; semper nitida glaberrima & fere lucida. Petioli eglandulosi.

Plumieri icones multum ab hac planta abludunt. *Jacq. l. c.* R.

* * * * *Foliis multifidis.*

24. PASSIFLORA cærulea. P. foliis palmatis integerrimis. f. 20. *monsp. Knorr. del.* 1. t. P. *Kniph. cent.* 2. n. 50.

P. foliis palmatis quinquepartitis integerrimis, involucris cordatis triphyllis. *Hort. cliff.*

Granadilla polyphyllus, fructu ovato. *Tournef. inst.* 241.

Granadilla pentaphyllos, flore cæruleo magno. *Barth. lugdb.* 1. p. 81. *Duham. arb.* 1. p. 272. t. 107.

Granadilla indica scandens & sempervirens, radice repente, foliis palmatis &c. *Mich. flor.* *

Clematis quinquefolia americana f. Flos passionis. *Rob. ic.*

Flos passionis major pentaphyllus. *Sloan. jam.* 104. *hist.* 1. p. 229. *Rai. suppl.* 339.

276 GYNANDRIA. HEXANDRIA.

Habitat in Brasilia. $\overline{\eta}$

Folia quinqueloba, obtusa. Petioli glandulis 2. Stipulae innares.

HEXANDRIA.

YIII. ARISTOLOCHIA.

Hexagyna. Cal. 6. Cor. 1. perala, lingulata, integra.
Caps. 6. locularis, infera.

14. ARISTOLOCHIA bœtica. A. foliis cordatis acuminatis
caule volubili, pedunculis subternis petiolo longio-
ribus.

A. Clematis serpens. Bauh. pin. 307. Moris. hist. 3. p. 109
f. 12. t. 17. f. 6.

A. Clematis bœtica. Clus. hist. 2. p. 71.

Habitat in Hispania, Creta, arbores scandens.

Stipulae ovato-rhombeæ, mucronatæ.

15. ARISTOLOCHIA sempervirens. A. foliis cordato-
oblongis acuminatis, undatis, caule infirmo, floribus
solitariis. Nil. dict. n. 5. Sabb. hort. 2. t. 82. Kniph.
cent. 3. n. 14.

A. caulibus infirmis, foliis cordato-oblongis undulatis,
floribus recurvis solitariis pendulis. Hort. cliff. 432.
Sauv. monsp. 111.

Pistolochia cretica. Bauh. pin. 307. Moris. hist. 3. p. 110
f. 12. t. 17. f. 16.

Pistolochia altera. Clus. hist. 2. p. 260.

Immergrüne Osterluzey. Linn. Pflanzensystem. 4. p. 482.

Habitat in Creta. $\overline{\eta}$

Flores curvi, foliis longiores. Caules angulati.

17. ARISTOLOCHIA Pistolochia. A. foliis cordatis cre-
nulatis subtus reticulatis petiolatis, floribus solitariis.
Hall. helv. n. 1029. * delph.

Aristolochia caule infirmo ramoso, foliis cordatis obsolete
dentatis, floribus solitariis erectis. Hort. cliff. monsp.

Aristolochia Pistolochia dicta. Bauh. pin. 307.

Pistolochia. Clus. hist. 2. p. 72. Dod. penit. 525.

Habitat in Hispania, G. Narbonensi, Helvetia. 24

Corollæ labium incurvum. Bractea cordata, sessilis,
parva, insidens basi pedunculi.

GYNANDRIA. HEXANDRIA. 479.

18. **ARISTOLOCHIA rotunda.** A foliis cordatis subtri-
libus obtusis, caule infirmo, floribus solitariis. *carn.*
Blackw. t. 256. Sabb. hort. 2. t. 80. delph.

A. caule infirmo ramoso, foliis cordatis integerrimis,
floribus solitariis erectis. *Hort. cliff. Sauv. monsp.*

A. rotunda, flore ex purpura nigro. *Bauh. pin. 307.*

A. rotunda. I. *Clus. hist. 2. p. 70.*

Aristolochia. Cam. epit. 419.

β. A. rotunda, flore ex albo purpurascens. *Bauh. pin. 307.*

Habitat in Italia, Hispania, G. Narbonensi. 24

Corollæ labium incurvum. Caulis angulatus. Varietas

β. *Foliis longius petiolatis, α vero vix manifestis.*

19. **ARISTOLOCHIA longa.** A. foliis cordatis petiolatis
integerrimis obtusiusculis, caule infirmo, floribus solitariis.
carn. Blackw. t. 257. Sabb. hort. 1. t. 81. delph.

A. caule infirmo ramoso, foliis cordatis integerrimis longe
petiolatis, floribus solitariis. *Sauv. monsp.*

A. longa vera. *Bauh. pin. 307.*

A. longa. *Clus. hist. 2. p. 70.*

β. A. longa hispanica. *Bauh. pin. 307.*

Habitat in Hispania, Italia, Gallia, Carniola. 24

Præcedenti nimis affinis.

21. **ARISTOLOCHIA Clematidis.** A. foliis cordatis, caule
erecto: floribus axillaribus confertis. *paris. Hall. helu.*
n. 1029. pal. sil. Ludw. ed. t. 105. Kniph. cent. 1.
t. 8. Knorr. del. hort. 2. t. A. 15. Blackw. t. 255. Regn.
bot. delph. lugd. burg.

A. caule erecto simplicissimo, foliis cordatis petiolatis,
floribus lateralibus confertis. *Hort. cliff. Sauv. monsp. 111.*

A. longa. *Zorn. ic. pl. med. cent. 1. t. 98. R.*

A. multiflora. *Riu. irreg. t. 116. Hall. R.*

A. clematidis recta. *Bauh. pin. 307.*

A. Clematidis vulgaris. *Clus. hist. 2. p. 71.*

A. rotunda. *Fuchf. hist. 90.*

Habitat in Austria, Gallia, Tataria, &c. 24

Tipula pennicornis facundat flores. Schreber.

478 GYNANDRIA POLYANDRIA

DODECANDRIA

III 5. CYTINUS.

Monogyna. Cal. 4. fidus, superus. Cor. 6. *Andr.*
16. fessiles. Bacca 8. locularis, polysperma.

1. CYTINUS *Hipocistis*.

Afarum (*Hipocistis*) foliis sessilibus imbricatis, floribus quadrifidis. *Spec. pl.*

Afarum aphyllum squamosum. *Sav. monsp.*

Hipocistis omnino lutea f. echræ colore. *Clas. hisp.* 143.

Hipocistis. *Bauh. pin.* 465. *Cam. epit.* 96. 97. *Duham. arb.* 1. p. 170. t. 68.

Habitas in Hispania, Lusitania, Mauritania, parasitica cisti.

POLYANDRIA.

III 9. ARUM.

Spatha monophylla, cucullata. *Spadix* supraculmatus,
inferne femineus, medio flamineus.

* *Acaulia*, foliis compositis.

1. ARUM *Dracunculus*. A. foliis pedatis: foliolis lanceolatis integerrimis æquantibus spatham spadice longiores. *car. Sabb. hort.* 2. t. 76. 77. *Regn. bot. delph.*

A. foliis palmatis: foliolis undecim lanceolatis integerrimis: intermediis majoribus, spadice clavato. *Her. cliff.* 344.

Dracunculus poliphyllus. *Bauh. pin.* 195. *Morif. hist.* 3. p. 548. f. 13. t. 5. f. 46.

Dracontium. *Dod. pempt.* 329.

Habitas in Europa australi. 2.

* * *Acaulia* foliis simplicibus.

3. ARUM *Colocasia*. A. acaule, foliis peltatis ovatis re-

GYNANDRIA. POLYANDRIA. 479

pandis : basi emarginatis. *Catesb. car.* 2. p. 45. t. 45.

Kniph. cent. 9. n. 11. & 12.

A. maximum ægyptium, quod vulgo Colocasia: *Bauh. pin.* 165.

A. ægypticum. *Col. ecphr.* 2. p. 1. t. 1. *Rumph. amb.* 5. p. 313. t. 109.

Colocasia. *Clus. hist.* 2. p. 75.

Habitat in Cretæ, Cypri, Syriæ, Ægypti *aquosis.* 24

6. ARUM esculentum. A. foliis peltatis ovatis integerrimis, basi emarginatis.

A. minus, nymphææ folio, esculentum. *Sloan. jam.* 62. *hist.* 1. p. 167. t. 106. f. 1.

Brassica brasiliara, foliis nymphææ. *Bauh. pin.* 111.

Caladium aquatile. *Rumph. amb.* 5. p. 318. t. 110. f. 11

Habitat in America 24

12. ARUM maculatum. A. acaule, foliis hastatis integerrimis : spadice clavato. *paris. carn. pal. sil. Fl. dan.* t. 505. *Blackw.* t. 228. *Sabb. hort.* 2. t. 74. *Knorr. del.* 2. t. A. 11. *Regn. bot.*

A. foliis sagittatis, spatha recta, clava cylindrica. *Hall. helv.* n. 1302.

A. vulgare non maculatum. *Bauh. pin.* 195.

Arum. *Fuchf. hist.* 69. *Lob. ic.* 597.

A. venis albis. *Bauh. pin.* 195.

β. A. maculatum, maculis candidis f. nigris. *Bauh. pin.* 195.

Habitat in Europa australiore. 24

Aliam adhuc varietatem pingit. *Sabb. hort. rom.* 2. t. 75. R.

14. ARUM proboscideum. A. acaule, foliis hastatis, spatha declinata filiformi-subulata. † *Sabb. hort.* 2. t. 78.

Arisarum flore in tenuem caudam abeunte. *Tournef. inst.* 161.

Arisarum minus proboscideum. *Bocc. mus.* 1. p. 61. t. 50.

Arisarum latifolium minus repens, cespitosum. *Barr. rar.* 182. t. 1150.

β. Arisarum rotundifolium f. potamogeiti folio. *Bocc. sic.*

Moris. hist. 3. p. 544. f. 13. t. 6. f. 19.

Habitat in Apenninis.

15 ARUM Arisarum. A. acaule, foliis cordatis oblongis, spathæ apertura ovata.

A. acaule, foliis cordato-oblongis, spatha bifida, spadice incurvo. *Spec. pl.* 2. *Sabb. hort.* 2. t. 79.

Arisarum latifolium majus. *Bauh. pin.* 196.

Arisarum latifolium alterum. *Bauh. pin.* 196.

Arisarum latifolium. *Clus. hist.* 2. p. 73. *Lob. ic.* 598.

480 GYNANDRIA. POLYANDRIA.

Arisarum serpentinum. Barr. ic. t. 573.

Habitas in Mauritania, Italia, Lusitania, Hispania, Galloprovincia nemoribus.

Spatha superne integra; subinflexa; inferne integra ac convoluta.

17. *ARUM tenuifolium. A. acaule, foliis lanceolatis, spadice setaceo declinato. Sauv. monsp.*

A. angustifolium. Bauh. pin. 196. Clus. hist. 2. p. 74.

Moris. hist. 3. p. 545. f. 13. t. 6. f. 21.

Habitas in Dalmatia, Romæ, Monspeli, Oriente. 24

Gouanus non indicat circa monspelim.

1121. C A L L A.

Spatha plana. Spadix tectus flosculis. Cal. o. Pet. o. Baccæ polyspermæ.

2. *CALLA palustris. C. foliis cordatis, spatha plana, spadice undique hermaphrodito. Fl. succ. ged. pal. fl. Fl. dan. t. 422. lithu.*

C. foliis cordatis. Fl. lapp.

Dracunculus palustris f. radice arundinacea. Bauh. pin. 199.

Dracunculus aquatilis. Dod. pempt. 330.

Provenzalina palustris. Petit. gen. 45.

Habitas in Europæ borealis paludibus.

1123. Z O S T E R A.

Spadix linearis, altero latere fructifero. Cal. o. Cor. o. Stam. alterna. Sem. solitaria, alterna.

1. *ZOSTERA marina. Z. pericarpis sessilibus. carn. Oeda, 15.*

Ruppia foliis linearibus obtusis. Marh. ad. angl. 1741 p. 217.

Fucus f. Alga marina graminea angustifolia seminifera ramosior. Rai. angl. 3. p. 52.

Habitas in mari Balthico, Oceano.

2. *ZOSTERA oceanica. Z. pericarpis pedicellatis oliviformibus.*

Alga angustifolia vitriariorum. Bauh. pin. 364.

Alga marina. Lob. ic. 251.

Alcyonium XIV. Bauh. pin. 368. radio est.

Habitas in Oceano. 4

Differt a precedente Fructibus pedunculatis; Radicibus in pilum complicatis.

CLASSIS

C L A S S I S X X I.

M O N O E C I A.

M O N A N D R I A.

1127. C H A R A.

♂. Cal. nullus. Cor. nulla.
♀. Cal. 4. phyllus. Cor. o.
Stigm. 3. fidum. Sem. 1.

1124. ZANNICHELLIA.

♂. Cal. nullus. Cbr. nulla.
♀. Cal. phyllus. Cor. o.
Pist. 4. Sem. 4.

1125. CERATOCARPUS.

♂. Cal. 2. partitus. Cor. o.
♀. Cal. 2. phyllus. Cbr. o.
Styl. 2. Sem. 1. inferum.

1126. CYNOMORIUM.

♂. Cal. amenti. Cor. nulla.
♀. Cal. amenti. Cor. o.
Styl. 1. Sem. 1. subrotundum.* *Callitriche verna.*

D I A N D R I A.

1130. L E M N A.

♂. Cal. 1. phyllus. Cor. o.
♀. Cal. 1. phyllus. Cor. o.
Styl. 1. Caps. 1. loculi.

T R I A N D R I A.

1133. Z E A.

♂. Gluma 2. flora. Glumid
2. valvis.
♀. Gluma. 1. flora. Glumid.
2. valvis. Styl. 1. Sem.
1. nudum subrot.*Linnaei spec. plant. Europ. Pars II.* H h

1135. COIX.

♂. Glum. 2. flora. Glum.
2. valvis.♀. Glum. 2. flora. Glum.
2. valvis. Styl. 2. fi-
dus. Sem. 1, tectum
nuce.

1137. CAREX.

♂. Ament. 1. florum. Cor.
nulla.♀. Ament. 1. florum. Cor.
1. Styl. 1. Sem. 1. rumi-
catum.

1132. SPARGANIUM.

♂. Cal. 3. phyllus. Cor. 6.

♀. Cal. 3. phyllus. Cor. 6.
Stigm. 2. Sem. 1. sper-
mum.

1131. TYPHA.

♂. Cal. 3. phyllus. Cor. 6.

♀. Cal. capillaris. Cor. 6.
Styl. 1. Sem. 1. pappi-
rui.* *Amaranthi vari.*

T E T R A N D R I A.

1149. URTICA.

♂. Cal. 4. phyllus. Cor. 6.
Nectarium cyathyfor-
me.♀. Cal. 2. valvis. Cor. 6.
Stigm. villosum. Sem. 1.
ovatum.

1150. MORUS.

♂. Cal. 4. partitus. Cor. 6.

♀. Cal. 4. phyllus. Cor. 6.
Styl. 2. Sem. 1, bacci-
catum.

1148. BUXUS.

♂. Cal. 3. phyllus. Cor.
2. petala.♀. Cal. 4. phyllus. Cor. 3.
petala. Stigm. 3. Cap.
3. locularis.

1147. BETULA.

♂. Ament. 3. florum. Cal.
4. partita.

MONOECIA.

483

♀. *Ament.* 2. *florum.* *Cor.* 0. *Styli* 2. *Sem.* 1. ovatum.

1145. LITTORELLA.

♂. *Cal.* 4. *phyllus.* *Cor.* 4. *fida.* *Stam.* longissima.

♀. *Cal.* 0. *Cor.* 4. *fida.* *Styl.* longissimus. *Sem.* nux.

PENTANDRIA.

1152. XANTHIUM.

♂. *Cal.* comm. *polyph.* *Cor.* 5. *fida.* *Filam.* connexa.

♀. *Cal.* 0. *Cor.* 0. *Styli* 2. *Drupa* 2. *locul.*

1157. AMARANTHUS.

♂. *Cal.* *propr.* 5. *phyll.* *Cor.* 0. *Stam.* 3. *f.* 5.

♀. *Cal.* *idem.* *Cor.* 8. *Styli* 3. *Caps.* *circumscissa.*

* *Rumex spinosus; alpinus.*

HEPTANDRIA.

POLYANDRIA. (plura quam 7.)

1164. SAGITTARIA.

♂. *Cal.* 3. *phyllus.* *Cor.* 3. *petala.* *Stam.* 24. *cir-* *citer.*

♀. *Cal.* & *Cor.* *maris.* *Pist.* 100. *Sem.* *numerosa.*

1163. MYRIOPHYLLUM.

♂. *Cal.* 4. *phyllus.* *Cor.* 0. *Stam.* 8.

♀. *Cal.* 4. *phyllus.* *Cor.* 0. *Pist.* 4. *Sem.* 4.

1162. CERATOPHYLLUM

♂. *Cal.* *sub* 7. *partitus.* *Cor.* 0. *Stam.* 18. *cir-* *citer.*

H B 2

1167. POTERIUM.

♀. Cal. sub 7. partitus. Cor.
o. Pist. 1. Sem. 1.

♂. Cal. 4. phyllus. Cor. 4.
partita. Stam. 32. cir-
citer.

♀. Cal. 4. phyllus. Cor. 4.
pet. Pist. 2. Sem. 2.
obducta.

1170. FAGUS.

♂. Cal. 5. fidus. Cor. o.
Stam. 12. circiter.

♀. Cal. 4. fidus. Cor. o.
Styl. 3. Caps. 2. sper-
ma.

1168. QUERCUS.

♂. Cal. 5. fidus. Cor. o.
Stam. 10. circiter.

♀. Cal. integer. Cor. o.
Styli 5. Nux coria-
cea.

1169. JUGLANS.

♂. Ament. imbricat. Cor.
6. partita. Stam. 18. cir-
citer.

♀. Cal. 4. fidus. Cor. 4.
pet. Styli 2. Nux co-
riacea.

1172. CORYLUS.

♂. Ament. imbricat. Cor. o.
Stam. 8.

♀. Cal. 2. phyll. Cor. o.
Styli 2. Nux nuda.

1171. CARPINUS.

♂. Ament. imbricat. Cor.
o. Stamina 10.

♀. Cal. 6. fidus. Cor. o.
Pist. 2. Nux nuda.

1173. PLATANUS.

♂. Ament. globos. Cor. ob-
soleta. Anth. circum-
nata.

♀. Ament. globos. Cor. 5.
pet. Styl. 1. Sem. 1.
papposum.

MONADELPHIA.

1175. PINUS.

♂. Cal. 4. phyllus. Cor. o.
Stam. plurima.

♀. Ament. strobilac. Cor. o.
Pist. 2. Nuces 2. ala-
tae.

1177. CUPRESSUS.

♂. Ament. Cor. o. Anth.
4. sessiles.

♀. Ament. strobilac. Cor. o.
Stigm. 2. Nux. angu-
lata.

1176. THUJA.

♂. Amentum. mor. o.
Anth. 4.

♀. Ament. strobilæ. Cor. o.
Pist. 2. Nux cincta
ala.

1181. CROTON.

♂. Cal. 5. phyll. Cor. 5.
pet. Stam. 15.

♀. Cal. 5. phyll. Cor. o.
Styli 3. Caps. 3. coc-
ca.

1184. RICINUS.

♂. Cal. 3. part. Cor. o.
Stam. multa.

♀. Cal. 5. part. Cor. o.
Styli 3. Caps. 3. coc-
ca.

SYNGENESIA.

1191. MOMORDICA.

♂. Cal. 5. fid. Cor. 5. fida.
Filam. 3.

♀. Cal. 5. fidus. Cor. 5. fi-
da. 3. fid. Pomum. elas-
ticum.

1193. CUCUMIS.

♂. Cal. 5. dentat. Cor. 5.
fida. Filam. 3.

♀. Cal. 5. dent. Cor. 5.
fida. Styl. 3. fid. Per-
mum, Sem. argutis.

1192. CUCURBITA.

♂. Cal. 5. dent. Cor. 5.
fida. Filam. 3.♀. Cal. 5. dent. Cor. 5.
fida. Styl. 3. fidus. Po-
mum. Sem. margina-
tis.

1194. BRYONIA.

♂. Cal. 5. dent. Cor. 5.
part. Filam. 3.♀. Cal. 5. dent. Cor. 5.
part. Styl. 3. fidus. Bo-
ca.

MONOECIA.

MONANDRIA.

1124. ZANICHELLIA.

Masculi. *Cal.* o. *Cor.* o.

Feminei. *Cal.* 1. phyllus. *Cor.* o. *Germina* circiter
4. *Sem.* totidem.

1. ZANICHELLIA *palustris*. Zanichellia. Hall. *helu. n.*
1604. *pal.* Fl. dan. t. 67. lithu. delph. lugd.

Z. *palustris* major, foliis gramineis. Mich. gen. 71. t. 34
f. 1. 2.

Graminifolia. Dill. *giss. app.* 168.

Algoides vulgaris. Vaill. *act.* 1719. p. 15. t. 1. f. 1.

Potamogeton capillaceum, capitulis ad alas trifidis. Bauh.
pin. 193. *prodr.* 101.

Potamogetoni similis graminifolia ramosa & ad, genicula
polyceratos. Pluk. *alm.* 305. t. 102. f. 7.

Habitat in Europæ, Virginiae fossis, fluviis. ☉

1126. CYNOMORIUM.

Masc. *Cal.* Amentum imbricatum. *Cor.* o.

Fem. *Cal.* In eodem amento. *Cor.* o. *Styl.* 1
Sem. 1. subrotundum.

1. CYNOMORIUM *coccineum*. Cynomorium. *Aman.*
acad. 4. p. 351. t. 2. *

C. erectum breve cylindricum nudum prima ætate squa-
matum. Brown. *jam.* 334.

C. purpureum officinarum. Mich. gen. 17. t. 12.

Fungus mauritanicus verrucosus ruber. Pet. *gaz.* t. 39. f. 8.

Fungus typhoides liburnensis. Til. *pis.* 64. t. 25.

H h. 4

Fungus typhoides concineus tuberosus, molliculus. *Rehm.*
musf. 2. p. 69. figc. 80. t. 81.

Habitat in Jamaica, Mauritania, parvula in Malta, Sicilia,
Gades, parasiticum radicum.

Tota planta est unicum Amentum.

1127. CHARA.

Masc. Cal. 0, Cor. 0. Anthera germinat subiecta,

Fem. Cal. 4. phyllus, Cor. 0. Stigma 1. fidem,
Sem. 1.

1. CHARA tomentosa. C. aculeis caulinis ovatis, delph.

C. caule subtomentoso fragili. *Hall. helu. n. 1683.*

Equisetum fragile majus subcinereum aquis immersum,
Morif. hist. 3. p. 621. f. 15. t. 4. f. 9.

Equisetum f. Hippuris lacustris, foliis mansu arenosa, Pluk.
alm. 135. t. 29. f. 4.

Habitat in Europæ stagnis, mari, lacubus.

2. CHARA vulgaris. C. caulibus lævibus, frondibus in-
 terne dentatis. *god. carn. gallob. fil. Fl. dan. t. 150. pel.*
lithu. lugd. delph. Fl. suec. 492. 1131.

C. caule levi fragili. *Hall. helu. n. 1681.*

C. caule herbaceo rigido, ramulis foliaceis patulis. *Wass.*
crypt.

Hippuris foetida, Dill. giff. 105. R.

Chara caulibus lævibus. Roy. lugd. 533.

C. vulgaris foetida. *Vaill. aq. 1719. p. 23. t. 3. f. 1.*

Equisetum foetidum sub aqua repens. Bauh. pin. 16. p. 25.
theatr. 251. Bauh. hist. 3. p. 731.

Habitat in Europæ aquis pigris.

Baccæ oblongæ polyspermæ, Weis.

3. CHARA hispida. C. aculeis caulinis capillaribus confertis.
*Fl. suec. * Fl. dan. t. 154. lugd. delph.*

C. caulibus aculeatis. *Hort. cliff.*

C. caule fragili spinoso. *Hall. helu. n. 1682.*

C. major, caulibus spinosis. *Vaill. aq. 1719. p. 18. t. 3. f. 3.*

Hippuris muscosa fetis per extremum stellatis. Dill. ephem.
N. C. cont. 8. app. p. 58. t. 13. f. 1. Hall. an. R.

Equisetum f. Hippuris muscosus sub aqua repens. Pluk. alm.
135. t. 193. f. 6.

Habitat in Europæ maritimis.

MONOECIA. DIANDRIA. 489

4. *CHARA flexilis*. C. caulium articulis inermibus diaphanis superne latioribus. *Fl. suec. pal. delph.*
C. caule levi pellucido flexili. Hall. helu. n. 1684.
C. translucens major & minor flexilis. Vaill. par. 18. t. 3. f. 8. 9.
C. translucens minor flexilis. Rai. angl. 3. p. 133.
Hippuris setis biturcis. Dill. ephem. N. C. cens. 6. app. p. 59. t. 13. f. 2. Hall. R.
Habitat in Europæ maritimis.

D I A N D R I A.

1130. L E M N A.

Masc. Cal. 1. phyllus. Cor. nulla.

Fem. Cal. 1. phyllus. Cor. 0. Styl. 1. Caps. unilocularis.

1. *LEMNA trisulca*. L. foliis petiolatis lanceolatis. *suec. ged. carn. sub Lenticula. pal. sil. delph. lithu. burg.*
L. foliis lanceolatis : extremis decussatis. Sauv. monsp.
Hydrophace caule filiformi, foliis elliptico-lanceolatis viviparis. Hall. helu. n. 1901.
Lenticularia ramosa monorrhiza, foliis oblongis pediculis longioribus donatis. Mich. gen. 16. t. 11. f. 5.
Lenticula aquatica trisulca. Bauh. pin. 362.
Hederula aquatica. Lob. ic. 2. p. 36.
Habitat in Europa sub aquis pigris, puris.
 2. *LEMNA minor*. L. foliis sessilibus utrinque planiusculis, radicibus solitariis. *Fl. suec. ged. carn. sub Lenticula. pal. sil. lugd. lithu. burg.*
Hydrophace monorrhiza, foliis planis ovatis. Hall. helu. n. 1900.
Lenticula minor monorrhiza, foliis subrotundis utrinque viridibus. Mich. gen. 16. t. 11. f. 3. Tozzet. app. 148.
Lens palustris. Cam. epit. 852. Rai. angl. 3. p. 129. t. 4. f. 1. Vallisn. oper 2. p. 81. t. 14. 15. Blackw. t. 380.
Lenticula palustris. Bauh. pin. 362. Vaill. 14. t. 20. f. 3.
Habitat in Europæ aquis quietis.
Floret quotannis sub canicula.

490 MONOCOTYLEDONATA. DICANDRIA.

3. LEMNA gibba. L. foliis sessilibus subrotundis hemisphaericis, radicibus foliaceis. Fl. succ. ped. fil. burg. lushu.

Hydrophace monorhiza, foliis pulvinatis, inferae cuneato-ventricosa. Hall. bot. n. 1898.

Lenticula palustris major: inferae magis convexa, fructu polyspermo. Mich. gen. 15. t. 11. f. 2.

Habitat in Europae aquis sequitur.

Varietas precedenti secundum Gerard. 122.

4. LEMNA polyrrhiza. L. foliis sessilibus, radicibus quatuor. Hort. cliff. 417. Fl. succ. ped. pal. fil. lushu. burg.

Hydrophace polyrrhiza, foliis ovatis planis. Hall. bot. 1899.

Lenticularia major polyrrhiza inferae sexoperma. Mich. gen. 16. t. 11. f. 1.

Lenticula palustris major. Rai. angl. 3. p. 129. t. 4. f. 2. Vaill. par. 114. t. 20. f. 2.

Habitat in Europae paludibus, fossis.

5. LEMNA arhiza. L. foliis geminis oppositis.

Lenticula omnium minima, arhiza. Mich. gen. 16. t. 11. f. 4.

Habitat in Italia, Galliae aquis, De Clafac.

Radix plane nulla. Folia quatuordecim dies calerent, citius minora.

TRIANDRIA.

1131. TYPHA.

Masc. Amentum cylindricum. Cal. obsoletum, triphyllum. Cor. o.

Fem. Amentum cylindricum, infra masculis. Cal. capillo villoso. Cor. o. Sem. 1. involucro capillari.

1. TYPHA latifolia. T. foliis subensiformibus, cula femineaeque approximatis. Fl. succ. ped. pal. fil.

Fl. dan. t. 643. Kniph. ent. 6. n. 97. leg. lushu. burg.

T. clava unica. Hall. helu. n. 1505.

T. palustris major. Bauh. pin. 20. Marif. 122. t. 2. f. 8. t. 13. f. 1.

Habitas in paludibus Europæ.

Spathæ 2 deciduæ : altera ad Spicæ masculæ basin ; altera ad ejus medium.

2. **TYPHA angustifolia.** C. foliis semicylindricis, spica mascula femineaque remotis. *carn. pal. lithu. burg.*

T. clava mascula a femina distincta. *Hall. helu. n. 1306.*

Typha. *Gmel. sib. 1. p. 133. Fuchs. hist. 822.*

T. palustris, clava gracili. *Bauh. pin. 20.*

T. palustris media. *Moris. hist. 3. p. 246. f. 8. t. 13. f. 2. Bauh. hist. p. 540.*

β. T. palustris minor. *Bauh. pin. 20.*

T. minor. *Bauh. hist. 2. p. 540. Rai. extr. 259.*

Habitas in Europæ paludibus. 24

1132. SPARGANIUM.

Masc. Amentum subrotundum. *Cal. 3. phyllus. Cor. o.*

Fem. Amentum subrotundum. *Cor. 3. phyllus. Cor. o. Stigma bifidum. Drupa exsucca. 1. sperma.*

1. **SPARGANIUM erectum.** S. foliis erectis triquetris. *Fl. suec. carn. herb. pal. sil. Kniph. cent. n. 95. lithu. delph. lugd. burg.*

S. caule foliisque erectis. *Hall. helu. n. 1303.*

S. ramosum. *Bauh. pin. 15. theatr. 228.*

Platanaria f. Buromon. *Dod. pempt. 601.*

β. Sparganium non ramosum. *Bauh. pin. 15. theatr. 231. Oed. dan. t. 260.*

Habitas in Zonæ frigidæ septentrionalis aquosis. 24

Maxima individua vidi *Valdstrenæ*, altitudine fere humana.

2. **SPARGANIUM natans.** S. foliis decumbentibus planis, *pal.*

S. foliis natantibus plano-convexis. *Fl. suec. delph. burg.*

S. foliis planis mollibus decumbentibus. *Hall. helu. n. 1304.*

S. non ramosum minus. *Dill. giff. 130. spec. 58.*

S. minimum. *Rai. hist. 1910. angl. 3. p. 437.*

Habitas in Europæ borealis lacubus, paludibus. 24

Tenera in paludibus foliis aut erectis aut procumbentibus ;

492 **MONOCIA. TRIANGULARIA.**
in lacubus extra arundines, imo extra Scirpus, ubi etiam
alta notant folia, quasi diversa.
 Tabulam 260 flor. den. huc trahit Linnæus, est peristylus
 ad Sp. erectum var. A. Hall. Pollich. R.

1133. **Z E A.**

Masc. in distinctis spicis. Cal. Gluma 2. flora;
 mutica. Cor. Gluma mutica.

Fem. Cal. Gluma 2. valvis. Cor. Gluma 2. valvis.
 Stylus 1, filiformis, pendulus. Sem. solitaria,
 receptaculo oblongo immersa.

1. ZEA Mays. Blackw. 1. 547. a. b. Kniph. cent. 3. p. 99.
 Frumentum indicum Mays dictum, Bauh. pin. 23. theatr.
 490.
 Frumentum indicum. Cam. epl. 186. Dod. pempt. 309.
 Habitat in America. ☉
 Varietates hujus plurimæ existunt.

1135. **C O I X.**

Masc. in spicis remotis. Cal. Gluma 2. flora;
 Cor. Gluma mutica.

Fem. Cal. Gluma 2. flora. Cor. Gluma reptica.
 Stylus 2. partitus. Sem. calyce ossificato
 tectum.

1. COIX Lachryma. C. seminibus ovatis. Kniph. cent. 4.
 n. 19. Regn. bot.
 Lithospermum arundinaceum. Bauh. pin. 258.
 Lachryma Jobi. Clus. hist. 2. p. 216.
 Ova piscium. Rumph. amb. 5. p. 191. t. 75.
 Catriconda. Rheed. mal. 12. p. 133. t. 70.
 Habitat in Indiis. 24
 Speciem aliam etiam commemorant Plumier & Rumph.
 mihi ignotam.

1137. CAREX.

Masc. *Amentum* imbricatum. *cal.* 1. *phyllus*.
Cor. 0.

Fem. *Amentum* imbricatum. *Cal.* 1. *phyllus*. *Cor.* 0.
Nectarium inflatum, 2. dentatum. *Stigm.* 3. *Sem.*
triquetrum, intra nectarium.

* * *Spica unica simplici.*

1. CAREX *dioica*. C. *spica simplici dioica*. *Fl. suec. gallob.*
Flor. dan. t. 369. *delph. lugd.*

Cyperoides parvum, caulibus & foliis tenuissimis triangularibus, spica longiore capsulis oblongis. *Mich. gen.* 56.
t. 32. f. 1.

Gramen cyperoides minimum, spica simplici crassa. *Morif. hist.* 3. p. 244. f. 8. t. 12. ult. (mas.)

Gramen cyperoides minus, ranunculi capitulo longiore. *Rai. angl.* 3. p. 425.

Habitat in Europæ pratis humidis. 2

2. CAREX *capitata*. C. *spica simplici androgyna ovata* :
superne mascula, capsulis imbricato-patulis. *Fl. dan.*
t. 372.

C. *spica unica subrotunda*, seminibus inflatis. *Hall. helv.*
n. 1351.

Cyperoides parvum, caulibus & foliis tenuissimis triangularibus, spica subrotunda. *Mich. gen.* 56. t. 32. f. 2.

Gramen Cyperoides minimum, ranunculi capitulo simplici asperiore rotundo. *Morif. hist.* 3. p. 245. f. 8. t. 12. f. 6.

Habitat in Lapponia, Anglia. 2

Simillima C. dioica, sed Spica exacte ovata : apice mascula ; nec cylindrica dioica.

3. CAREX *pulicaris*. C. *spica simplici androgyna* : superne mascula, capsulis divaricatis retroflexis. *carn. gallob. herb.* t. 14. f. 1. *delph.*

C. *spica unica*, terete, seminibus rostratis. *Hall. helv.* n. 1350.

C. *spica simplici androgyna*. *Fl. suec. paris.*

C. *spica unica*. *Fl. lapp.* 747.

C. *minima*, caulibus & foliis capillaceis, capitulo singulari

494 MONOECIA. TRIANDRIA:

tenuiori, capsulis oblongis utrinque acuminatis & deorsum flexis. *Mich. gen.* 66. t. 33. f. 1.

Gramen cyperoides spica simplici casta. *Scheuchz. gram.* 497. t. 11. f. 9. 10. *Hall. R.*

Gramen cyperoides minimum, spica simplici casta. *Morif. l. c. f.* 22. *planta mascula. Hall. R.*

Gramen cyperoides minimum, seminibus deorsum reflexis puliciformibus. *Rai. hist.* 1298. *Pluck. phyt.* 34. f. 10. *Morif. hist.* 3. p. 244. f. 8. t. 12. f. 21. *femina.*

Habitat in Europæ paludibus limosis. 2

Tabulam 166. *flor. dan. huc pertinere monet Hællerus. l. c. R.*

5. CAREX cyperoides. C. umbella subcapitata, floribus simplicissimis subulatis, involucrio tetraphyllo elongato.

C. (bohémica) spica globosa conglomerata androgyna, capsulis subulatis, involucrio triphyllo. *Schreb. gram.* t. 28. f. 3. *R.*

C. bohémica aquatica annua, cyperi facie, caule molli; capitulis in glomeratam spicam digestis. *Mich. gen.* 70. t. 33. f. 19.

Scirpus spica multipartita, seminibus caudatis. *Gmel. sib.* 1. p. 81.

Cyperus minor, capitulis inflexis. *Buxb. cent.* 4. p. 34. t. 61. *Habitat in Sibiria, Bohemia.* 2

Plantam misit D. Schreber Prof. Erlang. De genere n. novi. Radix perennis. Culmi pedales, triquetri. Capitulum terminale e glomerulis pluribus. Glumæ exteriores singuli glomeruli masculæ; interiores femineæ. Stamina 3. alba. Pistillum nectario longius, stylo bifido. *Schreb.*

* * Spicis androgynis.

6. CAREX baldensis. C. spinis ternis congestis sessilibus ovatis triquetris androgynis, involucrio diphylo. *Amæn. acad.* 4. * *delph.*

C. alpina, capitulis albis: squamis triquetris & pulchre lucentibus. *Segu. ver.* 1. p. 125.

Gramen junceum montanum, capite squamoso. *Bauh. pin.* 6. *prodr.* 13. t. 13.

Gramen junceum e monte baldo. *Bauh. hist.* 2. p. 509.

Habitat in Baldo. Seguiet. Allioni. 2

Involucri altera valvula spicis longior; altera brevior. Spicæ 3. f. 4. in capitulum congestæ, albæ oblongæ paleis lanceolatis flosculos masculos distinguuntibus, ad quarum basin flosculi feminei globosi: Stylo longo tripartito.

7. CAREX *arenaria*. C. spica composita, spiculis androgynis inferioribus remotioribus foliolo longiori instructis, culmo triquetro. *Fl. suec.* * *Leers. herb. n.* 706. t. 14. f. 2. *delph.*

C. spicis stipula brevioribus imis distinctis, superioribus congestis. *Hall. helv. n.* 1363. R.

Carex maritima humilis, radice repente, caule triquetro, spica spadicea. *Mich. gen.* 67. t. 33. f. 4.

Gramen cyperoides e monte ballon simile humilius, in arenosis nascens. *Pluk. alm.* 178. t. 34. f. 8.

Gramen cyperoides minus repens, spica divisa. *Lass. pruss.* 116. t. 31.

Habitat in Europæ arena præsertim mobili. 2x

Conf. Weigel *obs.* p. 34. & Carex disticha Schreb. *Spicil.* p. 63. vel. Car. spicata. *Pollich. pal. n.* 875. R.

8. CAREX *uliginosa*. C. spica composita, spiculis androgyni s. inferioribus remotioribus foliolo longiori instructis, culmo tereti. *Fl. suec. delph.*

Habitat in Sueciæ paludibus sylvaticis.

Synonyma Schoeni compressi huc referuntur. a variis sed parum refert. Spica totalis magis compressa, quam facile alia species.

9. CAREX *leporina*. C. spica composita, spiculis ovatis sessilibus approximatis alternis androgynis nudis. *Fl. suec. Leers. herb. n.* 707. t. 14. f. 6. *pal. fil. Flor. dan. t.* 294. *delph. burg. lugd.*

C. spinis ternis sessilibus confertis androgynis. *Fl. lapp.* 322.

C. spicis nudis subrotundis, distinctis, in summo caule congestis. *Hall. helv. n.* 1361. R.

C. angustifolia, caule triquetro, spicis pluribus elegantibus parum inter se distantibus. *Segu. ver. 1. p.* 124. t. 1. f. 3.

Gramen cyperoides, spica e pluribus spicis mollibus composita. *Scheuch. gram.* 456. t. 10. f. 15.

Gramen cyperoides palustre majus, spica divisa. *Bauh. pin.* 6. *Moris. hist.* 3. p. 244. f. 8. t. 12. f. 29.

Habitat in Europæ pratis udis. 2x

Spica e spiculis 3. f. 6. approximatis: Paleis flosculos distinguuntibus, griseis, seminibus ipsis longioribus. Styli incurvi.

Tab. 294. *fl. dan.* non videtur huc pertinere *Haller. l. c.* Synonymon *Seguieri* etiam non huc pertinet, monente *Hallero*; quod est Carex præcox Schreb. *spicil.* p. 63. n. 1010. R.

10. CAREX *vulpina*. C. spica supra decomposita inferne

496 **MONOECIA. TRIANDRIA.**

laxiore : spiculis androgynis ovatis glomeratis : superioribus masculis. *Fl. suec. paris. gallob. tub. Leers. herb. n. 708. t. 14. f. 5. pal. sil. Fl. dan. t. 308. delph. burg. lagd.*

C. spicis subrotundis, echinatis, supremis confluentibus; capsulis rostratis bifidis. *Hall. helu. n. 1364. R.*

C. culmo triquetro; spica supra decomposita; racemis imbricatis; spiculis androgynis, sessilibus, ovatis; foliis angulatis. *Scop. carn.*

Carex palustris major, radice fibrosa, caule exquisitè triangulari, spica brevi habitatiore compacta. *Mich. gen. 69. t. 33. f. 13. 14.*

Gramen Cyperoides palustre majus, spica compacta. *Bauh. pin. 6. Moris. hist. 3. p. 244. f. 8. t. 12. f. 14.*

β. Gramen spartium cyperoides, echinata spica, nemorum majus. *Barr. rar. 19. Hall. R.*

Habitat in Europæ paludibus.

Tabula 32. flor. *sik. non bona, si nostra est. Hall. l. c. R.*

11. CAREX *brizoides*. C. spica composita disticha nuda: spiculis androgynis oblongis contiguus, culmo nudo. *Darrh. nass. delph.*

C. spicis distichis, teretibus, repandis. *Hall. helu. n. 1358.*

C. culmo triquetro; spica composita; spiculis androgynis; alternis, acutis, teretibus, imbricatis. *Scop. carn.*

C. fibrata radice, angustifolia, caule exquisitè triangulari. *Mich. gen. 70. t. 33. f. 17.*

Gramen Cyperoides elegans, spica composita asperiore. *Rai. hist. 1297. Moris. hist. 3. p. 245.*

Habitat in Europa.

12. CAREX *muricata*. C. spiculis subovatis sessilibus remotis androgynis, capsulis acutis divergentibus spinosis. *Fl. suec. ged. Leers. herb. n. 709. t. 14. f. 8. pal. sil. gallob. Flor. dan. t. 284. delph.*

C. spicis ternis echinatis, glumis lanceolatis, capsulae mucrone simplici. *Hall. helu. n. 1366.*

C. spica androgyna, spiculis compactis obscure alternis, culmo nudo compresso. *Guett. stamp.*

C. nemorosa, fibrosa radice, angustifolia minima, caule exquisitè triangulari, spica brevi interrupta. *Mich. gen. 69. t. 33. f. 11.*

Gramen Cyperoides spicatum minimum, spica divulsa aculeata. *Scheuch. gram. 485.*

Gramen nemorosum, spicis parvis asperis. *Bauh. pin. 7. Moris. hist. 3. p. 244. f. 8. t. 12. f. 27.*

Habitat

Habitat in Europæ nemoribus humilibus. 24

13. CAREX loliacea. C. spiculis subovatis sessilibus remotis androgynis : capsulis ovatis tetetiusculis muticis divaricatis. *Fl. suec.*

C. nemorosa, fibrosa radice, caule triangulari, spica divulsa s. interrupta, capitulis solitariis præterquam ultimo. *Mich. gen. 69. t. 33. f. 10. sec. Schreb. R.*

Habitat in Suecia, Saxonia. Schreb. spicil. p. 64.

Radix repens. Folia graminea, tenera, lævia. Culmus lævis, superne nudus. Spiculæ 4. s. 8. ovatae, parvæ, ad apicem culmi sparsæ, albidæ. Semina ovata, obtusa, muticæ demto pistillo, subtus rotundata. Differt à C. muricata, cui smillima, quod dimidio minor, quod capsulæ minus divaricatae, obtusæ, nec margine acutæ, ad hanc nostram proprius accedunt figuræ *Moris. hist. f. 8. t. 12. f. 26. Pluk. phyt. 34. t. 4. quam ad alias.*

14. CAREX remota. C. spicis ovatis subsessilibus remotis androgynis, bracteis culmum æquantibus. *delph. burg.*

C. spiculis ovatis subsessilibus remotis androgynis, foliis caulinis culmum æquantibus. *Spec. pl. Scholl. barb. Leers. herb. n. 711. t. 15. f. 1. pal. sil. Fl. dan. t. 370.*

C. spicis foliorum in alis sessilibus. *Guett. stamp. 1. p. 151.*

C. spicis androgynis, imis in alâ prælongi folii sessilibus, supremis nudis. *Hall. helu. n. 1357.*

C. culmo triquetro; spicis androgynis, sessilibus, oblongis; inferioribus remotis; superioribus approximatis, nudis. *Scop. carn.*

C. angustifolia, caule triquetro, capitulis pulchellis; strigosioribus. *Mich. gen. p. 70. n. 3. t. 33. f. 16.*

Cyperoides angustifolium, spicis sessilibus in foliorum alis. *Rai. hist. 1295. Moris. hist. 3. p. 243. f. 8. t. 12. f. 17. Pluk. alm. 178. t. 34. f. 3. Scheuch. gram. 483.*

β. Carex (axillaris) spiculis ovatis sessilibus remotis androgynis, bracteis culmum æquantibus. *Spec. ed. 2. p. 1382.*

Habitat in Europæ umbrosis subhumidis. 24

15. CAREX elongata. C. spiculis oblongis sessilibus remotis androgynis, capsulis ovatis acutis. *Fl. suec. tub. Leers. herb. n. 711. t. 14. f. 7. pal.*

C. spicis ovatis, imis remotis, supremis congestis. *Hall. helu. n. 1359.*

Cyperoides polystachyon, spicis laxis paniculam veluti componentibus. *Scheuch. gram. 487. t. 11. f. 4.*

Linnaei spec. plant. Europ. Pars II. 11

238 **MONOECIA TRIANDRIA.**

Gramen Cyperoides angustifolium, spicis longis erectis.
Banh. pin. 6. theatr. 84.

Gramen Cyperoides medium angustifolium, spicis
 tribus erectis flavicantibus. *Morif. hist. 3. p. 242. f.*
8. a. 12. f. 8.

Habitas in Europa. 2.

16. **CAREX canescens.** C. spiculis subrotundis remotis
 sessilibus obtusis androgynis, capsulis ovatis obtusif-
 culis. *Fl. succ. tab. pal. Fl. dan. a. 285. delph.*

C. spicis ovatis, alternis, supremis se contingentibus. *Hall.*
helu. n. 1360. R.

C. spica androgyna, spiculis compactis obscure alternis,
 culmo nudo compresso. *Grev. fl. scot.*

C. spicis plurimis remotis sessilibus subrotundis angulis.
Fl. lapp. 332.

Gramen Cyperoides spicis curvis divulsis. *Less. prof.*
11752. 32.

Habitas in Europa septentrionali. 2.

Carex Halleri n. 1360. ad Caricem suam tinctorum
Pollichio pertinere videtur, non vero Symplocarum ab Hallero
adducta. R.

17. **CAREX paniculata.** C. racemo corymboso, spiculis
 androgynis. *gal. Leers. herb. n. 713. a. 14. f. 4. delph.*

C. spica paniculata. *Hall. helu. n. 1368. pal.*

C. angustifolia, caule triquetro, spica umbellata. *Grev.*
Mich. gen. 68. t. 33. f. 7.

Cyperus longus inodorus sylvaticus. *Banh. pin. 14.*
theatr. 221.

Cyperus alpinus longus inodorus, panicula brevior
 minus sparsa. *Scheuch. prodr. 7. a. 8.*

Gramen Cyperoides palustre, spica longiore lina. *Rai.*
hist. 1269. Morif. hist. 3. p. 244. f. 8. a. 12. f. 13.
Scheuch. gram. 499.

β. Cyperoides palustre, spica laxa, minus. *Scheuch. gram.*
801. Hall. Pollich. Leers. R.

Habitas in Europa australioris alpinis uliginosis. 2.

*** *Spicis sexu distinctis: femineis sessilibus.*

19. **CAREX flava.** C. spicis confertis subsessilibus sub-
 rotundis: mascula linearis, capsulis acutis recurvis. *Fl. pa-*
ris. gallob. carn. Leers. herb. n. 714. a. 15. f. 6. pal.
sil. delph. burg. lugd.

C. spicis ad apicem tribus sessilibus: femineis subrotundis,
 capsulis acutis recurvis. *Fl. lapp. 334.*

C. spicis femininis petiolatis, erectis, ovatis, echinatis:
Hall. helu. n. 1380.

Gramen Cyperoides palustre aculeatum, capitulo brevior:
T. Scheuch. gram. 426.

Gramen Cyperoides aculeatum germanicum s. minus. *Bauh. pin. 7. Moris. hist. 3. p. 243. f. 8. t. 12. f. 19.*

Habitat in Europæ paludibus. 24

20. CAREX pedata. C. spicis feminis sessilibus oblongis: inferiore axillari, foliis subfiliformibus. *Dærrh. nass. delph.*

C. spicis femininis raris, marem æquantibus, imâ petiolata. *Hall. helu. n. 1375.*

Gramen caryophylleum, angustissimis foliis, spicis sessilibus brevibus erectis non compactis. *Moris. hist. 3. p. 243.*

Gramen caryophyllatum nemorosum, spica multiplici.
Bauh. pin. 4. theatr. 47. prodr. 22. Scheuch. gram. 450. Mich. gen. t. 32. f. 14.

Gramen parvum sylvaticum paniculis forma pedum avium:
Bauh. hist. 2. p. 499. Hall. R.

Habitat in Helvetia, Anglia, Lapponia, Germania. 24

Folia pedalia, angustissima, rigidiuscula, inferne lævia:

Culmus solidior, læviusculus, hinc rotundatus, foliis longior. Spicæ tres, sessiles, Mascula terminalis, oblonga, ferruginea. Femineæ 2. alternæ, sessiles, ovato-oblongæ; fusco-ferrugineæ: superiore absque subjecto folio; inferiore ex ala folii, longitudine culmi. Flosculi minus arcte imbricati. Capsulæ obsolete pubescentes.

21. CAREX digitata. C. spicis linearibus erectis: mascula brevior inferioreque; bracteis aphyllis, capsulis distantibus. *Fl. suec. paris. ged. Leers. herb. n. 715. t. 16. f. 4. pal. Dærr. nass. delph. lithu.*

C. spica mascula terminali, quam feminea suprema brevior, capsulis distantibus. *Fl. lapp. 338.*

C. spicis femininis petiolatis raris, masculam superantibus.
Hall. helu. n. 1376:

C. culmo subtereti, nudo; spicis erectis, imbricatis; feminis supremis supra marem elatis. *Scop. carn.*

Cyperoides montanum nemorosum, caule triquetro compresso, spicis ferrugineis tenuioribus inter se distantibus:
Mich. gen. 65. t. 32. f. 9.

Gramen caryophyllatum montanum, spica varia. *Bauh. pin. 4. prodr. 9. Scheuch. gram. 448. t. 10. f. 14. Moris. hist. 3. p. 243. t. 12. f. 13.*

500 MONOECIA. TRIANDRIA.

Gramen caryophyllatum polycarpon, fructu triangulari.

Las. pruss. 112. t. 27.

Habitat in Europæ nemoribus. 24

22. CAREX montana. C. spicis femineis sessilibus subsolitariis ovatis masculæ approximatis, culmo nudo, capsulis pubescentibus. *Fl. suec.* * *Reyg. ged. Leers. herb. n.* 716. t. 16. f. 6. *pal. sil. Fl. dan. t.* 444. *delph. lithu.*

C. foliis caryophyllæis, spicis sessilibus approximatis, paucifloris, capsulis ovatis subhirsutis. *Hall. helu. n.* 1372. *Leers. Gouan. R.*

Cyperoides montanum, foliis angustis parvis, spica spadicea divisa. *Scheuch. gram.* 419. t. 10. f. 8. 9.

Cyperoides alpinum saxatile, capillaceo folio, caule rotundo triquetro, spica seminali nigricante plerumque unica. *Mich. gen.* 64. t. 32. f. 3.

Habitat in Europæ montanis apricis. 24

Folia mollia, quam congenerum magis flavescentia. Spica feminea evadit maturitate nigra. Semina vix conspiciuntur hirta sunt.

23. CAREX tomentosa. C. spicis femineis subpedunculatis, capsulis subglobosis tomentosis. *delph.*

C. angustifolius, spicis sessilibus, teretibus, approximatis, capsulis hirsutis. *Hall. helu. n.* 1373. R.

C. montana. *Scop. carn. excluso Linnaeo. R.*

Cyperoides montanum vernum, foliis angustis, &c. *Mich. gen.* 64. n. 68. R.

Cyperoides angustifolium montanum, folliculis feminum villosis. *Scheuchz. gram.* 423.

Gramen spicatum angustifolium montanum. *Bauh. pin. 4 prodr.* 20. *theatr.* 47. R.

Habitat in Austria. Jacq. Saxonia. Schreb. Helvetia, Craniolia. 24

Facias C. paniceæ, figura *Mich. gen. t.* 132. f. 13. hoc potius, quam C. pallescentem repræsentat; differt a Schenckiana. Bracteis spica longioribus. Culmus triquetus. Folia angusta. Spica mascula terminalis, oblonga, pallescenti-ferruginea. Sp. femineæ 2, alternæ, oblongæ, erectæ, breviter pedunculatæ. Bractæe longitudine culmi superioris. Folliculi, s. Capsulæ subglobosæ s. subovatæ, vix acuminatæ notabiliter tomentosæ, pilis brevissimis mollibus quasi muricatæ. Semina glabra, subglobosa cum lævi protuberantia apicis.

24. CAREX globularis. C. spica mascula oblonga, femina

sessili ovata, foliolo florali breviori approximata. *Fl. suec. Scholl. barb. delph.*

C. spicis sessilibus approximatis ovatis. *Hall. helu. n. 1371. Pollich.*

C. (*cricetorum*) spica mascula terminali obverse ovata, obtusa, femineis sessilibus, ovatis, capsulis pubescentibus. *Pollich. pal. n. 886. ex sentent. Pollichi. R.*

C. spica mascula a feminea in ala folioli remota. *Fl. lapp. 336.*

Habitat in Europa frigidiore.

25. CAREX *filiformis*. C. spica mascula oblonga, femineis sessilibus oblongis, inferiore foliolo proprio brevior. *Fl. suec. tub. norv. Leers. herb. n. 718. t. 16. f. 5. pal. delph.*

C. culmo foliisque firmis erectis; spicis femininis quaternis, longe petiolatis erectis. *Hall. helu. n. 1383. secundum Gouan. & Pollich. R.*

Cyperoides sylvaticum tenuifolium, spicis parvis tenuibus spadiceo-viridibus. *Scheuch. gram. 425. t. 10. f. 11.*

Gramen cyperoides flavicans, spicis brevibus prope summum caulis. *Morif. hist. 3. sect. 8. t. 2. f. 16. ic. elegans. sec. Gouan. R.*

Habitat in Europæ nemoribus.

26. CAREX *pilulifera*. C. spicis terminalibus confertis subrotundis: masculina oblonga. *Gunn. norv. delph.*

Gramen cyperoides tenuifolium, spicis ad summitatem caulis sessilibus globulorum æmulis. *Pluk. alm. 178. t. 91. f. 8.*

Habitat in Europa.

27. CAREX *saxatilis*. C. spicis tribus ovatis sessilibus alternis; mascula oblonga. *Fl. suec. Gmel. tub. Scop. Flor. dan. t. 159. delph.*

C. spicis tribus ad apicem sessilibus, femineis ovatis atris. *Fl. lapp.*

C. spicis tribus alternis sessilibus bracteatis, feminis ovatis acutis, mare oblonga. *Gmel. sib. 1. p. 134. n. 71. R.*

Cyperoides alpinum saxatile, caule cubitali rotundo-triquetro nudo, spicis brevibus binis s. ternis confertim nascentibus, squamis obtuse mucronatis. *Mich. gen. 63. t. 32. f. 4.*

Habitat in Europæ alpibus.

Spicæ atro-rubrae. Capsulae nigricantes. *Slop. l. c. R.*

102 MONOECIA. TRIANDRIA.

- **** *Spicis fern distinctis : femineis pedunculatis.*
28. CAREX *oputa*. C. spicis androgynis terminalibus pedunculatis : florentibus erectis ; fructiferis pendulis Fl. succ. carn. Fl. dan. t. 158. delph.
- C. spicis ad apicem culmi pendulis androgynis. Fl. lapp.
- C. spicis ovatis , petiolatis , congestis , senescento cunctantibus. Hall. helv. n. 1369.
- Gramen alpicum pulchrum , foliis caryophyllaris spicis atris & tumetibus. Schenck. gram. 481.
- Gramen Cyperoides alpinum majus , spicis atris plurimis simul junctis. Radb. elys. 1. p. 55. t. 23.
- Habitat in Alpibus Europæ.
29. CAREX *limosa*. C. spicis ovatis pendulis : mascula longiore erectiore , radice repente. Fl. succ. carn. u. b. Leers. herb. n. 719. t. 15. f. 3. demia varietate β. pal. Flor. dan. t. 646.
- Carex ~~spicis~~ versus apicem pendulis : masculis & femineis. Fl. lapp. 325.
- C. spicis femineis pendulis , capsulis ovatis compressis. Hall. helv. n. 1392.
- Cyperoides spica pendula brevior : squamis e spadiceo l. fusco rutilante viridibus. Schenck. gram. 442. t. 10. f. 15.
- Gramen cyperoides spica pendula minus. Beauh. fin. 35.
- Habitat in Europæ frigida paludibus sylvaticis.
30. CAREX *capillaris*. C. spicis pendulis : mascula erecta , femineis oblongis distichis , capsulis nudis acumina. Fl. succ. carn. Flor. dan. t. 168. Leers. herb. n. 720. (excepta varietate β cum icone , quæ est C. parula Scopoli f. sylvatica Schreberi. R.) sub.
- C. spicis femininis raris , pendulis , capsulis ovato-conicis mucronatis. Hall. helv. n. 1394.
- C. spicis ex apice pendulis : masculina & femineis albis. Fl. lapp. 326.
- Cyperoides alpinum , spicis seminiferis pendulis : bis in summo caule. Segu. veron. 3. p. 83. t. 3. f. 1.
- Gramen cyperoides pusillum , spica flavescente. Radb. elys. 1. p. 58. t. 40.
- Habitat in Sueciæ pratis humidis.
31. CAREX *pallefcens*. C. spicis pendulis : masculis erectis , femineis ovatis imbricatis , capsulis confertis obtusis. Fl. succ. carn. gallob. Leers herb. n. 721. t. 15. f. 4. delph.
- C. spicis femininis pendulis , capsulis confertis obtusis. Hall. helv. n. 1393.

C. spicis tribus pedunculatis erectis remotis : mascula & femineis. *Fl. lapp.*

Cyperoides polystachion flavicans, spicis brevibus prope summitatem caulis. *Pluk. alm.* 178. t. 34. f. 5. *Mich. gen.* 61. t. 32. f. 13. *Scheuchz. gram.* 430.

Habitat in Europæ paludibus. 24

Folia pubescentia.

32. **CAREX panicea.** **C.** spicis pedunculatis erectis remotis : femineis linearibus , capsulis obtusiusculis inflatis. *Fl. suec. paris. Leers. herb.* n. 722. t. 15. f. 5. *pal. sil. Fl. dan.* t. 261. *delph. lith.*

C. spicis remotis sessilibus , capsulis globosis. *Fl. lapp.* 333.

C. spica mare , unica pluribusve , feminis petiolatis , erectis , capsulis raris , maximis , ovato-triquetris bidentatis. *Hall. helu.* n. 1405.

Cyperoides foliis caryophyllæis , spicis e rarioribus & tumidi-
oribus vesicis compositis. *Mich. gen.* 61. t. 32. f. 11.
Pluk. alm. 178. t. 91. f. 7.

Habitat in Europæ uliginosis. 24

34. **CALEX Pseudo-cyperus.** **C.** spicis pendulis , pedun-
culis geminatis. *ged. gallob. pal. delph. lugd. burg. lithu.*

C. spicis femininis pendulis , asperis , capsulis longe aristatis ,
retroverfis. *Hall. helu.* n. 1397.

C. spicis pendulis : omnibus femineis : unica androgyna in-
ferne mascula. *Fl. suec. paris.*

C. spicis pluribus in summo caule tenuibus viridibus , præter-
masculam quasi aristatis. *Gmel. sib* 1. p. 141. n. 81.

Cyperoides spica pendula brevior. *Bauh. pin.* 6. *Moris. hist.*
3. p. 242. f. 8. t. 12. f. 5.

Pseudo Cyperus. *Dod. pempt.* 339.

Habitat in Europæ fossis. 24

35. **CAREX cæspitosa.** **C.** spicis erectis cylindricis ternis
subsessilibus : mascula terminali , culmo triquetro. *Fl. suec.*
tub. sil. delph. burg. lugd. lithu.

C. spicis sessilibus approximatis , teretibus folio insidentibus.
Hall. helu. n. 1383. R.

Cyperoides sylvaticum angustifolium , spicis parvis tenui-
bus spadiceo-viridibus. *Scheuchz. gram.* 425. t. 10. f. 11.
Hall. R.

Gramen Cyperoides caryophyllatum , folio longiore & an-
gustiore , spicis sessilibus compactis erectis. *Moris. hist.* 3.
p. 243. f. 8. t. 12. f. 14. *an potius f.* 3.

Habitat in Europæ paludibus turgosis. 24

504 MONOECIA. TRIANDRIA.

36. CAREX *distans*. C. spicis remotissimis subseffilibus
bractea vaginali, capsulis mucronatis. Pollich.
pal. delph.

C. spicis brevissime petiolatis distitis. Hall. helv. n. 1382.

C. culmo subtriquetro; spicis erectis, remotis; semineis
subseffilibus; capsulis acutis. Scop. carn.

C. spicis parvis longe distantibus, T. Rai, hist. 1295 Scheuch.
gram. 431.

Græmen cyperoides gracile alterum, glomeratis corula
spatio distantibus. Moris. hist. 3. p. 243. f. 8. t. 12. f. 18.

Habitat in Europæ australioris paludibus. 24

Differt a C. cæspitosa spicis longe distantibus, singulis
vagina folii, breviter pedunculatis, nec mutantibus; ca-
psulis acuminatis: squama calycina acuta.

* * * * * Spicis sexu distinctis: masculis pluribus.

37. CAREX *acuta*. C. spicis masculis pluribus, femineis sub-
seffilibus, capsulis obtusiusculis. Fl. succ. parisi. tab.
Leer. herb. n. 723. fil. pal. delph. burg. lugd. luth.

C. spicis erectis oblongis sessilibus alternis foliolo florali
brevioribus: superioribus masculis. Hort. cliff. 439. Roy.
lugdb. 74.

4. Carex *acuta*, *nigra*, Roy. ged. 1. p. 224. n. 6. Leer.
l. c. t. 16. f. 1. †.

C. spicis maribus ovatis, femininis gracilibus, erectis,
capsulis triquetris conicis. Hall. helv. n. 1406.

C. culmo subtriquetro; spicis erectis nudis masculis plu-
ribus; femineis subseffilibus, capsulis obtusis. Scop.
carn.

Cyperoides nigro-luteum, vernum, minus. Scheuch.
gram. p. 460.

Carex nigra verna vulgaris. Fl. lapp. 330.

Græmen Cyperoides, foliis caryophyllæis, vulgatissima.
Rai. hist. 1292. Bauh. hist. 2. p. 494.

β. Carex. *acuta*, *ruffa*. Leer. l. c. t. 16. f. 1. Pollich. l. c.

C. spicis masculis ternis, femininis numerosis erectis brevif-
sime petiolatis, capsulis b cornibus. Hall. helv. n. 1404.

Carex spicis masculis superioribus numerosis, femineis
inferioribus: intermedia unica androgyna. Roy. lugdb.

75

Carex maxima, spicis plurimis remotis longis. Fl. lapp. 329.

Cyperoides folus caryophyllæis, spicis habitioribus: squa-
mis curtis. Mich. gen. 62. t. 32. f. 12.

Cyperoides lausfolium, spica ruffa f. caule triangulo,
Scheuch. gram. 438.

MONOECIA. TRIANDRIA. 505

Gramen Cyperoides spica rufa L. caule triangulo. Bauh.
pin. 6.

Habitat in Europa ubique, α in siccioribus, β in aquosis. 24

38. CAREX vesicaria. C. spicis masculis pluribus, femineis pedunculatis, capsulis inflatis acuminatis. *Fl. suec. lib. norv. Leers. herb. n. 724. t. 16. f. 2. I. II. III. pal. sil. Oed. fl. dan. t. 647. carn. delph. burg. lugd. lithu.*

C. spicis plurimis: florescentibus tenuibus, fructiferis crassis. *Fl. lapp.*

C. spicis maribus una pluribusve, femininis erectis, teretibus, capsulis ampullatis, rostratis. *Hall. helu. n. 1401. R.*

C. spicis maribus una pluribus, cylindricis, femininis teretibus, pendulis, capsulis conicis, rostratis, bicornibus. *Hall. helu. n. 1409. R.*

Cyperoides vesicarium, spicis viridantibus & subfuscis. *Scheuch. gram. 450.*

Gramen cyperoides angustifolium, spicis longis erectis. *Bauh. pin. 6.*

β. Carex culmo longissimo, spicis tenuibus remotis. Fl. lapp. 328. Roy. lugdb. 75. Neck. gallab. 381. n. 10. a.

C. (sylvatica) spicis pendulis: mascula erecta, femineis filiformibus pedunculo brevioribus, capsulis acutis distantibus. *Huds. angl. Schreb. spicil. 62. Weber. spicill. p. 26. Distincta species.*

C. spicis femineis pendulis raris praelongis, capsulis rostratis bifidis. *Hall. helu. n. 1395.*

C. (patula) culmo triquetro, spica mascula albicante triquetra, femineis pedunculatis, pendulis; capsulis acutis glabris. *Scop. carn. 2. t. 59. Pollich. pal.*

C. capillaris. *Leys. fl. hall. n. 841. sec. Schreb.*

Icon flor. dan. t. 404. & Leers. herb. t. 15. f. 2. sub C. capillari. β. R.

Gramen Cyperoides sylvarum tenuius spicatum. *Rai. hist. 1295. Moris. hist. 3. p. 243. f. 8. t. 12. f. 9. Scheuch. gram. 418.*

γ. Carex spicis femininis oblongis pedunculatis brevioribus: masculis sessilibus, ultima longissima, capsulis inflatis acuminatis. Roy. lugdb. 76.

Gramen Cyperoides majus præcox, spicis turgidis teretibus flavescentibus. *Moris. hist. 3. p. 242.*

Habitat in Europæ udis sylvaticis.

39. CAREX hirta. C. spicis remotis: masculis pluribus,

506 MONOECIA. TRIANDRIA.

femineis subpedunculatis erectis, capitulis luteis. *Fl. sancti ped. gallob. Leers. herb. n. 7. t. 16. f. 3. pal. fl. flor. dan. n. 379. delph. burg. lithu. lugd.*

C. pilosa, spicis erectis: masculis tribus sessilibus: secundum minore: femineis duabus remotis. *Hort. cliff. 438.*

C. caule foliis capsulisque hirsutis. *Hall. helu. n. 1403.*

Cyperoides polystachion lanuginosum. *Morif. hist. 3. p. 244. f. 3. t. 12. f. 10. Pluk. alm. 178. t. 34. f. 6. Schenck. gram. 478.*

Habitat in Europæ sabulosis. 24

Capitula, villosa, inflata, apice bicornes. Folia exaequalia vaginæ albo-villosa.

1145. LITTORELLA.

Masc. Cal. 4. phyllus, Cor. 4. fida. Stam. longa

Fem. Cal. 0. Cor. obsoleta 3. fida (sub 4. fida. *Syst. Veg. p. 708.*) Stylus longus. Sem. Nucleus.

1. LITTORELLA lacustris. Littorella. *Berg. Ad. Stock. 1768. 341. Oed. dan. 170. Giseke ic. fasc. 1. t. 21.*

Plantago (uniflora) scapo uniflora. *Spec. ed. Fl. suec. Delib. parif. Hall. helu. n. 655.*

Plantago floribus femineis sessilibus ad exortum scapi usque maris. *Juss. oed. 1742. p. 131. t. 7. Guett. stamp.*

Germen junceum, capitulis quatuor longissimis filamentis donatis. *Pluk. alm. 180. t. 35. f. 2. Morif. hist. 3. f. 8. t. 9. f. 30.*

Habitat ad Europæ litora lacuum.

Femina ex eadem radice cum mare.

Sexus & imprimis Fructus videtur hanc a *Plantagine* distinguere, dum fructus non sit Bulbus.

1147. BETULA.

Masc. Cal. 1. phyllus, 3. fidus, 3. florus. Cor. 4. partita.

Fem. Cal. 1. phyllus, sub 3. fidus, 2. florus. Sem. utrinque membrana alata.

1. BETULA alba. B. foliis ovalis acuminatis serratis. B.

MONOECIA. TRIANDRIA. 507

succ. carn. herb. pal. sil. Kniph. cens. 12. n. 15. lugd. lith. burg.

B. foliis cordato lanceolatis ferratis. *Hall. helu. n. 1628.*

B. foliis cordatis ferratis. *Fl. lapp.*

Betula. Bauh. pin. 427. Cam. epit. 69. Blackw. t. 240.

β. *Betula pumila Broccembergensis. Thal. hercyn. 20. Bauh. pin. 427. Mapp. als. 139. Gmel. l. c. β. du Roi. l. c. β.*

Habitat in Europa frigidior. ¶

4. BETULA nana. **B.** foliis orbiculatis crenatis. *Fl. lapp. t. 6. f. 4. Fl. succ. Amæn. acad. 1. p. 4. t. 1. Flor. dan. 91.*

B. *pumila*, foliis subrotundis. *Amm. ad. 9. p. 314. t. 14. ruth. 259.*

Habitat in Alpibus Lapponicis, paludibus Sueciæ, Russiæ, Helvetiæ, Hercyniæ. ¶

6. BETULA Alnus. **B.** pedunculis ramosis. *pal. herb. sil. lith. lugd. burg.*

Alnus. Fl. succ.

α. *Betula glutinosa.*

Alnus foliis glabris, rotundis, rotunde crenatis spongiolis ad nervorum angulos. *Hall. helu. n. 1630. R.*

Alnus foliis obverse ovatis rugosis. *Mill. dict. n. 7. Du Roi, herbk. 1. p. 100. R.*

Alnus rotundifolia glutinosa viridis. Bauh. pin. 428.

Alnus. Cam. epit. 68. Læf. pruss. 10. t. 1. Matth. 140.

β. *Betula incana.*

Alnus foliis mucronatis, acute ferratis, subtus lanuginosis. *Hall. helu. n. 1631. Du Roi l. c. p. 109.*

Alnus secundus. Clus. 12.

Alnus incana & hirsuta. Bauh. hist. 1. p. 2. p. 154.!

Alnus folio incano. Bauh. pin. 428.

Habitat in Europa a Lapponia in Gades. ¶

Limites inter Alnum & Betulam nullos a natura positos esse docuere species 2 & 3.

Plantæ β datur varietas folio laciniato J. B. & alia varietas, quæ est Alnus alpina minor. Bauh. pin. Hall. l. c. R.

508 MONOECIA. TETRANDRIA.

1148. BUXUS.

Malc. Cal. 3. phyllus. Petals 2. Germinis rudimentum.

Fem. Cal. 4. phyllus. Petals 3. Styl. 3. Caps. 3. rostris, 3. locularis. Sem. 2.

1. BUXUS sempervirens. Buxus. Hall. Kets. n. 1610. ca. Lath. ed. t. 4. Kniph. cens. 1. t. 11. l. 1.

a. Buxus arborefcens.

B. (arborefcens) foliis ovatis. Mill. dict. n. 1. Black. 196.

B. arborefcens. Benth. pin. 471.

Buxus. Dod. penult. 782.

Buxus (arborefcens)

B. foliis orbiculatis. Mill. dict. n. 3.

Habitat in Europa australi. ☉

Inter omnes nulla est difformis specifica. D. 196. p. 35. B.

1149. URTICA.

Malc. Cal. 4. phyllus. Cor. o. Neckerum corollae cyathiforme.

Fem. Cal. 2. valvis. Cor. o. Sem. 2. minutum.

* Oppofiti folia.

1. URTICA pilulifera. U. foliis oppofitis ovatis lanceolatis fructiferis globosis. Mart. 1789.

Urtica urens pilulas ferens. Benth. pin. 232.

Urtica 1. doctore dis femine lili. Dod. mon. 4. p. 107.

U. romana. Lob. ic. 521.

Habitat in Europa australi. ☉

3. URTICA Dodartii. U. foliis oppofitis ovatis lanceolatis fructiferis globosis. Fl. 1711. n. 200.

U. altera pilulifera, parietariae facie. Benth. pin. 4.

U. pilulifera, parietariae facie, femine lili. Mart. 1789. 106.

Habitat . . . ☉

Affinis nimium præcedentibus duabus, etiam floribus masculis inferne locatis, sed Folia vix serrata.

6. *URTICA urens*. U. foliis oppositis ovalibus. *pal. Flor. dan. t. 739. lithu. burg. lugd.*

U. androgyna, foliis ovalibus. *fl. suec. paris.*

U. foliis ovatis, amentis cylindraceis androgynis. *fl. lapp.*

U. sexubus sede disjunctis, foliis ovato lanceolatis serratis, spicis oblongis. *Hall. helu. n. 1615.*

U. foliis ovalibus oppositis subduplicato-serratis petiolatis, racemis androgynis. *Neck. gallob.*

U. terria. *Matth. 127. Hall. R.*

U. minor acrior. *Lob. ic. 522. Hall. R.*

U. urens minor. *Bauh. pin. 232.*

U. urens minima. *Dod. pempt. 152.*

Habitat in Europæ cultis. ☉

7. *URTICA dioica*. U. foliis oppositis cordatis, racemis geminis. *Mat. med. 201. gallob. pal. sil. Blackw. 4, 12. mas. Flor. dan. t. 746. lith. burg. lugd.*

U. dioica, foliis oblongo-cordatis. *fl. suec. paris.*

U. foliis cordatis, amentis cylindraceis, sexu distincta. *fl. lapp.*

U. stirpibus sexu distinctis, foliis ovato-lanceolatis, productis serratis. *Hall. helu. n. 1614.*

U. vulgaris major. *Bauh. hist. 3. p. 445.*

U. urens maxima. *Bauh. pin. 232.*

U. urens altera. *Dod. pempt. 151.*

γ. *Urtica rubra*. *Tabern. 234. Gmel. l. c. Hall. R.*

Habitat in Europæ ruderatis. 24

In hac & U. urente Cal. fem. 4. phyllus; foliolis duobus ovato-cordatis duobus oppositis ovatis exiguis, quadruplo minoribus. Leers herb. n. 729.

* * *Alternifolia.*

16. *URTICA canadensis*. U. foliis alternis cordato-ovatis, amentis ramosis distichis erectis.

U. canadensis racemosa mitior s. minus urens. *Moris. hist. 3. p. 434. f. II. t. 25. f. 2.*

Habitat in Canada, Sibiria. 24

Dioica est.

Planta primum mascula, tandem eadem planta flores masculos & femineos produxit. Fabric. helmst. p. 365. R.

§10 MONOECIA. TRIANDRIA

1150. MORUS.

Masc. Cal. 4. partitus. Cor. o.

Fem. Cal. 4. phyllus. Cor. o. Styl. 2. Cal. bicatus. Sem. 1.

1. MORUS alba, M. foliis oblique cordatis lævibus. *parif. carn.*

M. foliis scabris, cordatis & semitrilobis. *Hall. helv. 1611.*

M. alba, fructu minori albo insulso. *Du ham. arb. 3.*

M. fructu albo. *Bauh. pin. 459.*

M. candida. *Dod. pempt. 810.*

Habitat in China, Persia sponte. Gmel. 5

Folia indivisa, nitentia.

2. MORUS nigra. M. foliis cordatis scabris. *parif. Lab. ed. t. 114. Kniph. cent. 3. n. 64. Regn. bot.*

M. fructu nigro. *Bauh. pin. 459. Blackw. t. 126.*

Morus. *Dod. pempt. 810.*

Habitat in Italia maritima; in Persia sponte. 5

Folia subquincqueloba, obtusiuscula, scabra. Dalm.

PENTANDRIA

1152. XANTHIUM.

Masc. Cal. comm. imbricatus. Cor. 1. petala, 5. fida infundibulif. *Recept. paleaceum.*

Fem. Cal. involucrum diphyllum, 2. florum. Cor. o. Drupa ficca, muricata, 2. fida. *Nuckel 2. locularis.*

1. XANTHIUM strumarium. Xanthium caule inermi, foliis ovatis trinervatis. *Fl. succ. parif. carn. pol. f. Regn. bot. Luth.*

2. foliis ovatis, aculeis uncinatis. *Hall. helv. 1611.*

MONOECIA. TRIANDRIA. 312
Xanthium. Fuchs. hist. 579. Blackw. t. 455.

Lappa minor f. *Xanthium discoridis. Bauh. pin. 198.*

Habitat in Europa. ☉

3. **XANTHIUM spinosum** X. spinis ternatis, foliis trilobis.
Kniph. cent. 2. n. 100.

X. spinis triplicibus. *Hort. cliff.*

X. spinosum, atriplicis folio. *Moris. hist. 3. p. 604. f. 15. t. 2. f. 3.*

X. lusitanicum spinosum. *Pluk. alm. 206. t. 239. f. 1. Herm. parad. 246. t. 246. Magn. hort. 208. t. 208.*

Volk. norib. 149. t. 149.

Habitat in Lusitania, Monspelii. ☉

Spinæ trifurcatae sunt stipulae, quarum altera fit fructus.

1153. AMBROSIA.

Masc. *Cal.* communis, 1. phyllus. *Cor.* 1. petalæ.
5. fidæ, infundibuliformes. *Recept.* nudum.

Fem. *Cal.* 1. phyllus, integer, ventre 5. dentato,
1. florus. *Cor.* 0. *Nux* e calyce indurato. 1.
sperma.

1. **AMBROSIA trifida.** A. foliis trilobis ferratis.

A. foliis palmatis: laciniis lanceolatis ferratis. *Hort. cliff. 443. Gron. virg. 188. 147. Roy. lugdb. 86.*

A. virginiana maxima, platini orientalis folio. *Moris. hist. 3. p. 4. f. 6. t. 1. f. 4.*

β. A. gigantea inodora, foliis asperis trifidis. *Rai. suppl. 109.*
Habitat in Virginia, Canada. ☉

4. **AMBROSIA maritima.** A. foliis multifidis, spicis solitariis pilosis subsessilibus. *Kniph. cent. 2. n. 3.*

A. foliis multifidis tripartitis. *Hort. cliff.*

A. maritima. *Bauh. pin. 138.*

Ambrosia. *Dod. pempt. 35.*

Habitat in Hetruriæ, Cappadociæ maritimis arenosis. ☉

Folia bipinnatifida, mollissima, pubescentia. Caules pilosi.

Racemi densi: Floribus masculis subsessilibus confertis, nec

Spica, ut in præcedenti, longo pedunculo insidet.

512 MONOECIA. TRIANDRIA.

1157. AMARANTHUS.

Masc. Cal. 3. f. 5 - phyllus. Cor. o. Sem. 3. f.

Fem. Cal. maris. Cor. o. Styli 3. Caps. 1. locularis
circumscissa. Sem. 1.

* Triandri.

1. AMARANTHUS albus. A. glomerulis triandris axil-
laribus bipartitis, bracteis subulatis, foliis ovatis emar-
ginatis lineatis strictis. Kniph. cent. 11. n. 2.

Blitum maritimum minus, calycibus aculeatis. Till. pis. 25.
Habitat in Pensylvania; migravit nunc in Italiam. ☉

Caulis erectus, tetragonus, lavis Ramis simplicibus.
Folia glabra, emarginata cum setula, subtus margine lacera.
Spicae axillares, brevissimæ, binæ, bipartitæ. Bractea
lanceolata, spinosa, calyce longiores.

4. AMARANTHUS tricolor. A. glomerulis triandris axilla-
ribus subrotundis amplexantibus, foliis lanceolato-ovatis
coloratis. Kniph. cent. 9. n. 6. Knorr. del. hort. 2.
t. A. 3. 4. 5.

A. capitalis subrotundis lateralibus, foliis lanceolatis ac-
minatis. Hort. upf.

A. folio variegato. Bauh. pin. 121.

A. tricolor. Lob. ic. 252.

Habitat in India, Russia. Gmel. it. 1. p. 182. ☉

Folia utrinque viridia: suprema purpureascentia.

11. AMARANTHUS Blitum. A. glomerulis lateralibus
floribus trifidis, foliis ovatis retusis, caule diffuso. Kniph.
cent. 11. n. 3. burg.

A. foliis ovatis emarginatis. Fl. suec. parif.

A. procumbens, foliis ovatis emarginatis. Hall. tab. 1606.

A. floribus triandris, glomeratis; foliis subovato-cuneatis;
caule diffuso. Scop. carn.

A. caule decumbente, foliis subrhomboideis
retusis, alternis spicis triandris. Neck. gallob.

Amaranthus sylvestris & vulgaris. Tournef. parif. 2. p. 24.

Blitum album minus. Rai. hist. 200. Cam. epit. 236. Phil.
phys. 212. f. 2.

MONOECIA. PENTANDRIA. § 13

Blitum rubrum minus. Bauh. pin. 118. Bauh. hist. 2. p. 967. & majus. Bauh. hist.

Blitum rubrum supinum. Lob. ic. 250.

Habitat in Europa temperiore. ☉

12. AMARANTHUS viridis. A. glomerulis triandris : floribus masculis trifidis, foliis ovatis emarginatis, caule erecto. *pal. ged. lithu. burg. lugd.*

Blitum album minus. Bauh. pin. 118. Bauh. hist. 2. p. 967.

Blitum minus album polyspermum folio subrotundo. Sloan. jam. 49. hist. 1. t. 92. f. 1.

Blitum sylvestre spicatum. Tournef. parif. 2. p. 273.

Cararu. Pis. bras. 241. t. 241.

Habitat in Europa, Brasilia. ☉

Planta Caule erecto, rubro striato. Folia oblongo-ovata margine membranaceo rubicundo undato. Spicæ ex glomerulis virides; Flores masculi triphylli.

* * Pentandri.

20. AMARANTHUS hypochondriacus. A. racemis pentandris compositis confertis erectis, foliis ovatis mucronatis.

A. sylvestris maximus novæ angliae, spicis purpureis. Tourn. inst. 245.

Habitat in Virginia. ☉

Caulis viridis. Folia valde acuta; disco ruffa lurida, præsertim subtus disco purpurascens. Spicæ erectæ flosculis purpuratissimis. Stamina 5. flava.

22. AMARANTHUS caudatus. A. racemis pentandris decompositis cylindricis pendulis longissimis.

A. maximus. Bauh. pin. 120.

Blitum majus peruvianum. Clus. hist. 2. p. 81.

Habitat in Peru, Persia, Zeylona, Russia. ☉

23. AMARANTHUS spinosus. A. racemis pentandris cylindricis erectis, axillis spinosis.

A. indicus spinosus, spica herbacea. Herm. lugdb. 31. t. 33.

Blitum spinosum. Rumph. amb. 5. p. 234. t. 83. f. 1.

Habitat in Indiis. ☉

114 MONOECIA. POLYANDRIA.

POLYANDRIA. Stamina ultra VII.

1162. CERATOPHYLLUM.

Masc. Cal. multipartitus. Cor. o. Stam. 16 - 20.

Fem. Cal. multipartitus. Cor. o. Pistil. 1. Styl. o. Sem. 1. nudum.

1. CERATOPHYLLUM *demersum*. C. foliis dichotomo-
bigeminis fructibus trispinosis. *succ. parif. ged. carn. pol.*
fil. lithu. lugd.

C. foliis subspinosis, capsulis tricornibus. *Hall. helv. 2*
1599.

Hydroceratophyllum folio aspero quatuor cornibus armato.
Vaill. aët. 1719. p. 21. t. 2. f. 1.

Dichotophyllum. *Dill. giff. 1. 7. Hall. R.*

Millefolium aquaticum cornutum. *Bauh. hist. 3. p. 784*
non verum. Tournef. parif. p. 290. non vero institut
Hall. R.

Equisetum sub aqua repens, foliis bifurcis. *Laf. pruss. 67.*
t. 12.

Habitat in Europæ fossis majoribus sub aqua.

2. CERATOPHYLLUM *submersum*. C. foliis dichotomo-
trigeminis, fructibus muticis. *herb. Flor. dan.*
t. 310.

Hydroceratophyllum folio lævi octo cornibus armato. *Vaill.*
aët. 1719. p. 21. t. 2. f. 2.

Habitat in Europæ aquis.

Affinis nimiam præcedenti, ut fere varietas.

1163. MYRIOPHYLLUM

Masc. Cal. 4. phyllus. Cor. o. Stam. 3.

Fem. Cal. 4. phyllus. Cor. o. Pist. 4. Styl. o. Sem. 4
nuda.

1. MYRIOPHYLLUM *spicatum*. M. floribus masculis
interrupte spicatis. *Fl. succ. parif. monsp. carn. and. 1719.*

MONOECIA. POLYANDRIA. 513
pal. fil. Flor. dan. t. 681. Kniph. cent. 5. n. 59. lugd. lith. burg.

Pentapteris spica nuda. Hall. helu. n. 993.

Millefolium aquaticum pennatum spicatum. Bauh. pin. 141. prodr. 73. t. 73.

Habitat in Europæ aquis quietis. 24.

Spica aphylla. Lin. Folia perpetuo quaterna. Gmel. R.

Verticilli quatuor superiores flores masculinos habent. Flores feminei infra masculos ponuntur; hic calyx nullus, corolla nulla visibilis erat. Pollich. l. c. R.

Confer. Potamogeton aquaticum pennatum minus, foliolis singularibus latiusculis flosculis subjectis. Moris. hist. 3. p. 622. f. 15. t. 4. f. 7. Rai. angl. 3. p. 151.

2. MYRIOPHYLLUM verticillatum. *M. floribus omnibus verticillatis hermaphroditis. Fl. suec. parif. monsp. pal. fil. lugd.*

M. aquaticum minus. clus. hist. 2. p. 252.

Pentapteris floribus alaribus. Hall. helu. n. 992.

Millefolium aquaticum, flosculis ad foliorum nodos. Bauh. pin. 141.

Habitat in Europæ inundatis. 24

Vidi plantas floribus masculis & femineis distinctis, vidi & rarius floribus hermaphroditis.

Folia quaterna. Flores hermaphroditi Turra.

1164. SAGITTARIA.

Masc. Cal. 3. phyllus. Cor. 3. petala. Filam. fere 24.

Fem. Cal. 3. phyllus. Cor. 3. petala. Pist. multa.

Sem. multa, nuda.

1. SAGITTARIA sagittifolia. *S. foliis sagittatis acutis. Fl. suec. parif. gallob. pal. fil. Fl. dan. t. 172. Knorr. del. hort. 2. t. f. 2. lithu. burg. lugd.*

Sagitta foliis acuminatis. Hall. helu. n. 1183.

Sagitta major. Scop. carn.

Sagitta major & minor. Tabern. p. 743.

Sagitta minor. Dod. pempt. 588.

Sagitta aquatica minor latifolia. Bauh. pin. 194.

β. Sagitta aquatica minor angustifolia. Bauh. pin. 194.

γ. Sagitta aquatica major. Bauh. pin. 94. lithu.

δ. Sagitta aquatica foliis variis. Læf. pruss. 234. t. 74. lithu.

516 MONOECIA. POLYANDRIA:

Gramen bulbosum aquaticum. *Bauh. prodr. 4. lichu.*

Habitat in Europæ, Americæ fluviiis, lacubus argillofis.

An sufficienter distincta SAGITTARIA foliis cordatis, quæ a Moris. hist. 3. f. 15. t. 4. f. 6. delineata.

1166. THELIGONUM.

Masc. Cal. 2. fidus. Cor. 0. Stamina fere 12.

Fem. Cal. 2. fidus. Cor. 0. Pist. 1. Caps. coriacea
1. locularis, 1. sperma.

1. THELIGONUM cynocrambe. Theligonum. *Kaiph. cat. 4. n. 89.*

Parietariæ affinis Cynocrambe dicta, floribus muscosis. *Plat. alm. 280.*

Conocrambe dioscoridis. *Bauh. pin. 122. prodr. 59. t. 59.*

Conocrambe alfine foliis. *Barr. ic. 335. bona.*

Alfine facie planta nova. *Col. phys. 121. t. 36.*

Alfina fœtida fabio columnæ dioscoridea. *Bauh. hist. 3. f. 365. Rai. hist. 206.*

Habitat in India orientali communis, inque Italia, Montepelii, Siciliæ umbrosis. ☉

Caules teretes, diffusi, flexuosi, succulenti. Rami oppositi, divaricati, ex alis foliorum oppositorum. Folia ovata, obtusiuscula, lævia, nervosa, sublineata, oleracea, margine scabra, inferiora opposita, superiora alterna deflexa in Petiolos, longitudine foliorum, connexos utrinque Superius tridentata, membranacea, lata, brevi. Flores masculini, oppositifolii; pedunculis propriis, brevissimis, mixtis. Calyx masculus bivalvis, obtusus, revolutus. Antheræ 13. 19, lineares, rectæ. Feminæ calyx bivalvis, solitarius, sessilis & alaris. Stylus 1, filiformis longus. Stigma simplex. Semen 1, rotundum, insidens receptaculo carnosio, cum quo decidit.

1167. POTERIUM.

Masc. Cal. 4. phyllus. Cor. 4. partita. Stam. 30-50

Fem. Cal. 4. phyllus. Cor. 4. partita. Pist. 2. Baccæ e tubo corollæ indurato.

1. FOTERIUM sanguisorba. P. inermis, caulibus subter-

MONOECIA. POLYANDRIA. 517

lofis. herb. pal. fil. Ludw. ed. t. 187. Sabb. hort. 2. t. 71.

Blackw. t. 413. lugd. burg.

P. inerme, filamentis longissimis. *Dalib. parif.*

Pimpinella polystemon. *Hall. helu. n. 706.*

Sanguisorba officinalis. *Regn. bot. tom. 1.*

Pimpinella Sanguisorba minor hirsuta. *Bauh. pin. 160.*

Pimpinella sanguisorba. *Camm. epit. 777.*

β. Pimpinella Sanguisorba minor lævis. *Bauh. pin. 160.*
Pollich. l. c. Kniph. cent. 1. n. 66.

γ. Pimpinella Sanguisorba inodora. *Bauh. pin. 160. prodr. 84.*
Habitat in Europæ australioris asperis. 24

2. POTERIUM hybridum. **P. inerme**, caulibus teretibus
strictis. *Sabb. hort. 2. t. 72.*

P. inerme filamentis florem vix superantibus. *Roy. lugdb.*

P. agrimonoides. *Hort. upf.*

Pimpinella agrimonoides. *Morif. bles. 294. umb. 57. hist. 3.*
p. 264. f. 8. t. 18. f. 9. Bocc. sic. 57. t. 29.

Pimpinella agrimonoides odorata. *T. Barr. rar. 18. t. 632.*
Habitat Monspeli. 24

3. POTERIUM spinosum. **P. spinis** ramosis. *Sabb. hort. 2.*
t. 73. Kniph. cent. 9. n. 74.

Pimpinella spinosa. *Morif. hist. 3. p. 262. f. 8. t. 18. f. 5.*

Poterio affinis, foliis pimpinellæ, spinosa. *Bauh. pin. 288.*

Stœbe legitima, dioscoridis. *Clus. hist. 2. p. 303.*

Habitat in Creta, Libano. 7

1168. QUERCUS.

Masc. Cal. 5. fidus fere. **Cor. o. Stam. 5. 10.**

Fem. Cal. 1. phyllus, integerrimus, scaber. **Cor. o.**
Styli 2 - 5. Sem. 1. ovatum.

3. QUERCUS Ilex. **Q. foliis** ovato-oblongis indivisis serratisque subtus incanis, cortice integro. *monsp. Scop. earn. Regn bot.*

Ilex oblongo serrato folio. *Bauh. pin. 424.*

Ilex arborea. *Bauh. hist. 1. p. 95. **

Ilex coccigera. *Blackw. t. 186.*

β. Quercus Smilax foliis oblongo-ovatis subtus tomentosis integerrimis. *Roy. lugdb. 81.*

Quercus (Ilex) foliis oblongo-ovatis subtus tomentosis integerrimis. *Mill. dict. n. 13. R.*

Ilex folio angusto non serrato. *Bauh. pin. 424.*

Smiax dalech. *Bauh. hist. 2. p. 101. fig.*

518 MONOECIA. POLYANDRIA.

7. *Quercus* (*gramuntia*) foliis oblonga - ovatis, sinuato-spinosis sessilibus subtus tomentosis, glandibus pedunculatis. *Spec. ed. 2. monsp. Gerard. prou.*

Ilex foliis rotundioribus & spinosis, e luco *gramuntio. May. monsp. 140.*

Habitat in Europa australi. H

4. *QUERCUS Suber. Q.* foliis ovato-oblongis indivisis serratis subtus tomentosis; cortice ramoso fungoso, *Mill. dict. n. 17. Du Roi. harpk. 2. p. 253. Reg. bot.*

Suber latifolium sempervirens. Bauh. pin. 424. Du Ham. arb. 2. t. 80. Blackw. t. 193.

Habitat in Europa australi. H

5. *QUERCUS coccifera. Q.* foliis indivisis spinoso-dentatis utrinque glabris. *Scop. carn.*

Q. foliis ovatis dentato - spinosis, glandibus sessilibus. *Sauv. monsp.*

Ilex aculeata cocciglandiflora. Bauh. pin. 423. Gail. air. 245. t. 53.

Ilex coccifera. Cam. epit. 774.

Habitat in G. Narbonensi, Hispania, Italia, Sicilia, Istria, Oriente, Judæa. H

10. *QUERCUS Esculus. Q.* foliis pinnatifidis: lacinias lanceolatis remotis acutis, postice angulatis. *Wag.*

Q. foliis pinnato - sinuatis lævibus fructibus sessilibus. *Spec. pl. 2.*

Q. parva s. Fagus græcorum & esculus. Bauh. pin. 40.

Habitat in Europa australi. H

Folia pinnato-partita fere ad costam marginatam; lacinias remotis, lanceolatis, acutis, postice imprimis angulatis; subtus etiam nuda.

11. *QUERCUS robur. Q.* foliis deciduis oblongis superne latioribus: sinubus acutioribus: angulis obtusis. *Fl. succ. parif. Scop. carn. pal. sil. Ludw. t. 69. burg. lithu. lapp.*

Q. foliis glabris semipinnatis, pinnis obtusis, superioribus majoribus. *Hall. helu. n. 1662.*

Q. foliis deciduis oblongis, superne latioribus, sinubus acutioribus, angulis obtusis, petiolatis, glandibus sessilibus. *Mill. dict.*

Q. foliis deciduis petiolatis oblongis superne latioribus, sinubus acutioribus, angulis obtusis, glandibus sessilibus. *Du Roi. harpk. 2. p. 237.*

Q. latifolia mas, quæ brevi pediculo est. Bauh. pin. 40. Du Ham. arb. 1.

MONOECIA. POLYANDRIA. 519

β. Quercus (femina) foliis deciduis oblongis obtusis, pinnato-sinuatis, petiolis brevissimis, pedunculis glandularum longissimis. *Mill. dict. n. 2.*

Q. foliis deciduis subpetiolatis oblongis obtusis pinnato-sinuatis : glandibus pedunculatis subsolitariis. *Dur. Roi. l. c. 240.*

Q. cum longo pediculo. *Bauh. pin. 420. Du Ham. arb. 3. Knorr. del. hort. I. t. E. 2.*

Quercus. *Fuchsf. 229. Læf. pruss. t. 69.*

Habitas in Europa. H

Var. β. Fructus omnes pedunculati sunt ; facie magis, quam charactere distinguenda. Varietas hæc adeo constans, ut qui distinguere velit, videtur inniti Ger.

12. QUERCUS Ægylops. **Q.** foliis ovato-oblongis glabris ferrato-dentatis. *Mill. dict. n. 7. ic. t. 215. Scop. carn.*

Q. calyce echinato, glande majore. *Bauh. pin. 420.*

Cerri glans Ægilops aspris. *Bauh. hist. I. p. 77. fructus.*

Habitas in Hispania. H

Noscitur calyce maximo.

13. QUERCUS Carris. **Q.** foliis oblongis lyrato-pinnatifidis : laciniis transversis acutis ; subtus subtomentosis. *Dur. Roi. harpk. 2. p. 259. t. 5. f. 1.*

Q. calyce hispido, glande minore. *Bauh. pin. 420.*

Phagus s. Esculus. *Bauh. hist. I. p. 74. fig.*

Habitas in Hispania, Austria. H

Ramenta multa linearia ex aliis foliorum.

1169. JUGLANS.

Masc. Cal. 1. phyllus, squamiformis. **Cor. 6.** partita.
Filamenta 18.

Fem. Cal. 4. fidus, superus. **Cor. 4.** partita. **Styli 2.**
Drupa nucleo sulcato.

1. JUGLANS regia. **J.** foliolis ovalibus glabris subserratis subæqualibus. *paris. Ludw. eccl. t. 188. Blackw. t. 247. Knorr. del. I. t. N. 7. Kniph. cent. I. n. 47. Regn. bot. lugd. barg.*

J. foliis septenis, ovato-lanceolatis, integerrimis. *Hall. helu. n. 1624.*

Nux juglans s. regia vulgaris. *Bauh. pin. 417.*

Nux juglans. *Dod. pempt. 816.*

320 MONOECIA. POLYANDRIA.

β. Nux juglans fructu maximo. *Bauh. pin. 417.*

γ. Nux juglans fructu tenero & fragili peramine, *Am. pin. 417.*

δ. Nux juglans bifera. *Bauh. pin. 417.*

ε. Nux juglans serotina. *Bauh. pin. 417.*

Habitat in Persia D. Lerche. H

America septentrionalis est Patria juglandum; in Persia hujus hodie primaria; Perside a Regibus Asiae Plin.

1170. FAGUS.

Masc. *Cal. 5. fidus, campanulatus. Cor. a. Stam. 11.*

Fem. *Cal. 4. dentatus. Cor. a. Styl. 3. Caps. (Calyx antea) muricata, 4. valvis. Sem. 2.*

1. *FAGUS Castanea. F. foliis lanceolaris acuminato-strictis subtus nudis. parif. Kniph. cent. 5. n. 31. Reg. bot. lugd. burg.*

Castanea foliis ovatae lanceolatis, serratis, densibus glabris. Hall. helv. n. 1623.

Castanea sativa. Mill. dict. n. 1. Scop. carn. ed. 2. n. 1187.

Castanea sylvestris. Bauh. pin. 419.

Castanea. Cam. epit. 118.

β. *Castanea sativa. Bauh. pin. 418. Blackw. 1. 33. Willd. ic. 1. 84.*

Habitat in Italiae & austriacae Europae montibus H

3. *FAGUS sylvatica. F. foliis ovatis obsolete serratis. Fl. suec. parif. gallob. pal. sil. lugd. burg. lishua.*

Fagus foliis ovato-lanceolatis, ovis undulatis. Hall. helv. n. 1622.

Castanea Fagus. Scop. carn. ed. 2. n. 1188.

Fagus. Bauh. pin. 419. Cam. epit. 12.

Stamina 4, 6, 8, 9, 11, 12. Willd. 1. c. R.

MONOECIA. POLYANDRIA. 521

1171. CARPINUS.

Masc. Cal. 1. phyllus : squama ciliata. **Cor.** o. **Stam.** 10.

Fem. Cal. 1. phyllus : squama ciliata. **Cor.** o. **Germina** 2. **Styli** singulis 2. **Nux** ovata.

1. **CARPINUS Betulus.** C. squamis strobilorum planis. *Fl. suec. paris. Hall. helu. n. 1627. carn. pal. gallob. sil. burg. lugd. lithu.*

C. vulgaris. *Mill. dict.*

Carpinus. *Dod. pempt. 841. Cam. epit. 71.*

Ostria ulmo similis, fructu in umbilicis foliaceis. Bauh. pin. 427.

Fagus sæpium vulgo. Bauh. hist. 1. p. 2. p. 146.

Habitat in Europa, Canada. H

Stamina flos gerit. 8. 16.

2. **CARPINUS ostrea.** C. squamis strobilorum inflatis.

Ostria carpinifolia. Scop. carn.

Ostrea ulmo similis, fructu racemoso lupulo simili. Bauh. pin. 427. Du Ham. arb. 5.

Ostrea italica, carpini folio, fructu longiore s. brevior habitiore. Mich. gen. 223. t. 204. f. 1. 2.

β. *Aceris cognata Ostria dicta flavescent. Pluk. alm. 7. t. 156. f. 1.*

Habitat in Italia, Virgin a.

Fructus feminei Humili similes, sed ex Squamis inflatis, undique clausis, ad basin villosis, includentibus Semen biloculare.

1172. CORYLUS.

Masc. Cal. 1. phyllus, 3. fidus, squamiformis, 1. florus. **Cor.** o. **Stam.** 8.

Fem. Cal. 2. phyllus, lacerus. **Cor.** o. **Styli** 2. **Nux** ovata.

1. **CORYLLUS Avellana.** C. stipulis ovatis obtusis. *Fl. suec. paris. carn. pal. sil. Blackw: t. 293. Kniph. cent. 1. n. 19. lugd. lithu. burg.*

522 MONOECIA. POLYANDRIA.

Corylus sylvestris. *Bauh. pin.* 418. *Hall. helu. n.* 1625.

Avellana nux sylvestris. *Fuchf. hist.* 398.

β. *Corylus sativa*, fructu albo majore, f. vulgaris. *Bauh. pin.* 418. *Kniph. cent.* 1. n. 20.

γ. *Corylus sativa*, fructu rotundo maximo. *Bauh. pin.* 418. *Knorr. del. hort.* 2. t. C. 5.

δ. *Corylus sativa*, fructu oblongo rubente. *Bauh. pin.* 418.

ε. (*maxima*) stipulis oblongis obtusis, ramis erectionibus. *Mili. dict. n.* 2. *Du Roi. herbk.* 1. p. 176.

ς. *Corylus nucibus* in racemum congestis. *Bauh. pin.* 418.

ζ. (*arborescens*) laciniis perianthii pinnatifidis. *Du Roi herbk.* 1. 178.

Habitas in Europæ sepibus. ☞

1173. PLATANUS.

Masc. *Cal.* Amentum globosum- *Cor.* vix manifestum
Antheræ filamentum circumnata.

Fem. *Cal.* Amentum globosum. *Cor.* polypetalum
Styli stigmate recurvo. *Sem.* subrotundo, stylo
mucronata, basi papposa.

1. *PLATANUS orientalis*. P. foliis palmatus.

Platanus. *Bauh. pin.* 431. *Clus. hist.* 1. p. 9. *Dech. hist.* 92. *Bell. itin.* 9.

P. *orientalis verus*. *Park. theatr.* 1427. *Du Ham. arb.* 1. tom 2. t. 33.

P. *orientalis Aceris folio*. *Tournef. Du Ham. arb.* 2. *Du Roi l. c.* 133. R.

Habitas in Asia, Tauro, Macedonia, Atho, Lemno, Creta locis riguis. ☞

Folia variant figura, sed sunt semper glabra; Pedunculi fusci. *Du Roi. R.*

2. *PLATANUS occidentalis*. P. foliis lobatis. *Du Roi herbk.* 2. p. 134.

P. *occidentalis*. *Catesb. car.* 1. p. 156. t. 56.

P. *occidentalis f. virginienfis*. *Park. theatr.* 1421. *Du Ham. arb.* t. 25.

Habitas in America septentrionali. ☞

Folia subtus lanata. *Du Roi l. c.* R.

Confer. Platanus orientalis. Pack. itin. 2. t. 89.

MONOECIA. MONADELPHIA. 523

1174. LIQUIDAMBAR.

Masc. *Cal.* communis 4. phyllus. *Cor.* o. *Filam.* numerosa.

Fem. *Cal.* in globum, 4. phylli. *Cor.* o. *Styli* 2. *Caps.* multæ in globum. 2. valves, polyspermæ.

1. LIQUIDAMBAR *Styracis flava*. L. foliis palmato angulatis : lobis indivisis acutis. *Du Roi herb.* 1. p. 369. *D'actw.* 1. 485.

L. foliis quinquelobatis serratis. *Mill. dict. n.* 1.

Liquidambar. *Bauh. pin.* 502.

Liquidambar arbor f. *Stryaciflua aceris folio.* *Plum alm.* 2. 24. 1. 42. f. 6. *Catesb. car.* 2. p. 65. 1. 65. *Du Ham. arb.* 1. 1. 139.

Syrax aceris folio. *Rai. hist.* 1681. 1799.

Habitat in Virginiae, Mexici uliginosis. H

MONADELPHIA.

1175. PINUS.

Masc. *Cal.* 4. phyllus. *Cor.* o. *Stam.* plurima. *Antheræ* nudæ.

Fem. *Cal.* strobili : squama 2. flora. *Cor.* o. *Pist.* 1. *Nux* ala membranacea excepta.

* *Foliis pluribus ex eadem basi vaginali.*

1. PINUS *sylvestris*. P. foliis geminis : primordialibus solitariis glabris. *Fl. suec. paris. carn. pal. norv. sil.* *Trew. in nov. act. A. N. C. III. app. p.* 452. 1. 15. f. 1. 3. 1. 16. f. 11. 25. *Mill. illustr. lugd. lithu.*

P. foliis binis, convexo-concavis, julis masculis solitariis alaribus. *Hall. helv. n.* 1660. *excluso Mugho. R.*

P. foliis geminis : conis pyramidalis, squamis oblongis obtusis. *Du Roy. herb.* 2. p. 13.

530 MONOECIA. SYNGENESIA.

C. verrucosa. Bauh. hist. 2. p. 222.

Melopepo verrucosus. Tournes. inst. 106.

Habitat . . . ☉

Planta, Flos, Semina, &c. C. Pepontis. Fructus minor cortice duriore fere lignoso.

5. CUCURBITA *Melopepo*. *C. foliis lobatis, caule erecto, pomis depresso nodosis.*

Melopepo clypeiformis. Bauh. pin. 312.

Cucurbita clypeiformis f. Siciliana. Bauh. hist. 2. p. 224.

Habitat . . . ☉

Cirrhi sunt in caule, quamvis non scandat, nec procumbat.

6. CUCURBITA *Citrullus*. *C. foliis multipartitis. cent. 9. n. 28.*

C. seminum margine basi dilatato. Hort. cliff.

Anguria indica. Rumph. amb. 5. p. 400. t. 46.

Anguria Citrullus dicta. Bauh. pin. 312. Blackw. t. 157.

Citrullus folio colocynthidis secto, semine nigro. Bauh. hist. 2. p. 235.

Habitat in Apulia, Calabria, Sicilia. ☉

1193. CUCUMIS.

Masc. Cal. 5. dentatus. Cor. 5. partita. Filam.

Fem. Cal. 5. dentatus. Cor. 5. partita. Pist. 3. fidum.

Pomi semina arguta.

1. CUCUMIS *Colocynthis*. *C. foliis multifidis, pomis globosis glabris. Blackw. t. 441. Sabb. hort. 1. t. 70.*

C. foliis multifidis. Roy. lugdb. 263.

Colocynthis, fructu rotundo, major. Bauh. pin. 313.

Colocynthis. Cam. epit. 982.

Habitat . . . ☉

2. CUCUMIS *prophetarum*. *C. foliis cordatis quinquelobis denticulatis obtusis, pomis globosis - spinoso - muricatis. Jacq. hort. t. 9. Blackw. t. 589.*

Colocynthis pumila echinata arabica, striis duodecim luteis & viridibus variegata. Shaw. afr. 164.

Habitat in Arabia.

5. CUCUMIS *Melo*. *C. foliorum angulis rotundatis, pomis torulosis. Sabb. hort. 1. t. 65. Blackw. t. 329. Kniph. cent. 10. n. 33.*

Melo vulgaris. Bauh. pin. 310.

MONOECIA. SYNGENESIA. 531

Melo. *Bauh. hist.* 2. p. 242.

Habitat in Calmucchia. ☉

8. CUCUMIS *fativus*. C. foliorum angulis rectis, pomis oblongis scabris. *Sabbat. hort.* 1. t. 63. *Blackw.* 1. 4.

Kniph. cent. 10. n. 34. *Regn. bot.*

C. *fativus vulgaris*. *Bauh. pin.* 310.

C. *vulgaris*. *Dod. pempt.* 66.

Habitat ☉

10. CUCUMIS *flexuosus*. C. foliis angulato - sub - lobatis, pomis cylindricis sulcatis curvatis.

C. *flexuosus*. *Bauh. pin.* 310.

C. *oblongus*. *Bauh. hist.* 2. p. 247. *Dod. pempt.* 662.

Habitat in India. ☉

1194. BRYONIA.

Masc. *Cal.* 5. dentatus. *Cor.* 5. partita. *Filam.* 3.

Fem. *Cal.* 5. dentatus. *Cor.* 5. partita. *Stylus* 3! fidus. *Bacca* subglobosa; polysperma.

1. BRYONIA *alba*. B. foliis palmatis utrinque calloso-scabris. *succ. paris. ged. gallob. sil. carn. Blackw.* 1. 533.

a. b. Knorr. del. 1. t. Z. 2. *lugd. lith. burg.*

B. *alba*, baccis nigris. *Bauh. pin.* 297.

Vitis alba, baccis nigris. *Fuchf. hist.* 94. *Cam. epit.* 987.

β. Bryonia *aspera* f. *alba* baccis rubris. *Bauh. pin. Regn. bot.*

B. foliis palmatis scabris. *Hall. helu. n.* 574.

B. *dioica*. *Jacq. austr.* 1. 199. *herb. pal. mæhofs. Blackw.* 1. 37. *Mill. ic.* 71. *Sabb. hort.* 1. 58. *lithu.*

B. *alba*. *Dod. purg.* p. 230.

Habitat in Europa ad pagos & sepes. 24

Baccis nigris semper monoica; α. baccis rubris semper dioica. *Mygind. Jacq. Reich.*

526 MONOECIA. MONADELPHIA.

lævibus bifariam versis. *Scop. carn. norv. Trew. nov. Ad. A. N. C. III. app. 451. t. 14. f. 5. 10. & t. 16. f. 1. 10. fl. Blackw. t. 198. Fl. dan. t. 193. Regn. bot. lund. lishu.*
P. foliis solitariis tetragonis mucronatis. Hall. helv. n. 1656.

P. (picea) foliis solitariis subulatis bifariam versis : conis oblongis pendulis, squamis ovalibus planis marginibus undulatis & laceris. Du Roi herbk. 2. p. 110.

Abies foliis solitariis apice acuminatis. Fl. suec. paris. De Ham. arb. 5.

Abies rubra. Trag. 1117. Hall. R.

Picea. Cam. epit. 47. Math. 97.

β. Picea major prima s. Abies rubra. Bauh. pin. 493.

γ. Abies alba s. femina. Bauh. pin. 505.

Abies minor. Bauh. pin. 493.

Habitas in Europæ, Asiæ borealibus humidiusculis. ἥ
Folia compressa, utrinque subcarinata, supra lutea.

1176. THUJA.

Masc. Cal. Amenti squama. Cor. o. Stam. 5.

Fem. Cal. strobili : squama 2. flora. Cor. o. Pist. 1.
Nux 1. cincta ala marginata.

1. *THUJA occidentalis. T. strobilis lævibus : squamis ob-*
rufis. Blackw. t. 210. Kniph. cent. 1. n. 91.

Thuja theophrasti. Bauh. pin. 488. lishu.

Arbor vitæ. Clus. hist. 1. p. 36. Dod. pempt. 630.

Habitat in Canada, Sibiria subhumidis. ἥ

Rami patuli.

2. *THUJA orientalis. T. strobilis squarrosus : squamis*
acutis.

T. strobilis squarrosus : squamis acuminatis reflexis. Du Roi
herbk. 2. p. 458.

T. strobilis uncinatus : squamis reflexo acuminatis. Mart.
cliff.

Habitat in China. ἥ

Rami erecti.

1203. OSYRIS.

♂. Cal. 3. fidus. Cor. o.
 ♀. Cal. 3. fidus. Cor. o.
 Styl. 1. Bacca 1. sperma.

* *Valeriana dioica. Carex dioica. Salix triandra.*

T E T R A N D R I A.

1210. HIPPOPHAE.

♂. Cal. 2. partitus. Cor. o.
 ♀. Cal. 2. fidus. Cor. o.
 Pist. 1. Bacca 1. sperma
 arillo truncato.

1209. VISCUM.

♂. Cal. 4. partitus. Cor. o.
 ♀. Cal. 4. phyllus. Cor.
 Stigm. obtusum. Bacca
 1. sperma, infera.

1211. MYRICA.

♂. Ament. squama. Cor. o.
 ♀. Ament. squama. Cor. o.
 Styli 2. Bacca 1. sperma.

* *Urtica varia. Morus nigra.*

P E N T A N D R I A.

1220. CANNABIS.

♂. Cal. 5. partit. Cor. o.
 ♀. Cal. 1. phyllus. Cor. o.
 Styl. 2. Nux.

1221. HUMULUS.

♂. Cal. 5. phyllus. Cor.
 nulla.
 ♀. Cal. 1. phyllus. Cor. o.
 Styl. 2. Sem. calyce
 alatum.

1212. PISTACIA.

♂. Cal. 5. fid. Cor. o.
 ♀. Cal. 3. fidus. Cor. o.
 Styli 3. Drupa sicca.

1218. SPINACIA.

♂. Cal. part. Cor. o.
 ♀. Cal. 4. fidus. Cor. o.
 Styl. 4. Sem. 1. calycis
 num.

*as alaternus.**alis pentandra.***ANDRIA.**

♂. Cal. 6. phyllus. Cor. o.
 ♀. Cal. 6. phyllus. Cor. o.
 Styli 3. *Bacca* supera, 3.
 locul.

♂. Cal. 6. phyllus. Cor. o.
 ♀. Cal. & Cor. maris.
 3. fid. *Bacca* infera,
 locul.

* *Rumex Acetosa, Acetofella, aculeata,*

OCTANDRIA.**POPULUS.**

♂. *Argent. lacina.* Cor. o.
Neder. ovum. Stam.
 8-16.

♀. *Argent. lacera.* Cor. o.
 Stigm. 4. fidum. Caps.
 2. valvis. Semen pap-
 posa.

1229. RHODIOLA.

♂. Cal. 4. partit. Cor. 4
 petala.

♀. Cal. 4. partit. Cor. o.
 Pist. 4. Caps. 4. poly-
 permæ.

* *Laurus nobilis, Acer rubrum.*

ENNEANDRIA.**1230. MERCURIALIS.**

♂. Cal. 3. phyll. Cor. o.
 Stam. 9. 12.

♀. Cal. 3. phyllus. Cor. o.
 Styli. 2. Caps. 2. coçca.

D I O E C I A.

133

1231. HYDROCHARIS.

♂. Cal. 3. phyll. Cor. 3.
petala.
♀. Cal. 3. phyllus. Cor. 3.
petala. Styli 6. Caps. in-
fera ; 6. locul.

* *Laurus*, an *omnis* ?

D E C A N D R I A.

1235. CORIARIA.

Cal. 5. phyllus. Cor. 5.
petala.
♀. Cal. & Cor. maris.
Styli 5. Bacca 5. sperma.
petalina.

* *Lychnis dioica*. *Cucubalus Orites*.

Gypsophila paniculata.

D O D E C A N D R I A.

I C O S A N D R I A.

* *Spiræa* *Aruncus*. *Rubus Chamæmorus*,

Myrtus dioica.

* *Thalictrum dioicum*.

M O N A D E L P H I A.

1240. JUNIPERUS.

♂. Ament. Cor. 0. Stam. 3.
♀. Cal. 3. partitus. Cor. 3.
pet. Styli 3. Bacca infera,
sperma, calycina.

1241. TAXUS.

♂. Cal. 4. phyllus Cor. 0.
♀. Cal. 4. phyllus. Cor. 0.
Stigm. 1. Bacca 1. sperma.
recutita.

L 1 4

* *Rhamnus** *Salix pentandra.*

H E X A N D R I A.

1225. SMILAX.

♂. Cal. 6. phyllus. Cor. o.
♀. Cal. 6. phyllus. Cor. o.
Styli 3. Bacca supera, 3.
locul.

1224. TAMUS.

♂. Cal. 6. phyllus. Cor. o.
♀. Cal. & Cor. maris. fl.
3. fid. Bacca inserta, 3.
locul.* *Rumex Acetosa, Acetofella, aculeata.*

O C T A N D R I A.

1228. POPULUS.

♂. Ament. lacinum. Cor. o.
Nectar. ovatum. Stam.
8-16.♀. Ament. lacerum. Cor. o.
Stigm. 4. fidum. Caps.
2. valvis. Semen pap-
pula.

1229. RHODIOLA.

♂. Cal. 4. partit. Cor. 4
petala.♀. Cal. 4. partit. Cor. o.
Pist. 4. Caps. 4. polys-
permæ.* *Laurus nobilis. Aer rubrum.*

E N N E A N D R I A.

1230. MERCURIALIS.

♂. Cal. 3. phyll. Cor. o.
Stam. 9. 12.♀. Cal. 3. phyllus. Cor. o.
Styli. 2. Caps. 2. cocca.

DIOECIA.

133

1231. HYDROCHARIS.

♂. Cal. 3. phyll. Cor. 3.
 ♀. Cal. 3. phyllus. Cor. 3.
 petala. Styli 6. Caps. in-
 fera ; 6. locul.

* *Laurus*, an *omnis* ?

DECANDRIA.

1235. CORIARIA.

Cal. 5. phyllus. Cor. 5.
 petala.
 ♀. Cal. & Cor. maris.
 Styli 5. Bacca 5. sperma.
 petalina.

* *Lychnis dioica*. *Cucubalus Orites*.

Gypsophila paniculata.

DODECANDRIA.

ICOSANDRIA.

* *Spiraea*. *Aruncus*. *Rubus Chamæmorus*.

Myrtus dioica.

* *Thalictrum dioicum*.

MONADELPHIA.

1240. JUNIPERUS.

♂. Ament. Cor. 0. Stam. 3.
 ♀. Cal. 3. partitus. Cor. 3.
 pet. Styli 3. Bacca infera,
 sperma, calycina.

1241. TAXUS.

♂. Cal. 4. phyllus Cor. 0.
 ♀. Cal. 4. phyllus. Cor. 0.
 Stigm. 1. Bacca 1. sperma.
 recutita.

1242. EPHEDRA.

♂. Anther. 2. fil. Cor. 9.
 Stam. 7.
 ♀. Cal. imbricatus. Cor. 9.
 Pist. 2. Bacca 2. sperma.
 calycina.

SYNGENESIA.

1246. RUSCUS.

♂. Cal. 6. phyllus. Cor. 9.
 Stam. 5.
 ♀. Cal. 6. phyllus. Cor. 9.
 Pist. 1. Bacca 3. hyl.
 2. sperma.

† Gnaphalium. dioicum. Bryonia dioica.

DIOECIA.

MONANDRIA.

1198. NAJAS.

Masc. Cal. cylindricus, 2. fidus. Cor. 4. fida.
Filamentum nullum.

Fem. Cal. 0. Cor. 0. Pist. 1. Caps. ovata. 1.
ocularis.

1. NAJAS marina. Fl. suec. monsp.

N. fructu monospermo. Hall. helv. n. 556.

Fluvialis vulgaris latifolia. Vail. ad. 1719. p. 17. t. 1. f. 2.

Fluvialis latifolia (& angustifolia) fructu monospermo.

Mich. gen. 11. t. 8. f. 1. 2.

Fluvialis Pisana, foliis denticulatis. Bauh. hist. 3. p. 579.

Potamogeton fluviatile sargazo simile lucens, foliis margine dentatis. Pluk. alm. 305. t. 216. f. 4. Rai. suppl. 121.

β. Fluvialis species, angusto brevique folio, undequaque spinis infecta. Rai. suppl. 132.

γ. Fluvialis species, folio angusto ad margines denticulis spinosis incincto, flagellum Christi dicta. Rai. suppl. 121.

Habitas in Europæ maribus, Basileæ.

DIANDRIA.

1199. VALISNERIA.

Masc. *Spatha* 2. partita. *Spadix* rectus flosculus.
Cor. 3. partita.

Fem. *Spathæ* 2. fida. 1. flora. *Cal.* 3. partitus,
 superus. *Cor.* 3. petala. *Styli* 3. *Caps.* 1.
 locularis, polysperma.

1. VALLISNERIA *spiralis*. Vallisneria. *Fl. lapp.* 371.
parif. Hall. helu. n. 1309. *norv. lithu.*

♀ *V. palustris*, algæ folio, italica, foliis in summitate
 denticulatis, flore purpurascens. *Mich. gen.* 12. t.
 10. f. 1.

Potamogeton, algæ folio, pisanum. *Boc. mus.* 1. t. 20.

♂ Vallisnerioides palustre, algæ folio, italica, foliis
 summitate tenuissime denticulatis, floribus albis vix conf-
 picuis. *Mich. gen.* 3. t. 10. f. 2.

Habitat in Pisæ aliisque Italiæ fossis, etiam vulgaris in
India orientali & Norvegia.

Folia tenuia apice tenuissime serrata & natantia Cl.
Alstræmer.

1201. SALIX.

Masc. *Amenti squamæ.* *Cor.* 0. *Glandulæ* baseos
 nectarifera.

Fem. *Amenti squamæ.* *Cor.* 0. *Stylus* 2. fidus.
Caps. 1. locularis. 2. valvis. *Sem.* papposa.

* *Foliis glabris serratis.*

1. SALIX *hermaphrodita*. S. foliis serratis glabris, flo-
 ribus hermaphroditis diandris. *Fl. suec.*

Habitat Upsaliæ. H

DIOECIA. MONANDRIA. 539

Convenit cum *S. pentandra* foliis glabris obtuse, glandulis serratis, convolutis, sentis, amentis & singulis ramis, rufescentibus; Amentis villosis, magnitudine & habitu iisdem. Differt vero ab eadem Ramis pallide rufescentibus, nec purpurascentibus; Foliorum pagina superiore venis exaratis, nec venis elevatis; Glandulis ad basin petiolorum extuberantibus; Tempore prius florendi & gallis copiosis, quam sequentis.

2. *SALIX triandra*. S. foliis serratis glabris, floribus triandris. ingr. carn. gallob. herb. pal. sil. lithu.

S. foliis glabris, elliptico-lanceolatis, serratis, stipulis dentatis, julis gracilibus, triandris. Hall. helu. n. 1420.

S. foliis elliptico-lanceolatis utrinque glabris serratis appendiculatis. Gmel. sib. I. p. 155. t. 34. f. 3.

S. folio auriculato splendente, flexilis. Rai. hist. 1637.

Habitat in Helvetia, Sibiria, Germania. &c. ¶

Perquam affinis *S. amygdalinae*.

3. *SALIX pentandra*. S. foliis serratis glabris, floribus pentandris. Fl. suec. sil. burg. lithu.

S. floribus glabris, ovato-lanceolatis, petiolis glandulosis, floribus hexastemonibus. Hall. helu. n. 1639.

S. vulgaris rubens. Bauh. pin. 473.

S. pentandra. Fl. lapp. 370. t. 8. f. 3.

S. folia laureo s. lato glabro odorato. Rai. hist. 1420.

S. spontanea, folio amygdalino, fragilis non auriculata. Bauh. hist. I. p. 214.

Habitat in Europæ paludibus montosis duris. ¶

Stamina 5, 6, 7 gerit. R.

4. *SALIX phylicifolia*. S. foliis serratis glabris lanceolatis: crenis undatis. Fl. lapp. 351. t. 8. f. D. Fl. suec.

β. S. foliis serratis glabris oblongo-ovatis. Fl. lapp. 350. t. 8. f. C.

Habitat in Sueciæ borealibus, circa Herbornam Leers. herb. n. 746. ¶

Variat staminibus 2, 3. rarius 4.

5. *SALIX vitellina*. S. foliis serratis ovatis acutis glabris: serraturis cartilagineis, petiolis calloso-punctatis. Hort. ups. 295. gallob. sil. burg.

Salix foliis lineari-lanceolatis acuminatis, Guett. stamp.

Salix sativa lutea, folio crenato. Bauh. pin. 473.

S. tenuifolia dentata. Munting. ic. 12. R.

S. lutea tenuior sativa viminea. Bauh. hist. I. p. 2. p. 214. Hall. R.

140 DIOECIA. DIANDRIA.

Habitat in Europa temperatiore. ♀

Forse sibi permiffa, nec culta, nec putata, evadit S. alba. Hall. gatt. 8. Cum S. alba jungit. Hallerus.

6. *SALIX amygdalina. S. foliis serratis glabris lanceolatis petiolatis, stipulis trapeziformibus. Fl. suec. parif. pal. fil. gallob. burg.*

S. stipulis serratis, foliis lanceolatis, serratis, glabris. Hall. helv. n. 1636.

S. spontanea fragilis, amygdalino folio, auriculata. Bauh. hist. 1. p. 2. p. 213. Hall.

S. stipulis serratis glabris acuminatis appendiculatis. Fl. lapp. 7.

S. folio amygdalino utrinque virente aurito. Bauh. pin. 473. (sec. Rajum.)

Die Weide mit dem Mandelbaumblatt. Linn. Pfl. syst. 2. p. 406.

Habitat in Europæ sylvis. ♀

Affinis nimium Salici triandra.

7. *SALIX hastata. S. foliis serratis glabris subovatis acutis sessilibus, stipulis subcordatis. Leers. herb.*

S. foliis serratis glabris subovatis sessilibus appendiculatis. Fl. lapp. 334. t. 8. f. G. Fl. suec. 797. 882.

P. foliis glabris, ovatis serratis, stipulis latissimis. Hall. helv. n. 1634.

S. latifolia non hirsuta. Bauh. hist. 1. P. 2. p. 216. Hall. R.

Die spondonformige Weide. Linn. Pflanzenfyst. 2. p. 407.

Habitat in Lapponia, Helvetia, Germania. ♀

9. *SALIX fragilis. S. foliis serratis glabris ovato-lanceolatis, petiolis dentato glandulosis. Fl. lapp. 349. t. 8. f. B. Fl. suec. herb. pal. fil. burg. lithu.*

S. stipulis dentatis, foliis glabris, ovato-lanceolatis, glandis ovatis pilosis. Hall. helv. n. 1638.

S. foliis serratis glabris lanceolatis acuminatis appendiculatis. Roy. lugdb.

S. fragilis. Bauh. pin. 474. prodr. 159. (Dill. sist. 43.) Du Ham. arb. 7.

S. folio amygdalino utrinque virente aurito. Bauh. pin. 473. (sec. Hallerum.)

S. folio longo latoque splendente, fragilis. Rai, angl. 3. p. 448. cantsarb. 143. It. scan. 200.

Habitat in Europæ borealibus. ♀

Arbor procera. Amenta ad basin pedunculis gerunt. foliola

2 f. 3. *sæpius caduca*. Rami digiso concessi ad genicula annotina diffiliunt.

10. *SALIX Babylonica*. S. foliis serratis glabris lanceolatis, ramis pendulis. *Gouan. illustr. p. 77. Gmel. is 3. p. 309. t. 34. f. 2.*

S. *orientalis*, flagellis deorsum pulchre pendentibus. *Tournef. cor. 41. Du Ham. arb. 20.*

S. *arabica*, foliis atriplicis. *Bauh. pin. 475.*

Garb arabum. *Rauw. itin. 183. Dalech. app. 30.*

Habitat in Oriente. ♀

Rami laxi, læves, juncei, purpurascetes. Folia angusta lanceolata, argute serrata, glabra, glauca costa albida; Stipulae rarius subrotundae, minimae: sæpius earum loco utrinque Punctum glandulosum.

11. *SALIX purpurea*. S. foliis serratis glabris lanceolatis inferioribus oppositis. *Fl. succ. Scholl. barb. pal. burg. lithu.*

S. folio longo subluteo non auriculata, viminibus rubris. *Rai. angl. 3. p. 450.*

S. *vulgaris nigricans*, folio non serrato. *Bauh. pin. 473.*

S. *rubra minima fragilis*, folio longo angusto. *Bauh. hist. I. p. 215.*

S. *monandra*. *Arduin mem. I. sp. 67. t. II. sec. Lin.*

Habitat in Europæ australioribus. ♀

Rami sæpe sanguineo - corallini. Gemmae univales promunt Folia 3, transversim cauli applicata (contra in plerisque) ex 2 exterioribus transversim positis Rami tenacissimi omnium.

12. *SALIX Helix*. S. foliis serratis glabris lanceolato-linearibus: superioribus oppositis obliquis. *paris. gallob. herb. pal. sil. burg.*

S. *monostemon*, foliis glabris; lineari-lanceolatis, serratis, superne conjugatis, julis tomentosis. *Hall. helv. n. 1640. R.*

S. *purpurea*. *Scop. carn.*

S. *monandra*. *Arduin. mem. I. sp. 67. t. II. secundum Hallerum. R.*

S. *humilior*, foliis angustis subcæruleis ex adverso binis. *Rai. hist. 1421. cant. 144. angl. 3. p. 448.*

S. *tenuior*, folio minore utrinque glabro, fragilis. *Bauh. hist. I. p. 213. f. 2.*

S. *Helice theophrasti*. *Dalech. hist. 277.*

Habitat in Europa australiori ad sepes, locis aquosis. ♀

342. **DIŌECIA. DIANDRIA**

Rami angulofi. Folia adulta fubtus glauca, fuperiora oppofita.
Stamen unicum. An fit a *S. purpurea* Lin. diverfa, du-
bitant Hallerus & du Roi. R.

13. **SALIX myrsinites.** S. foliis ferratis glabris ovatis
venofis. *Fl. lapp.* 353. t. 8. f. F. & t. 7. f. 6. *Fl.*
suec. herb.

S. foliis ovato-lanceolatis, venofis, ferratis, julis toment-
ofis. *Hall. helu. n.* 1645.

S. alpina pumila myrsinites. *Pont. comp.* 149.

S. alpina, foliis anguftioribus fplendentibus ferratis. *Schreuch.*
alp. 340.

S. alpina pumila repens, foliis oblongis exiguis: fuperas
fplendentibus, inferne cinereis creberrimis & tenuiffimis
crenis. *Till. pis.* 151.

Habitat in Alpibus Lapponiæ, Helvetiæ, Italiæ, in Ger-
mania. ¶

14. **SALIX arbuscula.** S. foliis fuferratis glabris fubdi-
phanis fubtus glaucis; caule fuffruticofa, *Fl. suec. Jacq.*
auftr. t. 408.

S. foliis ferratis glabris obovatis. *Fl. lapp.* 352. t. 8. f. E.

S. foliis ovato-lanceolatis, teneris, ferratis, *Stipulis lan-*
ceolatis. *Hall. helu. n.* 1647. R.

S. foliis integris glabris ovatis confertis pellucidis. *Fl. lapp.*

7. S. foliis ferratis glabris lanceolatis utrinque serratis. *Fl.*
lapp. t. 8 f. M.

Habitat in Lapponiæ campis arenofis. ¶

In Horto mirè mutata, altitudine humana, foliis creffo-
ribus aliena.

15. **SALIX herbacea.** S. foliis ferratis glabris orbiculatis.
355. t. 8. f. H. & t. 7. f. 3. 4. *Fl. suec. Jacq. vind. Fl.*
dan. t. 117.

S. foliis orbiculatis, ferratis, glabris, julis paucifloris. *Hall.*
helu. n. 1649.

S. saxatilis minima. *Bauh. pin.* 474. *prodr.* 159.

S. alpina lucida repens, alni rotundæ folio. *Bocc. maf. 2.*
p. 19. t. 1. f. ult. *Rai. suppl.* 12. *angl.* 3. *p.* 448.

Herba facie pyrolæ. *Mart. spitsb.* 47. t. G. f. B.

Habitat in Alpibus Lapponiæ, Helvetiæ. ¶

Arbor omnium minima.

16. **SALIX retusa.** S. foliis fuferratis glabris obovatis
obtusiffimis. *Gouan. illuftr.* p. 76.

S. foliis glabris ovatis, nitentibus, julis paucifloris. *Hall.*
helu. n. 1648.

S. foliis lævibus ovatis, spica rarissima. Hall. opusc. 301.

S. alpina angustifolia repens non incana. Bauh. pin. 474. prodr. 159.

S. alpina, serpilli folio lucido. Bocc. musc. 2. p. 18. t. 1.

S. pusilla humilis. Cam. epit. 104.

Habitat in Alpibus Helvetiæ, Austriæ, Italiæ. 2

Parvitate certat cum S. herbacea. Folia venis parallelis.

** * Foliis glabris integerrimis.*

17. SALIX reticulata. *S. foliis integerrimis glabris ovatis obtusis. Jacq. vind. 296. Gouan. illustr. p. 77. Fl. dan. t. 212.*

S. foliis integris glabris ovatis subtus reticulatis. Fl. lapp. 359. t. 8. f. L. & t. 7. f. 1. 2. Fl. suec. 801. 888.

S. foliis rotundis, integerrimis, subtus tomentosis, julis gracilibus longe petiolatis. Hall. helu. n. 1650. R.

S. pumila, folio rotundo. Bauh. hist. 1. p. 12. Scheuch. alp. 43. 340. Rai. hist. 1423.

Habitat in Alpibus Lapponiæ, Helvetiæ. 2

Gemmæ promunt Folia 3, ovata, obtusa, petiolata, integerrima, obsolete pilosa: supra viridia rugosa; subtus glauca reticulata: venis primum rubris, demum viridibus. Spica ex eadem gemma.

18. SALIX myrtilloides. *S. foliis integris glabris ovatis acutis. Pall. it 3. p. 33. Gouan. illustr. p. 77.*

S. foliis ovato-lanceolatis glabris, obscure serratis, venosis. Hall. helu. n. 1646 sec. Gouan. R.

S. foliis integris glabris ovatis alternis. Fl. lapp. 357. t. 8. f. f. K. Fl. suec. 804. 889.

Habitat in Suecia septentrionali. 7

19. SALIX glauca. *S. foliis integerrimis, subtus tenuissime villosis ovato-oblongis. Fl. lapp. 363. t. 7. f. 5. & t. 8. f. P. Fl. suec.*

S. alpina pyrenaica. Bauh. pin. 474. prodr. 159. Burf. XXIV. 120.

Habitat in Alpibus Lapponicis & Pyrenaicis. Sauv. 7

** * * Foliis integerrimis, villosis.*

20. SALIX aurita. *S. foliis integerrimis utrinque villosis obovatis appendiculatis. Fl. lapp. 369. t. 8. f. Y. gallob. pal. lithu.*

S. foliis rugosis, reticulatis, integerrimis, inferne tomentosis, julis ovatis. Hall. helu. n. 1652.

S. folio rotundo minore. Dill. app. 37. Rai. angl. 3. p. 450.

§44 DIOECIA DIANDRIA

Habitat in Europæ borealis sylvis. 7

21. *SALIX lanata*. S. foliis utrinque lanatis subrotundis acutis. *Fl. lapp.* 368. t. 8. f. 2. & 3. *Fl. suec.* S. foliis integerrimis, ovato-lanceolatis, subrotundis, foliis ovatis. *Hall. helu. n.* 1651. R.

S. humilis latifolia erecta. *Bauh. pin.* 474. *prodr.* 159. *Rai. hist.* 1426.

β. *Salix depressa* foliis integris subtus villosis lanceolato-ovatis utrinque acutis. *Fl. lapp.* 361. t. 8. f. N. *fl. suec.* 2. n. 899. *

Habitat in Alpibus Lapponicis. 7

Folia conferta, ovata, utrinque lanata, integerrima, subsessilia.

Varietatem β. eandem esse cultura docuit.

22. *SALIX Lapponum*. S. foliis integerrimis hirsutis lanceolatis. *Fl. lapp.* 366. t. 8. f. T. *fl. suec. Gouan. illust.* p. 78.

S. foliis crassis, elliptico-lanceolatis, integerrimis, utrinque sericeis. *Hall. helu. n.* 1643. *secundum Gouan.* R.

Habitat in Alpibus Lapponiæ ubique, Helvetiæ. 7

23. *SALIX arenaria*. S. foliis integris ovatis acutis: supra subvillosis, subtus tomentosis. *Fl. suec. gallob. pal. Fl. dan.* t. 197. *lithu.*

S. foliis integris subtus villosis ovatis acutis; *Fl. lapp.* 361. t. 8. f. O. Q.

S. foliis integerrimis, ovato-lanceolatis, subtus sericeis, julis tomentosis. *Hall. helu. n.* 1642. t. 14. R.

S. foliis integris ovatis acutis; supra subvillosis, stipulis ovatis acutis. *Du Roi herb.* 2. p. 408.

S. foliis integris utrinque hirsutis lanceolatis. *Hall. helu. ed. pr.* 155. t. 5. f. 2.

S. pumila, foliis utrinque candicantibus & lanuginosis. *Bauh. pin.* 474. *Rai. pin.* 474. *Rai. angl.* 3. p. 447. *Gmel. sib. I.* p. 164. t. 36. f. 1. *Du Ham. arb.* 10.

Habitat in Europæ paludibus. 24. 7

24. *SALIX incubacea*. S. foliis integerrimis lanceolatis: subtus villosis nitidis, stipulis ovatis acutis. *Fl. suec. Scholl. barb.*

S. foliis ovato-lanceolatis integerrimis, ramis decumbentibus, radice reptatrice. *Dalib. paris.*

S. foliis integris ovatis aliquot linearum longis, caulibus vix assurgentibus. *Guett. stamp.*

S. pumila, foliis ellipticis integerrimis, subtus glaucis, spica rotundiore. *Hall. helu. n.* 1. p. 153.

Habitat

Habitat in Europæ pascuis duriusculis, uliginosis. 24. ♀

25. *SALIX repens*. S. foliis integerrimis lanceolatis utrinque nudiusculis, caule repente. *Fl. suec. Scop. carn.*

S. alpina pumila rotundifolia repens, inferne subcinerea *Bauh. pin. 474. Rai. angl. 3. p. 448.*

Salix pumila latifolia L. *Clus. hist. 1. p. 83.*

Habitat inter montes Sueciæ, locis humidis. 24. ♀

Caulis crassitie digiti. Rami virgati, decumbentes. Ramuli dispersi villis albis. Folia ovata s. ovato-oblonga, integerrima, utrinque glabra, subtus glauca, petiolata stipulis nullis: infima opposita transversa & parum pilosa. Amenta lateralia. Feminis capsulae rufescentes.

26. *SALIX fusca*. S. foliis integerrimis ovatis; subtus pubescens. *Jacq. austr. 1. 409.*

S. foliis integerrimis ovatis; subtus villosis nitidis. *Sp. pl. Fl. lapp. 365. t. 8. f. R. Fl. suec. Jacq. vind. gallob. Scholl. herb.*

S. pumilla folio utrinque glabro. *Bauh. hist. 1. p. 217.*

Habitat in Europæ pascuis humidis. 24. ♀

Gemma univalvis, bipartita; Amentum absque foliis.

Affines sunt S. arenaria, incubacea, repens, fusca.

27. *SALIX rosmarinifolia*. S. foliis integerrimis lanceolato-linearibus strictis sessilibus subtus tomentosis. *Roy. lugdb. Fl. suec. lithu.*

S. procumbens; foliis lanceolatis, subtus sericeis. *Hall. helu. n. 1644. R.*

S. foliis lanceolato-linearibus, subtus villosis splendentibus. *Gort. ingr. sec. Hall. R.*

S. pumila angustifolia; inferne lanuginosa. *Bauh. hist. 1. p. 212. Hall. R.*

S. humilis angustifolia. *Bauh. pin. 474.*

S. humilis repens angustifolia. *Lob. ic. 2. p. 137. Bauh. hist. 1. p. 214.*

Habitat in Europæ campis depressis. 24. ♀.

Folia subtus nitida, sericea, villosa.

* * * * Foliis subserratis; villosis.

28. *SALIX caprea*. S. foliis ovatis rugosis; subtus tomentosis, undatis, superne denticulatis. *Fl. suec. paris. gallob. fil. Fl. dan. 1. 245. burg. lithu.*

S. foliis subcrenatis utrinque villosis ovato oblongis. *Fl. lapp. 365. t. 8. f. S. Dalib. paris.*

S. foliis ovatis, rugosis, subtus reticulatis, tomentosis, junctis ovatis, stipulis amplexicaulibus serr. *Hall. helu. n. 1643.*

Linnæi spec. plant. Europ. Pars II.

M. n.

§46 MONOECIA. DIANDRIA.

S. caprea latifolia. Tabern. 1038. Hall. R.

S. latifolia inferne hirsuta. Bauh. hist. 1. p. 215. Hall. R.

S. latifolia rotunda. Bauh. pin. 474. Rai. angl. 3. p. 449. Du Ham. arb. 28.

β. *S. foliis oblongis subtus villosis : inferioribus crenatis, superioribus integris. Fl. lapp. 367. t. 8. f. U.*

γ. *S. foliis elliptico-lanceolatis subtus sericeis appendiculatis. Gmel. sib. 1. p. 163. n. 19.*

Habitat in Europæ siccis. H

Gemmæ inferiores Folia, superiores Amenta non foliacea producunt. H

29. *SALIX viminalis. S. foliis subintegerrimis lanceolato-linearibus longissimis acutis subtus sericeis, ramis virgatis. Fl. suec. paris. gallob. pal. sil. burg.*

S. foliis prælongis, obscure dentatis; subtus tomentosis & albicantibus. Hall. helu. n. 1641.

S. foliis exstipulatis, lanceolato-linearibus, dentatis, rugosis, intra tomentosis; marginibus reflexis. Scop. carn.

S. foliis undulatis lineari-lanceolatis subtus tomentosis. Roy. lugdb. 84.

S. folio longissimo angustissimo utrinque albido. Bauh. pin. 474. Du Ham. arb. 4.

S. folio longissimo. Rai. angl. 3. p. 450.

S. foliis angustis & longissimis crispis : subtus albicantibus. Bauh. hist. 1. p. 212.

β. *S. foliis ex lineari lanceolatis integris subtus incanis. Gmel. sib. 1. p. 162. n. 16.*

Habitat in Europa ad pagos. H

Frutex junceus, altissimus. Gemmæ foliaceæ & floriferae distinctæ ut in caprea. Folia ante explicationem in hæc revoluta.

30. *SALIX cinerea. S. foliis subserratis oblongo-ovatis : subtus subvillosis, stipulis dimidiato-cordatis. Fl. suec. Leers. herb.*

S. foliis integris glabris lanceolato-ovatis. Fl. lapp. 358.

Habitat in Europæ nemoribus paludosis subhumidis. H

Amenta florum fusca, infra folia collocata, Pedunculo Foliolis aliquot minimis, lanceolatis. Stipulæ dimidiato cordatæ, utrinque ferratæ glandulis tribus.

31. *SALIX alba. S. foliis lanceolatis acuminatis serratis utrinque pubescentibus : ferraturis infimis glandulosis. Fl. suec. paris. gallob. carn. pal. sil. Blackw. t. 327. Reg. burg. lithu.*

DIOECIA. TRIANDRIA. 547

S. arborea, foliis ellipticis, lanceolatis, subtus sericeis, dentibus crassescens. *Hall. helv. n. 1635.*

S. vulgaris alba arborescens. *Bauh. pin. 473.*

Habitat ad pagos & urbes Europæ. ♀

Est Species maxima cum Salice fragili.

Species hujus generis difficillime extricantur. Solum pda lustre, arenosum, alpestre, calidum multavit mira metamorphosi species, ut de iisdem hæsitarint sæpius Botanici; accessit Descriptio ubique imperfecta & barbara. Incipiendâ itaque harum historia e novo; quæ sit, si rejiciantur veterum nomina, ne confundant synonyma dubia certas plantas; & in Floris & Hædioricis describantur. 1. Gemmæ univales s. multivales. 2. Foliatio intra gemmas sita & plicatione. 3. Amenta ex eadem vel distincta gemma, supra aut infra folia. 4. Folia serrata aut integerrima: paginis nudis aut villosis. 5. Staminum humerus. 6. Arbores, frutices; repentes.

T R I A N D R I A.

1202. E M P E T R U M.

Masc. Cal. 3. partitus. Cor. 3. petala. Stam. longa.

Fem. Cal. & Cor. maris. Styli 9. Bacca 9. sperma:

1. EMPETRUM album. E. erectum.

E. lusitanicum, fructu albo. *Tournef. inst. 579.*

Erica erecta, baccis candidis. *Bauh. pin. 486.*

Erica coris folio X. *Clus. hist. 1. p. 45.*

Weisse Rauchbeere. *Linn. Pfl. Syst. 4. p. 567.*

Habitat in Lusitania ♀

Differt a sequenti Ramulis pubescentibus. Foliis longioribus supra scabriusculis, subtus canaliculatis.

2. EMPETRUM nigrum. E. procumbens. *Fl. suec. Jacq. vind. Reyg. ged. norv. fil. Mill. illustr. lithu.*

E. procumbens, foliis ovato lanceolatis obtusis. *Hall. helv. n. 1605.*

Erica baccifera procumbens nigra. *Bauh. pin. 436.*

Erica baccifera. *Clus. pan. 29. Lob. ic. 621. Bauh. hist. 1. p. 526.*

Habitat in Europæ frigidissimæ montis paludosis. ♀

M m 2

48 DIOECIA. TETRANDRIA.

Plantas hermaphroditas nuper habuit Jacquin ; ego quondam unicam vidi Upsaliae , nec deinde myriadibus dioicis unicam.

1203. O S Y R I S.

Masc. *Cal.* 3. fidus. *Cor.* o.

Fem. *Cal.* 3. fidus. *Cor.* o. *Stylus* 1. *Stigma* subrotundum. *Bacca* 1. *ocularis.*

1. *OSYRIS alba. monsp. carn.*

O. foliis linearibus acutis. *Læfl. it.*

O. frutescens baccifera. *Bauh. pin.* 212.

Casia poetica monspeliensium. Cam. epit. 26. *Lob. ic.* 43.

Casia latinorum. Alp. exot. 41.

Casia monspelii dicta. Gesn. epit. 50.

Habitat in Italia , Hispania , Monspelii , Libano , Carniolia. Fr

Femina potius hermaphrodita , cum staminibus gaudet , tamen sine mare sterilis.

In mare vidit rudimenta styli & Germinis Scopoli. l. c. R.

T E T R A N D R I A.

1209. V I S C U M.

Masc. *Cal.* 4. partitus. *Cor.* o. *Filamenta* o. *Antheræ* calyci adnatæ.

Fem. *Cal.* 3. phyllus , superus. *Stylus* o. *Cor.* o. *Bacca* 1. sperma. *Sem.* cordatum.

1. *VISCUM album. V.* foliis lanceolatis obtusis , caule dichotomo , spicis axillaribus. *Hall. helu. n.* 1609. *ged. carn. pal. sil. lith. lugd. burg.*

Viscum. Fl. suec. paris. Du Ham. arb. 356. *Blackw. t.* 184.

V. baccis albis. *Bauh. pin.* 423.

Viscum. Cam. epit. 555.

Habitas in Europæ arboribus parasitica. Fr

1210. HIPPOPHAE.

Masc. *Cal.* 2. *partitus.* *Cor.* 0.

Fem. *Cal.* 2. *fidus.* *Cor.* 0. *Styl.* 1. *Bacca monosperma.*

1. HIPPOPHAE *Ramnoïdes.* H. foliis lanceolatis. *norv. Fl. dan. t. 265. Du Ham. arb. 2. t. 49. lugd.*

Hippophaës. Fl. suec.

H. foliis linearibus, subtus rubiginosis. *Hall. herb. n. 1603.*

Osyris rhamnoides. Scop. carn.

Rhamnus salicis folio angustiore, fructu flavescente. Bauh. pin. 477.

Rhamni species. Cam. epit. 81.

Rhamnoides florifera, salicis folio. Tournef. cor. 53.

Rhamnus. 2. Clus. hist. 110. R.

Oeaster germanicus. Cord. hist. 3. c. 24. p. 186. Hall. R.

Habitat in Europæ arenosis. H.

Folia adspersa squamis confertis, peltatis, ciliatis, pallidis. Flores Masculi solitarii, infra folia, inter ramum & Gemmam univalvem, persistentem, longitudine floris. Feminei solitarii, sessiles, in ala foliorum infimorum.

1211. MYRICA.

Masc. *Amenti squama lunata. Cor.* 0.

Fem. *Amenti squama lunata. Cor.* 0. *Styli* 2. *Bacca* 1. *sperma.*

1. MYRICA *Gale.* M. foliis lanceolatis subserratis, caule suffrutic. *ingr. Flor. dan. t. 327. Kniph. cent. 9. n. 70. 71.*

M. foliis lanceolatis, fructu sicco. *Fl. suec. parif.*

Rhus myrthifolia belgica. Bauh. pin. 414.

Chamæleagnus. Dod. pempt. 780. app.

Habitat in Europæ, Americæ septentrionalis uliginosis. H.

PENTANDRIA.

1212. PISTACIA.

Masc. Amenti. Cal. 5. fidus. Cor. 0.

Fem. distincta. Cal. 3. fidus. Cor. 0. Styli 3. Drupa monosperma.

1. PISTACIA *trifolia*. P. foliis subternatis simplicibusque. *meth.*

Pistacium mas siculum, folio nigricante. *Bocc. mus. 2. p. 139. t. 93.*

Terebinthus f. Pistacia trifolia. *Tournef. inst. 580.*

Habitat in Sicilia. $\bar{\eta}$

Folia pinnata, ternata, demum simplicia in eodem ramo; ubi simplicia, sunt majora ovato-subrotunda; ubi ternata, sunt foliola lateralia parva.

2. PISTACIA *narbonensis*. P. foliis pinnatis ternatisque suborbiculatis.

P. foliis sæpius quinatis orbiculatis. *Sauv. monsp.*

Terebinthus peregrina, fructu majore pistacis simili eduli. *Bauh. pin. 400. Tournef. inst. 579.*

Terebinthus indica major, fructu rotundo. *Bauh. hist. 1. p. 277.*

Terebinthus major, pistaciæ folio. *Lob. adv. 412.*

Habitat Monspeli, in Persia, Mesopotamia, Armenia. $\bar{\eta}$

2. PISTACIA *vera*. P. foliis impari-pinnatis: foliolis subovatis recurvis. *Blackw. 1. 461. Kniph. cent. 2. n. 58. 59. Regn. bot.*

Pistacia Terebinthus. *Mill. dict. n. 1.*

P. peregrina fructu racemoso f. Terebinthina indica. *Bauh. pin. 401.*

Pistacia. *Bauh. hist. 1. p. 275.*

Terebinthus indica theophrasti, Pistacia dioscoridis. *Lob. adv. 413. Tournef. inst. 380.*

Habitat in Persia, Arabia, Syria, India. $\bar{\eta}$

$\bar{\eta}$. Syria in Italiam intulit L. Vitellius, postea Cæsar factus, cum Legatus in ea provincia esset. *Plin. XIII. 5.*

MONOECIA. PENTANDRIA. 551

4. **PISTACIA** *Terebinthus*. P. foliis impari - pinnatis : foliolis ovato-lanceolatis. *monsp. carn. Blackw. t. 478.*

P. vera. *Mill. dict. n. 4.*

Terebinthus vulgaris. Bauh. pin. 400. Tournef. inst. 579.

Terebinthus. Clus. hist. 1. p. 15. Dod. pempt. 871.

Habitat in Europa australi, Africa boreali, India. ¶

Confer. Terebinthus peregrina, fructu minore cæruleo e Julii. C. B. 296.

5. **PISTACIA** *Lentiscus*. P. foliis abrupte pinnatis : foliolis lanceolatis. *monsp. Ludw. ed. t. 41. Blackw. t. 195.*

Kniph. cent. 1. n. 64. femina. cent. 5. n. 67. mas. Regn. bot.

Lentiscus vulgaris. Bauh. pinn. 399. Tournef. inst. 580.

Lentiscus. Clus. hist. 1. p. 14. Dod. pempt. 871.

Der Mastixbaum. Linn. Pfl. Syst. 2. p. 432.

Habitat in Hispania, Lusitania, Italia, Palæstina. ¶

Confer. Lentiscus ex Chio, ex qua fluit mastiche. Tournef. it. 1. p. 295.

Lentiscus mas demum flores hermaphroditos triandros pentagynos & fructus produxit. Fabr. helmst. 433.

1218. SPINACIA.

Masc. Cal. 5. partitus. Cor. 0.

Fem. Cal. 4. fidus. Cor. c. Styli 4. Sem. 1. intra calycem induratum.

1. **SPINACIA** *oleracea*. S. fructibus sessilibus.

S. foliis sagittatis, seminibus aculeatis. *Mill. dict.*

Spinacia mas. Dalech. hist. 543. ♀. Blackw. t. 49.

Spinacia semine spinoso. Moris. hist. 2. p. 598.

Lapathum hortense f. spinacia semine spinoso. Bauh. pin. 114.

Spinacia femina. Dalech. hist. 543. ♂

β. *Lapathum hortense f. Spinacia semine non spinoso. Bauh. pin. 115.*

Spinacia (glabra) foliis oblongo-ovatis, seminibus glabris. Mill. dict. n. 2.

Habitat ☉

152 DIOEGIA. PENTANDRIA.

1220. CANNABIS.

Masc. Cal. 5. partitus. Cor. o.

Fem. Cal. 1. phyllus, integer, latere hians.
Cor. o. Styl. 2. Nux 2. valvis, intra calycem
clausum.

1. CANNABIS *sativa*. C. foliis digitatis. *paris. hall. helu.*
n. 1616. *ged. sil. Blackw. t. 322. a. b. Regn. bot.*

C. *sativa*. *Bauh. pin. 320. ♀*

C. *mas*. *Dalech. hist. 497. ♀*

C. *erratica*. *Bauh. pin. 320. ♂*

C. *Femina*. *Dalech. hist. 497. ♂*

Habitas in Persia, ☉

1221. HUMULUS.

Masc. Cal. 5. phyllus. Cor. o.

Fem. Cal. 1. phyllus, oblique patens, integer.
Cor. o. Styl. 2. Sem. 1. intra calycem fo-
liatum.

1. HUMULUS *Lupulus*. *Humulus. paris. ged. gallo. pal.*
sil. Kniph. cent. 9. n. 49. Blackw. t. 536. a. & b. Knorr.
del. hort. 2. t. L. 5. mas. Mill. illustr. Regn. bot. lugd.
lith. burg.

Lupulus, *Hall. helu. n. 1618. Mill. dict.*

Lupulus mas. *Bauh. pin. 298. ♀*

Lupulus salictarius. *Fuchs. hist. 124.*

Lupulus femina. *Bauh. pin. 298. Cam. epit. 954. ♂*

Habitat in Europæ sepibus & ad radices montium. 2

Murmur electricum, quasi remotissimum tonitru vento
exagitante humili palos, quia?

H E X A N D R I A.

1224. T A M U S.

Masc. *Cal.* 6. partitus. *Cor.* o.

Fem. *Cal.* 6. partitus. *Cor.* q. *Stylus* 3. *fidus*.
Bacca 3. *locularis*, *infera*. *Sem.* 2.

1. TAMUS *communis*. T. foliis cordatis indivisis. *Hall. helu. n.* 1620. *carn. gallab. Blackw. t.* 457. *Sabb. hort. i.* t. 59. *mas. Regn. bot. lugd. burg.*

Bryonia lævis f. *nigra racemosa*, *cujus baccae rufescunt* f. *nigrescunt*. *Bauh. pin.* 297.

Bryonia sylvestris baccifera. *Bauh. prodr.* 135. ♀.

Bryonia lævis f. *nigra racemosa*. *Bauh. pin.* 297. ♂.

Bryonia lævis f. *nigra baccifera*. *Bauh. pin.* 297. ♀.

Vitis sylvestris f. *Tamus*. *Dod. pempt.* 400.

Habitat in Europæ australis, Orientis sepibus. 24

2. TAMUS *cretica*. T. foliis trilobis.

Tamus cretica, *trifido folio*, *Tournef. cor.* 3.

Habitat in Creta. 24

Varietas præcedentis. *Ger.* 136.

1225. S M I L A X.

Masc. *Cal.* 6. *phyllus*. *Cor.* o.

Fem. *Cal.* 6. *phyllus*. *Cor.* o. *Styli* 3. *Bacca* 3.
locularis. *Sem.* 2.

* *Caule aculeato, angulato.*

1. SMILAX *aspera*, S. *caule aculeato angulato, foliis dentato aculeatis cordatis novemnerviis.* *monsp. carn.*

S. *caule angulato aculeato, foliis cordato-oblongis acutis aculeatis.* *Hort. cliff.*

S. *aspera, fructu rubente.* *Bauh. pin.* 296.

S. *aspera, rutilo fructu.* *Clus. hist. i. p.* 122.

β. S. *aspera minus spinosa, fructu nigro.* *Bauh. pin.* 236.

S. *aspera, nigro fructu.* *Clus. hist. i. p.* 113.

554 DIOECIA. OCTANDRIA.

γ. S. viticulis aspera, foliis longis angustis mucronatis lœvibus : auriculis ad basin rotundioribus. *Pluk. alm.* 348. t. 110. f. 3.

Habitas in Hispaniæ, Italiæ, Palæstinæ, Carniolæ sepibus. ἥ

O C T A N D R I A.

1228. P O P U L U S.

Masc. Amenti. *Cal.* lamina lacera. *Cor.* turbinata, obliqua, integra.

Fem. Amenti. *Cal. & Cor.* maris *Stigma* 4. fidum. *Caps.* 2. locularis. *Sem.* multa, papposa.

1. **POPULUS alba.** P. foliis subrotundis dentato - angulatis subtus tomentosis *Fl. suec. paris. gallob. herb. pal. sil. Du Ham. arb.* t. 36. *Blackw.* t. 548. *lugd. burg. lithu.*

P. foliis angulosis, subtus tomentosis, subrotundis, dentato-angulatis. *Hall. helu.* n. 1634.

P. alba, majoribus foliis. *Bauh. pin.* 429.

P. alba. *Dod. pempt.* 835.

P. alba latifolia. *Lob. ic.* 2. p. 193. *Hall.*

β. B. alba folio minore. *Bauh. hist.* 1. p. 160. *Hall. R.*

Habitas in Europa temperatiori. ἥ

Flores omnino ut in P. tremula.

2. **POPULUS tremula.** P. foliis subrotundis dentato angulatis utrinque glabris. *Fl. suec. paris. gallob. herb. pal. sil. Blackw.* t. 248. *Kniph. cent.* 6. n. 73. *Fem. lith. burg. lugd.*

P. foliis glabris, orbiculatis, acuminatis, laxè serratis. *Hall. helu.* n. 1633.

P. tremula. *Bauh. pin.* 429.

P. lybica. *Dod. pempt.* 836.

Habitas in Europæ frigidioribus. ἥ

Folia involuta basi interne glandula duplici.

Petioli apice compressi, unde Folia tremula. Gouan.

3. **POPULUS nigra.** P. foliis deltoidibus acuminatis serratis. *Fl. suec. paris. gallob. carn. herb. pal. sil. Blackw.* t. 248. 1. *Mill. illustr. Syst. lith. burg. lugd.*

P. foliis glabris cordato-rhomboides serratis. *Hall. helu.* n. 1632.

DIOECIA. OCTANDRIA. 555

P. nigra. *Bauh. pin.* 429. *Dod. pempt.* 836. *Matth.* 137.
Loeb. ic. 794.

Habitat in Europa temperatiore. H

*Folia involuta absque glandulis ad basin, hinc diversa a
præcedentis. Stamina 16. Leers.*

1229. RHODIOLA.

Masc. Cal. 4. partitus. *Cor.* 4. petala.

Fem. Cal. 4. partitus. *Cor.* 0. *Nectaria* 4. *Pist.* 4.
Caps. 4. polyspermæ.

1. RHODIOLA rosea. *Rhodiola. Fl. suec. norv. sil. Fl. dan.*
t. 183. *Blackw. t.* 586. *Kniph. cent.* 2. n. 69. *Fabric.*
helmst. p. 271.

R. staminibus corolla duplo longioribus. Mill. dict. n. 1.

Sedum sexu distinctum, foliis serratis, umbellis densissimis.
Hall. helu. n. 953.

Sedum roseum. Scop. carn.

Rhodia Radix. Bauh. pin. 286. *Clus. hist.* 2. p. 65. *Rai.*
hist. 690.

Telephium luteum minus, radice rosam redolente. Moris.
hist. 3. p. 468. *f.* 12. *t.* 10. *f.* 8.

H. stat in Alpibus Lapponiæ, Austriæ, Silesiæ, Helvetiæ,
Britanniæ. 2

Nectaria. 5. Stam. 6. 8. 12. Fabric. Scop. R.

E N N E A N D R I A.

1230. MERCURIALIS.

Masc. Cal. 3. partitus. *Cor.* 0. *Stam.* 9. *f.* 12.
Antheræ globosæ, didymæ.

Fem. Cal. 3. partitus. *Cor.* 0. *Styli* 2. *Caps.*
dicocca, 2. locularis. 1. sperma.

1. MERCURIALIS perennis. *M. caule simplicissimo, foliis*
scabris. Fl. suec. parif. gallob. pal. sil. Flor. dan. t. 400.
Kniph. cent. 1. n. 57. *mas. Mill. illustr. lithu. burg.*

556 · DIOECIA. ENNEANDRIA.

M. caule perenni, indiviso, foliis hirsutis ovato-lanceolatis. *Hall. helu. n. 1601.*

♀. M. montana testiculata. *Bauh. pin. 122.*

Cynocrambe mas. *Cam. epit. 999.*

♂. Mercurialis montana spicata. *Bauh. pin. 122.*

Cynocrambe femina. *Cam. epit. 999.*

Habitat in Europæ nemoribus. 24

2. MERCURIALIS *ambigua*. M. caule brachiato, foliis glabriusculis, floribus verticillatis: femineo maculisque. *Habitat in Hispania.*

Simillima M. annua.

3. MERCURIALIS *annua*. M. caule brachiato, foliis glabris, floribus spicatis. *paris. ged. carn. gallob. pal. Blackw. t. 162. femina. Regn. bot. lugd. burg.*

M. caule annuo brachiato, foliis glabris, conjugatis, ovato-lanceolatis. *Hall. helu. n. 1600.*

M. testiculata f. mas. *Bauh. pin. 121. ♀*

M. mas. *Dod. pempt. 658. ♀*

♂. M. spicata f. femina. *Dod. pempt. 658.*

Habitat in Europæ temperate umbrosis ☉

4. MERCURIALIS *tomentosa*. M. caule suffruticoso, foliis tomentosis. *monsp.*

Phyllon testiculatum. *Bauh. pin. 122. ♀*

Phyllon marificum. *Clus. hist. 2. p. 48.*

♂. Phyllon spicatum. *Bauh. pin. 122.*

Phyllon seminificum. *Clus. hist. 2. p. 48.*

Habitat in G. Narbonensi, Hispania. h

1231. HYDROCHARIS.

Masc. *Spatha* 2. *phylla*. *Cal.* 3. *fidus*. *Cor.* 3. *petala*.
Filamenta 3. interiora stylifera.

Fem. *Cal.* 3. *fidus*. *Cor.* 3. *petala*. *Styli* 6. *Caps.* 6.
locularis, polysperma, infera.

1. HYDROCHARIS *Morsus Ranæ*. *succ. paris. Hall. helu. n. 1068. pal. fil. lith. burg. lugd.*

Nymphæa alba minima. Bauh. pin. 193.

Ranæ morsus. Dod. pempt. 583. f. 2. ♂. 1. ♀.

Nymphæa minor f. Morsus ranæ. Bauh. hist. 3. p. 173. cum icone aliena. Hall. R.

Morsus ranæ flore pleno odoratissimo. *Rai. cens. 101. ♀.*

DIOECIA. DECANDRIA. 557

γ. *Nymphæa alba minor*. *Bauh. pin.* 193.

Habitat in Europæ fossis limosis. ☉ 24

Gemmifera s. bullifera est.

DECANDRIA.

1235. CORIARIA.

Masc. *Cal.* 5. phyllus. *Cor.* 5. petala, calyci similima, connexa. *Antheræ* bipartitæ.

Fem. *Cal.* 5. phyllus. *Cor.* maris. *Styli* 5. *Sem.* 5. petalis succulento-baccatis tecta.

1. CORIARIA *myrtifolia*. *C.* foliis ovato-oblongis. *monsp.*

C. hermaphrodita. *Turr. farset.* 13. (*Femina.*)

Rhus myrtifolia monspeliaca. *Bauh. pin.* 414. *Angl. hortul.* 64. t. 20. f. 1.

Rhus plinii myrtifolia monspeliensium. *Lob. ic.* 2. p. 98.

Habitat Monspelii. ☿

Femina gaudet Antheris sterilibus. *Gerard.* 341.

Est planta polygama monoica, trioica. *Gouan. R.*

Plantam hermaphroditam describit Medicus in obs. Sacæcon. Lutr. 1774. p. 177. *R.*

DODECANDRIA.

1237. DATISCA.

Masc. *Cal.* 5. phyllus. *Cor.* 0. *Antheræ* sessiles, oblongæ, 15.

Fem. *Cal.* 2. dentatus. *Cor.* 0. *Styli* 3. *Caps.* 3. angularis, 3. cornis, 1. locularis, pervia, polysperma, infera.

1. DATISCA *cannabina*. *D.* caule lævi. *Kniph. cent.* 11. n. 38. *fem.*

558 DIOECIA. DODECANDRIA.

Cannabis foliis pinnatis. Hort. cliff. 457.

Cannabis lutea fertilis. Alp. exot. 300. t. 298. Moris. hist. 3. p. 432. f. 8. t. 25. f. 4.

Luteola herba sterilis. Bauh. pin. 100.

Cannabis lutea cretica. Alp. exot. 296. t. 295.

Cannabis lutea sterilis. Alp. exot. 301. t. 300.

Luteola herba foliis cannabinis. Bauh. pin. 100.

Habitas in Creta. 24

1238. MENISPERMUM.

Masc. Petala 4 exteriora; 8 interiora. Stam. 16.

Fem. Cor. maris. Stam. 8. sterilia. Baccæ binæ, monospermæ.

1. *MENISPERMUM canadense. M. foliis peltatis cordatis subrotundo-angulatis. Hort. upf. 91. Gmel. sib. 3. p. 107. n. 86. Willich. obs. 14. Du Roi. herbk. 1. p. 405.*

Menispermum canadense scandens, umbilicato folio. Tourm. ad. 1705. p. 311.

Hedera monophyllos virginiana, convolvuli foliis. Pluk. alm. 181. t. 36. f. 2.

Cissampelos. Rupp. ien. 67. Gmel. R.

Canadischer Mondsaamen. Linn. Pflanzensyst. 4. p. 611.

Habitas in Virginia, Sibiria. h

Caulis volubilis contra solem, lævis. Folia subrotundocordata, quinquangula, lævia, acuminata. Pedunculus béracemosus, supra axillaris, nutans, petiolo brevior.

MONADELPHIA.

1240. JUNIPERUS.

Masc. Amenti Calyx squamæ. Cor. 0. Stam. 3.

Fem. Cal. 3. partitus. Petala 3. Styli 3. Baccæ 3: sperma, tribus tuberculis calycis inæqualis.

1. *JUNIPERUS thurifera. J. foliis quadrifariam imbricatis acutis.*

DIOECIA. MONADELPHIA. 559
Cedrus hispanica procerior, fructu maximo nigro. *Tournef. inst.* 588.

Habitat in Hispania. $\bar{\eta}$

5. **JUNIPERUS Sabina.** J. foliis oppositis erectis decurrentibus : oppositionibus pyxidatis. *monsp. carn. Ludw. ed.* t. 93. *Kniph. cent.* 5. n. 46.

J. foliis cauli adpressis, lanceolatis, alterne conjugatis. *Hall. helu.* n. 1662.

J. (*lusitanica*) foliis oppositis patulis decurrentibus, ramis erectioribus. *Mill. dict.* n. 11.

J. foliis inferne adnatis : oppositionibus concatenatis. *Hort. upf.*

Sabina folio cupressi. *Bauh. pin.* 487. *Du Ham. arb.* 2. t. 62.

Sabina. *Dod. pempt.* 854. *Blackw.* t. 214.

α . **Sabina folio tamarisci dioscoridis.** *Bauh. pin.* 487.

Juniperus (Sabina) foliis oppositis erectis decurrentibus, ramis patulis. *Mill. dict.* n. 10.

Habitat in Lusitania, Italia, Sibiria, Oriente, Olympo, Ararat. $\bar{\eta}$

7. **JUNIPERUS communis.** J. foliis ternis patentibus mucronatis bacca longioribus. *Scop. carn. herb. pal. sil. Ludw.* t. 189. *Blackw.* 187. *Knorr. del. hort.* 1. t. W. 8. *Kniph. cent.* 1. n. 48. *mas. Mill. illustr. lich. burg. lugd.*

J. foliis convexo concavis, aristatis, baccis alaribus. *Hall. helu.* n. 1661.

J. foliis patentibus. *Fl. succ. binerviis. Sauv. monsp.*

J. vulgaris fruticosa. *Bauh. pin.* 488.

β . J. vulgaris arbor. *Bauh. pin.* 488. *Mill. dict.* n. 2.

γ . J. minor montana, folio latiore, fructu longiore. *Bauh. pin.* 489.

Habitat in Europæ frigidioris sylvis frequentissima. $\bar{\eta}$
Antheræ sub squamis lateralibus sæpe 5. *Leers. R.*

8. **JUNIPERUS Oxycedrus.** J. foliis ternatis patentibus mucronatis bacca brevioribus. *carn.*

J. foliis quaternis patentibus subulatis mucronatis. *Sauv. ad. monsp.* 1743.

J. foliis quaternis patentibus subulatis pungentibus. *Roy. lugdb.*

J. foliis ternis patentibus subulatis pungentibus trinerviis. *Sauv. monsp.*

J. major, bacca rufescente. *Bauh. pin.* 489. *Du Ham. arb.* 2. p. 326. t. 128.

560 DIOECIA. MONADELPHIA?

Oxycedrus. Clus. hist. 1. p. 39.

Habitat in Hispania, G. Narbonensi. H

Ramuli triquetri. Folia terna, patentia, sessilia (minime adnata) omnino ut in *J. communi*, sed majora omnia. Baccæ rufescentes magnitudine *Avellanæ*.

9. *JUNIPERUS phænicea*. J. foliis ternis oblitteratis imbricatis obtusis. *monsp.*

Cedrus folio cupressi major, fructu flavescente. *Bauh. pin. 487.*

Juniperus major. *Clus. hist. 1. p. 38.*

Habitat in Europa australi. Oriente. H

Hanc a J. lycia vix specie diversam judicat Rai. exot. 90; inquirens populares.

Folia ovata, convexa, obtusa, minutissima.

10. *JUNIPERUS lycia*. J. foliis ternis undique imbricatis ovatis obtusis. *monsp.*

Cedrus, folio cupressi, media majoribus baccis. *Bauh. pin. 487.*

Cedrus phœnicea altera Plinii & Theophrasti. *Lob. ic. 221: media ejusd. ic. 221.*

Habitat in Gallia, Sibiria. H

1241. TAXUS.

Masc. *Cal.* 4. phyllus gemmæ. *Cor.* 0. *Stam.* multa. *Antheræ* peltatæ octofidæ.

Fem. *Cal.* 4. phyllus gemmæ. *Cor.* 0. *Styl.* 0. *Sem.* 1, calyculo baccato, integerrimo.

1. *TAXUS baccata*. T. foliis approximatis. *Hall. helv. n. 1663. norv. sil. Du Ham. arb. 2. t. 86. Blackw. 1. 572. Kniph. cent. 1. n. 89. mas. lithu. Fl. suec. paris.*

Taxus. Bauh. pin. 505. Cam. epis. 840. Dod. pempt. 859:

Habitat in Europa, Canada. H

Dicotyledonis est.

1242. EPHEDRA.

Malc. *Amenti. Cal* 2. fidus. *Cor.* o. *Stamina* 7.
Antheræ 4. inferiores. 3. superiores.

Fem. *Cal.* bipartitus, 5. *tuplex. Cor.* o. *Pist.* 2.
Sem. 2. calyce baccato tecta.

1. EPHEDRA *distachya.* E. pedunculis oppositis : amentis
geminis. *monsp.*

E. petiolis masculis repetito ramosis. *Hall. helu. n.* 1664.

Polygonum bacciferum maritimum minus. *Bauh. pin.* 15.

Polygonum fruticans, botryoides Narbonense, minus. 2.

Clus. Barrel. ic. 731. n. 2. *Hall. R.*

Una marina monspeliensium. *Lob. ic.* 796. *Hall. R.*

Tragum. *Cam. hort.* 171. t. 46.

*Habitat in G. Narbonensis, Hispaniæ, Helvetiæ saxosis
collibus marinis. H*

SYNGENESIA.

1246. RUSCUS.

Malc. *Cal.* 6. phyllus. *Cor.* o. *Nectarium* centrale,
ovatum, apice perforatum.

Fem. *Calyx*, *Corolla* & *Nectarium* maris. *Styl.* 1.
Bacca 3. locularis. *Sem.* 2.

1. RUSCUS *aculeatus.* R. foliis supra floriferis nudis. *paris.
carn. Blackw. t.* 155. *femina. Sabb. hort. I. t.* 8. *Kniph.
cent. I. n.* 77. *mas. Regn. bot.*

R. foliis ovato lanceolatis, pungentibus, petiolis brevissi-
mis vaginosis. *Hall. helu. n.* 1238.

R. myrtifolius aculeatus. *Tournef. inst.* 79:

Ruscus. *Bauh. pin.* 470.

Ruscum. *Dod. pempt.* 474.

*Habitat in Galliæ, Italiæ, Helvetiæ, &c. nemorosis as-
peris. H*

Linn.

Europ. Pars II,

N a

564 DIOECIA. SYNGENETA.

2. RUSCUS *Hypophyllum*. R. foliis subtus floriferis nudis.
Laurus alexandrina, fructu folio infidente. *Bauh. pin.* 305.
Blackw. t. 194.

Laurus alexandrina, *Chamædaphne*. *Col. ecphr.* I. p. I. p.
 364. t. 165. f. 1.

β. *Ruscus latifolius*, fructu ex medio foliorum extra pen-
 dente. *Dill. elth.* 333. t. 251. f. 303.

Habitat in Italia ad latera collium. 24

3. RUSCUS *Hypoglossum*. R. foliis supra floriferis sub
 foliolo. *Sabb. hort.* t. 9. *Kniph. cent.* I. n. 78. *mas.*

R. caule simplici; floribus pedunculatis, foliolo rectis.
Scop. carn.

L. alexandrina, fructu pediculo infidente. *Bauh. pin.* 304

Hypoglossum dioscoridis; *Laurotaxa Plinii*. *Col. ecphr.* I. p.
 166. t. 165. f. 2. *Blackw. t.* 128.

Habitat in Hungariæ, Liguriæ, Italiæ montibus umbrosis. 24

5. RUSCUS *racemosus*. R. racemo terminali hermaphrodi-
 tico. *monsp.* *Sabb. hort.* I. t. 10.

R. angustifolius, fructu summis ramulis inascente. T. Mich.
pis. 82. Du Ham. arb. 4. *Tournef. inst.* 79.

Laurus alexandrina angustifolia ramosa, fructu ad extremum
 ramoso. *Morif. hist.* 3. p. 541. f. 13. t. 5. f. 14.

Laurus alexandrina, fructu e summitate caulium prodeunte.
Herm. lugdb. 679. t. 681. R.

Habitat in Insulis Archipelagi. 7

Calyx cum corolla coalitus fere in totum & deciduus. *Hort.*
cliff. 466. Fructus monospermus. *Hall. gatt.* 78. Flos herma-
 phroditus.

CLASSIS XXIII.

POLYGAMIA.

MONOECIA.

1252. HOLCUS.

♀. *Glum.* 1. *flora.* *Glum.*
2. *valv.* *Stam.* 3. *Styl.*
2. *Sem.* 1.

♂. *Glum.* 1. *flora.* *Glum.*
2. *valv.* *Stam.* 3.

1255. CENCHRUS.

♀. *Glum.* 2. *flora.* *Glum.*
2. *valv.* *Stam.* 3. *Styl.* 2.
fid. *Sem.* 1.

♂. *Involucr.* *idem.* *Glum.*
2. *valv.* *Stam.* 3.

1254. ISCHÆMUM.

♀. *Glum.* 2. *flora.* *Glum.*
2. *valv.* *Stam.* 3. *Styl.*
2. *Sem.* 1.

♂. *Glum.* *ead.* *Glum.* bi-
valv. *Stam.* 3.

1256. ÆGILOPS.

♀. *Glum.* 3. *flora.* *Glum.*
3. *arist.* *Stam.* 3. *Styli* 2.
Sem. 1.

♂. *Glum.* 3. *flora.* *Glum.*
3. *arist.* *Sem.* 3.

1251. ANDROPOGON.

♀. *Glum.* 1. *flor.* *Glum.*
basi arist. *Stam.* 3. *Styl.*
2. *Sem.* 1.

♂. *Glum.* 1. *flor.* *Glum.*
basi arist. *Stam.* 3.

1258. VALANTIA.

♀. *Cal.* 0. *Cor.* 4. *partita.*
Stam. 4. *Styl.* 2. *fid.*
Sem. 1.

♂. *Cal.* 0. *Cor.* 3. *f.* 4. *par-*
tita. *Styl.* 3. *f.* 4.

564

1267. CELTIS.

POLYGAMIA.

♂. Cal. 5. part. Cor. o.
Stam. 5. Styl. 2. Dru-
pa.

1249. VERATRUM.

♂. Cal. 6. part. Cor. o.
Stam. 6.

♀. Cal. o. Cor. 6. pet. Sta.
6. Pist. 3. Caps. 3.

♂. Cal. o. Cor. 6. petal.
Stam. 6.

1266. ACER.

♀. Cal. 5. fidus. Cor. 5. pet.
Stam. 8. Styl. 2. Caps. 2.
cocca alata.

♂. Cal. 5. fidus. Cor. 5.
pet. Stam. 8.

1259. PARIETARIA.

♀. Cal. 4. fid. Cor. o. Sta.
4. Styl. 1. Sem. 1.

♂. Cal. 4. fid. Cor. o. Styl.
1. Sem. 1.

1260. ATRIPLEX.

♀. Cal. 5. phyll. Cor. o.
Stam. 5. Styl. 2. fid.
Sem. 1.

♂. Cal. 2. phyllus. Cor. o.
Syst. 1. fid. Sem. 1.

* *Æsculus. Fraxinus excelsior.*

Euphorbia. Melothria. Ilex.

Silene saxifraga.

DIOECIA.

1273. FRAXINUS.

♀. Cal. o. f. 4. part. Cor. o.
f. 4. pet. Stam. 2. Pist. 1.
Sem. 1.

♂. Cal. o. f. 4. part. Cor. o.
f. 4. petala. Pist. 1.
Sem. 1.

* *Ilex aquifolium.*

Rhamnus Alaternus.

TRIOECIA.

1282. CERATONIA.

A ♀ *Cal.* 5. partit. *Cor.* o.
Stam. 5. *Styl.* 1. *Legum.* coriaceum, poly-
sperm.

B ♂ *Cal.* 5. partit. *Cor.* o.
Stam. 5.

♀. *Cal.* sub 5. dent. *Cor.* o.
Styli 1. *Legumen* co-
riaceum polyspermum.

1283. FICUS.

Recept. commune turbina-
tum, conniventi-clau-
sum, carnosum.

A ♀ *Cal.* 5. part. *Cor.* o.
Pist. 1. *Sem.* 1.

B ♂ *Cal.* 3. part. *Cor.* o.
Stam. 3.

C ♂ & ♀ intra idem *Re-*
ceptaculum commune,
distinctis fructificatio-
nibus partialibus.

POLYGAMIA.

MONOE CIA.

1249. VERATRUM.

Hermaphrod. *Cal.* 0. *Cor.* 6. petala. *Stam.* 6. *Pist.* 3. *Caps.* 3. polyspermæ.

Masc. *Cal.* 0. *Cor.* 6. petala. *Stam.* 6. *Pistilli* rudimentum.

1. VERATRUM *album*. V. racemo supradecomposito, corollis erectis. *norv. carn. Jacq. austr. t. 335. fl. Ludw. c. 3. t. 137. & 138. Mill. illustr. lithu.*

Veratrum pedunculis corolla erecta patente brevioribus. *Gmel. sib.*

V. spica paniculata, floribus sexu distinctis. *Hall. helv. n. 1204.*

Helleborum s. Veratrum album. *Dod. pempt. 383. Blackw. t. 74.*

Helleborus albus, flore subviridi. *Bauh. pin. 186.*

Helleborus albus, exalbido flore. *Clus. hist. 1. p. 274.*

Habitat in Russiæ, Sibirici, Austriæ, Helvetiæ, Italiæ, Græciæ montosis. 24

2. VERATRUM *nigrum*. V. racemo compacto, corollis patentissimis. *carn. Jacq. austr. t. 336. Krypt. carn. t. n. 91.*

V. pedunculis corolla patentissima longioribus. *Gmel. fr. 1. p. 76.*

V. flore atro rubente. *Tournesf. inst.*

Helleborus albus, flore atro rubente. *Bauh. pin. 186. Moris. hist. 3. p. 485. f. 12. t. 4. f. 1.*

Habitat in Hungariæ, Sibirici apricis siccis. 24

Præcedenti n. 1. æ affine, diversum loco, colore, pedunculis villosis, corolla patente nec erecta, racemo compacto, nec subdiviso in paniculam.

1251. ANDROPOGON.

Hermaphrod. *Cal. Gluma* 1. *flora. Cor. Gluma*
basi aristata. Stam. 3. Styli 2. Sem. 1.

Masc. *Cal. & Cor. prioris. Stam. 3.*

4. ANDROPOGON *gryllus*. A. paniculæ pedunculis simplicissimis trifloris, flosculo hermaphrodito aristato ciliato basi barbato. *monsp. carn.*

Phoenix spica laxa, locustis longe petiolatis, quadrifloris, duobus floribus sessilibus, duobus petiolatis. *Hall. helv. n. 1412.*

Festuca dumetorum, utriculis lanugine flavescentibus, Ægilops Dioscoridis. *Bauh. theatr. 149. icon, non vero flos seorsim pictus. Hall. R.*

Ægilops bromoides, juba purpurascens. *Bauh. hist. 2. p. 436. Scheuch. gram. 267.*

Gramen spartheum festuceum. f. Ægilops sparthea villosa. *Barr. ic. 18. f. 2.*

Habitat in Rhætia, Helvetia, Veronæ, Monspeli.

Lanugo brevissima baseos hermaphroditi.

Obs. Festuca dumetorum, utriculis lanugine flavescentibus. *Bauh. pin. 9. theatr. 149. vulgo ad Avenam fatuam relata, hujus synonymon secundum Hallerum.*

10. ANDROPOGON *distachyum*. A. spicis binis terminalibus, culmo indiviso. *Ger. prou. 106. t. 3. f. 2.*

A. spicis geminis, locustis bifloris, folliculis alterne muticis. *Hall. helv. n. 1415.*

Gramen dactylon spica gemina. J. R. H. *Scheuchz. gram. 95.*

Gramen bicornis f. distachyophorum. *Bocc. sicc. 20.*

Habitat in Helvetia Burser.

Culmi plusquam pedales. Folia plana. Rami nulli. Spicæ duæ, æquales, oblongæ, erectiusculæ, culmum terminantes: Singulæ ex flosculis viginti pluribus. Pedicelli archi, brevissimi, planiusculi, basi pilosi, superne latiores. Flores solitarii. Calyces uniflori: gluma exterior flore fere longior latiorque lævis subtruncata apice scarioso pubescente aristâ longa, articulata, patente. Gluma calycina interior minima.

568 POLYGAMIA. MONOECIA.

14. ANDROPOGON *hirtum*. A paniculae spicis conjugatis, calycibus hirsutis. *Læfl. it.* *

Festuca junceo folio, spica gemina. *Bauh. pin. 9. prodr. 19. theatr. 145.*

Gramen Dactylon spica gemina. *T. Scheuch. gram. 95.*

Gramen Dactylon ficulum, multiplici panicula, spicis ab eodem exortu geminis. *Pluk. alm. 175. t. 92. f. 1.*

Habitat in Lusitania, Sicilia, Smyrnae.

Pedunculi hirti, uti & rachis & glumæ. Hirsuties glumarum flore non longior.

19. ANDROPOGON *Ischæmum*. A. spicis digitatis plurimis, flosculis sessilibus: aristato muticoque, pedicellis lanatis. *monsp. Jacq. austr. t. 384. carn. gallob. pal. Knapf. cent. 12. n. 9.*

A. spicis villosis septenis, locustis bifloris, folliculis alternis sessilibus. *Hall. helu. n. 1414.*

Ischoemum panicula digitata, spiculis oppositis disjunctive, florum serie duplici, culmo tereti. *Guett. stamp. Delib. paris.*

Gramen Dactylon spicis villosis. *Bauh. pin. 8. theatr. 116. Scheuch. gram. 94.*

Gramen Dactylon spicis aristatis geniculatis. *Barr. ic. 753. f. 2.*

Habitat in Europæ australioris aridis, collibus, saxosis.

Flosculi 2, approximati in singulo dente rachæos. Hermaphroditi Calyx hirsutus: Petalo exteriore tenero interiore aristato. Masculi pedicellus hirsutus.

1252. HOLCUS.

Hermaphrod. Cal. Gluma 1. f. 2. flora. Cor. Gluma aristata. Stam. 3. Styli 2. Sem. 1.

Masc. Cal. Gluma 2. valvis. Cor. 0. Stam. 3.

6. HOLCUS *mollis*. H. glumis bifloris nudiusculis: flosculo hermaphrodito mutico, masculo arista geniculata. *monsp. gram. 149. t. 20. f. 2. spicill. fl. lips. p. 51. sub Aira. gallob. herb.*

Avena. *Hall. helu. n. 1485. lithu.*

Gramen paniculatum molle, radice graminis canini repente. *Moris. hist. 3. p. 202.*

Gramen miliaceum aristatum molle. *Rai. angl. 3. p. 404.*

Gramen caninum longius radicum. *Bauh. pin.* 1.

Gramen caninum paniculatum molle. *Rai. hist.* 1285. *Scheuch. gram.* 235.

Habitat in Europa.

Simile H. lanato, sed Calyces acutiores, acuminati magis, quam in reliquis.

7. *HOLCUS lanatus.* H. glumis bifloris villosis; hermaphrodito mutico masculo arista recurva. *ged. carn. gallob. Schreb. gram.* 145. t. 20. f. 1. *herb.* 70. t. 7. f. 6. *pal. fil. lith. burg. lugd.*

Aira foliis pubescentibus, panicula contracta: flosculo hermaphrodito mutico, masculo arista uncinata, calyce brevior. *Fl. suec.*

Aira flosculo masculo aristato, femineo mutico. *Hort. cliff. Avena triantha, floribus ovatis, perfecto mutico, imperfecto aristato. Hall. helu. n.* 1484.

Aira spiculis polygamis; masculo aristato; arista recurva. *Scop. carn.*

Gramen pratense paniculatum molle. *Bauh. pin.* 2. *prodr.* 3. *theatr.* 27. *Scheuch. gram.* 234. *Læs. pruss. ic.* 25.

Gramen lanatum. *Dalech. hist.* 425.

Habitat in Europæ pascuis arenosis. 24

10. *HOLCUS odoratus.* H. glumis trifloris muticis acuminatis: hermaphrodito diandro. *ged. gallob. lithu.*

Hierocloe. Gmel. sib. 1.

Avena triantha, locustis paniculatis, hiantibus, flosculo tertio imperfecto. *Hall. helu. n.* 1496.

Aira foliis planis, floribus trifloris aristatis: flosculo intermedio mutico diandro. *Fl. suec.*

Poa spiculis bifloris cordatis. *Fl. lapp.*

Gramen paniculatum odoratum. *Bauh. pin. prodr.* 7. *Scheuch. gram.* 236.

Gramen mariæ boruflorum. *Læs. pruss. III.* t. 26.

Habitat in Europæ frigidioris pascuis humensibus. 24

Flosculi duo laterales masculi triandri: intermedius s. tertius hermaphroditus diandrus.

Involucrum laciniatum, echinatum, 2. florum.
Cal. Gluma 2. flora; altero masculo, altero
 hermaphrodito.

Hermaphrod. *Cor.* Gluma mutica. *Stam.* 3. *Styl.* 1.

Masc. *Cor.* Gluma mutica. *Stam.* 3.

1. CENCHRUS *racemosus*. C. panicula spicata, glumis
 muricatis setis ciliaribus. *stamp. parif. monsp.*

Agrostis panicula spicata, glumis echinatis. *Schreb. gram.*
 45. t. 4. *Ger. prou. quod omnes flores sunt hermaphroditi.*

Phalaris muricata. *Forskæhl. ægypt. R.*

Tragus. *Hall. helu. n. 1413.*

Gramen caninum maritimum, spica echinata. *Bauh. pin. 2.*
Scheuch. gram. 76.

Gramen caninum maritimum asperum. *Bauh. prodr. 2. t. 2.*

Gramen caninum maritimum spicatum, echinatis glumis.
Barr. ic. 718.

Habitat in Europæ australioris maritimis, & in India orientali.

Calyx: Gluma bivalvis, triflora, ovato-oblonga, acuminata, cartilaginea, echinata. Corollæ bivalves, lanceolatae, membranaceæ, sessiles, longitudine calycis. Tertia s. intermedia Mascula calyce univalvi, priorum calyci simili, echinata, pedicellata. Culmus reclinatus. Folia ciliata.

Obs. Involucro destituitur, adeoque aut species Cenchri aut Melicæ erit. *Mant. 501.*

Glumæ absoluta florescentia rubent. Gouan.

4. CENCHRUS *capitatus*. C. spica ovata simplici. *monsp.*

Gramen spica subrotunda echinata. *Bauh. pin. prodr. 16.*
Scheuch. gram. 74.

Gramen montanum echinatum tribuloides capitatum. *Cal. ephr. 1. p. 340. t. 338. f. 1.*

Gramen minimum, spica globosa echinata. *Barr. rar. 1176.*
t. 28. f. 1. & t. 862. f. 2.

Habitat in G. Narbonensi, Italia.

Culmus geniculo unico. Gouan.

1256. ÆGILOPS.

Hermaphrod. *Cal.* Gluma subtriflora, cartilaginea.
Cor. Gluma terminata triplici arista, *Stam.* 3.
Styli 2. *Sem.* 1.

Masc. *Cal.* & *Cor.* Gluma hermaphrodit. *Stam.* 3.

1. ÆGILOPS *ovata*, Æ. spica aristata, calycibus omnibus triaristatis. *carn. sub Phleo.*

Ægilops spica ovata aristis brevioribus. *paris. Sauv. monsp.*

Ægilops. *Dod. pempt. 73. Cam. epit. 928. f. 2.*

Festuca altera, capitulis duris, *Bauh. pin. 10. theat. 151.*
Gramen spicatum, durioribus & crassioribus locustis, spica brevi. *Scheuch. gram. 11.*

Habitat in Europa australi. ☉

3. ÆGILOPS *triuncialis*. Æ. spica aristata : calycibus inferioribus biaristatis. *Kniph. cent. 11. n. 1.*

Æ. spica oblonga aristata : aristis geminis ternisve. *Schreb. gram. 80. t. 10. f. 1.*

Festuca altera, capitulis duris, spica triunciali. *Bauh. pin. 10.*
Gramen spicatum, durioribus & crassioribus locustis, spica longissima. *T. Scheuch. gram. 12.*

Habitat in Monspeli, Massiliæ, Smyrnæ aridis.

Spica oblonga, aristis flore multo longioribus. *Calycis valvula altera aristis 3, altera 2. Corollæ valvula altera aristis 3, altera 1. Radix perennis.*

5. ÆGILOPS *incurvata*. Æ. spica subulata mutica lævi incurva, calycibus unifloris. *Scop. carn. sub Agrostide. burg.*

Nardus spica subulata disticha. *Roy. lugdb.*

Gramen foliaceum, spicis articulosis erectis. *T. Schench. gram. 43.*

Gramen myuros erectum minimum arundinaceum. *Bocc. mus. 70. t. 59.*

Habitat in Angliæ, Hispaniæ, Italiæ paludibus maritimis. ☉

Spica filiformis, vix scapo crassior, teres. Calyx bivalvis, exterus. Corollæ valvulæ calyci contrariæ.

Hermaphrod. *Cal.* 0. *Cor.* 4. partita. *Stam.* 4.
Styl. 2. fidus. *Sem.* 1.

Masc. *Cal.* 0. *Cor.* 3. f. 4. partita. *Stam.* 3. f. 4.
Pistillum obsoletum.

1. *VALANTIA muralis*. V. floribus masculis trifidis germi-
mini hermaphroditici glabro insidentibus. *monsp. Sabb.*
hort. 1. t. 83.

V. procumbens glabra. *Hort. cliff.*

V. annua quadrifolia verticillata, floribus ex viridi pallescen-
tibus, fructu ecchinato. *Mich. gen.* 13. s. 7.

Rubeola echinata saxatilis. *Bauh. pin.* 334.

Cruciata muralis minima romana. *Col. ecphr.* 1. p. 298.
t. 297. *Moeis. hist.* 3. p. 328. f. 9. t. 21. f. 2.

Habitat in G. Narbonensis, Italiae muris. ☉

Caulis & Folia glabra.

2. *VALANTIA hispida*. V. floribus masculis trifidis germi-
mini hermaphroditici hispido insidentibus.

V. erecta hispida. *Hort. cliff.*

Galium floribus masculis trifidis, omnibus plantae partibus
hispidis. *Zinn. grett.* 233.

Habitat in Europa australi. ☉

Affinis maxime V. murali, sed major erecta. Caules
hispidi. Folia scabra. Fructus hermaphroditus rotundus,
undique hispidus; nec laevis aut restratus.

4. *VALANTIA Aparine*. V. floribus masculis trifidis pe-
dicellatis hermaphroditici pedunculo insidentibus. *paris.*
monsp. pal. Kniph. cent. 9. n. 97. *lugd.*

Galium foliis ferratis, petiolis trifloris, recurvis. *Hall. helm.*
n. 725.

Aparine femine laevi. *Vaill. paris.* 18. t. 4. f. 3.

Habitat inter Germaniae, Galliae, Siciliae segetes. ☉

Facies Galii Aparines. Caulis retrorsum scaber, scandens.
Folia sena, margine scabra. Pedunculi longitudine foliorum,
apice triflori: Flosculo Hermaphrodito sessili: lateralibus
Masculis pedicellatis, nunc 3. fidis, nunc 4. fidis.

6. *VALANTIA Cruciata*. V. floribus masculis quadrifidis,
pedunculis diphyllis. *monsp. paris. gallob. herb. pal. fl.*
lugd. lith.

POLYGAMIA. MONOECIA. 573

Galium foliis quaternis, flosculis in alis confertis. *Hort. cliff.*

Galium foliis quaternis, hirsutis petiolis octifloris. *Hall. helu. n. 769.*

Galium floribus polygamis; racemis lateralibus, ternatis, diphyllis; foliis quaternis, villosis. *Scop. carn.*

Cruciata hirsuta. *Bauh. pin. 335.*

Cruciata. *Dod. pempt. 257.*

Habitas in Germania, Helveria, Gallia. 24

7. VALANTIA glabra. V. floribus masculis quadrifidis, pedunculis dichotomis, aphyllis, foliis ovalibus ciliatis. *lithu.*

Galium floribus polygamis, racemis lateralibus binatis nudis, foliis quaternis lanceolatis glabris. *Scop. carn.*

Galium latifolium glabrum. *Bauh. prodr. 146.*

Cruciata glabra. *Bauh. pin. 335.*

Habitat in Austria, Italia. 24

Affinis nimis V. Cruciatæ.

1259. PARIETARIA.

Hermaphrod. Cal. 4. fidus. Cor. o. Stam. 4. Styl. 1. Sem. 1. superum, elongatum.

Fem. Cal. 4. fidus. Cor. o. Stam. o. Styl. 1. Sem. 1. superum, elongatum.

2. PARIETARIA officinalis. P. foliis lanceolato-ovatis, pedunculis dichotomis, calycibus diphyllis. *Flor. dan. 521. Regn. bot. lud. burg.*

P. foliis lanceolato-ovatis alternis. *S. pl. paris. gelr. carn. ged. gallob. pal. sil. Ludw. ed. t. 115. Kniph. cent. 4. n. 60.*

P. foliis hirsutis elliptico-lanceolatis. *Hall. helu. n. 1612.*

P. officinarum & dioscoridis. *Bauh. pin. 121.*

Parietaria Helxine. *Tabern. p. 550. Blackw. t. 136.*

Helxine. *Cam. epit. 849.*

Habitas in Europæ temperationis ruderatis. 24

Flores feminei tetragoni, pyramidati.

3. PARIETARIA judaica. P. foliis ovatis, caulibus erectis, calycibus trifloris, corollis masculis elongato-cylindricis. *Kniph. cent. 11. n. 83.*

P. foliis subhirsutis, ovato-lanceolatis, *Hall. helu. n. 1613.*

574 POLYGAMIA. MONŌECIA:

P. minor ocymi folio. *Bauh. pin* 121 ?

P. ocymi folio Tragi. *Boec. fic.* 47. t. 24. f. A.

Habitas Goettingæ, in Helvetia, Palæstina. 24

Florum *acervi ad ramos, vix ulli ad caulem*. Flores mascululi gemini, tubulosi, cylindrici, oris dentibus conniventibus. Flos femineus intermedius, ovatus.

4. *PARIETARIA lusitanica*. *P. foliis ovatis obtusis, caulibus filiformibus striatis lævibus procumbentibus*.

P. lusitanica annua minima. *Tournef. inst.* 509. *Rat. suppl.* 129.

P. ficula, alfine folio. *Boec. fic.* 47. t. 24.

Habitas in Lusitania, Hispania. ☉

1260. ATRIPLEX.

Hermaphrod. *Cal.* 5. phyllus. *Cor.* o. *Stam.* 5. *Styl.* 2. partitus. *Sem.* 1. depressum.

Fem. *Cal.* 2. phyllus. *Cor.* o. *Stam.* o. *Styl.* 2. partitus. *Sem.* 1. compressum.

1. *ATRIPLEX Halimus*. A. caule fruticoso, foliis deltoideis integris.

Halimus latifolius s. fruticosus. *Bauh. pin.* 120.

Halimus. 1. *Clus. hist.* 1. p. 53.

Habitas in Hispaniæ, Lusitaniæ, Virginie, Sibirie sepibus maritimis. ☿

2. *ATRIPLEX portulacoides*. A. caule fruticoso, foliis obovatis. *Fl. suec. Scop. carn.*

A. caule fruticoso, foliis lanceolatis obtusis. *Roy. lugb.* 218.

Halimus s. *Portulaca marina*. *Bauh. pin.* 120.

Portulaca marina. *Dod. pempt.* 771.

Habitas in Oceani Europæi septentrionalis litoribus. ☿

3. *ATRIPLEX glauca*. A. caule suffruticoso procumbente, foliis ovatis sessilibus integerrimis: inferioribus subdentatis.

A. maritima hispanica frutescens & procumbens. *Dill. elb.* 46. t. 40. f. 46.

Polygonum incanum rotundifolium halimoides fruticans Hispanicum. *Barr. rar.* 1145. t. 733.

Habitas in Gallie australis, Hispanie maritimis. ☿

4. *ATRIPLEX rosea*. A. caule herbaceo, foliis incanis

ratis, fructibus quadrangularibus dentatis. *Scholl. barb. pal.*

A. sylvestris, fructu roseo compresso. *Bauh. pin. 119. prodr. 58. Rupp. jenn. 341.*

Habitat in Europa australiore. ☉

7. *ATRIPLEX hortensis*. A. caule erecto herbaceo, foliis triangularibus. *Blackw. t. 99. & 552.*

A. hortensis alba f. pallide virens. *Bauh. pin. 119.*

A. hortensis. *Dod. pempt. 615.*

β. A. hortensis rubra. *Bauh. pin. 119.*

Habitat in Tartaria. ☉

8. *ATRIPLEX laciniata*. A. caule herbaceo, foliis deltoideis dentatis subtus argenteis. *Fl. suec. gallob. ged.*

A. maritima laciniata. *Bauh. pin. 120. Moris. hist. 2. p. 607. f. 5. t. 32. f. 17.*

A. marina. *Dod. pempt. 615.*

Habitat ad Europæ & Virginie septentrionalis littora marina. ☉

Planta tota recta quasi epidermide secedente. Caulis erectus, teres, nudus, virgatus. Folia exceptis infimis alterna, deltoidea, dentata, argentata laminulis. Spicæ terminales hermaphroditæ antheris late rubris. Flores Feminei axillares gemini. Calyx fructus compressus, 5. dentatus: dente intermedio majore.

9. *ATRIPLEX hastata*. A. caule herbaceo: calycinis valvulis femineis magnis deltoideis sinuatis. *Fl. suec. parif. gallob. pal. herb. burg. lithu.*

A. foliis triangularibus, basi producta, valvulis triangularibus subasperis. *Hall. helu. n. 1617.*

A. sylvestris, folio hastato f. deltoide. *Moris. bles. 237. Rai. hist. 192.*

A. sylvestris annua, folio deltoide triangulari sinuato & mucronato hastæ cuspidis simili. *Moris. hist. 2. p. 607. f. 5. t. 32. f. 14.*

Habitat in Europa frigidiori. ☉

Valvulæ calycinæ femineæ in hac haud raro etiam disco dentatæ. *Leers. l. c. R.*

10. *ATRIPLEX patula*. A. caule herbaceo patulo, foliis subdeltoideo-lanceolatis, calycibus seminum disco dentatis. *Ged. gallob. carn. pal. lith. burg.*

β. A. caule herbaceo, valvulis femineis seminum disco dentatis. *Fl. suec. parif.*

γ. A. caule herbaceo, foliis superioribus lanceolatis integerrimis. *Hort. cliff.*

576 POLYGAMIA. MONŒCIA.

A. foliis imis hamatis ; reliquis lingularis , valvulis seminibus dentatis. *Hall. helv. n. 1616. **

A. angusto oblongo folio. Bauh. pin. 219.

A. sylvestris , polygoni f. helxines foliis. Lob. ic. 257.

A. vulgaris angustifolia cum folliculis. Bauh. hist. 2. p. 973.

Habitat in Europæ culus , ruderalis. ☉

11. *ATRIPLEX littoralis. A. caule herbaceo erecto , foliis omnibus linearibus integerrimis. Fl. suec. gallob.*

A. angustissimo & longissimo folio. Herm. lugdb. 79.

Habitat in Europæ septentrionalis litoribus maris. ☉

Folia & linearia & lanceolata. Gmel. is. 2. p. 2. Folia interdum dentata , sineque recurvo ; color glaucus ; calis sulcatus. Willich. illustr. n. 6. R.

12. *ATRIPLEX marina. A. caule herbaceo erecto , foliis linearibus serratis. Hudf. angl.*

A. maritima angustifolia. Bauh. pin. 152.

A. angustifolia dentata. Pet. herb. 8. 7. f. 4.

Habitat in Angliæ , Sueciæ litoribus marinis. ☉

Herba pusilla , vix palmaris. Foliis crassiusculis.

13. *ATRIPLEX pedunculata. A. caule herbaceo divaricato , foliis lanceolatis obtusis integris , calycibus femineis pedunculatis. Hudf. angl. Flor. dan. 304.*

A. maritima , semine lato. Rai. angl. 3. p. 153.

A. maritima Halimus dicta humilis erecta , folliculis in latitudinem expansis & utrinque recurvis , longo pediculo insidentibus. Pluk. alm. 61. t. 36. f. 1.

β. A. maritima nostras , ocymi minoris folio. Rai. angl. 3. p. 153.

Habitat in Angliæ , Daniæ maritimis. ☉

1266. A C E R.

Hermaphrod. Cal. 5. fidus. Cor. 5. petala. Stam. 8.

Pist. 1. Caps. 2. f. 3. monospermæ , ala terminatæ.

Masc. Cal. 5. fidus. Cor. 5. petala Stam. 8.

2. *ACER Pseudo-Platanus. A. foliis quinquelobis inequaliter serratis , floribus racemosis. ged. herb. pal. fil. In Ham. arb. 1. t. 36. f. 1. gallob. burg. lich. lugd.*

A. foliis quinquelobis acutis obtuse serratis , petiolis cuneulatis. Sauv. monsp.

POLYGAMIA. MONOGAMIA. 577

A. foliis palmatis obtusis serratis, floribus subeperalis racemosis. *Hort. cliff.*

A. foliis quinquelobis, acute serratis, racemis pendulis. *Hall. helu. n. 1029. * * * **

A. foliis quinquelobis acute serratis, florum racemo compo-
sito pendulo. *Du Roi. herbk. 1. p. 4.*

A. montanum candidum. *Bauh. pin. 430.*

A. majus. *Dod. pempt. 840.*

Habitat in Helvetiæ, Austriæ montanis. ¶

Flores flavo-virentes. Racemi subovati ternati. Lærs. l. c. R.

6. ACER platanoides. **A.** foliis quinquelobis acuminatis
acute dentatis glabris; floribus corymbosis. *Fl. suec. carn.*
norv. pal. sil. Kniph. cens. 12. n. 1. burg. lugd.

A. foliis palmatis acute dentatis, floribus corollatis dichotomo corymbosis. *Hort. cliff.*

A. foliis quinquelobis, dentibus acutis, intervalis latis; racemis erectis. *Hall. helu. n. 1029. * * * **

A. montanum, tenuissimis & acutissimis foliis. *Bauh. pin. 431.*

A. major. *Cam. epit. 63.*

A. montanum, orientalis platani foliis atrovirentibus. *Pluk. alm. 7. t. 252. f. 1.*

*Habitat in Europa boreali: usque in montibus Stiria, Sa-
paulia. ¶*

*Flores masculi die ultimo mensis frondescentia decidua
late sparguntur per terram subjacentem pulchre lutei.*

8. ACER campestre. **A.** foliis lobatis obtusis emarginatis
*Fl. suec. gallob. carn. herb. pal. sil. Kniph. cens. 3. n. 1.
burg. lith. lugd.*

A. foliis tripartito-palmatis; laciniis utrinque emarginatis
obtusis, cortice sulcato. *Hort. cliff. getr.*

A. foliis semitrilobis obtusis, lobis lateralibus emarginatis.
Hall. helu. n. 1029.

A. campestre & minus. *Bauh. pin. 431.*

A. minus. *Dod. pempt. 840. Tabern. 973.*

9. A. campestre & minus mas f. sterile. *Vaill. parisi*

Habitat in Scania & australiori Europa.

Frutex.

9. ACER monspessulanum. **A.** foliis trilobis integerrimis,
glabris annuis. *Scop. carn.*

A. foliis trilobis cordatis integerrimis, lobis aequalibus di-
varicatis. *Du Roi. herbk. 1. p. 26.*

Linnaei spec. plant. Europ. Pars II.

G. S.

478 POLYGANIA. MONOCHETIA.

A. trifolium. Bauh. pin. 431. Flak. alm. 7. t. 251. f. 3.
Duham. arb. 1. p. 28. t. 10. f. 8.
Habitat Mospeli. H

1267. CELTIS.

Hermaphrod. *Cal. 5. paritius. Cor. o. Stam. 5.*
Styl. 2. Drupa 1. sperma.

Masc. *Cal. 6. paritius. Cor. o. Stam. 6.*

1. *CELTIS australis. C. foliis ovato-lanceolatis, pist. carn.*

C. foliis lanceolatis acuminatis serratis nervosis. Mill. del

C. foliis scabris, serratis, longissime lanceolatis. Mill. del
n. 1619.

Lotus fructu cerasi. Bauh. pin. 447.

Lotus f. Celtis. Cam. epic. 135.

Lotus arbor. Lob. ic. 186.

Habitat in Europa australi & Africa citeriore. H

1271. MIMOSA.

Hermaphrod. *Cal. 5. dentatus. Cor. 5. fida. Stam.*
5. f. plura. Pist. 1. Legumen.

Masc. *Cal. 5. dentatus. Cor. 5. fida. Stam. 5.*
10. plura.

15. *MIMOSA pudica. M. aculeata, foliis subdigitatis pinnatis, caule hispido.*

M. foliis digitatis. foliolis pinnatis, caule aculeato hispidoque, foliis pinnato-palmatis: singulis numerose pinnatis.
Hort. upf.

M. humilis frutescens & spinosa, siliquis conglobatis. Plin. spec. 17. ic. 202.

M. spinosa 3, siliquis parvis echinatis. Breyer. bot. 40. t. 18.

Æschynomene spinosa, flore globoso albido, siliquis articulatis echinatis. Comm. hort. 1. p. 57. t. 29.

Habitat in Brasilia. H

Calyx infundibuliformis, trifidus: lobo altero larva.
C. ...ula ...ula, Stamina filamenta 4, calyce sexies lo-

POLYGAMIA. DIOECIA. 979

floræ, erecta, filiformia. Germen subrotundum. Stylus subulatus, staminibus paulo brevior. Stigma acutum.

Evigilat apud nos mense Junio & Julio hora 3 matutina, obdormit hora 6 vespertina.

34. *MIMOSA farnesiana. M. spinis stipularibus distinctis, foliis bipinnatis, parvis octojugis, spicis globosis sessilibus.*

Acacia indica farnesiana. Ald. farnes. 2. Rai. hist. 977.

Habitat in Domingo. H

Flores polyandri capitulis sessilibus, fragrantissimis, luteis. Legumina fusiformia. Petioli glandula infra foliola.

Sata in horto Farnesiano primum 1611; nunc hortis communis ob flores ambrosiacas.

DIOECIA.

1272. GLEDITSCHIA.

Hermaphr. Cal. 4. fidus. Cor. 4. petala. Stam. 6. Pist. 1. Legumen.

Masc. Cal. 3. phyllus. Cor. 3. petala. Stam. 6.

Fem. Cal. 5. phyllus. Cor. 5. petala. Pist. 1. Legumen.

1. *GLEDITSCHIA triacanthos. G. caule spinis triplicibus axillaribus. Du Roi. herb. 1. p. 294.*

Gleditsia. Hort. upf. 298.

Gleditsia spinosa. Du Ham. arb. 1. p. 266. t. 104.

Melilobus. Mitch. gen. 15.

Cæsalpinoides foliis pinnatis ac duplicato-pinnatis. Hort. cliff. 489.

Acacia americana, abruæ folio, triacanthos. Pluk. mant. 1. t. 352. f. 1. Hort. angl. t. 21.

β. *Acacia abruæ folio, triacanthos capsula ovali unico semine claudente. Catesb. car. 1. p. 43. t. 43.*

Habitat in Virginia. H

In mascula arbore Cor. 0. Nectar. 0. Stam. 6.

Hermaphrod. Cal. o. f. 4. partitus. Cor. o. f. 4-
petala. Stam. 2. Pist. 1. Sem. 1. lanceolatum.

Fem. Sem. 1. lanceolatum.

1. FRAXINUS *excelsior*. F. foliolis serratis, floribus apetalis. *ged. pal. fil. Blackw. t. 328. lugd. lish. burg.*

F. floribus nudis. *Fl. suec. paris.*

F. apetala, foliis pinnatis serratis. *Hall. helv. n. 528.*

F. Ornus. *Scop. carn.*

F. excelsior. *Bauh. pin. 416.*

Fraxinus. *Dod. pempt. 771.*

Die hohe Esche. *Linn. Pflanzensystem. 2. p. 512.*

Habitas in Europæ septibus. ♀

Gemmæ laterales racemos, terminalis folia promit.

2. FRAXINUS *Ornus*. F. foliolis serratis, floribus corollatis.

F. floribus completis. *Hort. ups.*

F. florifera. *Scop. carn.*

F. paniculata. *Mill. dict. mar.*

F. humilior f. altera theophrasti, minore & tenuiore folio. *Bauh. pin. 416.*

F. florifera botryoides. *Morif. prat. 265. Hort. angl. 33-
t. 9. Du Ham. arb. 4.*

F. tenuiore & minore folio. *Bauh. hist. 1. p. 177.*

Habitas in Europa australi. ♀

Petoli hujus æquales: foliolis lanceolatis æqualibus;

Præcedentis marginati: foliolis ovatis: impari maiore.

POLYGAMIA. TRIOECIA. 581

TRIOECIA.

1282. CERATONIA.

Hermaphrod. *Cal.* 5. *partitus*. *Cor.* o. *Stam.* 5.
Stylus filiform. *Legumen* coriaceum, polysperm.

Dioica. Mas & Femina distincta.

1. CERATONIA *siliqua*. *Ceratonia*. *Blackw.* t. 209.
Knorr. flor. 2. t. 8. 7. *Zorn. ic. pl. med.* cent. 1. t. 59.

Siliqua edulis. *Bauh. pin.* 400.

♀. *Siliqua*. *Cam. epit.* 139.

Habitat in Apulia, Sicilia, Creta, Cypro, Syria, Palestina, totaque Oriente. *h*

Marem Feminamque ex Australi Europa habui; Hermaphroditum ex Austria hortis misit D. Jacquin.

Non dioica est, sed hermaphrodita hexandra, raro hexandra. *Forskæhl. ægypt. p.* 77. *Hermaphroditam pentandram pronunciat Fabric. helmst.* 431. R.

1283. FICUS.

Receptaculum commune turbinatum, carnosum, connivens, occultans flosculos vel in eodem vel distincto.

Masc. *Cal.* 3. *partitus*. *Cor.* o. *Stam.* 3.

Fem. *Cal.* 5. *partitus*. *Cor.* o. *Pistillum* 1. *Sem.* 1.

1. FICUS *Carica*. *P.* foliis palmatis. *monsp.* *Hall. helv.* n. 1607. *carn.* *Blackw.* t. 125. *Ludw. ed.* t. 109. *Knorr. del. hort.* 1. t. F. *Kniph. cent.* 2. n. 25. *Regn. bot.*

F. communis. *Bauh. pin.* 457.

Ficus. *Dod. pempt.* 812.

♂. *Caprificus.* *Bauh. hist.* 1. p. 134.

Ficus hyemalis. *Bauh. pin.* 457.

Linn. Pflanz. 2. p. 330.

, Asia. *h*

CLASSIS XXIV.

CRYPTOGAMIA.
FILICES.* *Frustrificationes spicatae.*

1284. **EQUISETUM.** *Spica sparsa. Frustrif. peltata, basi valvulata.*
 1288. **OPHIOGLOSSUM.** *Spica articulata. Frustrif. circumscissae.*
 1289. **OSMUNDA.** *Spica racemosa. Frustrif. 2-valves.*

* * *Frustrificationes frondosae, in pagina inferiore.*

1290. **ACROSTICHUM.** *Macula Discum totum occupans.*
 1296. **POLYPODIUM.** *Puncta disci distincta.*
 1293. **HEMIONITIS.** *Lineae disci decussantes.*
 1295. **ASPLENIUM.** *Lineae disci sub parallelae variae.*
 1294. **LONCHITIS.** *Lineae Marginis ad finem.*
 1291. **PTERIS.** *Lineae Marginis ad peripheriam.*
 1297. **ADIANTHUM.** *Macula apicem marginis reflexo obiectae.*
 1298. **TRICHOMANES.** *Frustrif. solutae Marginis ipsi inferiorae.*

* *Frustrificationes radicales.*

1299. **M.** *Frustrif. 4 capsularis.*
 1300. **M.** *Frustrif. 4. locularis.*
 1301. **M.** *Frustrif. 2. locularis.*

CRYPTOGAMIA.

MUSCI.

* *Acalypttrati*.

1302. LYCOPODIUM. *Anthera* bivalvis, sessilis.
 1304. SPHAGNUM. *Anthera* ore laevi.
 1305. PHASCUM. *Anthera* ore ciliato.

* * *Calyptrati dichini*.

1308. SPLACHNUM. *Anth.* cum apophysi maxima.
 1309. POLYTRICHUM. *Anth.* cum Apophysi minima, marginata.
 1310. MNIMUM. *Anth.* sine Apophysi.

* * * *Calyptrati monochini*.

1311. BRYUM. *Anth.* pedunculo terminali & tuberculo.
 1312. HYPNUM. *Anth.* pedunculo laterali & Perichætio.
 1306. FONTINALIS. *Anth.* sessilis, Perichætio imbricato obvoluta.
 1307. BUXBAUMIA. *Anth.* pedunculata, altero latere membranacea.

A L G E.

* *Terrestres*.

- 1315 MARCHANTIA. *Fl.* calyce communi polistato, subrus florido.
 1313. JUNGERMANIA. *Fl.* calyce simplici, 4. valvi.
 1314. TARGIONIA. *Fl.* calyce 2. valvi.
 1316 ANTHOCEROS. *Fl.* calyce tubuloso. *Anth.* subulata, 2. valvis.
Fr. cylindrica tubulosa.

A.

1315. RIGCIA.

Fr. Granulis frondi innatis.

1319. LICHEN.

Fr. receptaculo lævi nigro.

1324. BYSSUS.

Substantia lanuginosa.

* * *Aquaticæ.*

1320. TREMELLA.

A. gelatinosa.

1322. ULVA.

A. membranacea.

1321. FUCUS.

A. coriacea.

1323. CONFERVA.

A. capillaris.

F U N G I.

Pileati.

1325. AGARICUS.

Pileus subtrus lamellosus.

1326. BOLETUS.

Pileus subtrus porosus.

1327. HYDNUM.

Pileus subtrus echinatus.

1328. PHALLUS.

Pileus subtrus lævis.

* * *Pileo, destituti.*

1329. CLATHRUS.

Fung. cancellatus.

1330. HELVELLA.

F. turbinatus.

1331. PEZIZA.

F. campanulatus.

1332. CLAVARIA.

F. oblongus.

1333. LYCOPERDON.

F. globosus.

1334. MUCOR.

F. vesicularis stipitatus.

CRYPTOGAMIA.

FILICES.

1284. EUISETUM.

Spica fructificationibus peltatis, basi dehiscentibus multivalvi.

1. EUISETUM *sylvaticum*. E. caule spicato, frondibus compositis. *Fl. suec. ged. 1. herb. pal. burg. lithu.*

E. foliis repetito-ramosis, vaginis laxissimis. *Hall. helu. n. 1680.*

E. caule sulcato; frondibus racemosis. *Scop. carn.*

E. setis ramosis internodio multoties longioribus. *Roy. lugdb.*

E. *sylvaticum*. *Fl. lapp. Tabern. hist. 562.*

E. *sylvaticum*, tenuissimis setis. *Bauh. pin. 16. theat. 245.*

Habitat in Europæ septentrionalis pratis sylvaticis. 24

2. EUISETUM *arvense*. E. scapo fructificante nudo; sterili frondoso. *ged. carn. herb. pal. Kniph. cent. 1. n. 28. Burg. lithu. lugd.*

E. scapo fructificante nudo, caule sterili ramis compositis. *Fl. suec. paris.*

E. caule florifero nudo, sterili verticillato, radiorum duodecim. *Hall. helu. n. 1676.*

E. setis quadrangularibus internodio longioribus, caulibus septantibus. *Guett. stamp.*

E. setis simplicibus internodio multoties longioribus. *Roy. lugdb.*

E. *arvense*, longicribus setis. *Bauh. pin. 16.*

E. minus. *Fuchf. hist. 323.*

Scapus caudæ equinæ. *Blackw 1. 217. f. 3.*

Habitat in Europæ & Orientis agris, pratis. 24

3. EUISETUM *palustre*. E. caule angulato, frondibus simplicibus. *ged. herb. pal. carn. burg. lithu. lugd.*

286 CRYPTOGRAMIA. FULICES.

E. caule sulcato, ramis multifloris, foliis indivisis. Hall. helu. n. 1677. var. a.

E. sens simplicibus internodiis vix superantibus. Fl. succ. parif.

E. palustre brevioribus setis. Bauh. pin. 16. theatr. 242.

B. E. palustre minus polystachion. Bauh. pin. 16. prodr. 24. Rai. angl. 3. p. 131. t. 5. f. 3. Leers. l. c.

Habitas in Europæ aquosiss. 24

4. *EQUISETUM fluviale. E. caule striato, frondibus simplicibus. herb. pal*

F. caule non sulcato latissimo, verticillis densissimis. Fl. succ. parif. burg. lithu. lugd.

F. caule storigero nudo, sterili verticillato, radiorum quadraginta. Hall. helu. n. 1675.

F. caule lævi, subnudo. Scop. carn. n. 1.

E. palustre, longioribus setis. Bauh. pin. 15. theatr. 241.

Cauda equina. Blackw. t. 277. f. 2.

Hippuris. Lob. ic. p. 793. R.

Habitas in Europa ad ripas, lacuum, fluviorum. 24

Caulis striatus, non sulcatus Hall Scop. R.

Caulis floriferi a sterilibus distincti, ut E. arvensis. Hall.

5. *EQUISETUM limosum. F. caule subnudo lævi. succ. parif. herb.*

F. scapo nudo simplicissimo. Roy. lugdb.

F. nudum lævius nostras. Rai. angl. 3. p. 131. t. 5. f. 2.

Habitas in Europæ paludibus, turfosiss, profundis. 24

Varietas E. palustris creditur Hallero.

6. *EQUISETUM hyemale. E. caule nudo scabro basi subramoso. Fl. succ. parif. ged. herb. pal. burg. lith.*

F. caule subnudo, asperrimo, vaginis caulinis indivisis, ramis foliatis. Hall. helu. n. 1679.

F. scapo nudo simplicissimo. Roy. lugdb.

B. foliis nudum ramosum. Bauh. pin. 16.

E. nudum minus variegatum basilense. Bauh. pin. 16. prodr. 25.

Equisetum. Cam. epis. 770. f. A.

Habitas in Europæ sylvis, asperis, uliginosis. 24

Caulis viridis, scaber, rodens. Vaginz articulorum pallida, basi margineque atris, denticulis obsoletis.

1288 OPHIOGLOSSUM.

Spica articulata, disticha: articulis transversim debilitantibus.

1. **OPHIOGLOSSUM vulgatum.** O. fronde ovata. *Fl. succ. parisi. carn. pal. Weis. crypt. flor. dan. 147. Kniph. cent. 6. n. 64. Knorr. del. hort. 2. t. O. 3. Blackw. t. 416. burg. lithu. lugd.*

O. folio unico, ovato-lanceolato obtuso. *Hall. helu. n. 1685.*

O. folio ovato, spica disticha. *Hort. cliff.*

O. vulgatum. *Bauh. pin. 354. Plum. fil. XXXVI. t. B. f. 5. Moris. hist. 3. p. 545. f. 15. t. 5. f. 1.*

Ophioglossum. *Bauh. pin. 364. Cam. epit. 364.*

f. O. minus, subrotundo folio. *Bauh. pin. 354. prodr. 150.*

Habitat in Europæ pratis sylvaticis.

2. **OPHIOGLOSSUM lusitanicum.** O. fronde lanceolata.

O. pumilum autumnale. *Grisl. lusit.*

O. angustifolium minimum. *Barr. rar. 1280. t. 252. f. 2.*

Habitat in Lusitania.

1289. OSMUNDA.

Spica ramosa: Fructificanti. globosis.

* *Scapis insidentibus cauli ad basin frondis.*

2. **OSMUNDA Lunaria.** O. scapo caulino solitario, fronde pinnata solitaria. *Fl. succ. parisi. Weis. crypt. pal. carn. Fl. dan. t. 18. f. 1. Kniph. cent. 11. n. 82. Blackw. t. 420. Knorr. del. hort. 2. t. O. 3. Zorn. ic. pl. med. cent. 1. t. 65. lugd. burg. lithu.*

O. fronde pinnata caulina: pinnis lunulatis. *Hort. cliff.*

O. folio pinnatifido: pinnis lunulatis. *Fl. iapp.*

O. foliis pinnatis flabelliformibus lunatis. *Hall. helu. n. 1686.*

Lunaria racemosa minor & vulgaris. *Bauh. pin. 354.*

Lunaria minor. *Cam. epit. 643.*

f. Lunaria racemosa ramosa major. *Bauh. pin. 355.*

Lunariæ rarior specie. *Cam. epit. 644.*

γ. Lunaria racemosa minor, adianti folio. *Breyn. cent. 183. t. 93. Oed. flor. dan. t. 18. f. 3.*

588 CRYPTOGANIA. FILICES.

♀. *Lunaria minor*, rutaceo folio. *Bauh. pin.* 355.

Habitat in Europa. 2.

*** *Fronde ipsa fructificationes ferente.*

12. *OSMUNDA regalis*. O. frondibus bipinnatis : apice racemiferis. *Flor. dan.* t. 217. *Blackw.* t. 324. *Kniph. cent.* 2. n. 49. *burg.*

O. scapo paniculato polyphylo. *Fl. suec. paris.*

O. frondibus caulinis simpliciter pinnatis : pinnis lanceolatis. *Hort. cliff.*

O. regalis. f. *Filix florida*. *Plum. fil.* 35. t. B. f. 4.

Filix ramosa non dentata florida. *Bauh. pin.* 357.

Filix palustris. *Dod. pempt.* 473.

♀. *Filix florida* f. *Osmunda regalis*, foliis alternis, loba seminiferis. *Gron. virg.* 123.

Filix non dentata florida, foliis alternis & in summo cute seminibus occultatis. *Pluk. alm.* 156. t. 181. f. 4.

Filix botryoites f. *floridana major virginiana*, pinnis non dentatis alternatim positis. *Morif. hist.* 3. p. 593.

Habitat in Europa, Virginia ad fluvios.

*** *Frondibus aliis foliaceis, aliis fructificantibus.*

15. *OSMUNDA Struthiopteris*. O. frondibus pinnatis pinnatifidis, scapo fructificante disticho. *Fl. suec. nov. n. I. t. I.* f. 1. 2. 3. *Fl. dan.* t. 169.

Filix palustris altera, fusco pulvere hirsuta. *Bauh. pin.* 358.

Filicetum septentrionale & palustre. *Amm. rath.* 175. ed. petrop. 10. t. 18.

Filix norvegica minor. *Rai. app.* 68.

Habitat in Suecia, Helvetia, Russia, Norvegia, Ins. *Rupp.* 2.

17. *OSMUNDA Spicant*. O. frondibus lanceolatis pinnatifidis : laciniis confluentibus integerrimis parallelis. *Fl. dan.* t. 99. *lithu.*

Pteris fronde pinnata lanceolata, laciniis parallelis integerrimis. *Fl. suec. paris.*

Pteris fronte pinnata : foliolis linearibus parallelis. *Ag. luzdb.*

Struthiopteris. *Hall. helu.* n. 1687.

Struthiopteris spicant. *Scop. carn.* 2. n. 1258.

Struthiopteris (Spicant.) frondibus sterilibus pinnatis, pinnulis densis oblongis f. ; fructificantibus majoribus, laxius pinnatis, a

CRYPTOGAMIA. FILICES. 589

Filicastrum polypodii foliis brevioribus & crassioribus. *Ad. petrop.* 10. p. 280. *Hall. R.*

Polypodioides vulgaris foliis angustis. *Mich. florent.* 163. *Tozzet. it.* 1. p. 197.

Lonchitis minor. *Bauh. pin.* 359.

Lonchitis aspera minor. *Cam. epit.* 665.

Habitat in Europa, Suecia, Misnia, Pyrenæis. 24

18. *OSMUNDA crispa.* O. frondibus supradecompositis : pinnis alternis subrotundis incis. *Huds. angl. Oed. dan.* t. 496.

Pteris (heterophylla) frondibus bipinnatis : pinnis ovato-oblongis serratis obtusis ; fertilibus integerrimis. *Spec. ed.* 2.

Filix foliis duplicato-pinnatis , sterilibus obtusis dentatis , floralibus ovatis integerrimis. *Hall. helv. n.* 1659.

Ruta muraria major , foliis variis scilicet oblongis integris & subrotundis serratis. *Sloan. jam. 21. hist. 1. p. 92. t. 53. f. 2.*

Adiantum foliis minutim in oblongum scissis. *Moris. hist. 3. p. 585. f. 14. t. 5. f. 25. Hæc adultior.*

Adiantum foliis minutim in oblongum scissis, pediculis viridibus. *Bauh. pin.* 755.

Adiantum album tenuifolium, rutæ murariæ accedens. *Bauh. hist. 3. p. 743. Rai. hist. 147.*

Adiantum album alpinum crispum. *Rai. hist. 47. angl. 3. p. 126.*

Adiantum album floridum s. *Filicula petræa crispa.* *Pluk. alm. 9. t. 3. f. 2.*

Filix botryoides s. *filicula petræa florida anglica* , foliis plurifariam divisis. *Moris. hist. 3. p. 593. f. 14. t. 4. f. 4.*

Habitat in Jamaica, Anglia, Helvetia, Monspeli, Pyrenæis. 24

Planta quasi media inter Osmundas ultimas & Acrosticha.

Conf. Filicula alpina crispa. *Bauh. pin.* 358. e *Bursero.*

1290. A C R O S T I C H U M.

Frustrificationes discum totum frondis tegentes.

* *Fronde simplici divisa.*

6. **ACROSTICHUM** *septentrionale.* A. frondibus nudis li-

390 CRYPTOGAMIA. FILICES.

nearibus laciniatis. *Fl. succ. Oed. dan. t. 60. parif. Vaf. crypt. carn. pal. lisk. burg.*

A. caule bis bifurcato Hall. helu. n. 1714.

Holosteum petraeum. Tabern. 736.

Filicula saxatilis corniculata. Bauh. pin. 338.

Filix saxatilis tragi. Lob. ic. 47.

Habitat in Europae fissuris rupium. 2.

22. *ACROSTICHUM Marantæ. A. frondibus subbipinnatis: pinnis opposito-coadunatis subtus hirsutissimis: basi subdentatis.*

A. pinnis subtus pinnatis, squamosis, obtusis, basi utraque appendiculata. Hall. helu. n. 1715.

Lonchitis folio ceterach. Bauh. pin. 359. Bar. ic. 857.

Lonchitis aspera marantæ Cam. epit. 666.

Filicula crispa, lanugine hepatici coloris vestita. Fluk. alm. 150. t. 281. f. 4.

Habitat in Europa australi,

Stipites spithamei, atr. purpurei. Fronis ferme bipinnatis lanceolata, subtus dense imbricata squamis paleaceis, croceis, exsiccata contrahitur & occultat paginam internam. Foliola opposita, fere pinnata. Pinnæ ovatae, obtusae, crossae, integerrimae; at inferiores basi saepe utrinque dentatae. Fructificationes sparsae inter squamas foliaceas per totam paginam possicam.

23. *ACROSTICHUM iluense. A. frondibus subbipinnatis: pinnis oppositis - coadunatis obtusis subtus hirsutis; basi integerrimis. Flor. dan. t. 391. Kniph. cent. 10. n. 1.*

Polypodium fronde duplicato pinnata: pinnulis coadunatis obtusis subtus hispidis. Fl. succ.

Polypodium duplicato-pinnatum: pinnulis obtusis ramosis subtus villosum. Fl. lapp.

Polypodium lonchitidis marantæ species cambro-brinnica. Pluk. alm. 150. t. 89. f. 5.

Filicula alpina teretior, alis latiusculis brevioribus integris profunde dentatis. Moris. hist. 3. p. 576. f. 14. t. 3. j. 23.

Habitat in Europae frigidissima rupibus.

Maxime affinis precedenti, at vix digito longius, ut spithameum; stipite virescente, nec atropurpureo ut illud. Proximuni Polypodium ob praectis fructificationum, sed vob conseria.

1291. P T E R I S.

Fructificationes in lineis marginalibus.

15. **PTERIS aquilina.** P. frondibus supradecompositis : foliolis pinnatis : pinnis lanceolatis : infimis pinnatifidis : superioribus minoribus. *Fl. succ. paris. carn. norv. Weis. crypt. pal. 190. lugd. lithu. burg.*

Filix foliis triplicato-pinnatis, pinnis nervosis integerrimis, ultimis lanceolatis. *Hall. helu. n. 1688.*

Filix ramosa major, pinnulis obtusis non dentata. *Bauh. pin. 357. Plum. amer. 23. t. A. f. 1.*

Filix femina. *Fuchs. hist. 596. Cam. epis. 992. Matth. 291. Blackw. t. 325.*

Habitat in Europæ sylvis, præsertim cæduis.

1295. A S P L E N I U M.

Fructificat. in lineolis disci frondis sparsis.

* *Fronde simplici.*

1. **ASPLENIUM Hemionitis.** A. frondibus simplicibus cordato-hastatis quinquelobis integerrimis, stipitibus lævibus.

Hemionitis vulgaris. Bauh. pin. 353.

Hemionitis vera. Clus. hist. 2. p. 214.

Habitas in Italia, Hispania.

Maxime affine sequenti, at diameter frondis longitudinalis vix superat transversalem.

3. **ASPLENIUM Scolopendrium.** A. frondibus simplicibus cordato-lingulatis integerrimis, stipitibus hirsutis. *carn. Weis. crypt. pal. Kniph. cent. 10. n. 12. Regn. bot. Zorn. ic. pl. med. cent. 1. t. 47. burg. lugd.*

A. petiolis hirsutis, folio longe lineari lanceolato integerrimo, prostrato cordato. *Hall. helu. n. 1695.*

A. frondibus ensiformibus integris basi cordatis inflexis, petiolis hirsutis. *Fl. succ. paris. monsp.*

Lingua Cervina officinarum. Bauh. pin. 233. Plum. fil. 34. t. A. f. 4. Blackw. t. 138. Knorr. del. 2. t. L. 8.

Phyllitis. Com. epis. 579. Matth. 83. epis. 579.

β. Phyllitis crispa. Bauh. hist. 3. p. 757.

592 CRYPTOGAMIA: FILICES:

7. Phyllitis f. Lingua Cervina maxima, undulato folio auriculato per basin. *Pluk. phyt.* 243. f. 1.

8. Lingua Cervina medio folio nervo in aculeum abeundi. *Moris. hist.* 3. p. 557. f. 14. t. 1 f. 9.

9. Lingua Cervina multifido folio. *Bauh. pin.* 333.

10. Phyllitis f. Lingua Cervina crispa, folio multifido, ramosa. *Pluk. phyt.* 248. f. 2.

Habitat in Europæ umbrosis, nemorosis, saxosis. 2x.

* * Fronde pinnatifida.

8. ASPLENIUM *Ceterach*. A. frondibus pinnatifidis: obsoletis alternis confluentibus obtusis. *paris. monsp. angl. carn. Blackw.* t. 216. *Regn. bot. burg. lugd.*

A. foliis pinnatis, pinnis in basi dilatatis, obtusis, subtus muscosis. *Hall. helv.* n. 1694.

Asplenium f. *Ceterach*. *Bauh. pin.* 3. p. 479. *Moris. hist.* 3. p. 561. f. 14. t. 2. f. 1.

Barr. ic. 1043. 1044. 1051. *l. XXXIII.* t. 8. f. 3. 52. 604. & *forte* 601.

Ceterach officinarum. *Bauh. pin.* 4. *Habitat in Orientis, Monspeliæ, Italiae, Hercynia fissuris rupium humidis.*

Folia subtus adeo squamosa, ut fructificationes non conspiciantur.

* * * Fronde pinnata.

12. ASPLENIUM *trichomanes*. A. frondibus pinnatis: pinnis subrotundis crenatis. *paris. carn. Weis. crypt. pal. Flor. dan.* t. 119. *aph. cent.* 6. n. 14. *Blackw.* t. 370. *Regn. bot. Zorn. ic. pl. med. cent.* 1. t. 95. *burg. lithu. lugd.*

A. foliis pinnatis, pinnis ovatis, crenatis. *Hall. helv.* n. 1693.

Trichomanes f. *Polytrichum officinarum*. *Bauh. pin.* 356. *Plum. fil.* 26. t. V. f. i.

Trichomanes. *Fuchs. hist.* 796. *Cam. epic.* 925.

β. *Trichomanes* foliis eleganter incis. *Tournef. inst.* 539. t. 315. f. J. C.

Adiantum maritimum segmentis subrotundis. *Pluck. phyt.* 73. f. 6.

γ. *Trichomanes minus* & tenerius. *Bauh. pin.* 356.

Habitat in Europæ fissuris rupium, inque Oriente.

14. ASPLENIUM *marinum*. A. frondibus pinnatis: pinnis obovatis serratis superne gibbis obtusis: basi cuneatis.

Chamae Filix marina aeglea. *Moris. hist.* 3. p. 573. f. 14. t. 3. f. 25. *bon.*

Pluk.

Filicula maritima ex insulis stoechadis. Bauh. pin. 359. Bursf. XX. 39.

8. *Asplenium minus assurgens simplex, foliis oblongis margine inæquali crenato. Brown. jam 93.*

Adiantum jamaic. pinnulis auriculatis ad basin strictioribus. Pluk. alm. 9. t. 253. f. 5.

Lonchitis minor, pinnis latoribus leviter denticulatis, superiore latere auriculatis. Sloan. jam. 16. hist. 1. p. 78. t. 33. f. 1?

Habitat in Anglia, insulis Stochadum. Bursf. America.

19. *ASPLENIUM Ruta muraria. A. frondibus alternatim decompositis: foliolis cuneiformibus crenulatis. Fl. suec. parif. carn. Weis. crypt. 298. pal. Flor. dan. t. 190. Kniph. cent. 6. n. 13. Regn. bot.*

A. fronde laxa ramosa, ramis secundis trifoliis, superioribus semitrilobis rhomboideis, circumferratis. Hall. helv. n. 1691.

Ruta muraria. Bauh. pin. 356.

Adiantum album. Tabern. 796. Blackw. t. 219.

Baronychia. Cam. epit. 785. Matth. 1041.

Habitat in Europa ex rupium fissuris. 24.

20. *ASPLENIUM Adiantum nigrum. A. frondibus subtripinnatis: foliolis alternis pinnis lanceolatis inciso ferratis. carn. pal. Fl. dan. t. 250. Blackw. t. 220. Kniph. cent. 7. n. 3. Regn. bot.*

A. foliis triangularibus, pinnis pinnatis, pinnulis semipinnatis, lobulis ovatis, ferratis. Hall. helv. n. 1692.

A. frondibus duplicato-pinnatis, foliis inferioribus majoribus: foliolis ovatis superne crenatis. Dalib. parif.

Adiantum pulcherrimum. Bauh. hist. 3. p. 737.

Adiantum foliis longioribus pulverulentis, pediculo nigro. Bauh. pin. 355.

Dryopteris nigra. Dod. pempt. 466.

Onopteris major. Tabern. p. 796. R.

Habitat in Italia, Gallia, Hareynia, Anglia.

Pollen crocei est coloris.

21. *ASPLENIUM Trichomanes ramosum. A. frondibus duplicato-pinnatis: pinnis obovatis crenatis: foliolis inferioribus minoribus.*

Trichomanes ramosum majus & minus. Bauh. pin. 356.

Trichomanes ramosum. Bauh. hist. 3. p. 747. Rai. angl. 3. p. 119.

Habitat in Arvonie rupibus.

Linnaei spec. plant. Europ. Pars II.

P r

1296. POLYPODIUM.

Fruſif. in punctis ſubrotundis, ſparſis per diſcuti frondis.

* * *Fronde pinnatifida lobis cœdunatis.*

14. POLYPODIUM *vulgare*. P. frondibus pinnatifidis : pinnis oblongis ſubſerratis obtuſis, radice ſquamata. *carn. Veſf. crypt. pal. Blackw. t. 215. Kniph. cent. 6. n. 72. Regn. bot. Zorn. ic. pl. cent. 1. t. 46. burg. lugd.*

P. fronde pinnata, pinnis lanceolatis indiviſis ſerrulatis alternis connato-ſeſſilibus. *Fl. ſuec. pariſ.*

P. foliis pinnatis, lanceolatis, radice ſquamata. *Hall. herb. n. 1696.*

P. pinnatum pinnis, lanceolatis integris. *Fl. lapp. 380.*

P. *vulgare*. *Bauh. pin. 359. Plum. fil. 27. t. A. f. 2.*

P. *majus*. *Dod. pempt. 464.*

Filicula ſ. Polypodium. *Cam. epit. 993. Hall. R.*

β. Polypodium minus. *Bauh. pin. 359. Dod. pempt. 464. Habitat in Europæ rimis rupium.*

23. POLYPODIUM *cambricum*. P. frondibus pinnatifidis ; pinnis lanceolatis lacero pinnatifidis ſerratis.

P. fronde pinnatifida : foliolis lanceolatis ſinuato pinnatifidis. *Hort. cliſſ.*

P. cambro-britannicum, lobis foliorum profunde dentatis. *Moriſ. hiſt. 3. p. 565. f. 14. t. 2. f. 8. bene.*

P. cambro-britannicum, pinnulis ad margines laciniatis. *Raiſ. hiſt. 237. angl. 3. p. 117.*

Filix ampliſſima, lobis foliorum laciniatis, cambrica. *Pluk. alm. 153. t. 30. f. 1.*

Habitat in Anglia, Monſpelii.

* * * * *Fronde pinnata.*

27. POLYPODIUM *Lonchitis*. P. frondibus pinnatis ; pinnis lunulatis ciliato ſerratis declinatis, ſtipitibus ſtrigosis. *carn. Flor. dan. t. 497. norv. burg.*

P. foliis pinnatis, pinnis ciliatis, dentatis, appendiculatis. *Hall. helu. n. 1711.*

Lonchitis aspera. *Bauh. pin. 359.*

Lonchitis aspera major. *Moriſ. hiſt. 3. p. 566. f. 14. t. 2. f. 1.*

β. *Lonchitis mas*, ſ. *latifolia aspera minor*. *Barr. ic. 1121.*

Habitat in Alpinis Helvetiæ, Baldi, Arvonix, Walliæ, Monſpeli, Daliæ, Sueticæ, Virginix.

CRYPTOGAMIA. FILICES. 399

*Virginica plantæ pinnæ magis elongatæ & minus incurvatæ
serraturis minus ciliato-spinosis.*

37. **POLYPODIUM fontanum.** P. frondibus pinnatis lanceolatis : foliolis subrotundis argute incisiss stipite lævi. herb.

P. duriusculum pinnis laxè pinnatis , pinnulis bi & tridentatis. *Hall. helu. n. 1706.*

P. frondibus bipinnatis , foliolis argute incisiss, inferioribus subpalmatis , stipite lævi. *Gouan. illustr. 80.*

Acrostichum frondibus supinnatis : pinnis alternis cuneiformibus incisiss. *Ger. prou.*

Filicula saxatilis omnium minima. *Tournef. inst. 71.*

Adiantum album. *Lob. ic. 810. Hall. R.*

Filicula fontana. *Tabern. 792. Hall. R.*

Filicula fontana minor. *Bauh. pin. 258.*

Adiantum filicinum durius crispum minimum. *Bar. ic. 452. f. 1.*

Habitat in Sibiria , Galloprovincia , Helvetia , &c.

Habitus P. fragilis , at Foliola archiora , minus profunde subdivisa . punctis floriferis majoribus , prodeuntibus non ex squama subrotunda , sed ex rima s. valvula lineari , oblonga , alba.

***** *Fronde subbipinnata , cujus pinnæ confluent basi , ut semipinnata potius , quam perfecte duplicato-pinnata sit.*

38. **POLYPODIUM Phegopteris.** P. frondibus subbipinnatis : foliolis infimis reflexis : paribus pinnula quadrangulata coadunatis. *Fl. succ. norv. Vais. crypt. pal.*

P. foliis pinnatis , reflexis , pinnis ovatis , hirsutis primis cum nervo confluentibus. *Hall. helu. n. 1698.*

P. pinnatum : pinnis lanceolatis pinnatifidis integris : inferioribus nutantibus. *Fl. lapp.*

Phyllitis minor hirsuta , pinnis variis. *Amm. act. petrop. 10. t. 21.*

Filix non ramosa minor & sylvestris. *Petit. epist. 56. t. 504*

Filix minor britannica , pediculo pallidiore , alis inferioribus deorsum spectantibus. *Moris. hist. 3. p. 575. f. 14. t. 4. f. 17.*

Habitat in Europæ fagetis & in Virginia.

40. **POLYPODIUM fragrans.** P. frondibus subbipinnatis lanceolatis : foliolis confertis : lobis obtusis serratis , stipite paleaceo. *Huds. angl*

Dryopteris rubrum idæum spirans. *Amm. ruth. 251.*

596 CRYPTOGAMIA. FILICES.

Habitat in Sibiria, Anglia.

Habitus P. f. maris, at longe minus. Foliola densius congesta, lobis lateralibus obtusis, profundius serratis.

43. **POLYPODIUM cristatum.** P. frondibus sub-bipinnatis, foliolis ovato, oblongis: pinnis obtusiusculis apice acute serratis. *carn. burg. lugd. lithu.*

P. fronde duplicato-pinnata: pinnulis obtusis coadunatis crenatis. *Fl. suec.*

P. frondibus subbipinnatis: foliolis ovato-oblongis: pinnis obtusiusculis apice acute serratis: serraturis acuminatis. *Schreb. spicil. p. 71.*

P. pinnulis pinnatis; lobulis semipinnatis dentatis aristatis. *Hall. helv. n. 1075.*

P. (*Filix femina, cristata*) frondibus subtripinnatis; pinnis summis pinnatifidis, imis pinnatis; pinnulis repetito pinnatifidis, omnibus undique serratis, serraturis spinosis. *Weis. crypt. 317. f. R.*

P. frondibus subtripinnatis: pinnulis utraque serratis. *Müll. Fridr. p. 193. n. 845. t. 2. f. 4. R.*

Filix ramosa dentata, ramulis & pinnulis longius a se invicem distantibus. Mapp. als. 106. f. 8. b. R.

Filix non ramosa major pinnulis latiusculis longioribus in profundos denticulos divis. Moris. hist. 3. f. 14. t. 3. f. 11.

Filix mas ramosa, pinnulis dentatis. Pluk. phys. 181. f. 2. Habitat in Europa septentrionali.

florent huic foliola superiora, non inferiora.

Quæ ego adduxi synonyma, videntur quidem ad Poly-podium cristatum Lin. pertinere, licet denominatio Linneana non plane respondeat. R.

44. **POLYPODIUM filix mas.** P. frondibus bipinnatis: pinnis obtusis crenulatis, stipite paleaceo. *Fl. suec. paris. norv. t. 1. f. 4. carn. Weis. crypt. pal. Blackw. t. 323. Kniph. cent. 6. n. 71. Regn. bot. burg. lithu. lugd.*

P. pinnis pinnatis obtusis dentatis. *Hall. helv. n. 1701.*

Filix mas pinnulis cristatis. Veill. paris. t. 9. f. 2. Hall. R.

Filix mas non ramosa dentata. Bauh. pin. 358.

Filix mas. Fuchs. hist. 594. Tabern. 591.

Habitat in Europæ sylvis.

Flores reniformes.

45. **POLYPODIUM filix femina.** P. frondibus bipinnatis, pinnulis lanceolatis pinnatifidis acutis. *Fl. suec. paris. norv. carn. pal. Blackw. t. 325. burg. lithu. lugd.*

CRYPTOGAMIA. FILICES. 597

P. pinnis pinnatis, pinnulis lanceolatis, semipinnatis, lobis acute bisectis. *Hall. helu. n.* 1704.

P. frondibus bipinnatis; pinnis pinnatis confertis, pinnulis pinnatifidis, primis majoribus, lacinulis acuta bifidis. *Veis. crypt. β. R.*

Filix mas non ramosa, pinnulis angustis rarioribus profunde dentatis. *Moris. hist. 3. p.* 579. *f.* 14. *t.* 3. *f.* 8. *Pluk. phyt.* 180. *f.* 4.

Filix petræa femina 1. *Tabernæm.* 793. *Hall. R.*

Habitat in Europæ frigidioris subhumidis.

Flores ovati, subciliati, solitarii.

46. POLYPODIUM Thelypteris. **P.** fronde bipinnata: pinnis pinnatifidis integerrimis subtus undique polline rectis. *Dærr. nass. Flor. dan.* 760.

P. pinnis pinnatis, pinnulis acutis integerrimis. *Hall. helu. n.* 1697.

P. frondibus pinnatis; pinnarum laciniiis cum nervo connatis, ovatis, integerrimis, prima majore. *Veis. crypt.*

Acrostichum (Thelypteris) frondibus pinnatis: pinnis pinnatifidis integerrimis. *Spec. pl. ed. 2. fl. suec. Reyg. ged.*

Filix mollis f. glabra, vulgari mari non ramosæ accedens. *Bauh. hist.* 730. *t.* 731. *f.* 1.

Filix non ramosa minor, sylvatica, repens. *Moris. hist. 3. p.* 550. *n.* 17. *R.*

Filix petrea femina 3. *Tabern. hist.* 1182. *Hall. R.*

Filix fontis admirabilis ad marem vulgarem non ramosam accedens. *Bauh. hist. 3. p.* 731. *non figura.* *Hall. R.*

Filix tenuissime & profunde dentata. *Bauh. hist. 3. p.* 731. *figura, non descriptio.* *Hall. R.*

Filix minor non ramosa *Bauh. hist. 3. p.* 149. *Mapp. alsat. ic.* 7. *Hall. R.*

Thelypteris palustris non ramosa. *Rupp. jen.* 277. *Schmide. ic, p.* 45. *t.* 11. 13.

Habitat in Europæ septentrionalioris paludibus.

Habitus Polyd. Filicis maris. Stipes glaber. Folia sterilia latoria, obtusiora.

Filix mollis Bauhini non huc pertinet, ast ad Polypod. molle Schreberi *spicil. p.* 70. *R.*

47. POLYPODIUM aculeatum. **P.** frondibus bipinnatis: pinnis lunulatis ciliato-dentatis stipite strigoso. *paris. Hudf. angl. carn. Veis. crypt. pal.*

pinnis pinnatis, ciliatis serratis, appendiculatis. *Hall.*

998 CRYPTOGAMIA. FILICES.

Filix aculeata major. Bauh. pin. 358. prodr. 151.

Filix mas non ramosa, pinnulis latis articulatis spinosis Pluk. alm. 152. 179. f. 6. & 180. f. 1. Moriz. hist. 3. p. 580. f. 14. t. 3. f. 15.

Habitat in Europa.

48. *POLYPODIUM rhaeticum.* P. frondibus bipinnatis; foliolis pinnisque remotis lanceolatis: serraturis acuminatis. monsp. angl. carn. burg.

P. pinnis pinnatis, laxè divisis, pinnulis semipinnatis acutis. Hall. helv. n. 1708. R.

P. fumaroides laciniatum. Veis. crypt. an? R.

Filix rhaetica tenuissima denticulata. Bauh. hist. 3. p. 740.

Filicula fontana major s. *Adiantum album*, filicis folio. Bauh. pin. 358.

Habitat in Gallia, Helvetia, Anglia, Germania.

53. *POLYPODIUM fragile.* P. frondibus bipinnatis: foliolis remotis; pinnis subrotundis incis. et succ. parif. Fl. dan. t. 401. carn. lithu.

P. pinnis pinnatis; laxè divisis, pinnulis semipinnatis, lobulis subrotundis dentatis. Hall. helv. n. 1707. R.

P. (fumaroides, lobatum) frondibus bipinnatis: pinnis remotis suboppositis; pinnulis obtruse lobatis, fructificationibus dense confertis. Veis. crypt.

Adiantum filicinum aquaticum mollius, minimum. Barrel. ic. p. 432. f. 2. Hall. R.

Filix saxatilis, cauliculo tenui fragili. Pluk. alm. 150. t. 180. f. 5.

Filix pumila saxatilis. 2. Clus. pon. 706.

Filix saxatilis non ramosa nigris maculis punctata. Bauh. pin. 358?

Habitat in collibus Europæ frigidioris.

Synonyma Clusii & G. B. ad suum *Polypodium latissimum*, fumaroidem refert. Veis. crypt. p. 321. R.

55. *POLYPODIUM pteridioides.* P. (fragrans) frondibus bipinnatis: pinnis ovatis sublobatis obtusis subtus mucronatis margine reflexis, fructificationibus marginalibus.

Habitat in Gallia australi. Baro capacinus; ad muros Fuchsal. Kenig.

Satura P. fragilis. Recens siccatum fragantissimum. Stirps atropurpureus, lavis, basim versus paucis helvatis. Frons palmatis, bipinnata, lavis. Pinnæ oppositæ, bifides, ovatae, obtusæ, utrinque laeves, pinnatifidae, nigrae tenues reflexo fructificationes primum regente et

Pteride. Fructificationes marginales tantum, tectæ squama propria reniformi, omnino Polypodii.

Species Pteridi proxima, ad Polypodia ob squamam propriam relatum.

56. POLYPODIUM *regium*. P. frondibus bipinnatis : foliolis suboppositis; pinnis alternis laciniatis. *paris. carn. burg.*

Filix regia, fumaris pinnulis. Vaill. paris. 53. t. 9. f. 1. Habitat in Gallia.

Fructificationum pulvilli in hoc majores, quam in affinis.

Nomen Vaillantii ad P. fragile spectat. Hall. Veis. R.

57. POLYPODIUM ? *leptophyllum*. P. frondibus bipinnatis : sterilibus brevissimis : pinnis cuneiformi-lobatis. *Magn. monsp. 5. t. 5.*

Adiantum minimum, folia vario, Magn. hort. 5. t. 5.

Adiantum filicinum leptophyllum elatius hispanicum. Barr. rar. 1270. t. 431. opt.

Habitat in Hispania, Lusitania, Galloprovincia.

Medium inter Acrosticha, Polypodia, Osmundas.

* * * * * *fronde supradecomposita.*

67. POLYPODIUM *Dryopteris*. P. frondibus supradecompositis : foliolis ternis bipinnatis. *fl. suec. paris. ged. 1. p. 246. n. 3. Scop. carn. ed. 2. n. 1276. Veis. crypt. pal. flor. dan. t. 759.*

P. trifidum, ramis pinnatis : pinnis pinnatifidis. fl. lapp.

P. pinnis pinnatis, conjugatis, pinnulis ovatis, obtusis, crenatis. Hall. helv. n. 1699.

Thelypteris minor, pinnulis dentatis. Amm. ad. petrop. 10. p. 301. t. 22. f. 2.

Filix petraea 4. Tabern. p. 799. Hall. R.

Felix ramosa saxatilis, nigris maculis punctata. C. B. Moris. hist. 3. f. 14. t. 4. f. 19. Hall. R.

Felix ramosa minor, pinnulis dentatis. Bauh. pin. 358 Bauh. hist. 3. p. 733.

β. Filix pumila saxatilis. Clus. hist. 2. p. 212.

Habitat in Europæ nemoribus.

600 CRYPTOGAMIA. FILICES.

1297. ADIANTHUM.

Fructificationes in masculis terminalibus sub replicato margine frondis.

* * *Fronde composita.*

* * * *Fronde decomposita.*

12. ADIANTHUM *Capillus veneris*, A. frondibus decompositis : foliolis alternis : pinnis cuneiformibus lobatis pedicellatis. *Huds. angl. carn.*

A. periolis ramosis, foliis flabelliformibus lobatis, *Hall. helv. n. 1713.*

A. ramis pinnatis, foliis circuli sectorem referentibus, *Sammonsp.*

A. foliis coriandri. *Bauh. pin. 356. Mat. med. 480.*

Adiantum. Cam. epit. 924.

Habitat in Europa australi, Oriente.

1298. TRICHOMANES

Fructificationes solitariae, Stylo fetaceo terminatae, margini ipsi frondis insertae.

* * *Fronde composita.*

1. TRICHOMANES *pyxidiferum*. T. frondibus sub bipinnatis : pinnis alternis confertis lobatis linearibus. *Huds. angl.*

Filix pyxidifera. Plum. fil. 74. t. 50.

Filix humilis repens. Rai. angl. 3. p. 127. t. 3. f. 3. 4.

Darea tunbrigensis facie. Pet. fil. 104. t. 13. f. 13.

Habitat in America, Anglia.

6. TRICHOMANES *cambrigensa*. T. frondibus pinnatis : pinnis oblongis dichotomis currentibus dentatis *Hall. angl.*

Adiantum petrum perpusillum anglicum foliis bifidis trifidisque Rai. angl. 3. p. 123. suppl. 77.

Adiantum radicosum humisparsum f. Filicula pellucida nostrae coriandri foliis mollicellis, globuliferum. Pluk. atm. 10. 3. f. 5. & 6.

Darea tunbrigensis minor, Pet. mus. 761.

CRYPTOGAMIA. FILICES. 601

Muscus montanus italicus, *adianthi foliis*. *Bocc. musf.* 2. p. 24. t. 2. f. 1.

Habitas in Anglia, Italia, Jamaica.

Globulos seminales profert inter valvulas duas foliaceas.

10. *TRICHOMANES canariense*, T. frondibus supradecompositis tripartitis : foliolis alternis : pinnis alternis pinnatifidis.

Felix ramosa canariensis, rutæ murariæ pinnulis angustis altius incisis mediæ costæ alternatim alligatis. *Pluk. alm.* 156. t. 2. 291. f. 2.

β. *Polypodium (lusitanicum)* frondibus supradecompositis, foliolis alternis : pinnis oblongis longitudinaliter pinnatifidis. *Spec. ed.* 2. p. 1556. *Roy. lugdb.* 100.

Felix lusitana, polypodii radice. *Magnol. hort.* 79. t. 79.

Habitas in Canariis, Lusitania, ad latera montium.

1299. MARSILEA.

Flores masculi supra frondem.

Fructif. Feminea ad radicem, subrotunda, 4. capsularis.

1. *MARSILEA natans*. M. foliis oppositis simplicibus. *Schol. barb. Guenard. in Ad. gall.* 1762. p. 543. t. 29. f. 1.

M. (*salvinioides*) foliis oppositis, utrinque pilosis, longe radicatis, globiferis. *De Neck. in ad. pal. vol. 3. phys.* p. 297. t. 21.

Salvinia vulgaris aquis innatans, foliis subrotundis punctatis læte virentibus. *Mich. gen.* 107. t. 58. *Last. u.* 281.

Lenticula palustris latifolia punctata. *Bauh. pin.* 362.

Lens palustris patavina. *Bauh. hist.* 3. p. 785.

Habitat in Italiæ fossis paludosis stagnantibus lente fluentibus ; in America meridionali.

Rami natantes. Folia bisaria, ovata, complicata adpersa supra glomerulis et Filamentis 4, albis, apice cirrhosis ; inter radices globi 3. 7, stylo instructi, polyspermi. *Forster.*

2. *MARSILEA quadrifolia*. M. foliis quaternis integerrimis.

Zaluzianskia (marsiroides) foliis terminalibus quaternis utrinque glabris, capitulo peniolati solitario. *De Neck*

602 CRYPTOGRAMIA: FILICES:

ad. pal. vol. 3. phys. p. 303. cum charactere a Marsilea discrepante. R.

*Lemna. Juss. ad. 1740. p. 263. Guett. flam. 1. p. 62.
Lenticula palustris quadrifolia. Bauh. pin. 362. Mapp. alsat.
166. t. 166.*

*Lens palustris quadrifolia. Bauh. hist. 3. 789. Moris.
hist. 3. 629. f. 15. t. 4. f. 5.*

Lens palustris altera. Cam. epit. 853.

Habitat in Indiæ, Sibiria, Gallia, Alsatia fossis.

Capsulae edentulae.

1300. PILULARIA:

Flor. Masculi ad latus frondis.

Fructif. Feminea ad radicem, globosa, quadrilobularis.

1. PILULARIA globulifera. *Pilularia. Fl. suec. Flor. dan.
dan. t. 223. Pal. it. 1. p. 432.*

*P. palustris juncifolia. Daill. paris. 159. t. 15. f. 6. Dill.
musc. 538. t. 79. Juss. ad. 1739. p. 340. t. 11.*

*Graminifolia palustris repens, vasculis granorum piperis
æmulis. Rai. hist. 1325. angl. 3. p. 236. Moris. 3. p. 608.
f. 15. t. 7. f. 49.*

Gramen piperinum. Pet. herb. t. 29. f. 8.

*Musculus aureus capillaris, palustris inter folia folliculia
rotundis quadripartitis. Pluk. alm. 246. t. 48. f. 1.*

Habitat in Europæ inundatis.

1301. ISOETES.

Flor. Masculi Anthera, intra basin frondis.

*Flor. Feminei capsula bilocularis, intra basin
frondis.*

1. ISOETES lacustris. *Isoetes. It. Fl. suec. xl. dan. t. 191.
Marsilea foliis subulatis semicylindricis articulatis. F. suec.
996.*

*Camalaria folio longiore & graciliore. Dill. musc. 541.
t. 80. f. 2.*

*Subularia lacustris s. Calamistrum herba aquatica alpina
Rai. angl. 1. p. 210. t. 210.*

Habitat in Europæ frigida fundis lacuum.

MUSCI.

1302. LYCOPODIUM.

Anthera bivalvis, sessilis. *Calyptra* nulla.

4. LYCOPODIUM *clavatum*. L. foliis sparsis filamentosis, spicis teretibus pedunculatis geminis. *Reyg ged. Weis. crypt. fl. dan. t. 126. Blackw. t. 535. Zorn. ic. pl. med. cent. I. t. 54. lith. lugd. burg.*

L. caule repente, foliis patulis, pedunculis spica gemina terminatis. *Fl. suec. paris.*

L. ramis prolixis, spicis petiolatis gemellis, foliis linearibus pilo cristatis. *Hall. helv. n. 1722.*

L. (*officinale*) surculo vago reptante, ramoso: terminali erecto, imbricato, dichotomo, capitulifero. *De Neck. meth. 150.*

L. vulgare pilosum anfrigosum & repens. *Dill. musc. 441. t. 58. f. 1.*

Musculus terrestris repens, binis clavis foliosis se in altum erigentibus. *Pluk. alm. t. 47. f. 8. Moris. hist. 623. f. 15. t. 5. f. 2. Hall. R.*

Musculus terrestris clavatus. *Bauh. pin. 360. Pluk. phyt. 47. f. 8.*

Habitat in Europæ sylvis muscosis.

6. LYCOPODIUM *selaginoides*. L. foliis sparsis ciliatis lanceolatis, spicis solitariis terminalibus foliosis. *dan. t. 70.*

L. spicis sessilibus, foliis ovario lanceolatis? serratis, confertis. *Hall. helv. n. 1717. t. 56. f. 1.*

L. surculo decumbente sparso: terminali erecto, alis capitulifero, foliis lanceolatis ciliatis recurviusculis. *Neck. meth.*

L. spicis solitariis sessilibus, foliis aculeato-ciliatis, capsulis tetraspermis. *fl. suec.*

L. caule repente, ramis spica foliolis patula instructis. *fl. lapp. 421.*

Selaginoides foliis spinosis. *Dill. musc. 460. t. 68. f. 1. Habitat in Europæ pascuis muscosis.*

8. LYCOPODIUM *inundatum*. L. foliis sparsis integerrimis, spicis terminalibus foliosis. *fl. dan. t. 336.*

604 CRYPTOGRAMMA MUSCUL.

L. caule repente, spicis solitariis sessilibus foliosis, foliis brevibus. *Fl. suec. parif.*

L. spicis sessilibus, foliis linearibus confertissimis, *peruvia* Hall. *helu. n. 1721.*

L. furculo reptante sparso, terminali recto, foliis spiciformi. *Neck. meth.*

L. palustre repens, clava singulari. *Veist. parif. 133. t. 16. f. 11. Dill. musc. 452. t. 62. f. 7.*

Habitat in Europæ inundatis.

9. LYCOPODIUM Selago. L. foliis spicis octisariis, caule dichotomo erecto fastigiato, floribus sparsis. *Rey. can. Veis. crypt. pal. Fl. dan. t. 104. lithus.*

L. caule erecto dichotomo. *Fl. suec.*

L. spica sessili, foliis linearibus lanceolatis confertis. Hall. *helu. n. 1716.*

L. furculo erecto dichotomo capitulifero: primordialibus decedentibus sparsis. *Neck. meth. 148.*

Selago vulgaris abietis rubræ facie. *Dill. musc. 435. t. 56. f. 1.*

Muscus erectus ramosus saturate viridis. *Haub. pin. 360.*

Muscus erectus abietiformi. *Moris. hist. 3. p. 624. f. 3. t. 5. f. 9.*

Habitat in Europæ borealis sylvia acerofis.

11. LYCOPODIUM annotinum. L. foliis spicis quinquefariis subserratis; furculis annuino articulatis, spicis terminalibus glabris erectis. *Ingr. Veis. crypt. 11. t. 126.*

L. caule erecto, spica glabra terminali. *Fl. lapp. 419. Fl. suec.*

L. ramis prolixis, spicis sessilibus, foliis linearibus subserratis. Hall. *helu. n. 1720.*

L. furculo reptante adscendente repetito: terminali tereti solitario, capitulifero sessili, foliis serrulatis paulo densis. *Neck. meth.*

L. elatius juniperinum, clavis singularibus sine pediculis. *Dill. musc. 455. t. 63. f. 6. Giff. t. 2.*

Muscus clavatus, juniperinis foliis reflexis, clavis singularibus sine pediculis. *Moris. hist. 3. p. 624. f. 15. t. 5. f. 3.*

Muscus terrestris repens, clavis singularibus erectis. *Phil. alm. 248. t. 205. f. 5.*

Habitat in Europæ nemoribus.

15. LYCOPODIUM alpinum. L. foliis quadrifariis imbricatis acutis caubus erectis bifidis, spicis sessilibus, ramibus. *Veis. crypt. Fl. dan. t. 79.*

L. spicis sessilibus, foliis lanceolatis adpressis, utrinque ternatis. Hal. helu. n. 1719.

L. surculo adulto reptante, cæteris imbricato-fasciculato-fasciculatis. terminalibus capituliferis, erectis, dichotomis. Neck meth.

*L. caule repente, ramis tetragonis. Fl. lapp. 417. t. 11. f. 6. **

L. sabinae facie. Dill. musc. t. 58. f. 2. habitus accedit, sed floribus differt.

Muscus clavatus, foliis cupressi. Rai. hist. 121.

Habitat in Alpibus, Lapponiæ, Helvetiæ, Germaniæ.

16. *LYCOPODIUM complanatum. L. foliis bifariis connatis: superficialibus solitariis, spicis geminis pedunculatis. carn. pal. Fl. dan. t. 78. Kniph. cent. 10. n. 57. Vithu.*

L. caule repente, ramis triquetro-planis. Fl. lapp. 416. Fl. succ.

L. spicis petiolatis, quaternis, caulibus complanatis, foliis adpressis. Hall. helu. n. 1723.

L. surculo erecto, continuato, compresso, terminalibus capituliferis, dichotomo-spiciformibus. Neck. meth.

L. sabinae facie. Dill. musc. 445. synonyma.

Muscus clavatus, foliis cupressi. Bauh. pin. 360.

Muscus terrestris ramosus pulcher. Bauh. hist. 3. p. 759.

Lycopodium digitatum, foliis arboris vitæ, spicis bigemellis teretibus. Dill. musc. 448. t. 59. f. 3.

Savina sylvestris. Træg. 554. Hall. R.

Chamæcyparissus. Tabern. 945. Hall. R.

Habitat in Europæ & Americæ septentrionalis sylvis acrostit.

Confer. Lycopodium altissimum, spicis bigemellis quadrangulis. Dill. musc. 449. t. 60. f. 4.

18. *LYCOPODIUM helveticum. L. foliis bifariis petulis: superficialibus distichis, spicis geminis pedunculatis. Jacq. austr. t. 196. Scop. carn. burg.*

L. surculo reptante ramoso: terminali erecto, dichotomo, capitulifero. Neck. meth.

Lycopodioides foliis ellipticis alterne majoribus. Hall. helu. 93.

Muscus denticulatus major. Bauh. pin. 360.

Muscus repens, foliis alternis subrotundis per siccitatem falcatis. Moris. hist. 3. p. 626. f. 15. t. 6. f. 34.

Habitat in Alpibus Helvetiæ, Carniolæ.

19. *LYCOPODIUM denticulatum. L. foliis bifariis: superficialibus imbricatis, surculis repentibus, floribus sparsis.*

606 CRYPTOGAMIA. MUSCÆ.

L. caule plano, foliis superne quaternis, inferne binis, spicis obscure petiolatis. *Hall. helv. n. 1718.*

Lycopodioides imbricatum repens. Dill. musc. 462. t. 66. f. 1. A.

Muscus denticulatus minor. Bauh. pin. 360.

Muscus terrestris lusitanicus. Clus. hist. 2. p. 249.

Habitat in Lusitania, Hispania, Iberia.

Varietatem L. Helvetici suspicatur Hallerus.

1304. SPHAGNUM.

Anthera operculata: ore laevi. Calyptra nulla.

1. SPHAGNUM palustre. *S.* ramis deflexis. *Fl. suec. Veis. crypt. Leers herb. pal. Fl. dan. t. 474. litha.*

S. cauliferum ramis teretibus pendulis. *Hall. helv. n. 1724.*

S. ramis deflexis; pedunculis terminalibus; capsulis subrotundis. *Scop. carn.*

Hypnum (cubile) ramis lateralibus deflexis, primordialibus subrotundis, terminalibus congestis. *Neck. meth.*

Sphagnum palustre molle deflexum, squamis cymbiformibus. Dill. musc. 240. t. 32. f. 1.

Muscus palustris in ericetis nascens. Pluk. phyt. 101. f. 1. Vaill. paris. 139. t. 23. f. 3.

Muscus palustris terrestri similis. Pluk. alm. 254. t. 101. f. 4. Hall. Neck. R.

2. *Sphagnum palustre molle deflexum, squamis capillaceis. Dill. musc. 243. t. 32. f. 2.*

Habitat in Europæ paludibus profunde sylvaticis.

Antheræ diffiliunt cum sonu & crepitu.

3. SPHAGNUM alpinum. *S.* subramosum erectum.

Hypnum furculo erecto subdichotomo, foliis lineari-lanceolatis, trifariis, capitulis ovatis. *Neck. meth.*

S. subulatum viridissimum, capsulis ovatis. *Dill. musc. 245. t. 32. f. 3.*

Habitat in Angliæ summis alpium paludibus.

An Brii species? observante D. Schreber.

4. SPHAGNUM arboreum. *S.* ramosum repens, antheris lateralibus secundis. *Huds. angl.*

S. heteromallum polycephalon. *Dill. musc. 248. t. 32. f. 6.*

Hypnum (polycephalon) furculo ramoso-sparso, capitulis lateralibus secundis, margine ciliatis. *Neck. meth. 190.*

Habitat in Europa super truncos arborum.

1305. PHASCUM.

Anthera operculata : ore ciliato *Calyptra* minuta.

1. PHASCUM *pedunculatum*. P. acaule, anthera pedunculata. *Huds. angl.*

Bryum foliis ovato-lanceolatis, capsula pyriformi, brevissime papillata. *Hall. helu. n. 1831.*

Bryum ampullaceum, foliis serpilli pellucidis, collo crassiore. *Dill. musc. 334. t. 44. f. 5.*

Habitat in Europæ ericetis humidis.

Anthera inferne nigricans.

2. PHASCUM *acaulon*. P. acaule anthera sessili, foliis ovatis acutis conniventibus. *Fl. suec. ged. Veis. crypt. herb. pal. Flor. dan. t. 249. f. 3. Neck. gallob. lith.*

Sphagnum acaulon, foliis ovato-lanceolatis, pilo aristatis. *Hall. helu. n. 1726.*

Phascum (*cuspidatum*) caulescens, foliis ovatis cuspidatis patulis : terminalibus erectis conniventibus. *Schreb. de phasc. p. 8. t. 1. f. 1. 2. Weber. spicil. p. 125.*

Bryum (*bulbiforme*) capitulis globosis nitidis, calyptra basi dilatata, primordialibus bulbosis. *Neck. meth.*

Muscus trichodes acaulos minor latifolius. *Vaill. par. 128. t. 27. f. 2.*

Sphagnum acaulon bulbiforme majus. *Dill. musc. 251. t. 32. f. 11.*

Muscus trichodes acaulos minor latifolius. *Pet. gaz. 126. t. 95. f. 14.*

β *Sphagnum* acaulon bulbiforme minus. *Dill. musc. 252. t. 32. f. 12.*

Phascum (*muticum*) acaule, foliis ovatis muticis concavis conniventibus. *Schreb. de phasc. p. 8. t. 1. f. 11. 12. Weber. spicil. p. 126.*

Habitat in Europæ agris, hortis & ad fossas.

Calyptra cum Operculo.

3. PHASCUM *subulatum*. P. acaule, anthera sessili, foliis subulato-setaceis patulis. *angl. gallob. t. 1. f. 1. calyptra. Veis. crypt. herb. pal. n. 975. c. icone. Oed. fl. dan. t. 249. f. 1. 2. Hall. helu. n. 1727. lithu.*

P. caulescens, foliis lanceolato-linearibus patulis. *Schreb. de phasco. p. 8. Weber. spicil. 127.*

608 CRYPTOGAMIA. MUSCI.

Bryum (ericetorum) capitulis sessilibus ovoideis, calyptra conica, primordialibus plumosis. *Neck. meth.* 231.

Sphagnum acaulon trichodes. *Dill. musc.* 251. t. 32. f. 10.

Muscus trichodes minor acaulos, capillaceis foliis. *Pea. musc.* 87. *Vaill. parif.* 128. t. 29. f. 4.

Habitat in Europa. *Schreber.*

5. *PHASCUM repens.* P. caulescens repens, antheris lateralibus sessilibus.

Sphagnum vermiculare repens, capsulis intra foliorum squamulorum alas minimis. *Dill. musc.* 550. t. 85. f. 16.

Habitat in arborum truncis.

Obs. phascum acaulon. observante D. Schmiedelio, & subulatum, observante D. Schrebero gerens Calyptram, adeoque possent commode Bryis associari.

1306. FONTINALIS.

Anthera operculata. Calyptra sessilis, perichætio inclusa.

1. *FONTINALIS antipyretica.* F. foliis complicato-carinatis trifariis acutis, antheris lateralibus. *Fl. suec. ged. gallob. Veis. crypt. pal. Kniph. cent.* 12. n. 46. *burg. lichu.*

Hypnum foliis complicatis, ovato-lanceolatis, capsulis in alis petiolatis ovatis. *Hall. helv.* n. 1794.

Hypnum furculis trifariis, capitulis subsessilibus axillaribus; primordialibus oblongis teretibus. *Neck. meth.* 191. n. 48. a.

Fontinalis capitulis lateralibus. *Roy. lugdb.* 506.

Fontinalis triangularis major complicata e foliorum alis capsulifera. *Dill. musc.* 254. t. 33. f. 1.

Muscus squamosus, foliis acutissimis, in aquis nascens. *Vaill. parif.* 140. t. 33. f. 5. *Mich. gen.* t. 59. f. 9.

Muscus aquaticus, terrestris vulgari similis, major. *Buxb. cent.* 3. p. 39. t. 69. f. 2.

Muscus apocarpos aquaticus, foliis triangularibus complicatis. *Rup. jen.* 84. *Hall. R.*

Muscus aquaticus folio expanso. *Bauh. pr.* 152. *Hall. R.*

Muscus aquaticus, viticulis longis minus ramosis lucidis. *Morif. hist.* 3. p. 626. f. 15. t. 6. f. 32.

Muscus.

Muscus aquaticus denticulatus. Læs. pruss. 173. t. 52.

Habitat in Europæ fluviiis.

2. *FONTINALIS minor. F. foliis ovatis concavis trifariis acutis, passim geminis, antheris terminalibus. norv. n. 969. t. 3. f. 2. herb. Web. spicil. 35.*

Hypnum foliis ovato lanceolatis carinatis, apicibus capsuliferis. Hall. helu. n. 1795.

Hypnum (minus) surculis trifariis, apice capsuliferis, operculo capitulorum cuspidato. Neck. meth. p. 192. n. 48. β. cum præcedente conjungit. R.

Fontinalis triangularis minor carinata, e cymis capsu lifera. Dill. musc. 257. t. 33. f. 2.

F. capitulis terminalibus. Roy. lugdb. 506.

Habitat in Europæ fluviiis.

Folia complicato-carinata, in ramis crassioribus gemina.

3. *FONTINALIS squamosa. F. foliis imbricatis subulato lanceolatis, antheris lateralibus.*

Hypnum foliis ovato-lanceolatis, pilo terminatis capsulis per alas sessilibus. Hall. helu. n. 1796.

Fontinalis squamosa tenuis sericea atrovirens. Dill. musc. 259. t. 33. f. 3.

Fontinalis minor lucens. Bauh. hist. 3. p. 778.

Habitat in Britannia, Gallia, Helvetia.

4. *FONTINALIS pennata. F. foliis bifariis patentibus, antheris lateralibus. pal.*

Hypnum caulibus distichis, ramosis, foliis ovatis, transversim undosis. Hall. helu. ed. 2. n. 1797. t. 45. f. 2.

Hypnum (pennatum) surculo pennato, decumbente, foliis undato-rugosis, acutis, recurviusculis, capitulis axillari- bus sessilibus subcylindræis. Neck. meth.

Sphagnum pennatum undulatum, vagina squamosa. Dill. musc. 250. t. 32. f. 9.

Sphagnum cauliferum & ramosum, foliis crispis, crebris per caulem capitulis. Hall. helu. 96. t. 3. f. 2.

Muscus terrestris major, ramulis compressis, foliis super- ficie crispis. Vaill. paris. 129. t. 27. f. 4.

Muscus arboreus apocarpos, foliis pinnatis. Buxb. ephem. Nat. Cur. vol. 1. obs. 8. ex fide Halleri. R.

Habitat in Helvetiæ, Germaniæ arboribus.

Iconem Vaillantii ad Hypnum crispum amandat Weberus l. c. R.

Confer. Sphagnum pennatum undulatum, vagina squamosa. Dill. musc. 250. t. 32. f. 8.

Linnæi Spec. plant. Europ. Pars. II. Q q

Sphagnum pennatum undulatum, vagina pilosa. *Dill. musc.* 249. t. 32. f. 7.

1307. BUXBAUMIA.

Anthera operculata, hinc membranacea. *Calyptra* caduca (Schemiedel;) inter operculum sacculus pollinis.

1. BUXBAUMIA *aphylla*. Buxbaumia. *Hall. Mu. n.* 1728. *herb. pal. Fl. dan. t.* 44.

B. (*caulescens*) bulbo villis recto. *Schmied. diff. propr.* p. 25. n. 1.

Hippopodium. *Fabric. butisb. p.* 31. *

Musculus capillaceus aphyllus, capitulo crasso bivalvi. *Buxb. cent.* 2. p. 8. t. 4. f. 2. *Dill. musc.* 477. t. 68. f. 5.

Habitat in Suecia, Norvegia, Rutheno, Dania, Germania, Italia.

Calyptra reperitur medio octobris, valde caduca. R.

Antheram polliniferam *Millerum* 1764. primum detexisse perperam dicunt.

1308. SPLACHNUM.

Anthera infidens. *Apophysi* maxime colorata. *Calyptra* caduca. *Stella* feminea in distincto individuo.

1. SPLACHNUM *rubrum*. S. umbraculo orbiculari hemisphærico. p. 272. t. 3. f. 2. *

S. surculo simplicissimo, appendice capitulorum orbiculato hemisphærico. *Neck. meth.* 125.

Musculus norvegicus, umbraculo ruberrimo insignitus. *Pet. musc.* 70. t. 1. f. 10. *Dill. musc.* 547. t. 83. f. 9.

Musculus coronatus norvegicus, pediculo longissimo, umbraculo amplo. *Moris. hist.* 3. p. 630. f. 15. t. 7. f. 10.

Habitat in Norvegia, Finlandia, Russia, Sibiria.

Affine proximum ultimi.

2. SPLACHNUM *luteum*. S. umbraculo orbiculari plano. *Amœn. acad.* 2. p. 277. t. 3. f. 1. *

Habitat in Sueciæ paludibus sylvaticis.

Affine proximum sequentis.

CRYPTOGAMIA MUSCI. 611

3. **SPLACHNUM ampullaceum.** S. umbraculo ampullaceo obconico. *Fl. suec. carn. Flor. dan. t. 192.*

Bryum foliis ovato-lanceolatis, capsula piriformi, terminata cylindro. *Hall. helu. n. 1830.*

Splachnum furculo cespitoso, foliis lanceolatis nitidis, appendice capitulorum turbinato. *Neck. meth. p. 125.*

S. foliis linear-lanceolatis, nervo carinatis, acuminatissimis. *Zaga. flor. island. R.*

Bryum ampullaceum, foliis thymi pellucidis, collo strictiore. *Dill. musc. 343. t. 44. f. 3.*

Muscus capillaceus minor, capitulis geminatis. *Vall. paris. 130. t. 26. f. 4.*

Muscus aureus capillaris minor, capitulis geminatis erectis mutuo incubitu adnatis. *Pluk. alm. 246. Moris. hist. 3. p. 629. f. 15. t. 16. f. 10.*

Habitat in Europæ paludibus cespitosis nec nisi stercore conspurcatis; anne naturæ lusus totum hoc genus? an hoc Mnii fontani mera varietas.

4. **SPLACHNUM vasculosum.** S. umbraculo ampullaceo subgloboso.

S. foliis ovatis, nervo carinatis, apice obrufusculo. *Zag. flor. island. R.*

Muscus capillaceus, capitulis pyriformibus tubulosis. *Buxb. cent. 2. p. 1. t. 1. f. 1.*

Habitat cum præcedenti in Sueciæ borealis paludibus cespitosis, locis stercoreatis.

Simile præcedenti, sed illi umbraculum magis turbinatum & flavum; huic fere globosum sanguineum. Affine minimum Mnio annotino.

1309. POLYTRICHUM.

Anthera operculata, apophysi minimæ insidens. *Cal. villosa. Stella feminea in distincto individuo.*

1. **POLYTRICHUM commune.** P. caule simplici anthera parallepipeda. *Fl. suec. paris. ged. carn. gallob. Veiss. crypt. herb. pal. Blackw. t. 375. burg. lithu. Ingd.*

P. capsulis parallepipedis, foliis serrulatis acutis. *Schreb. spicil. 73.*

R. furculo capitulifero simpliciusculo: stellifero annotino-articulato, capitulis adultis tetraëdri obliquis. *Neck. meth.*

Mnium calyptris villosis, foliis serratis, capsulis quadrangulis, disco insidentibus. Hall. helu. n. 1835.

612 CRYPTOGRAMIA. MUSCI.

Polytrichum quadrangulare vulgare, yuccæ foliis serratis.
Dill. musc. 420. t. 54. f. 1.

P. aureum majus. *Bauh. pin.* 356.

Muscus juniperifolius, capitulo quadrangulo. *Vaill. paris.* 131. t. 23. f. 8.

β. *Polytrichum quadrangulare*, juniperi foliis brevioribus & rigidioribus. *Dill. musc.* 424. t. 54. f. 2. *Hall. l. c. γ.* *Flor. dan.* t. 295.

P. aureum medium. *Bauh. pin.* 356.

Muscus coronatus humilis. *Morif. hist.* 3. p. 630. f. 15. t. 7. f. 6. 8.

Muscus erectus, juniperi folioglauco. *Vaill. paris.* 131. t. 23. f. 6.

γ. *Polytrichum quadrangulare minus*, juniperi foliis pilosis.
Dill. musc. 426. t. 54. f. 3.

P. (piliferum) capsula parallepipeda, foliis integerrimis piliferis, caule simplici. *Schreb. spicil.* p. 74.

P. (pilosum) furculo simplicissimo, foliis piliferis linearibus integerrimis, capitulis obliquis tetraëdriis. *Neck. meth.* 123.

Mnium calyptris villosis, foliis serratis, pilo aristatis, capsulis quadrangulis, insidentibus disco. *Hall. helv. n.* 1836. R.

Muscus medicus, foliis juniperinis glaucis, non serratis, in longum & canescentem pilum desinentibus, capitulo tetragono, calyptra tomentosa rubra. *Mich. gen.* 109. n. 2. R.

Muscus capillaceus stellatus prolifer. *Vaill. paris.* t. 23. f. 7. R.

Habitat in Europæ α uliginosis, β. pascuis, γ. sterilissimis. 24

2. **POLYTRICHUM alpinum.** *P.* caule ramosissimo, pedunculis terminalibus. *Gunn. norv. Flor. dan.* t. 296.

P. furculo ramosissimo, setis terminalibus, capitulo ovato obliquo. *Neck. meth.*

Bryum foliis serratis, capsulis ovatis, basi turbinatis. *Hall. helv. n.* 1800. t. 46. f. 6.

Polytrichum alpinum ramosum, capsulis e summitate ellipticis. *Dill. musc.* 427. t. 55. f. 4.

Habitat in Alpibus Helvetiæ, Snovvdon, Germania.

3. **POLYTRICHUM umigerum.** *P.* furculo ramosissimo, setis lateralibus, capitulo erecto acuminato. *Neck. meth.*

P. (dubium) caule ramoso; capsulis oblongis, pedunculis lateralibus; foliis cartilagineo serratis. *Scop. carn. secundum Weberum.* R.

CRYPTOGAMIA. MUSCI. 613

Bryum caule ramoso, ex alis florigerum, foliis serratis, capsulis cylindricis. *Hall. helu. n. 1801.*

Bryum caule ramosissimo, pedunculis axillaribus calyptrisque pilosis. *Neck. gallob. 462. n. 20.*

Polytrichum ramosum, setis ex alis urnigeris. *Dill. musc. 427. t. 55. f. 5.*

Muscus ramosus erectus, calyptra villosa. *Vaill. paris. 131. t. 28. f. 13.*

Habitat in Europa, Jamaica.

1310. MNIUM.

Anthera operculata. Calyptra laevis.

Fem. capitulum nudum, pulverulentum, remotum.

1. **MNIUM pellucidum.** M. caule simplici, foliis ovatis, *Fl. succ. gallob. carn. herb. pal. Fl. dan. t. 300. Schmidel. ic. 1. p. 13. t. 3. optime. lithu.*

M. caule simplicissimo, foliis ovato lanceolatis, seta fungifera foliosa. *Hall. helu. n. 1853. t. 46. f. 8. exclusis synonymis Michellii & Buxbaumii. R.*

M. capitulo sphærophylloque distincto: surculis simplicibus, primordialibus plumulosis. *Neck. meth. 233. R.*

Bryum diaphanum. *Weber. spicil. p. 121. R.*

Mnium serpilli foliis tenuibus pellucidis. *Dill. musc. 232. t. 31. f. 2.*

Bryum alternans calyptra variegata & scyphulis foliosis. *Hall. helu. ed. pr. 118. t. 4. f. 8.*

Muscus coronatus minimus, capillaceis foliis capitulis oblongis. *Vaill. paris. 130. t. 24. f. 7.*

Habitat in Europæ pascuis succulentis umbrosis.

2. **MNIUM androgynum.** M. caule ramoso, androgynum. *Fl. succ. nerv. Veiss. crypt. herb. pal. Fl. dan. t. 299. lithu.*

M. caule simplici, capitulis rosiferis nudis. *Hall. helu. n. 1854.*

M. (*angustifolium*) capitulo pulverifero ovato, glabro: foliato globoso, surculis simplicibus ramosisque. *Neck. meth.*

Mnium perangustis & brevibus foliis. *Dill. musc. t. 31. f. 1. cat. giff. app. t. 1. f. A. B. C*

614 CRYPTOGRAMIA. MUSCI.

Muscoides : muscus capillaris minimus, capitulo minimo pulverulento. *Vaill. parif. 1. 29. f. 6.*

Muscus trichodes parvus, capitulo glomerato L. botryoide. *Mich. gen. 108. t. 59. f. 8. H. K.*

Muscus coronatus humilis minimus, foliolis pediculis & capitulis minimis erectis. *Morif. hist. 3. p. 631. f. 15. t. 7. f. 20.*

Musci parvi elegans species in truncis arborum. *Buxb. ad. petrop. 3. p. 273. t. 14. f. 2. Hall. R.*

Habitat in Europæ sylvis.

3. **MNIUM ramosum.** M. caule ramoso erecto, pedunculis femineis axillaribus. *Huds. angl.*

M. fluitans, prolixum, petiolis alaribus, sphaerulis pulverulentis. *Hall. helv. n. 1856.*

M. (prolixum) stipite setaceo axillari globulifero, furculo ramoso imbricato, primordiis plumulosis. *Neck. meth.*

M. majus, minus ramosum, capitulis pulverulentis crebris. *Dill. musc. 235. t. 31. f. 4.*

Habitat in Europæ paludibus.

An varietas alius speciei.

A. Mnio palustri vere differre vix credit cl. *Weberus spicil. fl. gatt. p. 123. Cum M. androgyno conjungunt. Neckerus & Leers. R.*

4. **MNIUM fontanum.** M. caule simplici geniculis inflexo. *Fl. suec. Vais. crypt. herb. pal. cum. ic. Weber. spicil. sub Bryo. Flor. dan. 298. lithu. burg.*

M. caulis teretibus, erectis, ex vertice multifidis. *Hall. helv. n. 1851.*

Bryum capsulis subrotundis obliquis, caule simplici geniculis inflexo. *Huds. angl.*

Mnium capsulis subrotundis; pedunculis radicalibus. *Scop. carn.*

Bryum furculo simplici tereti, capitulis obliquis subrotundis. *Neck. meth.*

Bryum palustre, scapis teretibus stellatis, capsulis magnis subrotundis. *Dill. musc. 340. t. 44. f. 2.*

Muscus stellatus alpinus palustris major. *Mich. gen. 1. 59. f. 4. Hall. R.*

Muscus capillaceus tenuissimus, pediculo longissimo purpurescente, capitulo rotundiore. T. *Vaill. parif. 134. t. 24. f. 10.*

Muscus trichodes palustris erectus coronatus, capitulis sphaericis amplioribus. *Morif. hist. 3. p. 628. f. 15. t. 6. f. 8.*

CRYPTOGAMIA. MUSCI. 615

Muscus parvus stellaris. Pluck. alm. 255. t. 47. f. 6.

Habitat in Europa ad fontes paludosos frigidos.

5. *MNIUM palustre.* M. caule dichotomo, foliis subulatis.

Fl. suec. carn. Vais. crypt. gallob. pal.

M. caulibus rubiginosis bipartitis, foliis mollibus, lanceolatis capsulis cylindricis acuminatis Hall. helv. n. 1855.

*Bryum succulo erecto dichotomo, setis axillaribus capitulo-
liferis, capsulis obliquis.* Neck. meth.

M. majus, ramis longioribus bifurcatis. Dill. musc. 233. t. 31. f. 3.

Muscus palustris erectus flavescens, capillaceo folio. Vaill. parif. 135. t. 24. f. 1.

Muscus palustris, flagellis erectis luteolis raro divisis, capitulis oblongis adianthi. Moris. hist. 3. p. 629. f. 15. t. 6. f. 9.

Muscus stellaris ramosus palustris, pediculo aureo erecto, capitulo magno sphaerico. Ruxb. cent. 1. p. 45. t. 65. f. 1. 2. 3. 4.

Habitat in Europæ paludibus.

6. *MNIUM hygrometricum.* M. a caule, anthera nutante, calyptra reflexa tetragona. ged. gallob. Wais. cryptog. herb. pal. Flor. Dan. t. 648. f. 2. lithu. burg.

M. foliis ovato lanceolatis, conniventibus, capsulis ovatis reflexis. Hall. helv. n. 1852.

Bryum acaule, capitulis turbinatis, obtusis, siccitate tetragonis, seta arcuata, primordialibus aggregatis. Neck. meth.

Bryum capsulis pendulis, obovatis, surculis simplicibus, foliis piliferis, ovatis. Scop. carn.

Bryum capsula nutante turbinato-ovata, calyptra reflexa quadrangulati. Fl. suec. parif.

Bryum bulbiforme aureum, calyptra quadrangulati capsulis pyriformibus nutantibus. Dill. musc. 407. t. 52. f. 74.

Muscus foliis scutellatis, capitulo pyriformi nutante. Vaill. parif. 135. t. 26. f. 16.

Muscus coronatus minor, foliolis latiusculis ad caulem convolutis, capitulis cernuis & obtusis aureis. Moris. hist. 3. p. 631. f. 15. t. 7. f. 17.

Habitat in Europæ sylvis apricis sterilibus.

7. *MNIUM purpureum.* M. caule dichotomo: axillis pendunculiferis, anthera erecta, foliis carinatis. ged. gallob. herb. pal. lithu.

Bryum capsula erecta oblonga, capillum divisis

616 CRYPTOGAMIA. MUSCI.

pedunculiferis : summitatibus stellatis , foliis carinatis.
Fl. suec. paris.

Bryum foliis lanceolatis subulatis , capsulis erectis , operculis conicis. *Hall. helu. n. 1816.*

Bryum antheris subnutantibus axillaribus , caule dichotomo , foliis carinatis. *Veiss. crypt. Weber. spicil.*

Bryum capsulis inclinatis striatis , operculo conico , foliis lineari-lanceolatis carinatis , caule subdiviso. *Schreb. spicil.*

Bryum tenue stellatum , fetis purpureis. *Dill. musc. 386. t. 49. f. 51.*

Muscus trichodes parvus , foliis musci vulgaris , capitulis longis acutis. *Moris. hist. 3. p. 628. f. 15. t. 6. f. 4. 16.*

Habitat in Europæ pascuis,

8. *MNIUM setaceum.* M. antheris erectis , operculis filiformibus longitudine antherarum. *Fl. suec. cum pal. iconq.*

Hypnum cauliculis teretibus , foliis lanceolatis repandis , capsulis erectis , cylindricis , aristatis. *Hall. helu. n. 1788. Veis. crypt.*

Bryum seta capitulifera subterminali , tuberculo setarum foliolis tecto. *Neck. meth.*

Bryum stellare nitidum pallidum , capsulis tenuissimis. *Dill. musc. 381. t. 48. f. 44.*

Habitat in Europæ muris , aggeribus , sepibus.

Caules erecti , brevissimi. Pedunculi atropurpurei , flexosi. *Antheræ cylindricæ , lividæ.* Opercula & Calyptræ filiformes , longiores & angustiores , quam in reliquis.

9. *MNIUM cirrhatum.* M. foliis arefactione revolutis. *Fl. suec. pal. icon. Flor. dan. t. 538. f. 4.*

Hypnum foliis capillaceis congestis , crispescentibus , capsulis ovato-cylindricis , brevissime ciliatis. *Hall. helu. n. 1786. Veis. crypt.*

Bryum surculo ramoso , foliis linearibus siccitate contortis , capitulis erectis cuspidatis. *Neck. meth.*

Bryum cirrhatum & stellatum , tenuioribus foliis. *Dill. musc. 379. t. 48. f. 42.*

Muscus muralis minimus roseus stellaris. *Vaill. paris. 130. t. 24. f. 8.*

Habitat in Europæ sylvis & ad sepes.

10. *MNIUM annotinum.* M. foliis ovatis acuminatis pellucidis subradicalibus antheris nutantibus. *Fl. suec. gallob. norv. Veis. crypt. herb. pal.*

Bryum furculo capitulifero stelliferoque fragili erecto sterili reptante, primordialibus plumosis. *Neck. meth.*

Bryum annotinum lanceolatum pellucidum, capsulis oblongis pendulis. *Dill. musc.* 399. t. 50. f. 68.

Habitat in Europæ humidiusculis.

11. *MNIUM* *hornum*. *M.* antheris pendulis, pedunculo curvato, furculo simplici, foliolis margine scabris. *Fl. suec. gallob. Veis. crypt. pal. burg.*

M. foliis lanceolatis, imbricatis, capsulis pendulis, cylindricis, obtusis. *Hall. helu. n.* 1846.

Bryum furculo capitulifero ramosiusculo: stellifero simplici, primordialibus plumulosis. *Neck. meth. p.* 215.

Bryum stellare hornum sylvarum, capsulis magnis natantibus. *Dill. musc.* 402. t. 51. f. 71.

Muscus capillaceus major stellatus. *Vaill. paris.* t. 24. f. 4. 5. *Mich. gen.* 108. t. 59. f. 2.

Muscus polytrichoides humilior, foliis brevioribus raris pallide viridentibus & vix pellucidis. *Moris. hist.* 3. p. 630. f. 15. t. 6. f. 3. 4.

Habitat in Europæ sylvis aggeribus.

12. *MNIUM* *capillare*. *M.* antheris pendulis, foliis ovatis setiferis carinatis, pedunculis longissimis. *Fl. suec. Dalib. paris. sub. Bryo.*

Bryum capillare. *Spec. ed. pal.*

Hypnum foliis ovato lanceolatis, capsulis ovatis, pendulis operculis conicis. *Hall. helu. n.* 1791. *Veis. crypt.*

Bryum capsulis pendulis oblongis; furculis ramosis; ramis annotinis, pedunculiferis; foliis piliferis, carinatis, ovatis. *Scop. carn.*

Bryum furculo stellifero acaule: capitulifero caulescente: foliis oblongis nitidis mucronatis. *Neck. meth.*

Bryum foliis latiusculis congestis, capsulis longis nutantibus. *Dill. musc.* 398. t. 50. f. 67.

Mnium capillare. *Fl. suec.*

Muscus capillaceus major, capitulis crassioribus cylindraceis nutantibus. *Vaill. paris.* 134. t. 24. f. 6.

Muscus capillaris, foliis latiusculis congestis, capitulis longis reflexis. *Moris. hist.* 3. p. 629. f. 15. t. 6. f. 19.

Muscus capillaris, folio longiore. *Magn. hort.* 138. t. 138.

Habitat in Europæ siccis, muris glareosis.

13. *MNIUM* *crudum*. *M.* antheris pendulis, calyptra recurvata foliis pellucidis. *herb. pal. cum icone. lithu.*

618 CRYPTOGRAMIA. MUSCI.

Bryum capsulis pendulis, calyptra recurvata, foliis subulatis acutis integerrimis, caule simplicissimo. *Fl. suec.*

Bryum capsulis oblongis pendulis, foliis acutis pellucidis ovatis: summis longioribus lanceolatis. *Schreb. spicil.*

Bryum furculo simplici, capitulis nutantibus, foliis intermediis latioribus, summis linearibus. *Neck. meth.*

Bryum pendulum horum molle, foliis & lanceolatis & gramineis. *Dill. musc. 401. t. 50. f. 70.*

Habitat in Europæ fylvis & declivitatibus.

14. *MNIUM* pyriforme. *M.* antheris pendulis turbinatis, stipite filiformi, floribus femineis setiferis. *Huds. angl. sub Bryo. Veis. crypt.*

Bryum antheris nutantibus turbinatis - foliis ensiformibus nitidis. *Fl. suec.*

Bryum foliis capillaceis, capitulis pyriformibus pendulis. *Hall. helv. n. 1813. t. 46. f. 7.*

Bryum (*aureum*) foliis capillaceis, capsulis pyriformibus pendulis, operculo brevissimo rotundato. *Schreb. spicil. Web. spicil.*

Bryum (*rupeum*) subcaule, capitulis turbinatis cernuis, seta flexuosa, operculis obtusis. *Neck. meth.*

Bryum trichodes aureum, capitulis pyriformibus nutantibus. *Dill. musc. 391. t. 50. f. 60.*

Habitat in Europæ rupibus.

15. *MNIUM* polytrichoides. *M.* calyptra villosa, *Fl. suec. pal. burg.*

M. acaulon, calyptra villosa, capsula cylindrica, foliis integerrimis. *Hall. helv. n. 1838.*

Polytrichum nanum. *Schreb. spicil. Neck. meth. Veis. crypt. var. β.*

Politrichum aloefolium. *Scop. carn.*

Polytrichum nanum, capsulis subrotundis galerinis aloës folio non serrato. *Dill. musc. 428. t. 55. f. 6. Delih. paris. 315.*

Muscus capillaceus minor, calyptra tomentosa. *Vaill. paris. 131. t. 26. f. 15.*

Muscus coronatus rigidus minor & humilior capitulis villosis brevioribus. *Morif. hist. 3. p. 630. f. 15. t. 7. f. 7.*

β. *Polytrichum* parvum, aloës folio serrato, capsulis oblongis. *Dill. musc. 429. t. 55. f. 7.*

Mnium calyptra villosa, acaulon, foliis serratis, capsulis cylindricis, erectis, *Hall. helv. n. 1837.*

Polytrichum (nanum) capsula cylindrica erecta, furculis simplicibus, brevissimis, foliis serrulatis. *Veis. crypt.*

Polytrichum (mnioides) acaule, foliis serrulatis, capitulis adultis erectis subcylindricis. *Neck. meth.*

Adiantum aureum medium, in ericetis proveniens. *Vaill. parif. 131. t. 29. f. 11.*

Muscus, adianthi aurei facie, humilis, capitulo rotundo turgido, calyptra villosa incana. *Buxb. cent. 1. p. 43. t. 63. f. 1.*

γ. *Polytrichum acaulon capillaceum*, capsulis cylindricis. *Dill. musc. 434. t. 55. f. 12.*

Habitas in Europæ ericetis subudis.

Calyptra villosa patula est ipsa Anthera major.

16. *MNIUM serpillifolium*. M. pedunculis aggregatis, foliis patentibus pellucidis. *gallob. crypt. herb. pal. Scop. carn. burg. lithu.*

α. *Mnium punctatum*.

M. pedunculis aggregatis, foliis obovatis integerrimis obtusis punctatis. *Darr. nass.*

M. foliis ovatis, integerrimis, capsulis pendulis, ovatis, aristatis. *Hall. helu. 1845.*

Bryum punctatum. *Schreb. spicil. Weber. spicil.*

Bryum (serpillifolium) furculo simpliciusculo, capitulis subaggregatis cernuis, foliis obovatis obtusis nitidis integerrimis. *Neck meth.*

Bryum capsulis nutantibus, foliis subovatis obtusis pellucidis punctatis, caulibus sterilibus procumbentibus, pedunculis aggregatis. *Fl. succ. Dalib. parif.*

Bryum caule procumbente, foliis alternis ovatis pellucidis. *Fl. lapp.*

Bryum pendulum serpilli folio rotundiore pellucido, capsulis ovatis. *Dill. muse. 416. t. 53. f. 81.*

Muscus folio lato subrotundo, capitulo singulari nutante, pediculo longo subrubenti infidente. *Vaill. parif. 136. t. 26. f. 5.*

Muscus polytrichoides palustris, major serpilli latioris folio pellucido. *Morif. hist. 3. p. 627. f. 51. t. 6. f. 39. 40.*

β. *Mnium (cuspidatum)* pedunculis aggregatis, folio alternis acutis serratis. *Scop. var. 1. Leers. Darr. nass.*

M. foliis ovato lanceolatis serratis confertis, capsulis nutantibus obtusa. *Hall. helu. n. 1841.*

Bryum cuspidatum. *Schreb. Spicil. Weber. spicil.*

620 GRYOTOGAMIA. MUSCI.

- Bryum (polymorphum)** furculo reptante pennato - capitulifero erecto, simplici, foliis mucronatis ferratis. *Neck. meth.*
- Bryum pendulum**, foliis variis pellucidis, capsulis ovatis *Dill. musc. 413. t. 53. f. 79.*
- Muscus palustris**, foliis subrotundis. *Vaill. parif. 26. f. 18.*
- Muscus polytrichoides pellucidus**, fere, foliis denticularis ad margines veluti crispis. *Pluk. alm. 257. t. 98. f. 8.*
- Mnium (proliferum)** pedunculis aggregatis, foliis rosaceo-congestis lanceolatis acutis. *Leers. l. c. γ. Pollich. l. c.*
- M.** foliis ovato-lanceolatis congestis, patulis, setis numerosis, capsulis ovatis, acuminatis, pendulis. *Hall. helv. n. 1839.*
- Bryum roseum.** *Schreb. spicil. p. 84. Veb. spicil. p. 118.*
- Mnium (roseum)** furculis simplicibus & proliferis, foliis rosaceo congestis, lanceolatis, acutis tenuissime argute crenulatis, pedunculis aggregatis. *Veis. crypt.*
- Bryum (roseum)** furculo capitulifero acaule : repetito stellato, foliis ovato-lanceolatis, patulis, mucronatis. *Neck. meth.*
- Bryum stellare roseum majus**, capsulis ovatis pendulis. *Dill. musc. 411. t. 52. f. 77.*
- Muscus stellaris roseus.** *Bauh. pin. 361. prodr. 151. Buxb. cent. 2. p. 2. t. 1. f. 3.*
- Muscus coronatus humilis**, foliolis latioribus stellatim nascentibus donatus. *Morif. hist. 3. p. 630. f. 15. t. 7. f. 9.*
- Mnium (undulatum)** pedunculis aggregatis, foliis oblongis undulatis. *Leers. l. c.*
- M.** foliis cordatis, ellipticis, setis aggregatis, capsulis ovatis obtusis pendulis. *Hall. helv. n. 1844.*
- M.** furculis erectis, ramis fasciculatis, foliolis ellipticis, undulatis, argute dentatis, pedunculis aggregatis. *Veis. crypt.*
- M.** capsulis ovatis pendulis, pedunculis aggregatis, furculis floriferis erectis, sterilibus humifusis, foliis pellucidis patulis oblongis undatis. *Scop. carn. ed. 1. p. 133. n. 7. var. δ.*
- Bryum (ligulatum)** capsulis nutantibus, pedunculis aggregatis, foliis oblongis undulatis serrulatis *Schreb. spicil.*
- Bryum (dendroides)** furculo capitulato stellatoque erecto ramoso, primordialibus plumulosis. *Neck. meth. 218. Weber. spicil.*

CRYPTOGAMIA MUSCI. 621.

Bryum caule adscendente, foliis ovato-oblongis paulis pellucidis. *Guett. stamp. Dalib. paris.*

Bryum dendriodes polycephalon, phyllidis folio undulato pellucido, capsulis ovatis pendulis. *Dill. musc. 410. t. 52. f. 76.*

Muscus erectus, folio linearis major. *Buxb. cent. 5. t. 71. f. 1. & minor. f. 2. Hall. R.*

Muscus roseus polycephalus, linearis foliis undulatis. *Vaill. paris. 135. t. 24. f. 3. Mich. gen. 108. t. 59. f. 5.*

Habitat in Europa fere ubique.

Varietas has pro distinctis speciebus sumunt varii: ego ob suas causas ita conjunxi, ut qui distinguendas velit, facile distinctas tradat.

17. *MNIUM triquetrum*. *M.* caulibus prælongis rubiginosis, foliis ovato-lanceolatis, capsulis ovatis pendulis *Hall. helu. n. 1849.*

Bryum (triquetrum) furculo ramoso, foliis ovato-lanceolatis, trifariam exstantibus, capitulis obtusis, ventricosis, cernuis. *Neck. meth.*

Bryum annotinum palustre, capsulis ventricosis pendulis. *Dill. musc. 404. t. 51. f. 72.*

Muscus denticulatus lucens fluviatilis maximus ad ramorum apices adianthi capitulis ornatus. Vaill. paris. 135. t. 24. f. 2. 2.

β. *Bryum lanceolatum bimum*, setis & capsulis oblongis pendulis. *Dill. musc. 405. t. 15. f. 73. Hudf.*

Bryum bimum. Schreb. spicil.

Mnium caulibus rubiginosis, foliis lanceolatis alternis, capsulis pendulis pyriformibus obtusis. *Hall. helu. n. 1847. R.*

γ. *Bryum palustre complicatum rubens*, capsulis turbinatis pendulis. *Dill. musc. 406. t. 51. f. 74 Hudson.*

Habitat in Europæ udis, surfosis.

Caules (in a) plurimi, in cespitem congesti, digitales & longiores, tomentosi. Folia trifariam posita, unde ramuli triquetri patuli, recentia pallide viridia, antiqua fusca, parva, distantia carinata, acuminata. Pedunculi solitarii longissimi, terminales purpurei. Antheræ erectæ, oblique f. arcuatæ, obtusissimæ, flavæ. Calyptra subulata. Stellæ femineæ ramos terminans in eadem planta cum capsula, sed distinctis ramis.

18. *MNIUM trichomanis*. *M.* foliis distichis integerrimis. *Fl. suec. herb. pal. gallob. lithu.*

623 CRYPTO GAMIA. MUSCI.

Jungermannia foliis integerrimis glabris ovatis diffusis, extremo caule setifero. *Hall. helu. n.* 1860.

Jungermannia scalaris. *Schmiedel. de junger. p.* 20. f. 17. *Schreb. spicil. p.* 103.

Jungermannia Trichomanis. *Scop. carn. ed. 2. n.* 1344. *Weber. spicil. p.* 148.

Mnium (*jungermannoides*) furculo pennato repente, apice globulifero ascendente, foliis integerrimis. *Neck. meth.*

Mnium trichomanis facie, foliolis integris. *Dill. musc.* 236. t. 31. f. 5.

Habitat in Sueciae, Angliae, Germaniae &c. arboribus.

In hoc Perichæzia frondalis abscondita reperitur: ergo Jungermannia ad sociandum Leers. l. c.

19. **MNIUM fissum.** *M.* foliis distichis bifidis. *herb.*

Jungermannia foliis bidentatis, fragiferis. *Hall. helu. n.* 1864.

Jungermannia fissa. *Scop. carn.*

Mnium (*trichomanis*) furculo reptante pennato, apice globulifero, foliis bidentatis. *Neck. meth.* 236. n. 5.

Mnium trichomanis facie, foliolis bifidis. *Dill. musc.* 237. t. 31. f. 6.

Jungermannia terrestris repens, foliis ex rotunditate acuminatis bifidis, apertura pene visibili. *Mik. gen.* 8. t. 5. f. 14.

Habitat in Europæ riguis.

20. **MNIUM Jungermannia.** *M.* foliis distichis, pinnis subtus auriculatis. *pal. lithu.*

M. caule procumbente simplici, foliis imbricatis integerrimis alternis antice appendiculatis. *Fl. suec.*

Jungermannia (*cochleariformis*) foliis amplexicaulibus, subrotundis *Hall. helu. n.* 1861. *Weis. crypt. p.* 123. *Weber. spicil.* 145.

Jungermannia purpurea. *Scop. carn.*

Lichenastrum alpinum purpureum, foliis auritis & cochleariformibus. *Dill. musc.* 479. t. 69. f. 1.

Jungermannia alpina palustris purpurea cambrica, foliis rotundioribus auritis tenuissime denticulatis. *Mik. gen.* 6. t. 5. f. 16.

Habitat in Europæ subhumidis.

1311. BRYUM.

Anthera operculata Calyptra lævis Filamentum e tuberculo terminali ortum.

* *Antheris sessilibus.*

1. BRYUM *apocarpum*. B. antheris sessilibus terminalibus, calyptra minima. *Veis. crypt. gallob. herb. pal. Fl. dan. 480. burg. lithu.*

Hypnum caulibus ramosis, foliis lanceolatis, hirsutis, operculis aristatis. *Hall. helu. n. 1793.*

Bryum antheris erectis, terminalibus, subsessilibus; surculis ramosis. *Scop. carn.*

Bryum surculo ramoso, foliis carinatis acutis, capitulis sessilibus ovatis terminalibus. *Neck. meth.*

Fontinalis apocarpa. *Weber. spicil.*

Sphagnum caulibus ramosis, foliis undique imbricatis capsulas obtegentibus. *Fl. suec.*

Sphagnum foliis pilo terminatis. *Ray. lugdb.*

Sphagnum subhirsutum obscure virens, capsulis rubellis *Dill. musc. 245. t. 32. f. 4.*

Muscus apocarpos hirsutus saxi adnascens, capitulis obscure rubris. *Vaill. paris. 129. t. 21. f. 15.*

β. Sphagnum nodosum hirsutum incanum. *Dill. musc. 246. t. 32. f. 5. Veis. l. c. p. 180. Leers. l. c. β.*

Hypnum caulibus teretibus ramosis, foliis ovato lanceolatis, pilo aristatis, operculo conico. *Hall. helu. n. 1792.*

Fontinalis albicans. *Weber. spicil. p. 38.*

Muscus squamosus saxatilis tortuosus ac nodosus. *Vaill. paris. 123. t. 27. f. 18.*

Habitat in Europæ saxi, arboribus.

2. BRYUM *striatum*. B. antheris subsessilibus sparsis, calyptris striatis sursumve pilosis. *ged. gallob. pal. Flor. dan. t. 537. f. 3. Schreb. spicil. 74. sub. Polytricho. lithu.*

Bryum. *Hall. helu. l. c. var. β.*

Polytrichum (*apocarpum*) caule ramoso: pedunculis brevissimis, sparsis; calyptris sursum pilosis. *Scop. carn.*

Polytrichum bryoides, α *striatum*. *Veis. crypt.*

Polytrichum capitulo laterali ovato subsessili, foliis integerrimis, ex appendiculatis, primordialibus gemmosis. *Neck. meth.*

624 CRYPTOGAMIA. MUSCI.

Polytrichum calyptra striata sursumve pilosa. Fl. suec. Dalib. paris.

Polytrichum bryi ruralis facie, capsulis sessilibus, majus. Dill. musc. 430. t. 55. f. 8.

Muscus apocarpus arboreus ramosus. Vaill. paris. 129. t. 25. f. 5. 6.

β. *Polytrichum bryi ruralis facie, capsulis, sessilibus minus. Dill. musc. 431. t. 55. f. 9. Veis. l. c. p. 176. Pollich. l. c. β. Neck. meth. β.*

Bryum setis brevissimis alaribus, calyptris villosis conicis: Hall. helu. n. 1799. α.

Muscus capillaceus minimus acaulos, calyptra striata. Vaill. paris. 30. t. 27. f. 10.

Muscus humilis tectorum subfuscus, capitulis brevibus pileolis erectis, pedunculis curtis. Moris. hist. 3. p. 629. f. 15. t. 6. f. 13.

γ *Polytrichum capsulis sessilibus, foliis brevibus rectis carinatis. Dill. musc. 432. t. 55. f. 10. Hall. l. c. γ.*

δ *Polytrichum capillaceum crispum, calyptris acutis pilosissimis. Dill. musc. 433. t. 55. f. 11. Neck. meth. γ. Pollich. l. c. γ. Fl. dan. t. 648. f. 1. Scop. carn. var. b.*

Polytrichum bryoides, capillaceum. Veis. crypt. p. 177.

Bryum setis brevissimis alaribus calyptris cylindricis villosissimis. Hall. helu. n. 1798.

Muscus capillaceus minimus, calyptra villosa. Vaill. paris. 130. t. 26. f. 9.

Habitat in Europæ α, arboribus, β saxis, γ. rupibus. δ. sylvis.

Proximum Polytrichis, sed absque apophysi sub anthera.

* * *Antheris pedunculatis erectis.*

3. *BRYUM pomiforme. B. antheris erectis sphaericis. Fl. suec. paris. ged. Veis. crypt. herb. pal. Weber. spicil. p. 119. lithu.*

B. caule erecto, foliis setaceis, capitulis globosis. Fl. lapp.

B. foliis subulatis, setis verticalibus, capsulis sphaericis. Hall. helu. n. 1803.

B. capsulis erectis sphaericis: pedunculis lateralibus; surculis ramosis; foliis linearibus. Scop. carn. exclusa varietate 1. secundum Weberum. R.

B. surculo erecto, capitulis lateralibus globosis. Neck. meth.

B. capillaceum, capsulis sphaericis. Dill. musc. 339. t. 44. f. 1.

Muscus capillaceus major, capitulis globosis. Buxb. cent. 5. app. n. 4. Hall. R.

Muscus

Muscus trichodes minimus sericeus capillaceus, capitulis sphaericis. *Morif. hist. 3. p. 628. f. 15. t. 6. f. 6. Vaill. parif. 129. t. 24. f. 9. 12.*

Habitat in Europæ scopulis humosis.

4. *BRYUM pyriforme*. B. antheris erectis obovatis ; calyptra subulata, surculis acaulibus foliis ovatis multatis. *gallob. Weis. crypt. pal. Fl. dan. t. 537. f. 1. lithu.*

B. caule brevissimo, foliis alato-lanceolatis, capsulis erectis, ovatis, operculo brevi. *Hall. helu. n. 1832.*

B. capsulis erectis, obovatis; surculis simplicibus; foliis pellucidis ovatis glabris. *Scop. carn.*

B. capitulis erectis turbinatis: operculo secedente truncatis. *Neck. meth.*

B. antheris erectis turbinato-ovatis: operculo obtusiusculo. *Fl. suec. Dalib. parif.*

B. serpillifolium pellucidum, capsulis pyriformibus. *Dill. musc. 345. t. 44. f. 6.*

Muscus capillaceus minimus, capitulis pyriformibus turgidis. *Vaill. parif. 129. t. 29. f. 3.*

Muscus minimus latifolius, capitulis erectis turgidis. *Buxb. cent. 1. p. 44. t. 64. f. 1.*

Muscus coronatus humilis stellaris, foliis latiusculis, capitulis pyriformibus erectis turgidiusculis. *Morif. hist. 3. p. 631. f. 15. t. 7. f. 16.*

Habitat in Europæ pratis ad aggeres.

3. *BRYUM extincitorium*. B. anthera erecta oblonga minor calyptris laxis æqualibus. *Fl. suec. parif. gallob. Weis. crypt. herb. pal. lithu.*

B. caule brevissimo, calyptra cylindrica, longe aristata. *Hall. helu. n. 1829.*

B. surculo simplici, foliis ovatis: supremis patulis, capitulis erectis, subcylindricis, calyptra basi dilatata. *Neck. meth.*

B. calyptra extincatorii forma; minus. *Dill. musc. 349. t. 45. f. 8.*

Muscus capillaceus minimus, calyptra longa conoide nitida. *T. Vaill. parif. 137. t. 26. f. 1.*

Muscus capillaris minimus cucullatus. *Magn. hort. 139. t. 82.*

Muscus coronatus humilis tenuifolius, pileolo magno acuto infra aperto extincitorium referente. *Morif. hist. 3. p. 630. f. 15. t. 7. f. 12.*

Bryum calyptra extincatorii figura, majus & ramosum, *Linnaei spec. Plant. Europ. Pars. II. R 2*

626 CRYPTOGAMIA. MUSCI.

Dill. musc. 350. t. 45. f. 9. Neck. l. c. var. β. Weis. l. c. β. Leers. l. c. β.

B. caule folioso, calyptra cylindrica ciliata, longe aristata, *Hall. helu. n. 1828.*

Habitat in Europæ arenosis.

6. **BRYUM turulatum.** B. antheris erectis subulatis, furculis acaulibus. *Fl. succ. ged. gallob. Weis. crypt. pal. Weber. spicil. p. 99. lithu.*

B. cauliculo brevissimo, foliis lanceolatis, capsulis & operculis longissimis. *Hall. helu. n. 1827.*

B. antheris erectis, subulatis; furculis simplicibus; foliis imberbibus. *Scop. carn.*

B. capitulis subulatis, foliis superioribus acutis, patentibus, primordialibus subradicalibus. *Neck. meth.*

B. capsulis longis subulatis. *Dill. musc. 350. t. 45. f. 10.*
Muscus capillaris, cauliculis longissimis & acutissimis. *Morif. hist. 3. p. 631. t. 7. f. 13. Hall. R.*

Muscus capillaris, corniculis longissimis incurvis. *R. Vaill. paris. 133. t. 25. f. 8.*

Habitat in Europæ aggeribus humidiusculis.

7. **BRYUM rurale.** Bryum antheris erectiusculis, foliis piliferis recurvis. *Fl. succ. paris. gallob. pal. lithu.*

B. caule erecto, foliis reflexis seta terminatis, capitulis falcatis. *Fl. lapp.*

Hypnum caulibus erectis foliis lanceolatis, pilo fluitante terminatis, capsulis erectis, longe rostratis. *Hall. helu. n. 1789. Weis. crypt.*

Bryum capsulis subnutantibus; pedunculis annotinis lateralibus; foliis supra medium dilatatis, piliferis, patulis. *Scop. carn. ed. 1. p. 145. n. 17. ed. 2. n. 1308.*

B. rurale unguiculatum hirsutum elatius & ramosius. *Dill. musc. 352. t. 45. f. 12.*

Muscus capillaceus, foliis Linariæ, capitulis corniculatis longissimis. *Buxb. cent. 5. t. 44. f. 1. Hall. R.*

Muscus capillaris rectorum, densis cæspitibus nascens, capitulis oblongis, foliis in pilum oblongum desinentibus. *Vaill. paris. 133. t. 25. f. 3.*

Muscus erectus hirsutus, capitulis longis acutis unguiculis avicularum similibus, vulgarissimus. *Morif. hist. 3. p. 628. f. 15. t. 6. f. 12.*

Habitat in Europæ muris, rusticorum tectis arborum truncis.

8. **BRYUM murale.** B. antheris erectis, foliis piliferis erectiusculis, furculis simplicibus cæspitosis. *Fl. succ. gallob. crypt. pal. herb. burg. lithu.*

B. caulibus brevissimis, foliis ovato-lanceolatis patulis, pilo terminatis, capsulis aristatis. *Hall. helu.* 1825.

B. capsulis erectis oblongis, surculis simplicibus, foliis rectiusculis; piliferis ovatis. *Scop. carn.*

B. capitulis oblongis erectis cuspidatis, foliis oblongis concavis piliferis. *Neck. meth.* 196.

B. regulare humile pilosum & incanum. *Dill. musc.* 355. t. 45. f. 1. *Muscus muralis*, omnium vulgarissimus, villosus. *Vaill. paris.* 135. t. 24. f. 15.

Muscus capillaris minor, capitulis erectis. *Mich. gen.* 108. t. 59. f. 7. *Hall. R.*

Muscus minimus; foliis acutis, in pilum canescentem desinentibus. *Buxb. cent.* 1. p. 45. t. 64. f. 4. *R.*

β. *Bryum* capitulo brevissimo, foliis ovatis lanceolatis patulis, capsulis aristatis. *Hall. helu. n.* 1826. *Leers. l. c. β.* *Neck. meth.* 197. β. *Pollich. l. c. var. β.*

B. humile, pilis carens, viride & pellucidum. *Dill. musc.* 356. t. 45. f. 15.

Muscus minimus latifolius, capitulis erectis longioribus. *Buxb. cent.* 1. p. 44. t. 25. f. 4. *Hall. R.*

Muscus muralis omnium vulgarissimus non villosus. *Vaill. paris.* 133. t. 24. f. 14.

Muscus capillaris minor, capitulis erectis, vulgarissimus, non villosus. *Vaill. l. c. t. 25. f. 4.*

Habitat in Europæ saxis, muris tectis, tegulis.

9. *BRYUM scoparium*. B. antheris erectiusculis, pedunculis aggregatis, foliis secundis recurvatis, caule declinato. *Fl. suec. paris. pal. ged. Kniph. cent.* 6. n. 77. *burg. lithu.*

B. caule declinato, foliis arrectis subulatis, capitulis erectiusculis. *Fl. lapp.*

Hypnum foliis falcatis, heteromallis, vaginis multifloris. *Hall. helu. n.* 1777. *Weis. crypt.*

Hypnum surculis erectiusculis; foliis secundis linearibus recurvis. *Scop. carn.*

Bryum surculo declinato, ramoso, foliis secundis, recurvatis, primordialibus plumulosis. *Neck. musc.*

B. reclinatum, foliis falcatis, scoparum effigie. *Dill. musc.* 357. t. 46. f. 16.

Muscus coronatus medius, foliis tenuissimis, pallidis, longiusculis, capitulis erectis, acutis. *Morif. hist.* 3. p. 630. f. 15. t. 7. f. 11. *Hall. R.*

Muscus coronatus humilis, corniculis longissimis acutissimis. *Morif. l. c. f. 13. Hall. R.*

628 CRYPTOGAMIA. MUSCI.

Muscus capillaceus major, pediculo & capitulo tenuioribus.
Vaill. parif. 132. t. 28. f. 12.

Muscus capillaceus, foliis pallidis, pediculis pluribus ex
communi theca egredientibus. *Buxb. cent. 2. p. 8. s. 4. f. 1.*

Habitat in Europæ sylvis arenosis, in truncis putridis. 24

10. BRYUM undulatum. B. antheris erectiusculis, pedun-
culis subsolitariis, foliis lanceolatis, carinatis, undulatis,
patentibus, serratis. *Fl. suec. parif. ged. gallob. Veis.
crypt. pal. Fl. dan. t. 477. burg. lithu.*

B. foliis lanceolatis, serratis, capsulis cylindricis inclinatis
aristatis. *Hall. helu. n. 1823.*

B. capsulis subnutantibus; furculis simplicibus; foliis lan-
ceolatis undatis serratis. *Scop. carn.*

B. (*phyllitifolium*) furculo simplici, foliis undato-serrulatis,
primordialibus plumulosis. *Neck. meth. 203.*

B. phyllidis folio rugoso acuto, capsulis incurvis. *Dill.
musc. 360. t. 46. f. 18.*

Muscus erectus, linariæ folio major. *Vaill. parif. 132. n. 1.*

Muscus capillaceus minor, capitulo longiore falcato. *Vaill.
l. c. t. 26. f. 17.*

Habitat in Europæ sylvis, aggeribus, pratis umbrosis.

11. BRYUM glaucum. B. antheris erectiusculis operculo
arcuato, foliis erectis imbricatis, furculis ramosis. *Fl.
suec. pal. burg. lithu.*

Hypnum caulibus erectis, teretibus, foliis imbricatis, fragi-
libus, capsulis inclinatis, aristatis. *Hall. helu. n. 1785. Veis. crypt.*

Bryum capsulis subnutantibus; operculo arcuato; furculis
ramosis; foliis fragilibus, erectis, glaucis. *Scop. carn.*

B. furculo ramoso, fragili, imbricato, foliis lanceolatis,
operculo capitulorum rostrato. *Neck. meth.*

B. albidum & glaucum fragile majus, foliis erectis, setis
brevibus. *Dill. musc. 362. t. 46. f. 20.*

Muscus trichodes montanus albidus fragilis. *Morif. hist. 3.
p. 630. f. 15. t. 6. f. 22.*

Muscus erectus capillaceus densissimus, glauco folio. *Vaill.
parif. 131. t. 26. f. 13.*

Muscus saxatilis ericoides. *Bauh. pin. 362. prodr. 151.*

Habitat in Europæ ericetis apricis.

Stirps facie *Sphagni palustris*, foliis albicantibus,
operculo ad modum subulato.

13. BRYUM pellucidum. B. antheris erectiusculis, foliis
acutis recurvis, caule hirsuto. *Fl. suec. herb. pal.*

B. foliis ovato-lanceolatis, aristatis, reflexis, capsulis ovatis
obliquis. *Hall. helu. n. 1824.*

B. furculo ferrugineo-tomentoso, foliis recurvatis nitidis carinatis, capitulis obliquis, cuspidatis. *Nech. meth.*

B. palustre pellucidum, capsulis & foliis brevibus recurvis. *Dill. musc.* 364. t. 46. f. 23.

Muscus polytrichodes elatior, foliis angustis pellucidis & fere membranaceis. *Pluk. alm* 247. t. 44. f. 7.

Muscus polytrichodes angustifolius pellucidus ramosus. *Pluk. alm.* 247.

B. Bryum erectus capitulis brevibus, foliis reflexis. *Dill. musc.* 345. t. 46. f. 24. *Pollich. l. c. β.*

Habitat in Europæ paludibus.

14. **BRYUM imberbe.** **B.** antheris erectis, ore dilatatis, foliis carinatis. *Huds. angl.*

B. foliis lanceolatis imbricatis, capsulis erectis cylindricis aristatis. *Hall. helv. n.* 1814. R.

B. tenue imberbe pallidum, foliis crebrioribus. *Dill. musc.* 382. t. 48. f. 46.

Habitat in Europæ arenosis graminosis, ad sepes, muros.

15. **BRYUM unguiculatum.** **B.** antheris erectis oblongis, pedunculis axillaribus, foliis erectis, acutis carinatis. *Huds. angl. Pollich. pal.*

B. capsulis urceolatis, operculo longo obliquo, foliis lanceolatis tortilibus, cauliculis erectis. *Schreb. spicil.* 78.

B. unguiculatum & barbatum, furculis in summitate crassioribus. *Dill. musc.* 383. t. 48. f. 47.

Muscus capillaris, capitulis erectis lageniformibus. *Buxb. cent.* 2. p. 9. t. 2. f. 9.

β. Bryum unguiculatum & barbatum tenuius & stellatum. *Dill. musc.* t. 48. f. 48. *Pollich. l. c. β. R.*

Habitat in Europæ muris arenosis.

Folia luteo-virentia. Pedunculi breves, axillares foliorum.

16. **BRYUM aciculare.** **B.** antheris erectis: operculo aciformi. foliis erectis subsecundis. *Scop. carn. sub Hypno.*

B. foliis imbricatis lanceolatis, setis alaribus, capsulis ovatis aristatis. *Hall. helv. n.* 1820.

Hypnum aciculare. *Weber. spicil.* p. 82.

B. montanum hemiheterophyllum, operculis acutis. *Dill. musc.* 366. t. 46. f. 25.

Habitat in Anglia, Helvetia, Germania, in montibus.

17. **BRYUM flexuosum.** **B.** antheris erectis, foliis setaceis, pedunculis flexuosis. *Fl. suec. Neck. gallob. pal.*

B. foliis capillaceis, capsulis cylindricis, operculis aristatis. *Hall. helv. n.* 1804.

612 CRYPTOGAMIA. MUSCI.

- P. ...*, filis sericeis, stipite flexuoso sub-
 ... Neck. meth. 205.
- P. ...* Dill. musc. 3-3. t. 47. f. 33.
 ...
12. *SPERMATOPHYTES*. B. antheris erectis, foliis ser-
 ... Fl. dan. gal. 1. p. 1. Fl. dan. 1. 479.
- P. ...*, capsulis ovatis, operculis
 ... Hall. bot. n. 1837. Crypt. f. 2. Hymn.
- P. ...* ; foliis sericeis
 ... Scop. carn.
- P. ...* capitulo annulo dorso capitulo
 ... Neck. meth. 205.
- P. ...* Dill. musc. 3-3. t. 47. f. 37.
- P. ...*, foliis ... spectantibus. Vill.
 ... f. 7.
- M. ...*, foliis ... capsulis minoribus
 ... f. 7.
- P. ...*, ...
13. *SPERMATOPHYTES*. B. antheris erectis, foliis sericeis
 ... Neck. meth. 205.
- P. ...*, foliis sericeis, ...
- H. ...*, ...
- P. ...*, operculo
 ... Neck. meth.
- I. ...*, ... Dill. musc. 3-3.
 ...
- M. ...* ...
- M. ...* ...
- H. ...* Helvetia, Suedia, ...
14. *SPERMATOPHYTES*. B. antheris erectis ...
 ... Fl. dan. gal. 1. p. 1. Fl. dan. 1. 479.
- P. ...* ovato lanceolatis, capsulis erectis, ovatis, ...
 ... Hall. bot. n. 1833.
- P. ...* erectis ovatis, ... ; ...
 ... Scop. carn.
- P. ...* erectis subrotundis : operculo ...
 ... Neck. meth. 195.

CRYPTOGAMIA. MUSCI. 631

B. exiguum, ereberrimis capsulis rufis. *Dill. musc.* 347.

t. 45. f. 7.

Muscus capillaceus, omnium minimus. *Vaill. paris.* 130.

t. 26. f. 2.

Muscus stellaris minimus, pediculo brevissimo, capitulis turgidis. *Buxb. cent.* 2. p. 4. t. 2. f. 2.

Muscus capillaris, capitulis rotundis, angustifolius. *Buxb. cent.* 2. p. 6. t. 2. f. 7.

Muscus coronatus minimus, foliis & capitulis oblongis in pediculis brevissimis. *Morif. hist.* 3. p. 631. f. 15. t. 7. f. 18.

Hyplopus Salomonis. *Hasselquist.*

Habitat in Europa ad agros, fossas, & sepes.

Capsulae amisso operculo apparent valde truncatae, obovatae & rusae, quibus manifeste differunt a sequente.

21. **BRYUM viriculum**. B. antheris erectis ovatis, foliis lanceolatis acuminatis imbricato-patulis. *Fl. suec. gallob. Weis. crypt. pal. lithu.*

B. capsulis capillar. capsulis ovatis aristatis. *Hall. helu. n.* 1809.

B. capsulis erectis ovatis; furculis simplicibus; foliis imbricatis, lanceolatis. *Scop. carn.*

B. furculo simpliciusculo, foliis linearibus exsiccatione crispis: capitulis erectis; operculo subrostrato tectis. *Neck. meth.*

B. capillaceum breve, pallide & laete virens, capsulis ovatis. *Dill. musc.* 380. t. 48. f. 43.

Muscus capillaceus omnium minimus, foliolis longioribus & angustioribus. *Vaill. paris.* 130. t. 29. f. 5.

Habitat in Europae agris, aggeribus.

Folia viridissima. Capsulae flavescentes, ciliatae.

Cum Bryo paludoso conjungit Weber. R.

22. **BRYUM paludosum**. B. acaulon, foliis setaceis, antheris obtusissimis patulis. *Fl. suec. gallob. Weis. crypt.*

B. foliis capillaribus, cauliculo sessili, capsulis ovatis aristatis. *Hall. helu. n.* 1810.

B. foliis setaceis, primordialibus solitariis. *Neck. meth.* 195.

B. trichodes acaulon palustre minimum, setis & capsulis brevissimis. *Dill. musc.* 387. t. 49. f. 53.

Habitat in Europae humidis.

Differunt a viridulo, capsulis fuscis & foliis siccitate minime crispis.

23. **BRYUM hypnoides**. B. antheris erectis, furculo erectiusculo, ramis lateralibus brevibus fertilibus. *Fl. suec. paris. gallob. herb. pal. burg.*

632 CRYPTOGAMIA. MUSCI.

Hypnum ramis alternis brevioribus, foliis pilosis, petiolis brevibus, flexuosis. *Hall. helu. n.* 1780. t. 46. f. 4.

Hypnum (*canescens* & *hirsutum*) caulibus suberectis, ramis alternis brevibus, foliis piliferis, pilis longis, vagis, setis brevibus, erectis. *Weis. crypt.*

Bryum furculo declinato, ramoso: ramis fertilibus, capitulis ovatis, erectis, operculo cuspidato. *Neck. meth.*

B. hypnoides polycephalon lanuginosum montanum. *Dill. musc.* 372. t. 47. f. 32.

Muscus alpinus ramosior erectus, flagellis brevioribus, lanuginosus. *Pluk. alm.* 255. t. 47. f. 5.

Muscus terrestris vulgari similis, lanuginosus. *Morif. hist.* 3. p. 625. f. 15. t. 5. f. 7.

B. Bryum hypnoides hirsutie virescens fasciculare alpinum, *Dill. musc.* 370. t. 47. f. 28.

γ. Bryum hypnoides, ericæ facie, capsulis barbatis, alpinum. *Dill. musc.* 371. t. 47. f. 31. *Leers. l. c. γ.*

Alia varietas est:

δ. Hypnum (canescens, pilosum) caulibus suberectis, ramis sparsis longioribus, foliis pilosis, setis longioribus erectis. *Weis. crypt. Leers. l. c. var. β. R.*

Bryum hypnoides, hirsutie canescens, vulgare. *Dill. musc.* 368. t. 47. f. 27. *Neck. gallob. var. α.*

Habitat in Europæ saxis, rupibus, sabulosis.

24. *BRYUM verticillatum.* *B. antheris erectis, pedunculis arefactione contortis, foliis piliferis, furculis fastigiatis.* *herb.*

B. pilosum verticillatum. *Dill. musc.* 374. t. 47. f. 35.

Habitat ad latera collium.

Cum sequenti conjungit Hallerus.

25. *BRYUM æstivum.* *B. antheris erectis subrotundis axillaribus, foliis subulatis distantibus.* *Huds. angl. gallob.*

B. cauliculis subnudis, ramosis, foliis subulatis brevibus, capsulis ovatis. *Hall. helu. n.* 1811.

B. furculo ramosiusculo, capitulis erectis subrotundis, foliis subulatis remotiusculis. *Neck. meth.*

B. palustre æstivum, conservæ facie. *Dill. musc.* 375. t. 47. f. 36.

Habitat in Europæ paludibus.

26. *BRYUM Celsii.* *B. antheris erectiusculis, pedunculis longissimis, foliis setaceis, furculis acaulibus.* *Fl. suec. norv.*

B. trichodes acaulon minimum, setis & capsulis oblongis. *Dill. musc.* 388. t. 49. f. 54.

Habitat Upsaliæ.

27. **BRYUM trichodes.** B. antheris erectis usculis : ore ciliato absque annulo, pedunculo longissimo. *Fl. suec. pal.*

B. foliis capillaceis erectis, setis longissimis, capsulis sursum lateſcentibus, operculo brevissimo. *Hall. helv. n. 1812.*

B. capitulis operculo decidente ciliatis, exanulatis, primordialibus reconditis. *Neck. meth.*

B. trichodes aureum, capsulis incurvis obtusis in setis longis, *Dill. musc. 389. t. 49. f. 58.*

Habitat in Suecia, Germania, Helvetia.

28. **BRYUM squarrosum.** B. antheris obliquis, foliis quinqueſariis imbricatis recurvis, caule ferrugineo tomentoso. *Fl. suec.*

Muscus palustris, foliis reflexis cauliculos quinquangulos formantibus. *Buxb. cent. 4. p. 36. t. 65. f. 1.*

Habitat in Europæ borealis paludibus apricis. *Læff.*

Muscus distantissimus ab Hypno squarroso. Surculi plures, conferti, erectiusculi, undique fusco-pubescentes : Ramo una alterove brevi. Folia minima, sessilia, ovata, acuminata, subcarinata, læte viridia, imbricata revoluta, quinqueſariam exstantia. Pedunculus terminalis, purpureus, longus. Capsula oblonga, obliqua.

*** *Antheris nutantibus.*

29. **BRYUM argenteum.** B. antheris pendulis, surculis cylindricis imbricatis lævibus. *Fl. suec. paris. gallob. crypt. 199. herb. pal. burg. lithu.*

B. cauliculis teretibus, capsulis ovatis, acuminatis, pendulis. *Hall. helv. n. 1821.*

B. capsulis pendulis, pedunculis subradicalibus, surculis teretibus ramosis. *Scop. carn.*

B. surculo nitido arcte imbricato, foliis ovato-concavis, capitulis ovatis nutantibus, stipite subradicali. *Neck. meth.*

B. pendulum julaceum argenteum & sericeum. *Dill. musc. 392. t. 50. f. 62.*

Muscus squamosus argenteus, ericæ folio. *Vaill. paris. 134. t. 26. f. 3.*

Muscus minimus e viridi argenteus, capitulis oblongis cernuis. *Moris. hist. 3. p. 627. f. 15. t. 6. f. 17.*

β. Bryum pendulum, surculis teretibus viridibus. *Dill. musc. 394. t. 50. f. 63.*

B. argenteum var. β. viride. *Leers. l. c.*

634 CRYPTOGRAMMA MUSCI.

B. trichoides, capitulis reflexis viridissimum, holosericeum;
Dill. giff. app. 86. t. 2. Hall. R.

Habitat in Europæ scdis, muris, rupibus.

30. BRYUM pulvinatum. B. antheris subrotundis, pedunculis reflexis, foliis piliferis. Fl. succ. parif. gallob. Weis. crypt. 201. pal. lithu.

B. foliis setaceis, pedunculo reflexo, capitulis subrotundis.
Fl. lapp.

B. foliis lanceolatis, pilo terminatis, capsulis pendulis ovatis aristatis. Hall. helv. n. 1822.

B. capsulis pendulis ovatis, pedunculis reflexis, furcula ramosis, foliis piliferis. Scop. carn.

B. furculo caespitose aggregato, foliis piliferis, setaceis, capitulis ovatis inter folia sepulcis. Neck. meth.

B. orbiculare pulvinatum hirsutie canescens, capsulis immerfis.
Dill. musc. 395. t. 50. f. 65.

Musci trichoides hirsutie canescens, capsulis subrotundis reflexis, in perbrevibus pediculis. Moris. hist. 3. p. 629. t. 6. f. 21.

Musci capillaceus lanuginosus minimus. T. Vaill. parif. 133. t. 29. f. 2.

Musci, capitulo inflexo, pene aculos. Pat. parif. 127. t. 95. f. 15.

Habitat in Europæ muris, rupibus.

Varietatem capsulis erectis describit, Pollich. l. c. R.

31. BRYUM caespitium. B. antheris pendulis, foliis lanceolatis acuminato-sericeis, pedunculis longissimis. Fl. succ. parif. gallob. Weis. crypt. sub Hypno herb. sub Mnio pal. c. icon. lithu.

B. foliis subulatis, pedunculo longissimo, capitulo reflexo.
Fl. lapp.

Hypnum foliis ovato-lanceolatis aristatis patulis, capsulis ovatis obtusis pendulis. Hall. helv. n. 1790.

Bryum foliis lanceolatis acuminatis nudis, stipite subdicali, capitulis ovatis obtusis. Neck. meth.

Bryum pendulum ovatum caespitium & pilosum, sub bicolori. Dill. musc. 396. t. 50. f. 66.

Musci capillaceus minimus, capitulo nutante, pediculo purpureo. Vaill. parif. 134. t. 29. f. 7.

Musci trichodes, capitulo parvo reflexo, pediculo longe medietate rubro, summa luteo-viridi. Moris. hist. 3. p. 629. f. 15. t. 6. f. 15.

Habitat in Europæ muris, scdis, glareosis.

Dantur individua stellifera : ergo.

Mnium. *Leers. l. c.*

32. **BRYUM carneum.** B. antheris pendulis subglobosis, foliis acutis alternis. *Fl. suec. herb. pal.*

B. foliis lanceolatis rarioribus; capsulis ovatis pendulis. *Hall. helu. n. 1834.*

B. furculo annotino stipitifero, operculis capitulorum obtusis, capitulis subrotundis, carnis. *Neck. meth.*

B. lanceolatum pellucidum, capsulis subrotundis, pedunculis carneis. *Dill. musc. 400. t. 50. f. 69.*

Muscus stellaris humilis, capitulis pendulis. *Buxb. cent. 2. p. 5. t. 2. f. 6. nimis humilis.*

Habitat in Europæ udis.

33. **BRYUM simplex.** B. anthera nutante oblonga, foliis subulatis, furculo simplicissimo: medio pedunculifero. *angl. sub rubro. gallob. herb. pal.*

B. rubrum. *Mant. 2. p. 309.*

B. cauliculis cernuis, foliis capillaribus heteromallis, capsulis curvulis cylindricis, operculis conicis. *Hall. helu. n. 1808.*

B. furculis apici dorsoque capituliferis simplicissimis, capitulis ovatis, foliis imbricato-patulis. *Neck. meth.*

B. trichoides, capsulis rubris carnis. *Dill. musc. 390. t. 50. f. 59.*

Muscus coronatus minimus. *Morif. hist. 3. 631. f. 15. t. 7. f. 19. Hall. R.*

Habitat in Europæ pascuis.

34. **BRYUM alpinum.** B. anthera pendula oblonga, foliis ovatis acuminatis carinatis, furculis ramosis, axillis pedunculiferis. *Huds. angl. 414.*

B. hynoides pendulum, sericeum, coma insigni atro rubente. *Dill. musc. 394. t. 50. f. 64.*

Habitat in Europæ rupibus, tenui terra tedis.

1312. HYPNUM.

Anthera operculata. Calyptra lævis. Filamentum laterale ortum e Perichætio.

* *frondibus pinnatis.*

2. **HYPNUM taxifolium.** H. fronde simplicissima pinnata lanceolata basi pedunculifera. *Fl. suec. gallob. crypt. 216. val. fl. danic. 17. f. 2. lithu.*

636 CRYPTOGAMIA. MUSCI.

H. foliis pinnatis contiguis linearibus, pedunculis radicalibus, capitulo subrotundo erecto. *Forst. fugh.*

H. foliis planis pinnatis, parium duodecim, setis radicalibus. *Hall. helu. n. 1775.*

H. fronde pinnata, simplici; pedunculis subradicalibus. *Scop. carn.*

H. furculo simplicissimo basi stipitifero capitulis incurvis, apice subulatis, foliis imbricatis acutis. *Neck. meth. 151.*

H. taxiforme minus: basi capsulifera. *Dill. musc. 263. t. 34. f. 2.*

Muscus pennatus, capitulis adianti. *Vaill. paris. 136. t. 24. f. II.*

Habitat in Europæ umbrosis.

3. HYPNUM denticulatum. H. fronde pinnata simplici: pinnis duplicatis basi pedunculifera. *Fl. suec. ged. gallob. pal. burg.*

H. foliis laxis pinnatis ovato-lanceolatis aristatis retro versis. *Hall. helu. n. 1772.*

H. furculo simpliciusculo decumbente basi setifero, capitulis obliquis, foliis ordine duplici, acutis, recurvatis. *Neck. meth.*

H. denticulatum pennatum, pinnulis duplicatis recurvis. *Dill. musc. 266 t. 34. f. 5.*

Muscus pennatus denticulatus minor. *Vaill. paris. 137. t. 29. f. 8.*

Muscus pennatus, capitulis adianti. *Morif. hist. 3. p. 626. f. 15. t. 6. f. 36.*

Habitat in Europa super terram umbrosam.

4. HYPNUM bryoides. H. fronde simplicissima pinnata, lanceolata apice pedunculifera. *Fl. suec. ged. Veis. crypt. 217. pal. Fl. dan. t. 473. f. I. lithu.*

H. foliis planis pinnatis, parium septem, setis ex summitate. *Hall. helu. 1776.*

H. fronde pinnata simplici, pedunculo terminali. *Scop. carn.*

H. fronde simplicissima pinnata, tuberculis pedunculorum nudis. *Neck. gallob. 464.*

H. furculo simplicissimo decumbente, apice stipitifero, capitulis erectis, ovoideis, foliis imbricatis. *Neck. meth. 152.*

H. taxiforme exiguum versus summitatem capsuliferum. *Dill. musc. 262. t. 34. f. I.*

Muscus polytrichoides, exiguis capitulis in summis furculis & foliis subrotundis erectis. *Vaill. paris. 136. t. 24. f. 13. Morif. hist. 3. p. 629. f. 15. t. 6. f. 11.*

Muscus terrestris minimus, foliis pennatim dispositis. *Buxb. cent. I. p. 44. t. 64. f. 3.*

Habitat in Europæ umbrosis.

Muscus minimus, capituli ora magna ciliis rubris spectabilis.

6. *HYPNUM adiantoides*. H. fronde pinnata ramosa erecta medio pedunculifera. *gallob. herb. pal. Scop. carn.*

H. foliis planis, pinnarum paribus quadraginta. *Hall. helu. n. 1774.*

H. surculo ramoso erecto, medio stipitifero, capitulis obliquis subulatis, foliis imbricatis acutis. *Neck. meth.*

H. taxiforme palustre ramosum majus & erectum. *Dill. musc. 264. t. 34. f. 3.*

Muscus taxiformis ramosus. Vaill. paris. 136. t. 28. f. 4.

Muscus filicifolius f. pennatus minor, pinnulis plurimis ad mediam costam adnexis. *Buxb. cent. 2. p. 3. t. 1. f. 4.*

Habitat in Europæ paludosis. 2c

7. *HYPNUM complanatum*. H. fronde pinnata ramosa 1 foliolis imbricatis acutis complicatis compressis. *Fl. suec. Veis crypt. herb. pal. lithu.*

H. caule disticho, foliis ovatis planis, capsulis ovatis aristatis. *Hall. helu. n. 1771.*

H. fronde ramosa: foliis distichis imbricatis, acinaciformibus acuminatis. *Schreb. spicil. 87.*

H. pennatum compressum & splendens, capsulis ovatis. *Dill. musc. 268. t. 34. f. 7.*

Muscus trichomanoides filicifolius splendens. Vaill. paris. 139. t. 23. f. 4.

Muscus terrestris repens primæ speciei similis. Moris. hist. 3. p. 625. f. 15. t. 5. f. 5.

Habitat in Europa ad truncos arborum.

α. *Hypnum complanatum. Neck. meth. musc. p. 155. non huc pertinet, sed ad H. trichomanoiden weberi. R.*

9. *HYPNUM sylvaticum*. H. fronde pinnata ramosa procumbente, foliolis acutis, medio pedunculifera. *Huds. angl.*

H. denticulatum pinnatum, pinnulis simplicibus rectioribus. *Dill. musc. 267 t. 34. f. 6. Hall. helu. n. 1773.*

Habitat in Europæ sylvis ad arborum radices.

* * *Surculis vagis.*

10. *HYPNUM lucens*. H. surculis ramosis, frondibus subpinnatis: foliolis punctatis.

H. fronde compressa ramosa; ramis vagis, obtusis; foliis

638 CRYPTOGRAMMA MUSCI.

ovatis imbricatis reticulatis plantis. Scop. carn. 1. p. 151. n. 7. ed. 2. n. 1319.

H. Pennatum aquaticum lucens, longis ~~lucens~~ ~~foliis~~ ~~foliis~~ Dill. musc. 270. t. 34. f. 10.

Habitat in Europæ paludosis.

11. HYPNUM rugosum. H. surculis vagis erectiusculis: foliis secundis recurvatis basi rugosis. Pollich. pal.

H. caulibus teretibus, foliis cincinnatis heteromallis. 1735.

H. (licopodioides) ramis apice recurvis decumbentibus, foliis secundis, antheris rectiusculis. Neck. gallob. p. 479. n. 28.

H. α. (licopodioides) surculo adscendente subpinnato, ramis recurvis: foliis dilatatis, secundis rugosis. Neck. meth. p. 168. Ad. pal. 2. p. 453. t. 7, f. 2.

H. lutescens crispum, lycopodii facie. Dill. musc. 289. t. 37. f. 24.

Habitat in Helvetia, Germania, Hollandia.

Musci rarius ramosus. Folia ad alterum latus flexa recurvataque basi dilatata & valde rugosa, sed vix sine oculo armato manifesta. Rami inter folia rubri. Pedunculi straminei in sicco contorti. Capsulæ obovatæ, breves absque ciliis.

Neckeri aliorumque descriptiones ab hac recedunt, ut de uno eodemque Musco loquantur? Weberus & Neckerus cum H. scorpioide L. conjungunt hunc Muscum. Vid. Weber. spicil. p. 66. R.

12. HYPNUM undulatum. H. surculis ramosis, frondibus subpinnatis: foliolis undulatis complicatis. Weis. Crypt. 220. herb.

H. caule plano, foliis distichis; obiter rugosis, nitidis, operculo conico. Hall. helv. n. 1770.

H. Fronde subpinnata; foliis undatis, crispis; pedunculis subradicalibus, longissimis; capsulis oblongis. Scop. carn.

H. surculo vago basi medioque setifero, capitulis oblongis, foliis distichis, complicatis, lanceolatis, undatis. Neck. meth. 156.

H. pennatum undularum, lycopodii instar sparsum. Dill. musc. 271. t. 36. f. 11.

Musci Adianti capitulis, Lycopodii in modum per terram se spargens. Moris. hist. 3. p. 626. n. 33. f. 15. t. 6. f. 33. Hall. R.

Habitat in Europæ locis saxosis cavernosis.

13 HYPNUM *crispum*. H. furculis ramosis, frondibus subpinnatis: foliolis undulatis planis. *ged. crypt. 221. pal. lithu.*

H. caule plano, foliis distichis, ovatis, transversim undulatis, operculis aristatis. *Hall. helu. ed. 2. n. 1769. t. 45. f. 5.*

H. repens *crispum*, ramulis compressis filicinorum more dispositis. *Hall. helu. ed. pr. 105. t. 3. f. 5.*

H. fronde subpinnata, ramosa; foliis undulatis, crispis; capsulis ovatis. *Scop. carn.*

H. furculo ramoso vago erectiusculo stipitifero, capitulis obliquis acuminatis, foliis undatis, obtusis, distichis. *Neck. meth. 153.*

H. pennatum undulatum crispum, setis & capsulis brevibus. *Dill. elth. 237. t. 36. f. 12.*

Musculus terrestris major, ramulis compressis, foliis superficie crispis. *Morif. hist. 3. p. 625. f. 15. t. 5. f. 10.*

Habitat in Europæ locis saxosis.

Musculus pennatus major. *Bauh. prodr. 151. Hall. R.*

Habitat in Europæ locis saxosis.

12 HYPNUM *triquetrum*. H. ramis vagis recurvis, foliis ovatis recurvatis patulis. *Fl. suec. ged. gallob. weif. crypt. pal. burg.*

H. foliis hispidis, ovato lanceolatis, aristatis. *Hall. helu. n. 1729.*

H. furculis repentibus; ramis vagis suberectis: apice dilatatis, ramulis recurvis. *Scop. carn.*

H. furculo adscendente; ramis supremis recurviusculis setiferis: infimis reptantibus, foliis imbricato patulis. *Neck. meth.*

H. ramis inæqualibus, foliis laxè imbricatis triangularibus acutis. *Fl. lapp.*

H. vulgare triangulum maximum & pallidum. *Dill. musc. 293. t. 38. f. 28.*

Musculus terrestris, maximus, erectus, ramosus, latioribus foliis & pallidioribus. *Buxb. cent. 4. t. 63. Hall. R.*

Musculus ramosus major spermatophorus. *Vaill. paris. 137. n. 11. t. 28. f. 9.*

Habitat in Europæ pratis, sylvis, sepibus.

15 HYPNUM *rutabulum*. H. ramis vagis subrepentibus, foliis ovatis mucronatis imbricatis. *gallob. Weif. crypt. 224. herb. pal. burg.*

H. foliis ovato-lanceolatis, patulis, striatis, breviter aristatis, operculis conicis. *Hall. helu. n. 1730.*

640 CRYPTOGAMIA. MUSCI.

H. ramis vagis erectiusculis, foliis ovatis acutis mucronatis parulis, operculis conicis. *Schreb. spicil.*

H. surculis repentibus; ramis vagis, suberectis; foliis acutis, ovatis; capsulis obtusis. *Scop. carn.*

H. surculo repente, ramis erectiusculis, teretibus, setiferis; foliis concavis, capitulo incurvo obtusiusculo. *Neck. meth.*

H. dentatum vulgatissimum, operculis obtusis. *Dill. musc.* 295. t. 38. f. 29.

Muscus myosuroides, rutabili fructu. *Vaill. paris.* t. 27. f. 8. & t. 23. f. 2.

Muscus denticulatus minor fericeus nostras, capitulis adiantis. *Moris. hist.* 3. p. 626. f. 15. t. 6. f. 35.

Habitat in Europæ septis, sylvis, ad arborum radices.

* * * *Surculis pinnatis.*

16. HYPNUM *filicinum*. H. surculis pinnatis: ramulis distantibus foliolis imbricatis incurvis acutis secundis. *Fl. suec. gallob. herb. pal.*

H. (*filicinum*, a. *spicatum*) surculis pinnatis: pinnis foliosis, simplicibus, distantibus, versus apicem in spicam approximatis: foliis imbricatis, falcatis. *Weiss. crypt.*

H. caulibus pinnatis, pinnis distantibus alterna longioribus, ex apice confertis, foliis cinnatis. *Hall. helv.* n. 1766.

H. surculis & ramis pinnatis, seta brevioribus, ramulis remotis, apicibus foliisque recurvis. *Scop. carn.*

H. surculo juniore erecto, simpliciter pinnato: adulto subrepente capitulifero, capitulis oblongis obliquis. *Neck. musc.* 164.

H. repens *filicinum* crispum. *Dill. musc.* 282. t. 36. f. 19.

Muscus *filcinus* palustris. *Vaill. paris.* 138. t. 29. f. 9. cum synonymis & descriptione aliena. *Weiss.*

Muscus pinnatus minor, cauliculis ramosis, in summitate veluti spicatus. *Lxj.* 167. ic. 43. sec. *Hall. R.*

Habitat in Europæ humidiusculis.

17. HYPNUM *proliferum*. H. surculis proliferis, plano-pinnatis, pedunculis aggregatis. *Fl. suec. pal. burglishu.*

H. caule suberecto, flexuoso, ramis teretibus, pinnatis, pinnulis pinnatis, operculis aristatis. *Hall. helv.* n. 1765.

H. *parietinum*. *Weiss. crypt.* *Neck. gallob.*

H.

CRYPTOGAMIA: MUSCI. 641

H. (parietinum) furculo pinnato, ramis arcuato subulatis teretibus, capitulis aggregatis obliquis, operculo subulato. *Neck. meth.* 157.

H. filicinum. Tamarisci foliis majoribus splendidibus. *Dill. musc.* 274. t. 35. f. 13.

Muscus filicinus flavescens major, tamosus. *Vaill. paris.* 140. n. 3.

Muscus filicinus major sericeus. *Ejusd.* t. 29. f. 1. bene.

Muscus vulgaris pennatus major. C. B. *Ejusd.* t. 28. f. 1.

Muscus filicinus major. *Morif. hist.* 3. p. 625. n. 20. f. 15. t. 5. f. 20. *Buxb. cent.* 2. p. 2. t. 1. f. 2.

Habitat in Europæ pratis, sylvis ad terram.

18. HYPNUM delicatulum. H. furculis subproliferis plano-pinnatis cuspidatis, pedunculis aggregatis. *Fl. suec.*

H. filicinum, tamarisci foliis minimis non splendidibus, setis capsulis brevioribus. *Dill. musc.* 546. t. 83. f. 6.

Habitat in America, Europa septentrionali.

Præcedenti simillimum, sed longe tenerius & forte sola varietas; antheras non vidi.

Cum H. parietino conjungit Pollichius, & Neckerus. R.

19. HYPNUM parietinum. H. furculis plano-pinnatis continuatis, pedunculis aggregatis. *Fl. suec. pal. burg. lithu.*

H. proliferum. *Weis. crypt. I. ex gallob.*

H. ramis teretibus, pinnatis, pinnulis pinnatis, foliis adpressis. *Hall. helv.* n. 1764.

H. furculis rugosis, nudis, frondibus bipinnatis: ramulis decurrentibus. *Scop. carn. sub H. prolifero.*

H. (proliferum) furculis proliferis, plano-pinnatis, stipitibus setaceis confertis. *Neck. in Ad. palat. 2. gallob.*

H. (tamariscifolium) furculo repente, ramis planiusculis proliferis, capitulis confertis incurvis. *Neck. meth.*

H. filicinum; Tamarisci foliis minoribus non splendidibus. *Dill. musc.* 276. t. 35. f. 14.

Muscus filicinus minor repens. *Læs. pruss.* 167. ic. 44. *Hall. R.*

Muscus filicinus minor. *Vaill. paris.* 140. t. 23. f. 9. *Morif. hist.* 3. p. 625. f. 14. t. 23. f. 9.

Habitat in Europa ad arborum truncos imos, & in sylvis ad terram.

Linnaei spec. plant. Europ. Pars II.

S 1

642 CRYPTOGRAMIA. MUSCI.

20. HYPNUM *prælongum*. H. furculis subpinnatis decumbentibus : ramulis remotis, foliolis ovatis, antheris cernuis. *Fl. suec. gallob. Weis. crypt. pal. lithu.*

H. foliis distichis, lanceolatis, pilo terminatis, capsulis ovatis, inclinatis, aristatis. *Hall. helu. n. 1758.*

H. furculo reptante, ramis subæqualibus, setis axillaribus, capitulis subcernis, foliis serrulatis, patulis. *Neck. meth.*

H. repens filicinum, triangularibus parvis foliis, prælongum. *Dill. musc. 278. t. 35. f. 15.*

Muscus terrestris minor, foliis crispis. *Euxb. cens. 4. p. 37. t. 63. f. 3.*

Habitat in Europæ terris, truncis, lignis.

Hæc species diff. ilius determinatur, quam dignoscitur.

21. HYPNUM *crista castrensis*. H. furculis pinnatis : ramulis approximatis, apicibus recurvis. *Fl. suec. gallob. crypt. 233. pal.*

H. foliis pinnatis, pinnis simplicibus, æqualibus foliis cinctis. *Hall. helu. n. 1768.*

H. furculis & ramis pinnatis : ramulis approximatis apicibus foliisque recurvis. *Scop. carn.*

H. furculo pinnato, ramis supremis contiguis apice deflexis, capsulis subrotundis obliquis. *Neck. meth.*

H. ramulis plano pinnatis : pinnis contiguis apicibus deflexis. *Fl. lapp.*

H. filicinum, cristam castrensem repræsentans. *Dill. musc. 284. t. 36. f. 20.*

Muscus cristam castrensem repræsentans flavescens nemorosus ramosus cassubicus. *Morif. hist. 3. p. 625. f. 15. t. 5. f. 8. icon. abludit.*

Muscus terrestris repens subflavus, foliis crispis minoribus, ramulisque densius confertis. *Vaill. paris. 141. t. 27. f. 14.*

Muscus pennatus major, caulibus ramosis in summitate veluti pinnatus. *Læf. pruss. 167. t. 42. bene.*

Habitat in Europa, Pennsylvania, ad radices Abietis.

22. HYPNUM *abietinum*. H. furculis pinnatis teretiusculis remotis inæqualibus. *Fl. lapp. 408. Fl. suec. gallob. Weis. crypt. lithu.*

H. ramis duris, teretibus, foliis adpressis, ovato-lanceolatis, pilo aristatis. *Hall. helu. n. 1762.*

H. furculis & ramis pinnatis : ramulis distantibus, subulatis, rectis. *Scop. carn.*

CRYPTOGAMIA. MUSCI. 643

H. furculis primordialibus obtusis, ramis teretibus subulatis; foliis ovatis acutis rachi siccata adpressis. *Neck. meth.* 163.

H. lutescens, alis subulatis tenacibus. *Dill. musc.* 280. t. 35. f. 17.

Muscus pennatus minor, caulibus ramosis, in summitate veluti spicatus. *Læs. pruss.* 167. t. 43. *Vaill. parif.* 1. 29. f. 12.

Muscus terrestris, furculis filamentosis tenacibus abietinis semel tantum divisis. *Morif. hist.* 3. p. 625. f. 15. t. 3. f. 22.

Habitat in Europæ sylvis abietinis.

Pedunculi & media rachi furculi, solitarii, purpurei & recti, longitudine; Calyptra erecta, subulata, pallida; Anthera ruffa, recurvata magis quam in ullo alio, margine oris integro, intra quem Cilia brevia, patula.

23. HYPNUM plumosum. **H.** furculis pinnatis repentibus; ramis confertis, foliis imbricatis subulatis, atheris erectis; *Huds. angl. Weis. cryptog. Scholl. barb. herb.*

Hypnum repens filicinum plumosum. *Lill. musc.* 280. t. 35. f. 16.

Habitat in Europa ad radices arborum.

Cum **H.** sericeo jungit Neckerus, sed ab eo differt habitu vere pinnato, & foliis patentissimis. *Weis. l. c. R.*

*** *Foliis reflexis.*

24. HYPNUM cupressiforme. **H.** furculis subpinnatis; foliis secundis recurvis apice subulatis. *Fl. suec. gallob. Weis. crypt. ged. pal. Fl. dan.* 535. f. 2. *burg. lithu.*

H. caule compresso, foliis imbricatis, falcatis; pilo aristatis; *Hall. helv. n.* 1737.

H. furculis & ramis pinnatis, foliis secundis: apicibus recurvis. *Scop. carn.*

H. furculo pinnato, ramis apice deflexis; remotis, capitulis ovatis, erectis. *Neck. meth.*

H. crispum cupressiforme, foliis aduncis. *Dill. musc.* 287. t. 37. f. 23.

Muscus squamosus ramosus minor & crispus. *Vaill. parif.* 139. t. 27. f. 13.

Muscus terrestris aureus minor, flagellis clavique brevioribus. *Morif. hist.* 3. p. 624. f. 15. t. 5. f. 8.

Habitat in Europæ sylvis ad arborum radices.

25. HYPNUM aduncum. **H.** furculis erectiusculo subramosis, foliis secundis recurvis subulatis, ramis recurvatis; *gallob. Weis. crypt. pal. lithu.* 9 & 2

644 CRYPTOGRAMMA MUSC.

H. caule erecto, foliis filicis, pila ciliata, speculo
conito. Hall. helv. n. 1776.

H. fuscato erecto simplicifolium, rami recurvi, capite
obliquo apice obtuso. Neck. musc.

H. palustre erectum, summitas obtusa. Dillen. musc.
292. t. 27. f. 26.

Habiat in Europe alpibus.

26. HYPNUM compressum. H. fuscis pinnatis compressis,
foliis acuminatis recurvis, antheris erectis ovatis.
Hedf. angl.

H. ramis pinnatis, pinnis ramosis, foliis ciliatis. Hall.
helv. n. 1767.

H. filicinum var. β . ramoso pinnatum) fuscis pinnatis;
pinnis filicis, subramosis, laxis, multinatis, apice
crispis; foliis imbricatis, filicis. Weis. crypt.

H. filicinum fuscum molle & pelliculosum, mucronis
obtusum. Dill. musc. 286. t. 36. f. 22.

Habiat in Europe truncis arborum.

27. HYPNUM scorpioides. H. ramis vagis procumbentibus
recurvis, foliis fuscis acuminatis. FL. suec.

H. scorpioides majus. Weber. spicil.

H. scorpioides palustre magnum, lycopodi inder. Dillen.
Dill. musc. 290. t. 37. f. 25.

Habiat in Anglia, Saccia palustris profunda.

Musci procumbens, rami vagis fuscis, apicibus obtusis.

Cum H. ramoso L. conjungit Neckerus & Weberus. H.

28. HYPNUM viticulosum. H. fuscis repensibus, ramis
vagis terribus, foliis parvis acuminatis. Gal. gall.
Weis. crypt. pol.

H. foliis lanceolatis crispis, capsulis erectis, speculo ar-
nico. Hall. helv. n. 1739.

H. fuscato germinante crassiore, adulto graciliore, ap-
tante capitalifero, primordiis ramosis. Neck. musc.

H. sabulosum, viticulis gracilibus erectis, capsulis ter-
ribus. Dill. musc. 307. t. 39. f. 43.

Musci squamosus, viticulis longioribus glabris. Hall.
parif. 227. t. 23. f. 1.

Musci montanus, flagellis longis terribus capillis
et plurimum indivis. Morf. 3. p. 624. f. 15. t.
5. f. 7.

Musci montanus gracilis,
glabris. Pluk. aln. 255.

Muscus densis foliis juniperinis in cespitem aggestis. Læf. pruss. 49. t. 49.

Habitat in Europæ montosis aridis ad arbores.

29. *HYPNUM squarrosum. H. ramis vagis, foliis lanceolatis, complicato carinatis quinquefariam recurvatis.*

*H. ramis vagis, foliis ovatis apice subulatis undique reflexis. Fl. suec. * Sp. pl. gallob. Weis. crypt. Scop. carn. pal. Fl. dan. t. 535. f. 1. lithu.*

H. foliis patulis subulatis, operculis conicis. Hall. helu. n. 1732.

H. surculo reptante setifero, ramis vagis, foliis recurvis subulatis striatis, capitulo ovali obliquo. Neck. meth.

H. repens, triangularibus reflexis foliis, majus. Dill. musc. 303. t. 39. f. 38.

Muscus terrestris erectior, foliis reflexis. Moris. hist. 3. p. 524. f. 15. t. 6. f. 2.

♂. *Hypnum repens, triangularibus reflexis foliis, minus. Dill. musc. 304. t. 39. f. 39.*

H. foliis cincinnatis, patulis, reflexis. Hall. helu. n. 1733. Flor. dan. t. 648. f. 3.

Muscus erectus, foliis reflexis. Vaill. paris. 139. t. 27. f. 5. Habitat in Europæ subhumidis.

30. *HYPNUM palustre. H. surculis repentibus, ramis confertis erectis, foliis ovatis secundis, antheris erectiusculis. Hudsf. angl. lithu.*

H. foliis ovato-lanceolatis falcatis, capsulis ovatis. Hall. helu. n. 1738.

H. (heterophyllum) surculo reptante simplici, ramis erectis setas subæquantibus: operculo mucronato. Neck. meth.

H. heterophyllum aquaticum polycephalum repens. Dill. musc. 293. t. 37. f. 27.

Habitat in Europæ udis.

31. *HYPNUM loreum. H. surculis reptantibus, ramis vagis erectis, foliis secundis, antheris subrotundis. Weis. crypt. herb.*

H. surculo repente, ramis erectis, foliis subsecundis, capitulo oblique subrotundo. Neck. meth. 170.

H. loreum montanum, capsulis subrotundis. Dill. musc. 304. t. 39. f. 40.

Muscus squamosus major, foliis angustioribus acutissimis. Vaill. paris. 138. t. 25. f. 2.

Muscus repens major, foliis & flagellis longis & tenuibus donatus. Moris. hist. 3. p. 626. f. 15. t. 5. f. 24.

Muscus montanus. Tabern. ic. 809.

448 CRYPTOGRAMIA MUSCI

Musco denticulato similis. *Bauh. pin.* 360.

Habitat in Europæ montosis.

Cum *H. squarroso* L. coniungit *Weber. spicil.* 69. R.

* * * * * *Surculis dendroidibus s. fasciculatis.*

32. *HYPNUM dendroides*. H. surculo erecto, ramis fasciculatis terminalibus simpliciusculis, antheris erectis. *Fl. suec. Vais. crypt. herb. pal. burg.*

H. caule erecto terne undique ramoso. *Fl. lapp.* 411.

H. caule nudo innomato, ramis teretibus confertis, capsulis acutis arisanis. *Hall. helv. n.* 1747.

H. surculis erectis, ramis terminalibus, teretibus, imbricatis, caule repente. *Scop. carn.*

H. surculo pristino reptante, ramis erectis congregatis simpliciusculis, operculo cuspidato. *Neck. meth.*

H. dendroides sericeum, setis & capsulis longioribus erectis. *Dill. musc.* 313. t. 40. f. 48.

Muscus squamosus ramulos erectus alopecuroides. *Tournef. inst.* 554. t. 326. f. 5. *Vaill. paris.* 137. t. 26. f. 6.

Muscus dendroides elatior, ramulis crebris minus furculosis, capitulis pedunculis brevibus insidentibus. *Morif. hist.* 3. p. 626. f. 15. t. 5. f. 31.

Muscus abietis facie. *Bauh. prodr.* 151. n. 3. *sec. Hall. R.*

Habitat in Europæ prarif. sylvisque subhumilis.

33. *HYPNUM alopecurum*. H. surculo erecto, ramis fasciculatis terminalibus subdivisis, antheris subnutantibus. *Fl. suec. gallob. Vais. crypt. herb. pal.*

H. caule nudo, confertim ramoso, ramis inclinatis, capsulis aristatis. *Hall. helv. n.* 1748.

H. surculo ramosissimo erecto, ramis spatio-pinnatis, primordialibus plumulosis. *Neck. meth.*

H. dendroides obicurus, setis & capsulis brevioribus subnutantibus. *Dill. musc.* 315. t. 41. f. 49.

Muscus dendroides sylvarum erectus, ramulis kali æmulis radice repente. *Morif. hist.* 3. p. 626. f. 15. t. 5. f. 30.

Muscus squamosus alopecuroides, flagellis recurvis. *Vaill. paris.* 137. t. 23. f. 5.

Habitat in Europæ sylvis humidiusculis.

* * * * * *Surculis teretiusculis.*

34. *HYPNUM curtispiculum*. H. surculis vagis teretibus, foliis ovatis acutis parulis, antheris pendulis. *Fl. suec. Vais. crypt. pal.*

H. ramis teretibus, foliis ovato-lanceolatis, pilis antheris, capsulis ovatis pendulis, operculis aristatis. *Hall. helv. n.* 1740.

H. furculo germinante crassiore, operculis capitulorum cuspidatis. Neck. meth.

H. dentatum curtispiculum, viticulis rigidis. Dill. musc. 333. t. 43. f. 69.

Habitat in Europa, America, ad radices arborum saxorum.

35. *HYPNUM purum. H. furculis pinnato-sparsis subulatis, foliis ovatis obtusis conniventibus. Fl. suec. ged. gallob. carn. pal. Flor. dan. t. 706. f. 2. lithu.*

H. ramis inæqualibus sparsis, foliis ovatis imbricatis, capitulis obliquis. Fl. lapp.

H. ramis teretibus, foliis ovato-lanceolatis, setis prælongis, capsulis inclinatis aristatis. Hall. helu. 1743.

H. furculo adscendente ramoso setifero: ramis incurvo-subulatis teretibus, foliis conniventibus obtusis mucronatis. Neck. meth.

H. cupressiforme vulgare, foliis obtusis. Dill. musc. 309. t. 40. f. 45.

Musculus squamosus cupressiformis. Vaill. paris. 138. t. 28. f. 3.

Habitat in Europæ pascuis, sylvis.

36. *HYPNUM filifolium. H. furculis vagis ramosissimis, ramis filiformibus, antheris obliquis. Hudf. angl. 421.*

H. (polyanthos) furculis ramosissimis: ramis rectis teretibus multifloris, capsulis erectis. Schreb. spicil. p. 97. R.

H. myosuroides, filiforme. Weber. spicil.

H. sericeum ramosius & tenuius, capsulis acuminatis. Dill. musc. 327. t. 42. f. 62.

Habitat in Europæ truncis arborum.

Est varietas H. myosuroidis secundum Weberum. Non huc videtur pertinere Hypnum foliis perangustis filo terminatis, capsulis inclinatis breviter aristatis. Hall. helu. n. 1755. monente Webero R.

37. *HYPNUM illecebrum. H. furculis ramisque vagis teretibus erectiusculis obtusis. Fl. suec. gallob. Fl. dan. t. 706. f. 1.*

Bryum caule erecto ramoso, foliis ovatis undique imbricatis. Fl. lapp.

H. ramis teretibus, foliis ovato-lanceolatis, concavis. Hall. helu. n. 1744.

H. furculo vago procumbente, ramis teretibus arcte imbricatis, foliis concavis obtusis, mucrone terminali. Neck. meth.

H. cupressiforme rotundius vel illecebræ æmulum. Dill. musc. 311. t. 40. f. 46.

648 CRYPTOGRAMMA MUSCI.

Muscus terrestris, furculis kali geniculati aut illecebræ
stimulis, foliis subrotundis squamatis incumben-
tibus. *Vaill. parif. 137. t. 25. f. 7.*

Habitat in Europæ & Americæ septentrionalis pascuis.

38. HYPNUM riparium. H. furculis teretibus ramosis,
foliolis acutis patulis distantibus. *Fl. succ. gallob. crypt. 250. pal. Flor. dan. 649. f. 1.*

H. ramis teretibus, foliis ovato-lanceolatis, pilo terminatis,
setis caulinis brevibus. *Hall. helu. n. 1746.*

H. furculo ramoso, tereti: immerso subpennato, foliis
patentibus lanceolaris alternis, capitulo acuminato.
Neck. meth.

H. aquaticum, flagellis & teretibus & pennatis. *Dill. musc. 308. t. 40. f. 44.*

H. ramosum fluitans pennatum. *Rai. angl. 3. p. 81.*

Habitat in Europa, ad fluviorum ripas.

39. HYPNUM cuspidatum. H. furculis vagis: apice foliis
convolutis acuminatis. *Fl. succ. ged. gallob. pal. lithu.*

H. foliis ovato-lanceolatis ramis teretibus, supremis
conicis, acuminatis. *Hall. helu. n. 1741.*

H. cuspidatum pungens. *Weis. crypt.*

H. furculo erectiusculo, ramis summis, pungenti-cuspidatis,
setis axillaribus longissimis. *Neck. meth. 163.*

H. palustre extremitatibus cuspidatis & pungentibus. *Dill. musc. 300. t. 39. f. 34.*

Muscus squamosus palustris, foliis flagellibusque rigidius-
culis, incurvis. *Vaill. parif. t. 23. f. 11. Hall.*

Muscus saxatilis repens, coma sparsa. *Buxb. cent. 2. p. 6. t. 3. f. 1. 2. 3. 4.*

Habitat in Europæ paludibus muscosis, quas replet.

***** Sursulis confertis.

40. HYPNUM sericeum. H. furculo repente, ramis con-
fertif erectis, foliis subulatis, antheris erectis. *Fl. succ. parif. ged. carn. gallob. Weis. crypt. pal. burg. lithu.*

H. ramis teretibus, foliis pilo præpilatis, capsulis cylin-
dricis erectis aristatis. *Hall. helu. n. 1750.*

H. repens, ramis annotinis ramosis erectis, capitulo erecto
acuminato, primordialibus acuminatis. *Neck. meth.*

H. vulgare sericeum recurvum, capsulis erectis cuspidatis.
Dill. musc. 323. t. 42. f. 59.

Muscus terrestris luteo-viridans sericeus repens. *Morif. hist. 3. p. 626. f. 15. t. 5. f. 25.*

Muscus arboreus splendens sericeus. *Vaill. parif. 132. t. 27. f. 3.*

Muscus terrestris vulgaris. Buxb. cent. 4. p. 34. t. 62. f. 1.

Muscus terrestris, vulgari similis, cauliculis cistis. Buxb. cent. 4. p. 37. t. 64. f. 1. Hall. R.

Habitat in Europa ad truncos, muros, inque campis.

41. *HYPNUM velutinum.* H. furculo repente, ramis confertis erectis, foliis subulatis, antheris subnatis. Fl. suec. ged. gallob. Vais. crypt. pal. Fl. dan. t. 475. burg. lithu.

H. ramis teretibus, foliis pilo terminatis, capsulis ovatis inclinatis mucronatis. Hall. helu. n. 1754.

H. reptans, ramis erectis teretibus seta duplo brevioribus, capitulis ovatis subnatis. Neck. meth.

H. velutinum, capsulis ovatis cernuis. Dill. musc. 326. t. 42. f. 61.

Muscus squamosus ramosus tenuior, capitulis incurvis. Vaill. paris. 138. t. 26. f. 9.

Muscus terrestris minor omnium vulgatissimus. Buxb. cent. 4. p. 35. t. 62. f. 23.

Habitat in Europa ad radices arborum umbrosas.

42. *HYPNUM serpens.* H. furculis repentibus, ramis filiformibus, foliis oblitteratis. Scop. carn. gallob. Weis. crypt. herb. excepta var. 2. pal. lithu.

H. caulibus prostratis, foliis minimis, pilo terminatis, capsulis cylindricis erectis mucronatis. Hall. helu. n. 1756.

H. furculo reptante, ramis æqualibus filiformibus, capitulo obliquo. Neck. meth.

H. furculis repentibus, ramis filiformibus & foliis oblitteratis, capsulis cylindraceis inclinatis, mucrone brevissimo recto. Schreb. spicil.

H. trichodes serpens, setis & capsulis longis erectis. Dill. musc. 329. t. 42. f. 64. A.

Muscus terrestris omnium minimus, capitulis majusculis oblongis erectis. Moris. hist. 3. p. 625. f. 15. t. 5. f. 14. 21. Vaill. paris. 138. t. 28. f. 2. 6. 7. 8.

Habitat in Europa, Virginia. ad caudices, ligna, saxa.

Synonymon Moris. & Vaill. huc non pertinere mones Weberus, l. c. R.

43. *HYPNUM sciuroides.* H. furculis erectis ramosis incurvatis. gallob. Weis. crypt. pal. Kniph. cent. 6. n. 46. herb. lithu.

H. caulibus teretibus ramosis, foliis appressis, pilo aristatis, operculis conicis. Hall. helu. n. 1749.

H. furculis erectis ramosis, ramis subradicalibus, foliis subovatis, capsulis erectis. Scop. carn.

650 CRYPTOGAMIA MUSCI.

H. furculo germinante crassiore; capitulifero tenuiore; ore capitulorum coarctato, primordialibus pilosis. Neck. meth.

H. arboreum sciuroides. Dill. msc. 319. t. 41. f. 54.

Muscus arboreus splendens myosuroides. Vaill. parif. t. 27. f. 12.

Muscus repens, serici modo lucens, viticulis longioribus erectis. Moris. hist. 3. p. 620. f. 15. t. 5. f. 27.

Habitas in Europæ truncis arborum.

Pedunculi maturi funis instar torti; feminae furculi distincti crescunt.

Hypnum palatinum Neck. meth. msc. hujus varietas est secundum de Leers. R.

44. HYPNUM gracile. H. furculis repentibus, ramis fasciculatis teretibus erectiusculis, antheris erectis ovatis. Hudf. angl.

H. ornithopodioides. Flor. dan. t. 649. f. 2. sec. Linneum.

H. gracile ornithopodioides. Dill. msc. 328. t. 41. f. 55. Habitat in Europæ segetis.

45. HYPNUM myosuroides. H. furculis ramosissimis: ramis subulatis utrinque attenuatis teretibus. gallob. Weis. crypt. pal. lukh.

H. ramis teretibus, foliis pilo præpilatis; capitulis ovatis mucronatis inclinatis. Hall. helu. n. 1752.

H. myosuroides tenuius, capitulis stantibus. Dill. msc. 317. t. 41. f. 51.

Muscus trichodes humilis ramosus, capitulis oblongis trimidiuſculis donatum. Moris. hist. 3. p. 628. f. 15. t. 6. f. 3.

Muscus squamosus minor myosuroides, capitulis incurvis. Vaill. parif. 137. t. 27. f. 6.

β. Hypnum myosuroides, crassius. Weber. spicil. p. 94. R.

H. (myosuroides) furculis ramosissimis, subulatis incurvis, utrinque attenuatis teretibus. Schreb. spicill. p. 97. n. 1069. Weber. R.

H. (myosuroides) furculo germinante crassiore; ramis teretibus subulatis; primordialibus sparsis. Neck. meth.

H. viviparum. Neck. gallob.

H. foliis ovato-lanceolatis, ramis teretibus, compressis. Hall. helu. n. 1745.

H. myosuroides crassius, capsulis erectis. Dill. msc. 317. t. 41. f. 50.

Habitas in Europa supra lapides arborum

CRYPTOGAMIA. ALGÆ. 651

46. **HYPNUM clavellatum.** H. repens, ramis erectis confertissimis, antheris incurvis; operculis inflexis. *Huds. angl. pal. c. icone.*

H. furculo repente, ramis contiguis teretibus erectis, duplo seta brevioribus, capitulo incurvo acuminato. *Neck. meth.*

H. clavellatum parvum repens, setis brevibus. *Dill. musc. 551. t. 35. f. 17.*

Habitat in Europæ lignis putridis.

ALGÆ.

1313. JUNGERMANNIA

Masc. Pedunculatus, nudus. *Anthera* 4. valvis.

Fem. Sessilis, nudus, feminibus subrotundis.

Frondeb. pinnatis secundis.

1. JUNGERMANNIA asplenoides. J. frondibus simpliciter pinnatis: foliolis ovatis subciliatis. *gallob. Weis. crypt. carn. pal. n. 1056. burg. lithu.*

J. foliis pinnatis, subrotundis, ferratis, ex apice florifera. *Hall. helu. n. 1868.*

J. furculo pinnato apice pedunculifero, foliis obtusis recurviusculis margine subciliatis. *Neck. meth.*

J. major, foliis brevioribus & obtusioribus non dentatis. *Mich. gen. 8. t. 5. f. 3. o. o.*

Lichenastrum asplenii facie, pinnis laxioribus. *Dill. musc. 482. t. 69. f. 5.*

Muscus bifolius procumbens, foliis subrotundis. *Læs. pruss. 167. t. 47.*

β. *Jungermannia* caule ramoso repente, foliis imbricatis ovato-crenulatis: alternis minoribus. *Roy. lugdb.*

J. major, foliis subrotundis tenuissime denticulatis. *Mich. gen. 7. t. 5. f. 1. 2.*

Lichenastrum asplenii facie, pinnis confertioribus. *Dill. musc. 483. t. 69. f. 6.*

γ. *Hepaticoides polytriche* facie. *Vaill. paris. 19. t. 19. f. 7.*

651 CRYPTOGAMIA: ALGÆ.

Milvus trichomanoides, foliis rotundioribus palmatis spon-
matim conjuncte sibi incumbentes. *Mich. gen. 8. t. 6. f. 42.*

Habitas in Europæ, Indiæ udis umbrosis.

2. *JUNGERMANNIA viticulosa*. J. frondibus pinnatis :
foliolis planis nudis linearibus. *prou. Weis. crypt. pal.*

J. foliis pinnatis subrotundis, ex caule florifera. *Hall. helv. n. 1869.*

J. furculo pennato basi medioque capitulifero, foliis ovatis
integerrimis. *Neck. meth.*

J. terrestris, viticulis longis, foliis perexiguis densissimis ex
rotunditate acuminatis. *Mich. gen. 8. t. 5. f. 4.*

Lichenastrum trichomanis facie, e basi & medio florent.
Dill. musc. 484. t. 69. f. 7.

Habitas in Europæ udis umbrosis sylvis.

3. *JUNGERMANNIA polyanthos*. J. frondibus simpliciter
pinnatis : foliolis integerrimis imbricatis convexis. *Weis. crypt. pal.*

J. foliis pinnatis ovatis, ex caule florifera. *Hall. helv. n. 1870.*

J. trichomanis. *Neck. gallob. p. 491.*

J. furculo ramoso pennato, ex lateribus globulifero, foliis
subrotundis integerrimis. *Neck. meth.*

J. palustris repens, foliis densissimis ex rotunditate acuminatis.
Mich. gen. 8. t. 5. f. 5.

Lichenastrum trichomanes facie, polyanthos, breve &
repens. *Dill. musc. 486. t. 70. f. 9.*

Habitas in Europæ palustribus.

4. *JUNGERMANNIA lanceolata*. J. frondibus simpliciter
pinnatis lanceolatis apice floriferis, foliolis integerrimis.
Weis. crypt. pal. lithu.

J. foliis pinnatis ovatis confertis, ex apice florifera. *Hall. helv. n. 1871.*

J. furculo pennato apice globulifero, foliis ovatis obtusis
integerrimis. *Neck. meth.*

J. palustris minima repens, foliis subrotundis densissimis
hæte virentibus. *Mich. gen. 8. t. 5. f. 67.*

Lichenastrum trichomanis facie minus, ab extremitate florens.
Dill. musc. 486. t. 70. f. 10.

Habitas in Europæ humidis umbrosis.

5. *JUNGERMANNIA bidentata*. J. frondibus simpliciter
pinnatis apice floriferis, foliolis bidentatis. *carn. gallob. Weis. crypt. pal. lithu.*

3. foliis bifidis, in ramo florifero majoribus. *Hall. helv. n. 1877. sec. Schreb. R.*

3. furculis procumbentibus ramosis, apicibus floriferis, foliis ovatis bicuspidatis, stipulis lanceolatis quadridentatis. *Schreb. Spicil.*

3. furculo pennato apice globulifero, foliis bidentatis. *Neck. meth. 130.*

3. major repens, foliis bifidis. *Mich. gen. 8. t. 5. f. 12. Seguiet. ver. suppl. 31.*

Lichenastrum pinnulis acutioribus & concavis bifidis, majus. *Dill. musc. 487. t. 70. f. 11.*

Hepaticoides, polytrichi facie, foliis bifidis, major. *Vaill. parif. 99. t. 19. f. 8.*

Musculus lichenoides, foliis pennatis bifidis, minor. *Morif. hist. 3. p. 627. J. 15. t. 6. f. 47.*

Habitas in Europæ ericetis umbrosis.

Cum Mnio fillo L. conjungis Weber. R.

6. JUNGERMANNIA *bicuspidata*. J. frondibus simpliciter pinnatis medio floriferis, foliolis bidentatis. *gallob. Weis. crypt. herb. pal.*

J. furculis procumbentibus, foliis distichis quadratis bidentatis, stipulis nullis. *Schreb. Spic.*

J. foliis bifidis, ex medio caule crebro florifera. *Hall. helv. n. 1878.*

J. furculo pennato lineari-ramoso, medio capitalifero, foliis bidentatis acutissimis. *Neck. meth.*

J. minima repens, foliis bifidis, vagina florum cylindracea. *Mich. gen. 9. t. 6. f. 17.*

Lichenastrum, pinnulis acutissime bifidis, minimum. *Dill. musc. 288. t. 70. f. 13.*

Habitas in Europæ umbrosis humidis.

7. JUNGERMANNIA *quinquedentata*. J. frondibus pinnatis ramosis apice floriferis, foliolis quinquedentatis. *Hudf. angl.*

J. furculo pennato ramoso, apice globulifero, foliis quinquedentatis. *Neck. meth. p. 132.*

Lichenastrum multifidum majus ab extremitate florens. *Dill. musc. 494. t. 71. f. 23.*

Habitas in Europa.

* * Frondibus pinnatis, pinnulis auriculatis superne tectis.

8. JUNGERMANNIA *undulata*. J. frondibus supra bipinnatis apice floriferis, foliolis subrotundis integerrimis undulatis. *Weis. crypt. pal. c. fig.*

651 CRYPTOGRAMIA: ALGÆ.

Muscus trichomanoides, foliis rotundioribus pellucidis squar-
matim conjuncte sibi incumbentibus. *Marif. hist. 3. p.*
627. f. 15. t. 6. f. 42.

Habitat in Europæ, Indiæ udis umbrosis.

2. *JUNGERMANNIA viticulosa*, J. frondibus pinnatis :
foliolis planis nudis linearibus. *prou. Weis. crypt. pal.*

J. foliis pinnatis subrotundis, ex caule florifera. *Hall. helv.*
n. 1869.

J. furculo pennato basi medioque capitulifero, foliis ovatis
integerrimis. *Neck. meth.*

J. terrestris, viticulis longis, foliis perexiguus densissimis ex
rotunditate acuminatis. *Mich. gen. 8. t. 5. f. 4.*

Lichenastrum trichomanis facie, e basi & medio florens.
Dill. musc. 484. t. 69. f. 7.

Habitat in Europæ udis umbrosis sylvis.

3. *JUNGERMANNIA polyanthos*. J. frondibus simpliciter
pinnatis : foliolis integerrimis imbricatis convexis. *Weis.*
crypt. pal.

J. foliis pinnatis ovatis, ex caule florifera. *Hall. helv. n.*
1870.

J. trichomanis. *Neck. gallob. p. 491.*

J. furculo ramoso pennato, ex lateribus globulifero, foliis
subrotundis integerrimis. *Neck. meth.*

J. palustris repens, foliis densissimis ex rotunditate acumina-
tis. *Mich. gen. 8. t. 5. f. 5.*

Lichenastrum trichomanes facie, polyanthemum breve &
repens. *Dill. musc. 486. t. 70. f. 9.*

Habitat in Europæ palustribus.

4. *JUNGERMANNIA lanceolata*. J. frondibus simpliciter
pinnatis lanceolatis apice floriferis, foliolis integerrimis.
Weis. crypt. pal. lithu.

J. foliis pinnatis ovatis confertis, ex apice florifera. *Hall.*
helv. n. 1871.

J. furculo pennato apice globulifero, foliis ovatis obtusis
integerrimis. *Neck. meth.*

J. palustris minima repens, foliis subrotundis densissimis
late virentibus. *Mich. gen. 8. t. 5. f. 67.*

Lichenastrum trichomanis facie minus, ab extremitate florens.
Dill. musc. 486. t. 70. f. 10.

Habitat in Europæ humidis umbrosis.

5. *JUNGERMANNIA bidentata*. J. frondibus simpliciter
pinnatis apice floriferis, foliolis bidentatis. *carn. gallob.*
Weis. crypt. pal. lithu.

J. furculo pennato ramofiusculo , apice capitulifero , foliis linearibus recurvis ; fupra auriculatis. *Neck. meth.*

Lichenaſtrum auriculatum : pinnulis anguſtis planis recurvis.
Dill. muſc. 492. t. 71. f. 20.

Hepaticoides albescens , foliis pinnatis. *Vaill. pariſ.* 100. t. 19. f. 5.

Habitat in Europæ umbroſis.

12. JUNGERMANNIA trilobata. **J.** frondibus ſubtus bipinnatis , foliolis quadratis obſolete trilobis. *Veſſ. crypt. herb.*

Jungermannia furculis planis pinnatis : foliolis obſolete trilobis auriculatis. *Fl. ſuec.*

J. caulibus convexis , foliis deſcendentibus imbricatis tridentatis. *Hall. helv. n.* 1866.

J. furculo pennato , ſimpliciuſculo reptante , foliis ſubquadratis , obtuſe crenatis , infra appendiculatis. *Neck. meth.*

Jungermannia foliis latusculis obtuſis undulatis & veluti anguloſis. *Mich. gen.* 7. t. 5. f. 10.

Lichenaſtrum multifidum, ab extremitate florens. *Dill. muſc.* 494. t. 7. f. 23. ſecundum *Hallerum* , &

Lichenaſtrum pinnulis obtuſe trifidis nervo geniculato. *Dill. muſc.* 493. t. 71. f. 22.

Habitat in Suecia , Anglia , Italia , Germania , Helvetia.

13. JUNGERMANNIA reptans. **J.** frondibus ſubtus bipinnatis apice reptantibus , foliolis quadridentatis. *Veſſ. crypt. lithu.*

J. minima , ramoſa , polyanthos , foliis imbricatis trifidis. *Hall. helv. n.* 1879.

J. furculis decompoſitis , medio floriferis , foliis quadratis antice inciſis , ſtipulis quadrifidis. *Schreb. ſpicil.*

J. furculo pennato , ramoſo , apice radicato , foliis ſubtus auriculatis ſubquadridentatis. *Neck. meth.*

Lichenaſtrum multifidum exiguum ad baſin florens per ſiccitatem imbricatum. *Dill. muſc.* 494. t. 71. f. 24.

Habitat in Europæ ericetis udis.

14. JUNGERMANNIA multiflora. **J.** fronde repente ramoſa : foliolis alternis geminis ſetaceis æqualibus. *pal. cum fig.*

J. fronde ſimpliciter pinnata , baſi florifera ; foliolis acutis. *Huff. angl.*

Lichenaſtrum multiflorum exile , foliis anguſtiſſimis. *Dill.* 481. t. 69. f. 4.

656 CRYPTOGAMIA. ALGÆ.

Habitat in Europæ sylvis.

Stirps filiformis, depresso planiuscula, repens. Folia alterna, ubique geminata, setacea, aqua hyalina, vix articulata, oculo armata videntia.

* * Frondibus imbricatis.

15. JUNGERMANNIA *complanata*. J. furculis repentibus: foliis inferne auriculatis duplicato imbricatis, ramis æqualibus. *Fl. succ. ged. norv. gallob. Weis. crypt. pal. Scop. carn. lithu.*

J. foliis rotundis, alterne imbricatis, caule plano, multifloro, setis brevissimis. *Hall. helu. n. 1860.*

J. furculo reptante, foliis subrotundis serie duplici ordinatis, subtus appendiculatis; vaginis ramorum plano-truncatis. *Neck. meth.*

J. foliis circinatis imbricatim dispositis ex viridi flavescens. *Mich. gen. 7. t. 5. f. 21.*

Lichenastrum imbricatum majus, squamis compressis & planis. *Dill. musc. 496. t. 72. f. 26.*

Hepaticoides foliis & furculis Thuyæ instar compressis major. *Vaill. paris. t. 19. f. 9. bene. Hall. R.*

Habitat in Europa ad arborum truncos.

Auriculas non invenit Scopoli, nec habet figuræ Dillenii. R.

16. JUNGERMANNIA *dilatata*. J. furculis repentibus, foliis inferne auriculatis duplicato-imbricatis, ramis apice latioribus. *Fl. succ. gallob. Weis. crypt. pal. burg.*

J. foliis imbricatis, alternis orbiculatis, setis brevissimis. *Hall. helu. n. 1874.*

J. furculo reptante imbricato, foliis serie duplici orbiculatis: ramorum apice latiusculo. *Neck. meth.*

Lichenastrum imbricatum minus, squamis convexo-concavis. *Dill. musc. 497. t. 72. f. 27.*

Hepaticoides, furculis & foliolis thuyæ instar compressis minor. *Vaill. paris. 99. t. 19. f. 10.*

Muscoides minimum, foliis alternis: superioribus circinnatis, inferioribus cucullatis, flore pediculo ferme carente. *Mich. gen. 10. t. 6. f. 6.*

Habitat in Europa, America, ad arborum truncos.

J. dilatata Scopoli carn. ed. 2. n. 1341. potius ad J. tamariscifoliam spectat, monente Webero. R.

17 JUNGERMANNIA *tamariscifolia*. J. foliis imbricatis serie duplici: superioribus subrotundis convexis obrutis quadruplo-majoribus. *Fl. succ. gallob. Weis. crypt. pal. lithu.*

γ. foliis imbricatis rotundis, superne alternis, convexis, inferne quadratis. *Hall. helv. n. 1873.*

J. (*tamariscifolia*) furculis decompositis apice floriferis, foliis subrotundis subtus auriculatis, stipulis ovatis emarginatis. *Schreb. spicil.*

J. furculo lineari reptante, foliis convexiusculis duplici ordine suborbiculatis: inferioribus quadruplo minoribus subbifidis. *Neek meth.*

Lichenastrum imbricatum, tamarisci narbonensis facie. *Dill. musc. 499. t. 72. f. 31.*

Hepaticoides, quæ Muscus trichomanoides terrestris minor floridus. *Vaill. paris. 100 t. 23. f. 10.*

Muscoides squamosum saxatile nigro - purpureum, furculis angustioribus, foliis circinatis minoribus. *Mich. gen. 10. t. 6. f. 5.*

Habitat in Europa ad truncos arborum; rupes.

J. rubiginosa. *Necker. in Act. pal. 2. p. 447. meth. p. 139: huc pertinere videtur. Weber l. c. R.*

18. JUNGERMANNIA *platyphyla*. J. furculis procumbentibus subtus imbricatis, foliis cordatis acutis. *gallob. Weis. crypt. pal.*

J. foliis imbricatis lanceolatis, superne planis, alternis, inferne concavis, quinquefariis. *Hall. helv. n. 1872.*

J. furculo decumbente imbricato, appendicibus subtus ordine duplici cordatis acutis. *Neck. meth. 142.*

Lichenastrum arboris vitæ facie, foliis minus rotundis. *Dill. musc. 501. t. 72. f. 32.*

Hepaticoides, furculis & foliolis thuyæ instar compressis, major. *Vaill. paris. t. 19. f. 9.*

Muscoides squamosum, atrovirens, foliis subrotundis. *Mich. gen. 9. t. 6. f. 3. 4.*

Muscus muralis floridus, foliis subrotundis creberrime imbricatis dispositis. *Moris. hist. 3. p. 627. f. 15. t. 6. f. 44.*

β. Muscoides squamosum saxatile maximum compressum, ex obscuro virescens, foliis subrotundis, nonnihil denticulatis. *Mich. gen. t. 6 f. 1. ex Hall. R.*

Lichenastrum Arboris vitæ facie, foliis rotundioribus. *Lill. musc. 502. t. 72. f. 33.*

Habitat in Europæ & Americæ septentrionalis sylvis.

19. JUNGERMANNIA *ciliaris*. J. furculis repentibus, foliolis duplicato imbricatis: inferne auriculatis ciliatis. *Fl. suec. pal. burg.*

Linnaei spec. Plant. Europ. Pars. II. Fc

658 CRYPTO GAMIA. ALGÆ.

J. caule pinnato, folus crispis, capillariter multifidis. *Hall. helv. n. 1881.*

J. furculo repente, foliis serie duplici ordinatis, subtus appendiculatis, ciliatis: vagina axillari, cylindracea, villosa. *Neck. meth.*

J. caule ramoso, furculis compressis, foliis imbricatis serie duplici ovato-acutis ciliatis. *Fl. lapp. 426.*

Lichenastrum filicinum pulchrum villosum. *Dill. musc. 303. t. 73. f. 35.*

Muscus palustris, absinthii folio. *Vaill. parif. 140. t. 26. f. 10.*

Habitat in Europa passim.

Folia superiora saepe bifida sunt.

20. JUNGERMANNIA varia. J. furculis erectiusculis bifariam imbricatis, foliolis bipartitis. *pal. lithu.*

J. furculis adscendentibus, foliis distichis bilobis subrotundis integerrimis, lobulis inaequalibus. *Schreb. spicil.*

J. repens, foliis cordatis carinatis. *Mich. gen. 8. t. 5. f. 9.*

Lichenastrum foliis variis. *Dill. musc. 303. t. 73. f. 36.*

Habitat in Europæ sylvis, ericetis.

**** *Surculis undique imbricatis, foliolis sparsis.*

22. JUNGERMANNIA rupestris. J. furculis teretibus; foliolis subulatis secundis.

J. furculosa erectiuscula, foliolis undique imbricatis hinc reflexis. *Il. jucc.*

J. furculo tereti, foliis subulatis secundis patulis. *Neck. meth.*

Lichenastrum alpinum nigricans, foliis capillaceis reflexis. *Dill. musc. 307. t. 73. f. 40.*

Habitat in Europæ fixatis rupibus humentibus.

23. JUNGERMANNIA trichophylla. J. furculis teretibus, foliolis capillaceis aequalibus. *Weis. crypt. herb.*

J. furculis erectiusculis, foliolis imbricatis acuminatis undique erectis. *Fl. jucc.*

J. minima, foliis capillariter multifidis. *Hall. helv. n. 1880.*

J. furculo tereti adscendente, apice globulifero, foliis subulato capillaceis sparsis. *Neck. meth. 144.*

Lichenastrum trichodes minimum in extremitate florae. *Dill. musc. 307. t. 73. f. 37.*

Habitat in Europa fixatis rupibus.

24. JUNGERMANNIA *alpina*. J. furculis teretibus, foliolis ovatis patentibus, calycibus imbricatis.

Lichenastrum alpinum atrorubens teres, calycibus squamosis. *Dill. musc.* 506. t. 73. f. 39.

Habitat in Alpibus Britanniae, inque rupibus Sueciae, Germaniae.

* * * * * *Acaules frondibus simplicibus.*

25. JUNGERMANNIA *epiphylla*. J. aculis, foliolo frondi innato. *Fl. suec. Weis. crypt. pal. Fl. dan.* t. 359. lithu.

J. fronde foliacea, obtuse lobata, ex medio florifera. *Hall. helu. n.* 1883.

J. acaulis; foliis laciniatis crispis; thecis dorsalibus, subovatis glabris. *Scop. carn.*

J. sublobata apice latior capitulo globoso: primordialibus frondosis obtusissimis marginalibus. *Neck. meth.*

Marsilea major atrovirens, floribus albicantibus e foliorum medio egredientibus. *Mich. gen.* 5. t. 74. f. 1.

Lichenastrum capitulis rotundis e foliorum medio enascentibus. *Dill. musc.* 508. t. 74. f. 41.

Lichen petræus, caliculo calceato. *Bauh. pin.* 362.

Lichen alter minor, folio calceato. *Col. ecphr.* 1. p. 331. t. 332.

Habitat in Europæ ripis elatioribus umbrosis udis.

Stamina circa germen primus observavi. *Fl. suec.* 1. n. 930.

26. JUNGERMANNIA *pinguis*. J. acaulis, fronde oblonga sinuata pingui. *Fl. suec. gallob. Weis. crypt.*

J. fronde foliosa lacera, ex oris florifera. *Hall. helu. n.* 1884.

J. sinuata obtusiuscula, ex lateribus capitalifera: primordialibus convolutis linearibus frondosis. *Neck. meth.*

Marsilea media pinguis pallide virens, floribus majoribus nigricantibus ad foliorum latera egredientibus. *Mich. gen.* 5. t. 4. f. 2.

Lichenastrum capitulis oblongis juxta foliorum divisuras enascentibus. *Dill. musc.* 509. t. 74. f. 42.

Hepatica saxatilis undulata, seminifera. *Vaill. parif. t.* 19. f. 6. *Hall. R.*

Lichen parvus erectus, foliis profunde laciniatis. *Pluck. alm.* 216. t. 42. f. 2.

Habitat in Europæ paludibus.

27. JUNGERMANNIA *multifida*. J. acaulis, fronde bipinnatifida. *Fl. suec. gallob. Weis. crypt. herb.*

450 CRYPTOGAMIA ALGAE.

1. General Information: Name, Address, City, State, Zip, Country, Telephone Number, E-mail Address, Date of Birth, Sex, Marital Status, Religion, Race, Ethnicity, Nationality, Language, Education, Occupation, Income, Social Security Number, Driver's License Number, Passport Number, Military Service, Other Identifiers, etc.

1. 1944-1945 1946-1947 1948-1949 1950-1951 1952-1953 1954-1955 1956-1957 1958-1959 1960-1961 1962-1963 1964-1965 1966-1967 1968-1969 1970-1971 1972-1973 1974-1975 1976-1977 1978-1979 1980-1981 1982-1983 1984-1985 1986-1987 1988-1989 1990-1991 1992-1993 1994-1995 1996-1997 1998-1999 2000-2001 2002-2003 2004-2005 2006-2007 2008-2009 2010-2011 2012-2013 2014-2015 2016-2017 2018-2019 2020-2021 2022-2023 2024-2025 2026-2027 2028-2029 2030-2031 2032-2033 2034-2035 2036-2037 2038-2039 2040-2041 2042-2043 2044-2045 2046-2047 2048-2049 2050-2051 2052-2053 2054-2055 2056-2057 2058-2059 2060-2061 2062-2063 2064-2065 2066-2067 2068-2069 2070-2071 2072-2073 2074-2075 2076-2077 2078-2079 2080-2081 2082-2083 2084-2085 2086-2087 2088-2089 2090-2091 2092-2093 2094-2095 2096-2097 2098-2099 2100-2101 2102-2103 2104-2105 2106-2107 2108-2109 2110-2111 2112-2113 2114-2115 2116-2117 2118-2119 2120-2121 2122-2123 2124-2125 2126-2127 2128-2129 2130-2131 2132-2133 2134-2135 2136-2137 2138-2139 2140-2141 2142-2143 2144-2145 2146-2147 2148-2149 2150-2151 2152-2153 2154-2155 2156-2157 2158-2159 2160-2161 2162-2163 2164-2165 2166-2167 2168-2169 2170-2171 2172-2173 2174-2175 2176-2177 2178-2179 2180-2181 2182-2183 2184-2185 2186-2187 2188-2189 2190-2191 2192-2193 2194-2195 2196-2197 2198-2199 2200-2201 2202-2203 2204-2205 2206-2207 2208-2209 2210-2211 2212-2213 2214-2215 2216-2217 2218-2219 2220-2221 2222-2223 2224-2225 2226-2227 2228-2229 2230-2231 2232-2233 2234-2235 2236-2237 2238-2239 2240-2241 2242-2243 2244-2245 2246-2247 2248-2249 2250-2251 2252-2253 2254-2255 2256-2257 2258-2259 2260-2261 2262-2263 2264-2265 2266-2267 2268-2269 2270-2271 2272-2273 2274-2275 2276-2277 2278-2279 2280-2281 2282-2283 2284-2285 2286-2287 2288-2289 2290-2291 2292-2293 2294-2295 2296-2297 2298-2299 2300-2301 2302-2303 2304-2305 2306-2307 2308-2309 2310-2311 2312-2313 2314-2315 2316-2317 2318-2319 2320-2321 2322-2323 2324-2325 2326-2327 2328-2329 2330-2331 2332-2333 2334-2335 2336-2337 2338-2339 2340-2341 2342-2343 2344-2345 2346-2347 2348-2349 2350-2351 2352-2353 2354-2355 2356-2357 2358-2359 2360-2361 2362-2363 2364-2365 2366-2367 2368-2369 2370-2371 2372-2373 2374-2375 2376-2377 2378-2379 2380-2381 2382-2383 2384-2385 2386-2387 2388-2389 2390-2391 2392-2393 2394-2395 2396-2397 2398-2399 2400-2401 2402-2403 2404-2405 2406-2407 2408-2409 2410-2411 2412-2413 2414-2415 2416-2417 2418-2419 2420-2421 2422-2423 2424-2425 2426-2427 2428-2429 2430-2431 2432-2433 2434-2435 2436-2437 2438-2439 2440-2441 2442-2443 2444-2445 2446-2447 2448-2449 2450-2451 2452-2453 2454-2455 2456-2457 2458-2459 2460-2461 2462-2463 2464-2465 2466-2467 2468-2469 2470-2471 2472-2473 2474-2475 2476-2477 2478-2479 2480-2481 2482-2483 2484-2485 2486-2487 2488-2489

Environ. Sci. Technol. 1991, 25, 1111-1115.

[illegible]

1. U.S. DEPARTMENT OF COMMERCE, Bureau of Census, P.O. 25,
WASH.

1. Statement of the problem: problem

W. J. ...
...
W. J. ...

1. Introduction - What is the purpose of the document?
 2. Background - What is the context of the document?
 3. Scope - What are the boundaries of the document?
 4. Methodology - How was the information gathered?
 5. Results - What were the findings?
 6. Conclusion - What are the main takeaways?
 7. References - What sources were used?
 8. Appendices - What additional information is provided?
 9. Index - How can the document be navigated?
 10. Summary - What is the overall message?

H. J. ...
 ...
 ...

Highway Engineering Department, Texas & Pacific

29. NON GERMANIA sicut? unde? unde
fuerit non minus, perinde quod. R. de.
et hoc videtur.

J.
... ..

J. Edgar Hoover, Director, Federal Bureau of Investigation
Washington, D. C.

Lichenzium *sligum*, capitulis a pileis loricatis e comis
 nervis distinctis. *Det. m. l. 1712. = L. 16.*

Edna is Extra mother, and is the only one

1314. T A R G I O N I A.

Cal. bivalvis, incl. sens. globum.

1. TARGIONIA *hypoleuca* T. *mauvea* & *viridis*.
Mauiensis T. *viridis*

Lotes permutados en lotes, fecha cont. E-2-A pta. 362
Diciembre 1972.

Deben also earlier hypothesis accepted. C... 475. 1. f. 33.
f. 33.

Leben interessanter Persönlichkeiten. Ein. u. d. f. d.
n. 61. f. 4

Habitus in Italia. Hirs. in Constantinopoli, Galibey, circa D

1315. MARCHANTIA.

Masc. *Cal.* peltatus, subtus tectus. *Cor.* 1. petalis.
Antheræ multifidæ.

Fem. *Cal.* sessilis, campanulatus, polyspermus.

1. MARCHANTIA *polymorpha*. *M.* calyce communi decemfido. *Fl. suec. ged. gallob. Veis. crypt. pal. burg. lithu.*

M. feminis analogo scobiformi : primordialibus marginalibus. *Neck. meth.*

M. linea frondem bipartiente, cyathis scobiferis ferratis, setis umbellatis & stellatis decemfidis. *Hall. helu. n. 1891.*

M. (*umbellata*) pileo crenato plano; capsulis glabris. *Scop. carn.*

Marchantia major, capitulo stellato : radiis teretibus. *Mich. gen. 2. t. 1. f. 1.*

Lichen fontanus major stellatus æque ac *umbellatus* & *cyathophorus*. *Dill. musc. 523. t. 76. f. 6.*

Lichen petræus latifolius, s. *Hepatica fontana*. *Bauh. pin. 362.*

Lichen. *Fuchsf. hist. 170. Gam. epit. 782.*

β. *Marchantia stellata*. *March. Añ. Paris. 1713. p. 307. t. 5.*

M. fronde æquabili, cyathis ferratis, bacillis umbellatis & stellatis octofidis. *Hall. helu. n. 1892.*

M. (*stellata*) pileo laciniato; laciniis teretibus, apice dilatatis. *Scop. carn.*

M. capitulo stellato : radiis teretibus. *Mich. gen. 2. t. 1. f. 2.*

Lichen domesticus minor stellatus æque ac *umbellatus* & *cyathophorus*. *Dill. musc. 527. t. 77. f. 7.*

Lichen petræus stellatus. *Bauh. pin. 362.*

Hepatica secunda. *Tabern. ic. 815.*

γ. *Marchantia* calyce communi octopartito : laciniis plano convexis. *Fl. lapp. 423. Roy lugdb. 507.*

Marchantia capitulo non dissecto. *Mich. gen. 1. t. 1. f. 5. Hall. n. 1892. var. β.*

Lichen petræus umbellatus. *Bauh. pin. 362.*

Hepatica tertia. *Tabern. ic. 815.*

642 CRYPTOGAMIA ALGAE

Hab. in Europa pars aquarum : litoris umbrosi.

Ex America septentrionali : Lich. linn. ex parte nomen in America, cum P. in arcto septentrionali exhibetur.

3. *MARCHANTIA crenata*. M. calyce communi quadriloculari subovato. *Fl. dan.* 1762.

M. foliis lanatis, capitulis quadrifidis. *Hall. helv.* n. 1889.

M. periclypeo masculino multilobato : femineo lunato. *Neck. germ.*

M. limbo calyce magis vix latiusculi : cavitationibus inter lobos. *Neck. meth. p.* 116.

M. foliis majoribus cruciformibus, femineis sessilibus lunatis. *Swed. musc.*

Lichen feminae lobulatus, florifer pileatus tandem cruciatus. *Dill. musc.* 511. t. 71. f. 4.

Lichen hepaticus, capitulis quadrifidis. *Burb. cent.* 1. t. 62. f. 2.

Lichen hepaticus. *Mich. gen.* 4. t. 4.

Hab. in Europa umbrosi.

5. *MARCHANTIA hemisphaerica*. M. calyce communi quinquelobato hemisphaerico, periclypeo nullo. *Fl. suec.* 1762. *Idem* 1762.

M. de crenata, capitulis conicis hirsutis. *Hall. helv.* n. 1890.

M. feminis analogo nullo : primordialibus marginalibus. *Neck. meth. rusc.*

Hemica media, capitulo hemisphaerico. *Mich. gen.* 3. t. 2. f. 2.

Lichen pileatus parvus, foliis crenatis. *Dill. musc.* 519. t. 71. f. 2.

Lichen hepaticus, capitulis stellatis crassis, pediculis brevibus infidentibus. *Burb. cent.* 2. p. 5. t. 5. f. 1.

Hab. in Europa paludosis.

Periclypea in hac specie nulla, nec calices seminiferi.

Capitulum hemisphaericum subius globulis 5. difflicentibus pollensque effundentibus.

6. *MARCHANTIA conica*. M. calice communi quinqueloculari subovato. *Strep. carn.* *Fl. dan.* t. 274. *Idem.*

M. verucis asperis, capitulis conicis. *Hall. helv.* n. 1889.

M. calice communi semiovario, periclypeo femineo orbiculari marginato. *Idem.*

CRYPTOGAMIA. ALGÆ. 663

- M. feminis** analogo subrotundo : primordialibus subart. cullatis. *Neck. meth.*
- Lichen vulgaris major pileatus & verrucosus.** *Dill. musc.* 516. t. 75. f. 1.
- Lichen petræus**, caliculo pileolum sustinente. *Bauh. pin.* 362. *Dill. giss. app.* 78. t. 1.
- Lichen plinii primus & pileatus.** *Col. exphr.* t. p. 330. t. 331.
- Hepatica reticulata & verrucosa.** *Vaill. par.* 97. t. 33. f. 8. *Hall. R.*
- Hepatica vulgaris major** f. officinarum Italiæ. *Mich. gen.* 3. t. 2. f. 1.
- Habitat in Europæ locis umbrosis, ripis elevatis.*
Feminei flores in fronde, tanquam verrucæ.
- 7. MARCHANTIA androgyna.** M. calyce communi integro hemisphærico.
- M. (angustifolia)** pileo hemisphærico, frondibus partialibus lineari dichotomis. *Neck. meth.*
- Lichen pileatus angustifolius dichotomus.** *Dill. musc.* 520. t. 75. f. 3.
- Habitat in Italia, Jamaica, inter saxorum crepidines; in introitu speluncæ Baumannianæ.* *Web.*
- Monoica f. Androgyna est hæc species.*

1316. BLASIA.

Masc. Cal. cylindricus granis repletus.

Fem. Cal. nudus; Fructu subrotundo, foliis immerso, polyspermo.

- 1. BLASIA pusilla.** *Blasia. Fl. suec. Schmied. diff. propr. herb. burg.*
- B.** fronde lobata, lobis subrotundis, nervis setiferis. *Hall. helu. n.* 1857.
- B. pusilla**, lichenis pixidati facie. *Mich. gen.* 14. t. 7. *Oed. dan.* t. 45.
- Mnium lichenis facie.** *Dill. musc.* 237. t. 31. f. 7.
- Habitat in Europa ad latera fossarum, solo ex arena sterili.*

1317. RICCIA.

Cal. o. nisi cavitatem vesicularem intra folii substantiam. *Cor.* o. *Anthera* cylindrica, sessilis, germiui composita. *Pist.* *Germen* turbinatum. *styl.* filiformis, antheram pertorans. *Peric.* sphericum, coronatum anthera emarcida. *Sem.* hemisphærica, pedicellata. Observante J. C. D. Schrebero.

1. *RICCIA crystallina*. R. frondibus superficie papillofis. *gall.b. Palat.* 2. t. 2. f. 9. *lithu.*

R. solus aspergine crystallina perfusus, margine incrassatis. *Fl. suec.*

R. fronde reticulata æquabili, lobis denticulatis, obtusis, simplicibus, & cordatis. *Hall helu. n.* 1896.

R. fronde superficie papillosa: primordialibus iecernentibus. *Neck meth.*

R. minima & minor pinguis aspergine crystallina perfusa. *Mich. gen.* 107. t. 57. f. 7. 3.

Hepatica palustris, lobis inflatis. *Vaill. paris.* 98. t. 19. f. 2.

Lichen palustris, rutæ folio. *Ceif. ups.* 28. *Dill. musc.* 355. t. 78. b. 12.

Habitat in Europæ locis subhumidis.

2. *RICCIA minima*. R. frondibus glabris bipartitis acutis. *Fl. suec. herb.*

R. minima nitida, segmentis angustioribus acutis. *Mich. gen.* 107. t. 57. f. 6.

Lichen omnium minimus, foliolis fissis super terram expansis. *Dill. musc.* 534. t. 78. f. 11.

Habitat in Europæ inundatis.

3. *RICCIA glauca*. R. frondibus glabris canaliculatis bilobis obtusis. *herb. pal. gall.b. burg. lithu.*

R. fronde sulco divisa, lobis biturcatis, lobulis lanceolatis obtusis. *Hall. helu.* 1897.

R. fronde bilobata obtusa superne sulcata. *Neck. meth. p.* 42.

R. minima pinguis, foliis latiusculis ample sulcatis e glauco virrescentibus. *Mich. gen.* 107. t. 57. f. 4.

Hepatica palustris bifurca, locis brevioribus carinatis. *Vaill. paris.* 98. t. 19. f. 1.

Lichen minimus, foliis venosis bifariam, vel trifariam se

CRYPTOGAMIA. ALGÆ. 665

dividendo progredientibus. *B. Dill. musc.* 533. t. 78. f. 10.

Lichen terrestris minor, foliis rutæ. *Buxb. cent.* 2. p. 10. t. 5 f. 5.

Habitat in Anglia, Italia, Gallia.

4. *RICCIA fluitans*. R. frondibus dichotomis lineari-filiformibus. *It. scan.* 241. *Fl. succ. ged. gallob. Flor. dan.* t. 275. *burg. lithu.*

R. fronde dichotoma, lineari ramosa, plana: primordialibus obtusis. *Neck. meth.*

Lichenastrum aquaticum fluitans tenuifolium furcatum. *Dill. musc.* 231. t. 74. f. 47.

Hepatica palustris dichotoma, segmentis angustioribus. *Vaill. parif.* 98. t. 10. f. 3.

Habitat in Europæ fossis, piscinis, ad ripas supra aquam extensa.

5. *RICCIA natans*. R. frondibus obcordatis ciliatis. *meth. pal. burg.*

Lichen parvus vernus cordiformis ima parte fimbriatus, lentis palustris modo aquis innatan. *Dill. meth.* 536. t. 78. f. 18. *Rai. angl.* 116.

Lens palustris, roris solis foliis cordatis. *Pet. musc.* 652.

Habitat in stagnis Angliæ, Germaniæ. *W. Schimielel.*

Margines ciliatæ revera non sunt, sed tantummodo radiculis marginalibus sæpius explanatis illam faciem debent. *Weber. l. c. R.*

1318. ANTHOCEROS.

Masc. Cal. sessilis, cylindricus, integer. *Anthera* subulata, longissima, 2. valvis.

Fem. Cal. 6. partitus. *Sem.* 3.

1. *ANTHOCEROS punctatus*, A. frondibus indivisis sinuatis punctatis. *gallob. he. b. pal. Fl. dan.* t. 396.

A. fronde crispa sinuato laciniata. *Schreb. spicil.*

A. fronde integra superne punctata. *Neck. meth.*

A. foliis minoribus magis laciniatis. *Dill. musc.* 476. t. 68. f. 1.

Anthoceros minor, foliis magis carinatis atque eleganter crenatis, subtus incurvatis. *Mich. gen.* 11. t. 7. f. 2.

666 CRYPTOGRAMIA. ALGÆ.

Habitat in Angliæ, Italiæ, Germaniæ uliginosis, umbrasis.

2. **ANTHOCEROS lævis.** A. frondibus indivisis sinuatis lævibus. *Leers. herb. Pollich. pal. Dærr. nass.*

A. frondibus indivisis, sinuatis, levibus. *Hall. helu. n. 1893.*

A. fronde sinuata integra obtusa. *Neck. meth.*

A. foliis majoribus minus laciniatis. *Dill. musc. 476. t. 68. f. 2.*

A. major. *Mich. gen. II. t. 7. f. 1.*

Lichen hepaticus, pediculis gramineis. *Buxb. cent. p. 40. t. 61. f. 1.*

Habitat in Europa & America boreali.

3. **ANTHOCEROS multifidus.** A. frondibus bipinnatifidis linearibus

A. fronde lineari triplicato pinnata. *Hall. helu. n. 1894.*

A. fronde lineari bipinnatifida: primordialibus linearibus simplicissimis. *Neck. meth.*

A. folio tenuissimo multifido. *Dill. musc. 477. t. 88. f. 4.*

Habitat in Germania.

1319. LICHEN.

Masc. *Receptaculum* subrotundum, planiusculum, nitidum.

Fem. *Farina* foliis adspersa.

* A. *Leprosi tuberculati.*

1. **LICHEN scriptus.** L. leprosus albicans, lineolis nigris ramosis characteriformibus. *Fl. suec. gallob. Veis. crypt. pal.*

L. bivalvis, de rimis efflorescens, ater characteristicus. *Hall. helu. n. 2079.*

L. crustaceus; tuberculis sessilibus, in lineas ordinatis. *Scop. carn.*

L. cinereo-albidus: lineolis characteriformibus divaricatis. *Neck. meth.*

Lichenes. *Mich. gen. p. 102. n. 9. 10. 11. t. 56. f. 3.*

Lichenoides crusta tenuissima, peregrinis voluti litteris inscripta. *Dill. musc. 128. t. 18. f. 1.*

Habitat in Europæ corticibus arborum.

2. **LICHEN geographicus.** L. leprosus flavescens, lineolis

nigris mappam referens. *Fl. suec. gallob. Weis. crypt. pal. Flor. dan. t. 468. f. 1. lithu.*

L. crusta inseparabili flava, scutellis nigris confluentibus.
Hall. helu. n. 2063.

L. flavicundus mappiformis : lineolis ramosis, punctis atris intermixtis. *Neck. meth.*

Lichen. Mich. nou. gen. p. 97. n. 19.

Lichenoides nigro-flavum tabulæ geographicæ instar pictum.
Dill. musc. 126. t. 18. f. 5.

Habitat in Europæ rupibus altis.

3. LICHEN atrovirens. *L. leprosus viridis marginis tuberculisque atris.*

L. leprosus ater, tuberculis viridibus confertis. *Fl. suec. Spec. plant.*

Habitat in Europæ rupibus.

Tubercula parva, flavo-virescentia, conferta ut tota area flavo-virens conspiciatur, quam cingit margo niger.
Sp. pl.

4. LICHEN biffoides. *L. leproso-farinaceus, peltis stipitatis subglobosis.* *Leers. fl. herb. born.**

Coralloides fungiforme ex ungula equina, livide rubescens.
Dill. musc. 78. t. 14. f. 5.

Fungi parvi globosi ex ungue equino putrescente. *Rai. angl. 3. p. 13.*

Habitat in Europæ glareosis.

Crusta farinosa, viridi-cinerea. Fungi pro ratione magni.

Lichenis ericetorum L. vel. Lich. fungiformis Sc. notabilis varietas esse videtur cl. Webero. spicil. p. 197. R.

5. LICHEN lacteus. *L. leprosus albus, tuberculis concoloribus hemisphæricis.* *fl. herb.* pal.*

Habitat ubique in rupibus & saxis. Zæga.

Tubercula majuscula sunt.

6. LICHEN rupicola. *L. leprosus albidus, tuberculis pallidis albo-marginatis.* *Leers fl. herb.*

Habitat supra rupes planiusculas nudas apricas in sylvis.
Zæga.

Crusta parvissima. Tubercula interdum sunt virescentia, subconvexa absque margine, unde diversa apparet.

7. LICHEN pertusus. *L. leprosus : verrucis subtessellatis lævigatis pertusis poro uno alterove cylindrico.* *herb. pal.*

Flor. dan. t. 766. v.

668 CRYPTOGAMIA. ALGÆ.

L. (*verrucosus*) crustaceus cinereus , scutellis virescentibus.
Huds. angl.

L. (*carpineus*) cinereus : tuberculis pertusis confertis.
meth. musc. cum. L. carpineo Lin. conjungit. R.

Sphæria (*pertusa*) subcomposita aggregata irregularis perforata virescens. *Weig. obs. Weber spicil.*

Lichenoides verrucosum & rugosum cinereum glabrum.
Dill. musc. 128. t. 18. f. 9.

Lichen crustaceus , verrucosus , alpinus , arboribus adnascens , cinereus , receptaculis florum infescentibus. *Mich. gen. 95. ord. 32. tab. 52. optimo habitum plantæ exprimit sec. Weberum. R.*

Lichenoides crustaceum arboreum tenuissimum cinereorum, capsulis exiguis crebris. *Mich. gen. 106. t. 56. f. 2.*

Habitas in Saxis & arboribus Europæ.

Crusta leprosa , alba , tenuis ; ubi vero floret crassa , fissa in verrucas angulatas , læves , poro uno alterove pertusas.

8. LICHEN *rugosus*. L. leprosus albicans , lineolis simplicibus punctisque nigris confertis. *Fl. suec. gallob. Weis. crypt. carn. pal.*

L. albidus : punctis lineolisque aggregatis atris. *Neck. meth. 46. . . .*

Lichenoides punctatum & rugosum nigrum. *Dill. musc. 125. t. 18. f. 2.*

Habitat in Europæ sylvis supra arborum truncos.

Lichen *Hall. stirp. helv. n. 2077. non potest esse Lichen rugosus L. quem in saxis nunquam legi , sed videtur esse Lichen Hall. l. c. n. 2079. var. γ. Neque videtur huc pertinere Lichen Hall. n. 2083. Weigel. l. c. R.*

9. LICHEN *sanguinarius*. L. leprosus cinereo virescens , tuberculis atris. *Fl. suec. gallob. Scholl. barb. pal. herb. lithu.*

L. crusta farinacea virescente , scutellis atris. *Hall. helv. n. 1065.*

Lichenoides leprosum , crusta cinereo virescente , tuberculis integerrimis. *Dill. musc. 126. t. 18. f. 3.*

Habitat in Europæ rupibus truncisque arborum.

Crusta tenuissima , subcinerea. Tubercula atra , subglobosa , magna , non nitida , margine obsoleto angulato , non membranaceo.

10. LICHEN *fusco-ater*. L. leprosus fuscus , tuberculis atris. *Fl. suec. pal.*

L. crusta farinacea inseparabili cinerea, scutis nigerrimis confertis. *Hall. helu. n. 2062. sec. Weis. R.*

Habitat in Europæ rupibus.

11. LICHEN vernalis. **L.** leprosus albidus, tuberculis subrotundis ferrugineis. *herb.*

L. crusta tenuissima cinerea, scutellis fuscis. *Hall. helu. n. 2070. R.*

Lichenoides leprosum tuberculis fuscis & ferrugineis. *Dill. musc. 126. t. 18. f. 4. & t. 55. f. 8.*

Habitat primo vere in aridissimo collo Polonico Upsaliæ, & passim in Europa.

Crusta cinerascens - alba. Tubercula subglobosa, gelatinosa, conferta, sessilia, absque margine calycino.

12. LICHEN calcarius. **L.** leprosus candidus, tuberculis atris. *Fl. suec. Weis. crypt.*

Lichenoides tartareum tinctorium candidum, tuberculis atris. *Dill. musc. 128. t. 18. f. 8.*

Habitat in Europæ rupibus marmoreis.

13. LICHEN cinereus. **L.** leprosus tuberculis nigris albo-marginatis. *Mant. 122. Leers. fl. herb.*

L. crusta inseparabili, reticulata, cinerea, scutellis nigris minutissimis. *Hall. helu. n. 2059. R.*

Habitat ubique in rupibus & saxis.

Similis L. fusco atro, sed tuberculis minoribus. Margo Lichenis niger est.

14. LICHEN atroalbus. **L.** leprosus niger, tuberculis atro alboque mixtis.

L. leprosus niger, tuberculis albis. *Fl. suec.*

Habitat in Alpium rupibus.

Difficile dictu, quinam color constituat crustam.

15. LICHEN ventosus. **L.** leprosus flavus, tuberculis rubris. *Fl. suec. Neck. meth. musc. pal. Fl. dan. 472. f. 2.*

Habitat in Alpium rupibus.

16. LICHEN fagineus. **L.** leprosus albus, tuberculis albis farinaceis. *Neck. gallob. Weis. crypt. pal.*

L. verrucosus, crustaceus, albus, orbiculis villosis concoloribus. *Hall. helu. n. 2086.*

L. (*orbiculatus*) crustaceus albicans, tuberculis albis farinaceis. *Schreb. spicil.*

L. crustaceus albus; tuberculis sessilibus, farinaceis. *Scop. carn.*

L. crustaceus, arboribus adnascens, farinaceus, albus,

omnium crassissimus, receptaculis florum nigricantibus. *Mich. gen. 96. t. 52. f. 6.*

Habitat in Europæ ad parietes rupium.

23. LICHEN *italiensis*. L. crustaceus albicans, scutellis pallidis. *Fl. suec. ged. gallob. Weis. crypt. pal.*

L. fronde laciniata appressa cervina concolore. *Hall. helu. n. 2024.*

L. fronde germinante suborbiculari: scutellifera difformi, primordialibus nudis granulosis viridulis. *Neck. meth.*

L. (*muralis*) imbricatus flavo virescens, scutellis concoloris demum lutescentibus: margine pallido. *Schreb. spicil. p. 120. R.*

L. pulmonarius saxatilis glauco-virescens angustioribus segmentis, receptaculis florum griseis. *Mich. gen. 94. t. 51. f. 4. Schreb.*

Lichenoides crustosum orbiculare incanum. *Dill. musc. 135. t. 18. f. 17.*

Habitat in Europæ lapidibus, assribus, trabibusque putridis.

Synonymon *Dillenii* excludere jubet *Schreberus*.

24. LICHEN *subfuscus*. L. crustaceus albicans, scutellis nigris: junioribus urceolatis cavis. *pal.*

L. crustaceus albicans, scutellis subfuscis: margine cinereo subcrenato. *Fl. suec. gallob. Weis. crypt. norv. ged.*

L. crusta tenace verrucosa cinerea, scutellis rufis marginatis. *Hall. helu. 249.*

L. crustaceus verrucosus albicans; scutell.² subfuscis, sessilibus, planis; margine cinereo, subcrenato. *Scop. carn.*

L. rugosus subcinereus: scutellis subfusci^s margine crenulatis. *Neck. meth.*

L. nigricans. *Ejusd. gallob. p. 504. Varietas est.*

L. crustaceus, arboribus adnascens, ex cinereo albicans, receptaculis florum crebris, subfuscis, limbo albo, crenato cinctis. *Mich. gen. 97. t. 29. Hall. R.*

Lichenoides crustaceum & leprosum, scutellis subfuscis. *Dill. musc. 134. t. 18. f. 16.*

Habitat in Europa, arboribus & rupibus innascens.

25. LICHEN *Parellus*. L. crustaceus albus, pelvis concavis obtusis pallidis. *flor. herb.*

L. verrucosus tenax, albus, scutellis excavatis marginatis, concoloribus. *Hall. helu. n. 2048.*

L. albidus rugoso-calcareus: scutellis irregularibus concoloribus. *Neck. meth.*

Lichenoides

Lichenoides leprosum tinctorium, scutellis lapidum cancri figura. *Dill. musc.* 130. t. 18. f. 10.

Pareille d'Auvergne. Tournef. it. 1. p. 233.

Habitat in Muris.

26. *LICHEN upsaliensis*. L. crustaceus, foliolis subulatis striatis. *Fl. suec.*

Habitat in summis Iugis supra terram sterilissimam campi polonici, Upsaliæ, nec alibi nobis visus.

Magnitudo Lichenis saxatilis, cinereo-albus, constans Setis rectis, albis, emarcidis, prostratis, inæqualibus, confusis, simplicissimis, longitudine unguis, fragilissimis, sæpe pluribus basi coalitis. Scutella albæ margine obtuso, majuscula, radicalia, nec setis insidentia.

C. Imbricati.

27. *LICHEN centrifugus*. L. imbricatus foliolis obsolete multifidis lævibus albidis centrifugis, scutellis rufo-fuscis.

Fl. suec. Weis. crypt. norv. pal.

L. foliis planis multifidis obtusis: laciniis linearibus, calycibus concavis. *Fl. lapp.* 448. t. 11. f. 2.

L. fronde semipinnata, rotunde lobata, crasse villosa, scutellis peramplis. *Hall. helu. n.* 2015.

L. fronde germinante pulverulenta, subrotunda, subtus villoso-atra: frondulis interioribus centrifugis. *Neck. meth.*

L. pulmonarius saxatilis, tenuiter laciniatus, elegantis sulphurei coloris. *Buxb. cent.* 2. t. 7. f. 3. *Hall. R.*

L. imbricatus viridans, scutellis badiis. *Dill. musc.* 180. t. 24. f. 75.

Habitat in Europæ frigida rupibus.

28. *LICHEN saxatilis*. L. imbricatus, foliolis sinuatis scabris lacunosis, scutellis badiis.

L. imbricatus foliolis sinuatis scabris lacunosis, scutellis folio concoloribus. *Fl. suec. gallob. Weis. crypt. pal.*

L. foliis planis acute laciniatis rubiginoso-albidis. *Fl. lapp.*

L. subviridis, fronde semipinnata, rotunde lobata, lineis pulverulentis reticulata, inferne scabra, carbonaria. *Hall. helu. n.* 2011.

L. foliaceus, repens, reticulato-lacunosus, inferne villosus, ater; scutellis rubris. *Scop. carn.*

L. fronde lacunata lævi scutellifeta; germinante asperiuscula, primordialibus pulverulentis. *Neck. meth.* 90.

Linnaei Spec. plant. Europ. Pars. II.

V v

674 CRYPTOGAMIA. ALGÆ.

L. opere phrygio ornatus. Vaill. paris. t. 21. f. 5.

Lichenoides vulgatissimum cinereo-glaucum lacunosum & cirrhosum. Dill. musc. 188. t. 24. f. 83.

Musco-fungus Lichenoides minor cinereus vulgatissimus. Moris. hist. 3. p. 534. f. 15. t. 7. f. 6.

Habitat in Europæ rupibus.

rupestris.

29. *LICHEN omphalodes. L. imbricatus, foliolis multifidis glabris obtusis incanis, punctis vagis eminentibus. Fl. suec. pal.*

L. fronde multifida, supra nitida, subtus villosa - atra: scutellis concavis fronde concoloribus. Neck. meth.

L. nigricans omphalodes Vaill. paris. 116. t. 20. f. 10.

Lichenoides saxatile tinctorium, foliis pilosis purpureis. Dill. musc. 185. t. 24. f. 80.

Habitat in Europæ rupibus.

arboreus & rupestris.

30. *LICHEN olivaceus. L. imbricatus: foliolis lobatis nitidis lividis. Fl. suec. gallob. Weis. crypt. pal. lithu.*

L. foliis planis multifidis obtusis: laciniis apice latioribus. Fl. lapp.

L. fronde semipinnata, rotunde lobata olivacea pulla, scutellis crenatis. Hall. helu. n. 2014.

L. fronde supra nitida scutellifera: margine scutellarum crenato. Neck. meth.

L. olivaceus. Scop. carn.

L. crustæ modo arboribus adnascens, olivaceus. Vaill. paris. t. 20. f. 8.

Lichenoides olivaceum, scutellis amplioribus verrucosis. Dill. musc. 184. t. 24. f. 78.

Habitat in Europæ rupibus.

arboreus & rupestris.

Peltæ orbiculatæ, elevatæ, membranaceæ, undique liberæ.

Cl. Weis. ac Scopolius hunc Lichenem cum L. pullo Schreb. spicil. p. 131. confundere videntur. R.

31. *LICHEN sablunensis. L. imbricatus: foliolis linearibus dichotomis planiusculis acutis nigris, scutellis atris. Fl. suec.*

L. fronde laciniata, undique nigerrima, scutellis concoloribus. Hall. helu. n. 2020.

L. fronde utrinque anthracina lineari-multifida scutellis concoloribus. Neck. meth.

Lichenoides tinctorium atrum, foliis minimis crispis. Dill. musc. 188. t. 24. f. 81.

Habitat in Europæ rupibus nudis.

rupestris.

32. LICHEN *stygius*. L. imbricatus, foliolis palmatis recurvis atris. *Fl. suec.*

Habitat in Suecia, imprimis in Insula Baltici Blåkulla: *rupestris.*

Scutella orbiculata, nigra, concava, margine obtuse crenulato.

33. LICHEN *crispus*. L. imbricatus, foliis lobatis truncatis crenatis atro-viridibus, scutellis concoloribus. *Hallf. anaf. herb. pal. lithu.*

L. (*murorum*) fronde imbricata: frondulis extimis latioribus obtuse crenatis, scutellis concoloribus. *Neck. meth. musci*

Lichenoides gelatinosum atrovitens crispum & rugosum: *Dill. musc. 139. t. 19. f. 23.*

Habitat ad Muros.

Figuram *Dillen. t. 26. f. 26. ad hanc speciem potius pertinere credit cl. Schreberus. R.*

34. LICHEN *cristatus*. L. imbricatus; dentato-ciliatus; scutellis folio majoribus. *gallob.*

L. gelatinosus ramosus, laciniatus, scutellis rubris. *Hall. helu. n. 2039. R.*

L. (*multifidus*) foliaceus, gelatinosus; foliis multifidis; scutellis rubris concavis. *Scop. carn.*

Lichenoides gelatinosum, fuscum, Jacobaeae maritimae divisa. *Dill. musc. 140. t. 19. f. 25. R.*

β. Lichenoides gelatinosum, foliis imbricatis & cristatis. *Dill. musc. 140. t. 19. f. 26.*

Habitat in Europa australi.

35. LICHEN *parietinus*. L. imbricatus, foliis crispis fulvis, peltis concoloribus fulvis. *Fl. suec. Neck. gallob. herb. pal. lithu.*

L. fronde rugosa, rotunde lobata, flava; concolore. *Hall. helu. n. 2021.*

L. foliaceus repens crispus; foliis & scutellis fulvis. *Scop. carn.*

Lichenoides vulgare sinuosum; foliis & scutellis luteis. *Dill. musc. 180. t. 24. f. 76. C.*

Habitat in Europae parietibus, rupibus, lignis.

36. LICHEN *physodes*. L. imbricatus: laciniis obtusis subinflatis. *gallob. pal. lithu.*

L. foliaceus subdepressus laevis: laciniis integerrimis truncatis: subtus concavis lacunosis atris. *Fl. suec.*

676 CRYPTOGAMIA. ALGÆ.

L. fronde semipinnata, cava, cornubus pulverulentis. *Hall. helu. n. 2018.*

L. fronde scutellifera, lævi: frondulis subtubulosis apice dilatiusculis fariniferis. *Neck. meth.*

L. foliaceus, laciniatus, glaber; laciniis tubulosis obtusis clausis. *Scop. carn.*

L. crustæ modo arboribus adnascens tenuiter divisus. *Buxb. cent. 3. p. 272. t. 14. f. 1.*

L. pulmonarius arboribus adnascens: desuper cinereus subrus anthracinus: segmentis teretibus tubulosis. *Mich. gen. 91. t. 50. f. 1. 2.*

Lichenoides ceratophyllum obtusius & minus ramosum. *Dill. musc. 154. t. 20. f. 49.*

Habitat in Europæ corticibus Betulæ, & Saxis ac rupibus.

Dignoscitur a reliquis apicibus foliorum tumidis, & si transversim secantur, inflatis.

Cum L. saxatili conjungit cl. Weisius. (crypt. 57.)

37. LICHEN stellaris. **L.** imbricatus: foliolis oblongis laciniatis angustis cinereis, scutellis pullis. *Fl. succ. Neck. gallob. pal. lithu.*

L. foliis laciniatis cinereis: laciniis obtusiusculis sæpius apice trifidis. *Roy. lugdb. 510.*

L. fronde semipinnata, cinerea, obtuse lobata, scutellis glaucis nigrescentibus. *Hall. helu. n. 2017.*

L. foliaceus, repens, coriaceus, crispus; scutellis atris. *Scop. carn.*

L. fronde cinerascete, lineari-multifida: scutellis fuscis. *Neck. meth.*

L. pulmonarius vulgarissimus superne albo cineretus, inferne nigricans, segmentis angustis, receptaculis nigricantibus. *Mich. gen. 91. t. 43. f. 2.*

Lichenoides cinereum, segmentis argutis stellatis, scutellis nigris. *Dill. musc. 176. t. 24. f. 70.*

Habitat in Europæ ramis arborum.

arboreus.

D. Foliacei.

40. LICHEN ciliaris. **L.** foliaceus erectiusculus: laciniis linearibus ciliaris, scutellis pedunculatis crenatis. *Fl. succ. norv. gallob. Weis. crypt. var. a. pal. Flor. dan. t. 711. burg. lithu.*

L. corniculatus, planus, ramulis capillaribus, scutellis petiolatis. *Hall. helu. n. 1980.*

L. foliaceus, laciniatus, cinereus; laciniis erectiusculis; acutis, pilosis; scutellis pedunculatis. *Scop. carn.*

L. fronde erectiuscula, laciniis linearibus ciliatis: scutellis concaviusculis stipitatis subcrenatis. *Neck. meth.*

L. foliis laciniatis cinereis; laciniis linearibus apice pilosis. *Roy. lugdb.*

L. cinereus arboreus, marginibus pilosis, major. *Vaill. parif. 115. t. 20. f. 4.*

Lichenoides hispidum majus & rigidius, scutellis nigris. *Dill. musc. 150. t. 20. f. 45.*

Musco fungus arboreus cinereus scutellatus; marginibus pilosis. *Meris. hist. 3. p. 634. f. 15. t. 7. f. 6.*

Musculus arboreus pyxoides pilosus. *Læs. pruss. 171. t. 50.*

Habitat in Europæ arboribus. *arboreus.*

41. LICHEN islandicus. **L.** foliaceus adscendens laciniatus; marginibus elevatis ciliatis. *Fl. suec. gad. carn. norv. Weis. crypt. pal. Flor. dan. t. 135. burg. lithu.*

L. foliis oblongis laciniatis; marginibus conniventibus ciliatis. *Fl. lapp.*

L. fronde convexa ciliata pustulata, obtuse ramosa, utrinque lævi, ramis brevissime bicornibus. *Hall. helu. n. 1978.*

L. fronde erectiuscula laxa, margine ciliata: scutellis subrotundis terminalibus planis. *Neck. meth.*

L. pulmonarius minor, &c. *Mich. gen. 85. t. 44. f. 4. Hall.*

L. terrestris, foliis eryngii. *Buxb. cent. 2. p. 11. t. 6. f. 1. 2.*

Lichenoides rigidum, eryngii folia referens. *Dill. musc. 209. t. 28. f. 111.*

Musculus Islandicus purgans. *Borrich. act. Hassn. 1674. p. 126.*

β. Lichenoides eryngii folia referens, tenuioribus & crispioribus foliis. *Dill. musc. 212. t. 28. f. 112.*

γ. Lichen islandicus, tenuissimus.

Borallioides tenuissimum nigricans, mundi muliebris instar textum. *Dill. musc. 113. t. 17. f. 32.*

Habitat in Europæ sylvis sterilissimis, Pinnetis, terrestribus.

Varietas **γ. habitat frequens in sterilissimis collibus Sueciæ, Ramulis omnino teretibus, intus cavis, nec ciliatis, adeoque diversissimus; at color luridus basis rubra. Varia specimina & gradus, unam eandemque speciem esse evincunt.**

42. LICHEN nivalis. **L.** foliaceus adscendens laciniatus

678 CRYPTOGAMIA. ALGÆ.

crispus glaber lacunolus albus margine elevato. *Fl. succ. Scop. carn. Weis. crypt. Fl. dan. t. 227.*

L. niveus sinibus dardaleis laciniatus, ramis erectis calyce orbiculato. *Fl. lapp. 446. t. 11. f. 1.*

L. convexo concavus, lacunatus & glaber, ramis crispis subluteis. *Hall. helu. n. 1977.*

L. fronde laciniata glabra, erecta, nitida; punctis marginalibus obscuris. *Neck. meth.*

Lichenoides lacunosum candidum glabrum, endiviæ crispæ facie. *Dill. musc. 162. t. 21. f. 56.*

Habitas in Lapponiæ, Upsaliæ, Groelandiæ *alpinis, apricis, siccis, glareosis, terrestribus.*

Duplex est varietas 1) albida, quam pinxit *Oederus*; 2) lutea. *Weh. l. c. 2.*

43. *LICHEN pulmonarius. L. foliaceus* laciniatus obtusus glaber; supra lacunosus; subtus tomentosus. *Fl. succ. ged. Weis. crypt. Blackw. t. 353. pal. burg. lithu.*

L. foliis oblongis divitis, superficie reticulato-punctata. *Fl. lapp.*

L. lacunatus, inferne gibbosus, reticulo farinoso, scutellis lateralibus. *Hall. helu. n. 1986.*

L. foliaceus, laciniatus, repens, supra reticulato-lacunosus; scutellis sparsis. *Scop. carn.*

L. fronde obtusa glabra, supra lacunoso-reticulata, subtus tomentosa. *Neck. gallob.*

Lichenoides pulmoneum reticulatum vulgare, marginibus peltiferis. *Dill. musc. 212. t. 29. f. 113.*

Musculus pulmonarius. Bauh. pin. 361.

Pulmonaria. Fuchsf. hist. 631. Cam. epit. 683.

Habitas in Europæ sylvis umbrosis super arbores antiquas, præsertim in *Fagis & Quercubus.*

Monstrum hujus *Lichenis* est in *Ephem. n. C. dec. 1. n. 2. phs. 50.*

44. *LICHEN furfuraceus. L. foliaceus* decumbens furfuraceus: laciniis acutis: subtus lacunosus atris. *Fl. succ. Weis. crypt. pal. lithu.*

L. foliis planis laxis flaccidis inæqualiter laciniatis: supra albis, subtus atris. *Fl. lapp.*

L. cornubus planiusculis ovatis, superne farinosis, inferne atris & reticulatis. *Hall. helu. n. 1976.*

L. fronde ramosissima superne cinerea, subvillosa, inferne nigra; ramulis acutis. *Neck. meth.*

L. arboreus, cornua cervi referens, subtus anthracinus, desuper cinereus. *Mich. gen.* 76. t. 38. f. 1. *Hall. R.*

L. arboreus leucomelanus: ramulis altera parte anthracinis, altera candidissimis. *Buxb. cent.* 2. p. 12. t. 7. f. 1.

Lichenoides cornutum amarum: desuper cinereum; inferne nigrum. *Dill. musc.* 157. t. 21. f. 52.

Muscus amarus, absinthii folio. *Bauh. hist.* 3. p. 764.

Habitat in Europæ arboribus.

Cum L. prunastri a Scopolio conjungitur, a quo vero vere differt. Weber. l. c. R.

45. LICHEN ampullaceus. **L. foliaceus** planiusculus lobatus crenatus, peltis globosis inflatis.

L. foliis laciniatis, marginibus convolutis in vesiculam abeuntibus. *Roy. lugdb.*

Lichneoides tinctorum glabrum vesiculosum. *Dill. musc.* 188. t. 24. f. 82.

Habitat in Lancastria Angliæ.

47. LICHEN farinaceus. **L. foliaceus** erectus compressus ramosus: verrucis marginalibus farinosis. *Schreb. spicil.* 125. *Pollich. pal. burg. lithu.*

L. foliaceus erectus compressus ramosus farinaceus: lateribus verrucosis. *Fl. suec. gallob. ged.*

L. foliis multifidis, calycibus in margine foliorum sessilibus. *Roy. lugdb.*

L. lacunofus, complanatus, ramis acutissimis, orbiculis farinosis marginalibus frequentissimis. *Hall. helv. n.* 1981.

L. foliaceus erectus, laciniatus; laciniis acutis, rugosus, rigidis; verrucis lateralibus. *Scop. carn.*

L. rostratus. *Femina. Scop. carn. ed. 2. cum sequenti conjungit.*

L. fronde lineari ramosa, laterali verrucosa: verrucis planis farinosis. *Neck. meth.*

L. cinereus angustior, scutis in marginibus segmentorum. *Vaill. paris.* 115. t. 20. f. 14. 15. 13.

Lichenoides segmentis angustioribus ad margines verrucosis & pulverulentis. *Dill. musc.* 172. t. 23. f. 63.

Habitat in Europæ arboribus, præsertim Quercetis, Fraxineis.

48. LICHEN calicaris. **L. foliaceus** erectus linearis ramosus lacunofus convexus mucronatus. *Fl. suec. ged. gallob. Weis. crypt. pal. burg. lithu.*

680 CRYPTOGAMIA. ALGÆ.

- L. utrinque lacunatus, complanatus, farinosus & scutellifer. *Hall. helu. n. 1983.*
- L. foliaceus erectus laciniatus; laciniis linearibus acutis rigidis; scutellis terminalibus. *Scop. carn.*
- L. rostratus, mas. *Scop. carn. cum præcedente jungit.*
- L. fronde cartilaginea, glabra, ramosa, erecta; scutellis appendiculatis terminalibus. *Neck. meth.*
- L. cinereus latifolius ramosus. *Vaill.-parif. 115. t. 20. f. 6.*
- Lichenoides coralliforme rostratum & caniculatum. *Dill. musc. 170. t. 23. f. 162. A. B. C.*
- Musco-fungus arboreus capitulis rostratis. *Moris. hist. 3. p. 634. f. 15. t. 7. f. 5.*
- β. Lichen pyxidatus damæ cornu divisura, acetabulorum oris crispis. *Vaill. parif. 115. n. 6. t. 21. f. 2. Weis. R.*
- Habitat in Europæ arboribus, rupibus.*
49. LICHEN *fraxineus*. L. foliaceus erectus oblongus lanceolatus sublaciniatus lacunosus glaber, scutellis subpedunculatis. *Fl. suec. gallob. Weis. crypt. Kniph. cent. 6. n. 55. burg. lithu.*
- L. foliis oblongis planis rugosis sinuatis, calycibus orbiculatis petiolatis. *Roy. lugdb.*
- L. utrinque lacunatus, fronde latissima, scutellis creberrimis. *Hall. helu. n. 1985.*
- L. foliaceus, erectus, laciniatus; laciniis lanceolatis, obtusis, rugosis, rigidis; scutellis sessilibus, confertis, sparsis. *Scop. carn.*
- L. fronde oblonga erecta, utrinque lacunata scutellataque: scutellis sessilibus & stipitatis. *Neck. meth. 105.*
- L. pulmonarius cinereus mollior in amplas lacinias divisus. *Tournef. inst. 549. t. 325. f. A. B.*
- Lichenoides longifolium rugosum rigidum. *Dill. musc. 165. t. 22. f. 59.*
- Musco fungus quernus latifolius cinereus. *Moris. hist. 3. p. 634. f. 15. t. 7. f. 14. 3. 4.*
- β. Lichen pulmonarius, rufescens, durior, in amplas lacinias divisus. *Mich. gen. 74. t. 36. f. 1. Hall. Weis. R.*
- Habitat in Europæ arboribus, præsertim Fraxinis.*
51. LICHEN *prunastri*. L. foliaceus erectiusculus lacunosus: subrus tomentosus albus. *Fl. suec. ged. gallob. Weis. crypt. pal. Kniph. cent. 12. n. 68. burg. lithu.*
- L. foliis mollibus candicantibus dichotomis, calycibus orbiculatis sessilibus. *Roy. lugdb.*

L. complanatus, utrinque lacunatus, undique farinosus. *Hall. helu.* 1984.

L. foliaceus, suberectus, laciniatus: laciniis furfuraceis, subtus lacunosis. *Scop. carn.*

L. fronde verrucifera molliuscula, tomentosa: scutellifera cartilaginea glaberrima: scutellis terminalibus utrinque planis. *Neck. meth.*

L. cinereus vulgatissimus cornua damæ referens. *Vaill. paris.* 115. t. 20. f. 11. 12.

L. pulmonarius mollior dichotomus, superne cinereus. *Mich. gen.* 75. t. 36. f. 3. *Hall. Vais. R.*

Lichenoides cornutum, bronchiale molle subtus incanum. *Dill. musc.* 160. t. 21. f. 55.

β. Lichenoides corniculatum candidum molle, segmentis angustis. *Dill. musc.* 159. t. 21. f. 54.

Lichen cornua Damæ referens, angustifolius. *Vaill. paris.* t. 20. f. 7.

Habitat ad Europæ arbores siccas.

52. **LICHEN juniperinus**. **L. foliaceus** laciniatus crispus fulvus, peltis lividis. *Fl. suec.*

L. fronde germinante byssiformi: explicata subrotunda; frondulis erectiusculis, exterioribus latioribus. *Neck. meth.* 96. cum **L. parietino** conjungit.

L. fulvus sinubus dædaleis laciniatus. *Fl. lapp.*

Habitat in Europæ juniperetis.

Differt a L. parietino colore pallide flavo; foliis laxis erectiusculis; peltis disco brunneis.

53. **LICHEN caperatus**. **L.** pallide viridis rugosus margine undulatus. *stamp. ged. gallob. Weis. crypt. pal.*

L. fronde rotunde lobata, rugosa, inferne atra & aspera. *Hall. helu. n.* 2005.

L. foliaceus, repens, lobatus, supra luteo-virescens, infra lævis, ater; scutellis verrucosis, sessilibus, umbilicatis rubris. *Scop. carn.*

L. fronde juniore supra rugosiuscula: germinante pulverulenta, frondulis extimis lævibus, latioribus obtusis. *Neck. meth.*

Lichenoides caperatum rosacee expansum. *Dill. musc.* 193. t. 25. f. 97. *C. C.*

Musco-fungus lichenoides, crustæ modo adnascens major cinereus. *Morif. hist.* 3. p. 633. f. 15. t. 7. f. 1.

Habitat in Europa & America ad saxæ & arbores.

Glaucescenti - lateus parum foliaceus, raro scutella præferens.

55. LICHEN *glaucus*. L. foliaceus depressus lobatus glaber : margine crispo farinaceo. *Fl. suec. pal. Fl. dan. t. 598.*

L. fronde laciniata & crispa, inferne atra & aspera. *Hall. helu. n. 2007.*

L. foliaceus, repens, lobatus, supra cinereus, subrus ater; marginibus verrucosis. *Scop. carn.*

L. fronde laciniata nitida, margine crispo-farinosa subrus atra: punctis scutelliformibus bruneis. *Neck. meth.*

L. pulmonarius saxatilis cinereus minor, umbilicis nigricantibus. *Vaill. paris, tab. 2. f. 12. Weber. R.*

L. pulmonarius, crispus inferne nigerrimis & glaber, superne cinereus, receptaculis florum subobscuris. *Mich. gen. 91. t. 50. f. 1. Weber. R.*

Lichenoides endiviæ foliis crispis splendentibus subrus nigricantibus. *Dill. musc. 192. t. 25. f. 96.*

Musco-Fungus lichenoides arborum crispus cinereus subrus nigricans. *Morif. hist. 3. p. 633. f. 15. t. 7. f. 4.*

Habitat in Europæ frigida, præsertim Sueciæ, truncis Betulinis.

Folia papyro tenuiora, albo cinerascente glauca.

56. LICHEN *fascicularis*. L. foliaceus gelatinosus, tuberculis turbinatis fasciculatis, fronde majoribus. *Oed. dan. t. 462. g. 2.*

L. (*glomeratus*) fronde erecta, scutellis terminalibus conglobatis. *Neck. meth.*

Lichen (*glomeratus*) foliis rectis congestis, bruneis, apice scutelliferis. *Neck. gallob. R.*

Lichen foliaceus, gelatinosus, tuberculatus, ater; scutellis convexis, rubris. *Scop. carn. R.*

Lichenoides gelatinosum palmatum, tuberculis conglomeratis. *Dill. musc. 141. t. 19. f. 27.*

Habitat in Suffexia Angliæ. Dill. ad Stenbrohult Smolandia. Filius. In Germania. Weber.

Folia radicalia, minuta, uti totus, gelatinosa, hyalina Tremellæ instar. Tubercula ratione folii ad modum magna, numerosa, subpetiolata, turbinata, fasciculata, extimo truncata, limbo obtuso cincta.

E. Coriacei.

57. LICHEN *aquaticus*. L. coriaceus repens lobatus obtusus, peltis hemisphæricis maximis. *Fl. suec.*

Lichenoides scutellis amplis. *Dill. musc. 150. t. 20. f. 44.*

Habitat in Suecia sub aqua in paludibus.

serm. 2. 12.

Planta diversa a Lichene crinito. Dill. musc. t. 84. f. 8. Folia lobata, pallide viridia, margine integerrima. Peltæ subsessiles, primum planæ, demum dilatatæ, folia majores, incarnato fuscæ, margine reflexo, supra discum folii positæ,

58. LICHEN *resupinatus*. L. coriaceus repens lobatus, peltis marginalibus posticis. *Fl. suec. herb. Flor. dan. t. 764.*

L. fronde peltifera sinuata, latiore lobata distincta: verrucifera crispata, peltis posticis. *Neck. meth.*

L. pulmonarius major s. minor ex obscuro cinereus: inferne ex albo rufescens, receptaculis florum rubris ad latera oblongis. *Mich. gen. 86. t. 44. f. 1. 2.*

Lichenoides fuscum, peltis posticis ferrugineis. *Dill. musc. 206. t. 28. f. 105.*

Habitat in Europæ sylvis. terrestris.

59. LICHEN *venosus*. L. coriaceus repens ovatus planus subtus venosus villosus, peltis marginalibus horizontalibus. *Fl. suec. gallob. Pollich. pal.*

L. flabelliformis, subtus reticulatus, scutellis marginalibus planis. *Hall. helu. n. 1993.*

L. fronde simplici venosâ decumbente: peltis planis orbiculatis. *Neck. meth.*

L. pulmonarius, minimus, inferne albus & niger reticulatus, &c. *Mich. gen. p. 85. n. 12. t. 44. f. 3. Hall. R.*

Lichen pulmonarius. *Ejusd. l. c. p. 86. n. 13. t. 44. f. 5. Hall. R.*

Lichenoides parvum virescens, peltis nigricantibus planis. *Dill. musc. 208. t. 28. f. 109.*

Habitat in Europa ad margines scrobiculorum in sylvis. terrestris.

60. LICHEN *aphtosus*. L. coriaceus repens lobatus obtusus planus, verrucis sparsis, peltâ marginali adscendente. *Fl. suec. Veis. crypt. carn. pal. Flor. dan. t. 767. f. 1. lithu.*

L. fronde obtuse lobata, superne verrucosa, inferne villosa. *Hall. helu. n. 1992.*

L. fronde lobata repente supra verrucata: peltis oblongis marginalibus adscendentibus. *Neck. meth. 80.*

L. pulmonarius maximus, verrucosus, superne e cinereo virescens inferne obscurus, receptaculis florum rubris & circinatis. *Mich. gen. 85. sec. Hall. R.*

Lichenoides digitatum læte virens, verrucis nigris notatum. *Dill. musc. 207. t. 28. f. 106.*

684 CRYPTOGAMIA. ALGÆ.

Habitat in Europæ sylvis acerosis sterilibus sub juniperis: terrestres.

L. canini varietatem facit cl. Weifs. , quæ tamen non est. Weber l. c. R. Folia siccata sunt glauca , virentia vero saurate viridia.

61. LICHEN *arcticus*. *L. coriaceus repens lobatus obtusus planus lævis : subtus avenius villosus. Fl. suec.*

Lichen foliis planis subrotundis lobatis obtusis , calyce plano ovali lacinulæ propriæ adnato , niveus. Fl. lapp. 442.

Habitat in Suecia boreali , sub juniperis. terrestres.

Folia lata , integra , albida , subtus nivea Pelta subtus versus marginem , magna , livido incarnata . magnitudine pollicis , adeoque inter omnes Lichenes latissima. Supra læve est folium.

62. LICHEN *caninus*. *L. coriaceus repens lobatus obtusus planus subtus venosus villosus , pelta marginali adscendente. Fl. suec. ged. gallob. Weifs. crypt. pal. Blackw. t. 336. Flor. dan. t. 767. f. 2. lithu.*

L. foliis planis subrotundis lobatis obtusis calyce plano lacinia propriæ adnato. Fl. lapp.

L. fronde subrotunde lobata , inferne reticulata , peltis convexo-concavis. Hall. helu. n. 1988.

L. foliaceus , repens , subtus reticulato - venosus ; peltis marginalibus , suberectis. Scop. carn.

L. fronde simplici obtusa sinuato-lobata , subtus venoso-villosa : peltis ovatis adscendentibus. Neck. meth.

Lichen folio orbiculato margine integriusculo , superficiæ subtus pustulata. Fl. lapp.

L. pulmonarius saxatilis digitatus. Vaill. paris. 116. t. 21. f. 16.

Lichenoides digitatum cinereum , lactucæ foliis sinuosis. Dill. musc. 200. t. 27. f. 102.

Musco-Fungus terrestris latifolius cinereus , hepaticæ facie. Moris. hist. 3. p. 632. f. 15. t. 7. f. 1.

Habitat in Europæ sylvis , juxta lapides , in terra. terrestres.

Folia quasi farina cinerea sunt tecta.

63. LICHEN *sylvaticus*. *L. coriaceus repens laciniatus lacunosus , peltis marginalibus adscendentibus. Hudf. angl. fl. herbor. **

L. lacunatus exasperatus , inferne villosus & scutellifer. Hall. helu. n. 1987.

L. (polyschides) fronde rigidiuscula oblonga supra verrucata lacunataque : peltis marginalibus subrotundis. *Neck. meth.*

Lichenoides polyschides villosum & scabrum, peltis parvis: *Dill. musc.* 199. t. 27. f. 101. *Mich. gen.* 84. t. 43.

Habitat in Angliæ, Germaniæ, Helvetiæ *rupibus. arboribus.*

64. LICHEN horizontalis. L. coriaceus repens planus subtus avenius, peltis marginalibus horizontalibus. *Huds. angl. Flor. dan.* t. 765. *herb. gallob.*

L. fronde rotunde lobata, peltis planissimis. *Hall. helu. n.* 1991.

L. fronde repente sinuato-lobata peltata ac verrucata distincta; peltis marginalibus orbicularis planis. *Neck. meth.* 78.

Lichenoides subfuscum, peltis horizontalibus planis. *Dill. musc.* 205. t. 28. f. 104.

Lichen pulmonarius minor inferne obscurus. *Mich. gen.* 85. t. 44. f. 1. 6.

Habitat in Helvetia, Anglia, Italia, Suecia, Germania.

Madidus dum est, Pelta horizontalis undique marginata.

65. LICHEN perlatus. L. coriaceus repens lobatus glaber: subtus ater, scutellis peduncularis integris. *Huds. angl. herborn. gallob. pal.*

L. pulmonarius saxatilis cinereus minor, umbilicis nigricantibus. *Vaill. paris.* t. 21. f. 12.

Lichen pulmonarius crispus, inferne nigricans glaber, superne cinereus, receptaculis subobscuris. *Mich. nou. gen.* 91. t. 50. f. 51.

Lichenoides glaucum perlatus subtus nigrum & cirrhosum. *Dill. musc.* 147. t. 20. f. 39.

Habitat in Europa *ad arborum, potissimum Quercuum truncos.*

66. LICHEN faccatus. L. coriaceus repens subrotundus: peltis depressis subtus faccatis. *Fl. suec. Oed. dan.* t. 532. f. 3. *herb.*

L. (*immersus*) foliaceus repens subrotundus planus peltis immersis. *Hudson. angl.*

L. fronde coriacea, rotunde lobata, scutellis immersis. *Hall. helu. n.* 1995.

L. foliaceus repens, lobatus, glauco-virescens; scutellis immersis fuscis. *Scop. carn.*

L. fronde reptante subcrenata lobata : peltis immersis. *Neck. meth.*

L. pulmonarius ; alpinus , terrestris , glauco - virescens ; receptaculis florum fuscis. *Mich. gen.* 95. *ord.* 31. *t.* 52. *Lichenoides* lichenis facie , peltis acetabulis immersis. *Dill. musc.* 223. *t.* 30. *f.* 121.

Habitat in Alpibus Europæ.

67. *LICHEN croceus.* *L.* coriaceus repens subrotundus planus ; subtus venosus villosus croceus , peltis sparsis adnatis. *Fl. suec. Flor. dan.* *t.* 263. *terrestris.*

L. foliis subrotundis planis levissime incis , calycibus orbiculatis disco folii adnatis. *Fl. lapp.* 443. *t.* 11. *f.* 3:

L. fronde rotunde lobata , plana , subtus venosa , miniata. *Hall. helu. n.* 1994.

L. fronde reptante subrotunda subtus venoso-villosa crocea : peltis sessilibus orbiculatis. *Neck. meth.* 84:

Lichenoides subtus croceum , peltis appressis. *Dill. musc.* 221. *t.* 30. *f.* 120.

Habitat in Lapponia , Helvetia , Groenlandia. *terrestris.*

F. Umbilicati , squalentes quasi fuligine.

68. *LICHEN miniatus.* *L.* umbilicatus gibbus punctatus ; subtus fulvus. *Scop. carn. Fl. dan.* *t.* 532. *f.* 1:

L. pulmonarius alpinus cinereus punctatus. *Hall. helu.* 75. *t.* 2. *f.* 2.

L. fronde imbricata , rotunde lobata , cinerea ; punctata ; inferne ochrea subaspera. *Hall. helu. ed.* 2. *n.* 1999. *t.* 47. *f.* 2:

L. fronde complicata , lobata , supra punctata. *Neck. meth.*

L. pulmonarius , saxatilis , e cinereo fuscus , minimus. *Mich. gen.* 101. *ord.* 36. *t.* 54. *f.* 1. *Hall. R.*

Lichenoides coriaceum nebulosum cinereum punctatum , subtus fulvum. *Dill. musc.* 223. *t.* 30. *f.* 127.

Habitat in Angliæ , Helvetiæ , &c. rupibus alpinis.

69. *LICHEN velleus.* *L.* umbilicatus : subtus hirsutissimus. *Fl. suec.*

L. folio subrotundo peltato ; margine fere integro , subtus margine hirsutus. *Fl. lapp.*

L. fronde coriacea , inferne pustulata , villosa , scutellis atris. *Hall. helu. n.* 1997: *R.*

L. fronde perampla sinuato-crenata inferne villosissima. *Neck. meth.*

Lichenoides coriaceum , latissimo folio umbilicato & verrucoso. *Dill. musc.* 145. *t.* 82. *f.* 5.

Habitat in alpinis Lapponiæ, Sueciæ, Angliæ. rupestris.

70. LICHEN *pustulatus*. L. umbilicatus; subtus lacunosus, furfure nigro adpersus. *Fl. suec. pal. Fl. dan. t. 597. f. 2.*
L. fronde cinerea, inferne lacunata, arbusculis superne nigris. *Hall. helu. n. 1996.*

L. fronde lobata fusca subtus lacunata: furfure marginall concolore. *Neck. meth.*

L. pulmonarius, saxatilis, inferne reticulatus & lacunatus, superne cinereus, ac verrucosus, receptaculis florum, & seminibus nigricantibus & veluti deustis. *Mich. gen. 89. ord. 18. t. 47.*

Lichenoides crustæ modo saxi adnascens verrucosus cinereus & veluti deustus. *Vaill. paris. 116 t. 20. f. 9.*

Lichenoides pustulatum cinereum & veluti ambrustum. *Dill. musc. 227. t. 30. f. 131.*

Habitat in Europæ rupibus apricis.

rupestris.

Humidus virescit.

71. LICHEN *proboscideus*. L. umbilicatus, peltis turbinatis truncatis perforatis. *Fl. suec.*

L. *cylindricus* foliaceus ciliatus, peltis cylindricis rectis sparsis perforatis truncatis circulo duplici. *Amæn. acad.*

Lichenoides foliorum laciniis crinitis. *Dill. musc. 149. t. 20. f. 42.*

Habitat in Suecia, Lapponia, Pennsylvania.

Folium magnitudine pollicis, subrotundum, orbiculatum, margine inæqualiter & obtuse lobatum, centro radicum, subtus læve, hinc inde & rarius fibræ radicantem exserens; Pagina superior planiuscula vel nonnihil contorta, cinerea, punctis fuscis elevatis exasperata. Peltæ per discum sparsæ, atræ, minimæ turbinatæ, truncatæ; poro pertusæ usque ad folium, margine lato plano.

72. LICHEN *deustus*. L. umbilicatus undique lævis. *Fl. suec. burg.*

L. fronde rotunde lobata, inferne nigerrima globosa. *Hall. helu. n. 1998.*

L. fronde siccitate fragili lævi. *Neck. meth.*

Lichen folio orbiculato peltato margine integriusculo undique glaber. *Fl. lapp.*

Lichen pulmonarius saxatilis e cinereo fuscus minimus. *Dill. musc. 116. t. 21. f. 14.*

Lichenoides coriaceum cinereum, peltis atris compressis. *Vaill. paris. 219. t. 20. f. 117.*

Habitat in Sueciæ, Galliæ rupibus s. caulibus elevatis apricis.

rupestris.

688 CRYPTOGAMIA: ALGÆ

73. LICHEN *polyphyllus*. L. umbilicatus polyphyllus utrinque lævis atrovirens crenatus. *Fl. suec.*

L. lapideus crispus, coloris utrinque schistei. *Hall. helu. n. 2001. t. 47. f. 5.*

L. frondulis imbricatis utrinque lævibus margine crenatis. *Neck. meth. 77.*

Lichenoides tenue pullum, foliis utrinque glaucis. *Dill. musc. 225. t. 30. f. 129.*

Habitat in Europæ rupibus elatis apricis. rupestris.

74. LICHEN *polyrhizus*. L. umbilicatus polyphyllus utrinque lævis polyrhizos. *Fl. suec. Weis. crypt. Fl. dan. t. 597. f. 1.*

L. fronde imbricata subrotunda punctata, inferne villosa, scutellis atris. *Hall. helu. n. 2000. t. 47. f. 4.*

L. frondulis superne lacunatis, subtus villosis. *Neck. meth. 77.*

L. ad marginem radicans alpinus, ex cœruleo nigricans, scutellis nigerrimis. *Hall. enum. helu. t. 2. f. 4. R.*

Lichenoides pullum superne & glabrum, inferne nigrum & cirrhosum. *Dill. musc. 226. t. 30. f. 130.*

Habitat in Arvonix, Sueciæ rupibus apricis elatis.

rupestris.

Folia cinereo glauca, subtus hirta.

G. Scyphiferi.

75. LICHEN *cocciferus*. L. scyphifer simplex integerrimus, stipite cylindrico, tuberculis coccineis. *Fl. suec. gallob. herb. pal. lithu.*

L. caule simplici, calyce turbinato margine carnosso colorato prolifero. *Fl. lapp.*

L. pyxidatus, cocciferus. *Weis. crypt. var. f.*

L. scyphiger, infundibulorum fungulis coccineis. *Hall. helu. n. 1918. infundibulorum, fungis fuscis. ibid. n. 1919. infundibulis proliferis, fungulis coccineis. n. 1927. infundibulis proliferis, fungulis atrofuscis. n. 1928. sec. Weis & Leers. R.*

L. coralloides, caule simplici pyxidato, marginibus tuberculosis, tuberculis coccineis sessilibus. *Scop. carn.*

L. stipite cylindrico scyphifero: scyphis integerrimis. *Neck. meth. 57.*

L. pyxidatus, oris coccineis & tumentibus. *Vaill. paris. 115. t. 11. f. 4. Mich. gen. 82. t. 41. f. 3.*

Coralloides scyphiforme, tuberculis coccineis. *Dill. musc. 82. t. 14. f. 7.*

Habitat in Europæ sylvis sterilibus, ericetis, rupibus.

76. LICHEN *cornucopioides*. L. *scyphifer simplex* folio brevior, tuberculis coccineis. *Fl. succ. herb.*

L. caule simplici, calyce turbinato: centro multipliciter prolifero. *Fl. lapp.*

L. infundibulis proliferis; foliosis & fungosis. *Hall. helu. n. 1929.*

L. stipite simplici, scyphis margine tuberculatis foliatisque. *Neck. meth. cum precedente conjungit.*

Coralloides scyphiforme: marginibus radiatis & foliatis. *Dill. musc. 85. t. 14. f. 9.*

Muscus pyxoides. *Barr. rar. 1284. t. 1268. f. 2.*

Habitat in Lapponiæ, Sueciæ, Angliæ sylvis glareosis.

77. LICHEN *pyxidatus*. L. *scyphifer simplex crenulatus*, tuberculis fuscis. *Fl. succ. ged. gallob. pal. herb. burg. lithu.*

L. caule simplici, calyce turbinato margine tenui. *Fl. lapp.*

L. scypho infundibuliformi simplici. *Hall. helu. n. 1912.*

L. pyxidatus, tuberculatus. *Weis. crypt. var. b.*

L. coralloides; caule simplici, pyxidato: marginibus tuberculosis, tuberculis sessilibus, fuscis. *Scop. carn. ed. 1. p. 87. n. 8. tuberculis stipitatis fuscis. ibid. n. 9.*

L. pyxidatis major, acerbulo fimbriato & tuberculoso. *Vaill. paris. t. 21. f. 11.*

L. scypho crenulato, tuberculis sessilibus stipitatisque. *Neck. meth.*

L. pyxidatus major. *Tournef. inst. 349. t. 325. f. D. Mich. gen. 82. t. 41. f. 2. 4.*

Coralloides scyphiforme, tuberculis fuscis. *Dill. musc. 79. t. 14. f. 6. B. C. & fig. 8. B.*

Fungus terrestris pyxidatus. *Magn. hort. 83. t. 83.*

β. Lichen caule simplici, calyce turbinato: centro simpliciter prolifero. *Fl. lapp.*

L. pyxidatus, prolifer. *Weis. crypt. p. 87.*

L. infundibulis glabris proliferis. *Hall. helu. n. 1923. infundibulis serratis proliferis. n. 1924.*

L. coralloides, caule simplici pyxidato: marginalibus tuberculosis proliferisque. *Scop. carn. marginibus proliferis serratis nudis.*

Coralloides scyphiforme tuberculis fuscis. *Dill. musc. p. 79. t. 14. f. 6. D.*

Musco-fungus pyxidatus, calyce altero alteri innato. *Morif. hist. 3. p. 632. f. 13. t. 7. f. 4. Hall. Weis. R.*

γ. Lichen caule simplici, calyce turbinato: centro simpliciter prolifero. *T. Vaill. paris. 113. t. 21. f. 3. 9. t. Plant. Europ. Pars. 11. X x*

890 CRYPTOGAMIA. ALGÆ.

Lichen pyxidatus major alpinus, receptaculis copiosioribus & rufescentibus. *Mich. gen.* 82. t. 41. f. 7. 9.

Habitat in Europæ sylvis.

78. LICHEN *umbellatus*. L. scyphifer simplex denticulatus, stipite cylindrico. *Fl. suec. gallob. pal. burg. lithu.*

L. caule simplici, calyce turbinato: margine acuto prolifero. *Fl. lapp.* 436.

L. pyxidatus fimbriatus. *Weisf. crypt. var.*

L. scyphiger infundibulis serratis. *Hall. helu. n.* 1915. infundibulis serratis coronatis. *helu.* 1916.

L. coralloides, caule simplici pyxidato nudo: marginibus serratis. *Scop. carn.* pyxidato squamoso.

L. stipite cylindrico scyphifero: scyphis margine denticulatis. *Neck. meth.* cum L. pyxidato conjungit.

L. pyxidatus minor. *Vaill. parif.* 115. t. 21. f. 6. *Mich. gen.* 83. t. 41. f. 45.

Coralloides scyphiforme gracile, marginibus serratis. *Dill. musc.* 84. t. 14. f. 8. 9.

Habitat in Europæ silvis sterilibus.

79. LICHEN *gracilis*. L. scyphifer ramosus denticulatus: ramis *Fl. suec.* *Neck. gallob. herb. pal.* *Weber. spm.* 223. *Weisf. crypt. var. h. lithu.*

L. infundibulis proliferis & corniculatis. *Hall. helu. n.* 1946.

L. coralloides, caule simplici subulato nudo. *Scop. carn.*

L. stipite scyphigero filiformi simplici ramisque, scyphis denticulatis. *Neck. meth.*

L. pyxidatus & corniculatus ramosus alpinus e fascio cæreus, pyxidulis crenatis. *Mich. gen.* 81. t. 41. f. 5.

Coralloides scyphiforme ferratum elatius, caulibus gracilibus glabris. *Dill. musc.* 88. t. 14. f. 13. C.

Musco fungus pyxidatus gracilior ramosus, calycibus serratis. *Morif. hist.* 3. p. 362. f. 15. t. 7. f. 6.

Habitat in Europæ ericetis, montosis, sylvis arcticis.

80. LICHEN *digitatus*. L. scyphifer ramosissimus: ramis cylindricis calycibus integris nodosis. *Fl. suec. pal.* *Weisf. crypt.*

L. caule inordinate ramoso, ramis in calyces turbinatos margine carnosos desinentibus. *Fl. lapp.* 436.

L. infundibulis palmaris, fungulis coccineis. *Hall. helu. n.* 1939.

L. coralloides, caule ramoso; ramis in tubercula coccinea desinentibus. *Scop. carn.*

L. stipite crustaceo ramoso : ramulis terminalibus palmatis tubercularisque. *Neck. meth.* 59. n. 20.

Coralloides ramulosum ; tuberculis coccineis: *Dill. musc.* 96. t. 15. f. 19.

Habitat in Europæ sylvis sterilibus.

81. *LICHEN cornutus*. *L.* scyphifer simpliciusculus subventricosus , calycibus integris. *Fl. suec. gallob. pal. lithu.*

L. caule simplici subulato , rarius bifido. *Fl. lapp.*

L. pyxidatus cornutus. *Weis. crypt.*

L. cornubus simplicissimis acutis farinosis. *Hall. helu. n.* 1902. cornubus subramosis farinosis. *Hall. l. c. n.* 1903.

L. coralloides ; caule subulato bifido. *Scop. carn.*

L. stipite subnudo difformi subventricoso : scyphis proliferis. *Neck. meth.*

L. cinereus proboscideus & corniculatus ut plurimum non ramosus. *Mich. gen.* 81. n. 12. 13. 14.

Coralloides vix ramosum , scyphis obscuris. *Dill. must.* 90. t. 15. f. 14. scyphiforme cornutum. 92. t. 15. f. 16. *E.*

Musculus fistulosus corniculatus. *Barr. rar.* 1286. t. 1277. f. 1. *Bacc. musc.* 2. p. 149. t. 107.

Habitat in Europæ ericetis.

82. *LICHEN deformis*. *L.* scyphifer simpliciusculus subventricosus , calycibus dentatis. *Fl. suec. herb. burg.*

L. caule simplici , apice acuto s. calyce turbinato terminato. *Fl. lapp.* 433. t. 11. f. 3.

L. corniculo simplici , maximo , lævi. *Hall. helu. n.* 1904.

L. pyxidatus difformis ; acetabulis cornigeris. *Hall. helu. n.* 1938.

L. pyxoides teres , acetabulis minoribus repandis. *Mich. gen.* 80. t. 41. f. 1.

Coralloides crassius subincanum , calycibus dentatis. *Dill. musc.* 95. t. 15. f. 18. *mala.*

Habitat in Europæ ericetis.

H. fruticulosi.

83. *LICHEN rangiferinus*. *L.* fruticulosus perforatus ramosissimus , ramulis nutantibus. *Fl. suec. ged. Weiz. obs.* 38. n. 31. *gallob. Weis. crypt. pal. burg. lithu.*

Lichen rangiferinus , alpestris. *Flor. dan.* t. 180. *Kniph. cent.* 6. n. 56.

Lichen caule erecto tereti ramosissimo , alis perforatis. *Fl. lapp.* 437.

L. fruticosus , farinosus , surculis innumeris nutantibus. *Hall. helu. n.* 1957.

692 CRYPTOGAMIA. ALGÆ.

L. coralloides , caule ramoso candido , extremitatibus patulis rufescentibus radiatis. *Scop. carn.*

L. (*alpestris*) fronde tereti , adulta ramosissima lævi alis perforata : ramulis nutantibus furcatis tuberculiferis. *Neck. meth.*

Coralloides montanum fruticuli specie ubique candicans. *Dill. musc.* 107. t. 16. f. 29.

Musco-fungus coralloides montanus ramosissimus cinereus vulgaris. *M. ris. hist.* 3. p. 633. f. 15. t. 7. f. 9.

Muscus coralloides f. cornutus montanus. *Bauh. pin.* 361.

Muscus corallinus f. Corallina montana. *Bauh. hist.* 1198.

β. Lichen rangiferus , sylvaticus. *Oed. fl. dan.* t. 539.

L. caule fruticoso folioso , ramis subnutantibus , minute multifidis , fungiferis. *Hall. helu. n.* 1956.

L. caule ramoso , alis perforatis , ramis capitatis. *Roy. lugdb.* 511.

Coralloides fruticuli specie candicans , corniculis rufescentibus. *Dill. musc.* 110. t. 16. f. 30.

Muscus terrestris coralloides erectus , corniculis rufescentibus. *Bauh. pin.* 361. *prodr.* 152.

Habitat in Alpibus & Europæ frigida sylvis sterilissimis.

Alpestris differt a sylvatico , ut flos plenus a simplici.

84. LICHEN uncialis. **L.** fruticulosus perforatus , ramulis brevissimis acutis. *Fl. suec. ged. gallob. pol. herb. lithu.*

Lichen caule ramoso , alis perforatis , ramis brevissimis acutis. *Fl. lapp.*

L. fruticosus , levis , perforatus , cauliculis palmatis brevissimis. *Hall. helu. n.* 1960.

Coralloides perforatum minus molle & tenue. *Dill. musc.* 99. t. 16. f. 22.

β. Lichen (*ceraoides*) fronde tereti aculeata , alis perforata lævi ventricosa : ramulis terminalibus subradiatis. *Neck. gallob.*

L. fruticosus lævis perforatus spinosus , ramulis punctiferis. *Hall. helu. n.* 1959.

Coralloides perforatum majus molle & crassum. *Dill. musc.* 98. t. 16. f. 21.

L. coralloides tubulosus albidus major & mollior , caulibus crassioribus , minus ramosis , receptaculis florum perexiguus rufescentibus. *Mich. gen.* 79. n. 6. t. 40. f. 2.

γ. Lichen caule fruticoso , compresso , ramosissimo , ramulis punctiferis. *Hall. helu. n.* 1961. *Web. R.*

L. coralloides tubulosus alpinus major , & durior ex cinereo

virescens , caulibus primariis nonnihil compressis. *Mich.*
gen. 80. n. 9.

Habitat in Europæ ericetis.

85. LICHEN *subulatus*. L. fruticulosus subdichotomus ,
ramis simplicibus subulatis. *fl. suec. gallob. Weis. crypt.*
pal. burg. lithu.

L. caule erecto dichotomo , ramis subulatis. *fl. lapp.*

L. caule fruticoso erecto , ramis adscendentibus dichotomis.
Hall. helu. n. 1955.

L. fronde subulata subnuda ramosiuscula dichotoma ; ramis
subulatis tuberculiferis. *Neck. meth.*

Coralloides corniculis longioribus & rarioribus. *Dill. muse.*
102. t. 16. f. 26.

Musculus corniculatus. *Tabern. ic. 809. Bauh. hist. 3. p. 767.*

Habitat in Europæ sylvis , ericetis.

86. LICHEN *globiferus*. L. fruticulosus solidus lævis , ru-
berculis globosis cavis terminalibus.

L. (*globosus*) fruticulosus solidus ramosissimus , ramulis
teretibus acutis furcatis , tuberculis terminalibus globosis.
Huds. angl.

L. fronde terete ramosa ; tuberculis globosis cavis termina-
libus. *Neck. meth.*

L. fruticulosus coralloides non tubulosus cinereus ramosissi-
mus , receptaculis sphaericis concoloribus. *Mich. gen. 105.*
t. 39. f. 6.

Coralloides cupressiforme , capitulis globosis. *Dill. muse.*
117. t. 17. f. 35.

Habitat in Tingitana , Anglia , Stenbrohult Smolandia , &c.

*Similis L. paschali , sed lævior , aphyllus , ramis ter-
minatis globis concavis ore parvo : sphaericæ hiantibus
intus nigris.*

Pro varietate L. fragilis habet cl. Weber. spicil. p. 206.

& *Neck. l. c.*

87. LICHEN *paschalis*. L. fruticulosus solidus tectus fo-
liolis crustaceis. *Fl. suec. Weis. crypt. herb. pal. Fl. dan.*
t. 151. lithu.

L. caule ramoso solido foliolis crustaceis tecto. *Fl. lapp.*

L. caule fruticoso , fungifero , tecto verrucis calcareis. *Hall.*
helu. n. 1963.

L. fronde ramosa , crustaceis foliis tecta : ramulis tuber-
culatis. *Neck. meth.*

L. alpinus glaucus ramosus borryoides. *Scheuch. alp. 137.*
t. 19. f. 4. *Mich. gen. t. 53. f. 5. 8.*

694 CRYPTOGAMIA. ALGÆ.

Coralloides crispum & botryoïdes alpinum. Dill. musc. 114. t. 17. f. 33.

Corallina alpina valde crispa. Pet. musc. t. 65. f. 7.

Musco-fungus coralloides terrestris dense ramificatus cinereus & veluti incrustatus norvegicus. Moris. hist. 3. p. 633. f. 15. t. 7. f. 12.

Muscus cupressiformis ramosus. Læf. pruss. 168. t. 48.

Habitat in Helvetiæ, Italiæ, Cambriæ, Lapponiæ, Scaniæ, Groenlandiæ, Pennsylvania alpestribus.

88. LICHEN fragilis. L. fruticulosus solidus, ramulis teretibus obtusis. *Fl.*

L. erectus ramosissimus, ramulis teretibus nudis filiformibus obtusis. Fl. lapp. 440. t. 11. f. 4.

L. fronde tereti ramosa: ramis obtusis. Neck. meth. 67. cum L. globifero conjungit.

Coralloides alpinum, corallinæ minoris facie. Dill. musc. 116. t. 17. f. 34.

Habitat in Europæ alpibus alpinisque, per Sueciam in rupibus.

I. Filamentosi.

90. LICHEN plicatus. L. filamentosus pendulus, ramis implexis, scutellis radiatis. *Fl. suec. ged.*

L. ramis filiformibus ramosis pendulis confertis. Fl. lapp.

L. pendulus teres exasperatus, ramis descendens, scutellis radiatis. Hall. helu. n. 1971. sec. Weber. R.

L. fronde pendula ramosissima tereti, implexa: scutellis lateralibus margine radiatis. Neck. meth.

Usnea vulgaris, loris longis implexis. Dill. musc. 56. t. 11. f. 1.

Muscus arboreus, Usnea officinarum. Bauh. pin. 361.

Habitat in Europæ & Americæ borealis sylvis densis umbrisque fagetis.

91. LICHEN barbatus. L. filamentosus pendulus subarticulatus, ramis patentibus. *Fl. suec. ged. lithu.*

L. barbatus, trunco articulo lævi. Hall. helu. n. 1970.

L. fronde pendula subarticulo longissima: verrucis levissime sparsis. Neck. meth. 71.

L. caule tereti coriaceo, ramulis tenuissimis cinereis. Roy. lugdb. 513.

Usnea barbata, loris tenuibus fibrosis. Dill. musc. 63. t. 12. f. 6.

Muscus capillaceus longissimus. Bauh. pin. 361.

Habitat in Europæ & Americæ septentrionalis sylvis fagetis.

92. LICHEN *divaricatus*. L. filamentosus pendulus angulatus intus tomentosus, ramis divaricatis, peltis orbiculatis sessilibus.

L. ramosus, lacunatus, flaccidus. *Hall. helv. n. 1975.*

L. filamentosus pendulus, loris compressis lacunosis glabris. *Scop. carn.*

L. (*mollis*) fronde articulata, pendula intus tomentosa; ramis furcatis compressis; scutellis orbiculatis glabris. *Neck. meth. musc.*

Usnea mollis, ramis longis compressis. *Dill. musc. 62. t. 12. f. 5.*

Habitat in Helvetiæ, Misiæ, cæt. cacuminibus arborum acifoliarum. *D. Schreber.*

94. LICHEN *jubatus*. L. filamentosus pendulus: axillis compressis. *Fl. suec. ged. Weis. crypt. lithu.*

L. ramis filiformibus ramosis pendulis, alis compressis. *Fl. lapp. 456.*

L. barbatus, pendulus, filamentis nigris prolixis. *Hall. helv. 1968.*

L. filamentosus pendulus; tuberculis lateralibus ianatis. *Scop. carn.*

L. fronde pendula ramosa: filamentis compressis teretibusque verrucosis. *Neck. meth.*

L. capillaceus longissimus s. Muscus arboreus nigricans, usnea officinarum. *Mich. gen. 77. n. 7. R.*

Usnea jubata nigricans. *Dill. musc. 64. t. 12. f. 7.*

Muscus arboreus niger & tenuior. *Dod. pempt. 471.*

Habitat in Europæ sylvis & rupibus.

95. LICHEN *lanatus*. L. filamentosus ramosissimus, decumbens implicatus opacus. *Fl. suec. herb.*

L. fronde ramosissima tereti decumbenti - implexa siccitate rigida. *Neck. meth. β. cum L. jubato conjungit.*

Usnea lanæ nigræ instar saxis adhærens. *Dill. musc. 66. t. 13. f. 8.*

Habitat in Europæ frigida rupibus.

96. LICHEN *pubescens*. L. filamentosus ramosissimus decumbens implexus nudus. *Fl. suec.*

L. fruticosus minimus flexuosus, cauliculis nigris capillaceis. *Hall. helv. n. 1967. jec. Weber. R.*

Usnea caespitosa exilis capillacea atra. *Dill. musc. 66. t. 13. f. 9.*

Habitat in Europæ septentrionali, Lapponia, in rupibus humilis, quæ imbres diluunt.

CRYPTOGAMIA. ALGÆ. 697

101. LICHEN *floridus*. L. filamentosus erectus, scutellis radiatis. *Fl. suec.* " *car. ged. Weis. crypt. pal. Kniph. cent.* 6. n. 54. *burg. lithu.*

L. caule ramoso solido, foliis setaceis, receptaculis maximis orbiculatis peltatis ciliato radiatis. *Mort. cliff.*

L. barbatus, ramis teretibus, furculis parallelis simplicibus, scutellis radiatis. *Hall. helu.* n. 1972.

L. fronde erecta ramosa tereti solida apice scutellifera: scutellis ciliatis. *Neck. meth.*

Ufnea vulgarissima tenuior & brevior cum orbiculis. *Dill. musc.* 69. t. 13. f. 13.

Musco - fungus arborum capillaceus, scutellis amplis per ambitum pilis radiatis. *Morif. hist.* 3. p. 635. f. 15. t. 7. f. 14.

Muscus arboreus cum orbiculis. *Bauh. pin.* 361.

Muscus ramosus floridus. *Tabern. ic.* 808.

Habitat in Europæ fagets.

1320. TREMELLA.

Fructificationes vix manifestæ, in corpore gelatinoso.

1. TREMELLA *junipera*. T. sessilis membranacea auriformis fulva. *Fl. suec.*

Fucus Tremella juniperina. *Gmel. fuc. R.*

Agaricum gelatinosum, flavo rubrum, superne tuberculosum. *Hall. helu.* n. 2260.

Byssus gelatinosa fugax, junipero innascens. *Fl. lapp.*

Habitat in Juniperetis primo vere.

Membranacea, gelatinosa, lutescente - virens. Siccata fragilis, nigricans.

2. TREMELLA *Nostoc*. T. plicata undulata. *Fl. suec. carn. gallob. Weis. crypt. herb. pal. burg.*

T. terrestris sinuosa pinguis & fugax. *Dill. musc.* 52. t. 10. f. 14.

Fucus tremella *Nostoc*. *Gmel. fuc. R.*

Lichen gelatinosus, plicatus, undulatus, laciniis crispis, granulosis. *Hall. helu.* n. 2041.

Byssus gelatinosa fugax terrestris. *Fl. lapp.* 530.

Linkia terrestris gelatinosa membranacea vulgarissima. *Mich. gen.* 126. t. 67. f. 1.

Nostoc paracelsi. Ad. parif. 17. p. 228.

698 CRYPTOGAMIA. ALGÆ.

β. Tremella arborea nigricans, minus pinguis & fugax.
Dill. musc. 54. t. 10. f. 15. *Leers l. c.* R.

Habitat in pratis post pluvias.

3. TREMELLA *lichenoides*. T. erecta plana: margine crispo lacinulato.

T. frondibus erectis planis: margine crispo lacinulato. *Fl. suec. gallob. pal.*

Lichen tremelloides. *Weis. crypt. herb.*

Lichen atrocoeruleus laciniatus & ciliatus. *Hall. helu. n.* 2032.

Lichen foliaceus gelatinosus, foliorum segmentis capillaceis. *Scop. carn.*

Lichen (*tremelloides*) fronde erecta marginibus lacinulata, crispa; scutellis subrotundis. *Neck. meth.* 87.

Lichenoides pellucidum, endiviæ foliis tenuibus crispis. *Dill. musc.* 143. t. 19. f. 31. 32. 33. 34. 35.

Lichen terrestris membranaceus mollior fuscus. *Mich. gen.* 26. t. 38.

Musco fungus terrestris minor fuscus, foliis e latitudine crenatis musco innascens. *Moris. hist.* 3. p. 632. f. 15. t. 7. f. 4.

Habitat in Muscis, locis umbrosis ad montes.

4. TREMELLA *verrucosa*. T. tuberculosa solida rugosa. *Fl. suec.*

T. lobata & rugosa. *Hall. helu. n.* 2127.

Fucus Tremella verrucosa. *Gmel. fuc.* R.

Tremella subrotunda sinuosa difformis gelatinosa. *Is. w. goth.* 139.

T. fluviatilis gelatinosa & uterculosa. *Dill. musc.* 54. t. 10. f. 16.

Linkia palustris gelatinosa saxi adnascens ex obscuro fulva & concava vesicam referens. *Mich. gen. n.* 26. t. 67. f. 2.

Habitat supra Lapides in rivulis.

5. TREMELLA *difformis*. T. subrotunda sinuata difformis gelatinosa. *Fl. suec.*

Habitat in Oceani Conservis.

6. TREMELLA *hemisphærica*. T. hemisphærica sparsa. *Fl. suec.*

Fucus Tremella hemisphærica. *Gmel. fuc.* 225. R.

Habitat in Rupibus & Conservis submarinis.

7. TREMELLA *purpurea*. T. subglobosa sessilis solitaria glabra. *Fl. suec. pal.*

Sphæria (*tremelloides*) composita solitaria, subpeduncu-

lata subglobosa purpurea subgelatinosa. Weis, obs. 46. n. 8. t. 33. f. 1.

Evela purpurea. Schreb. spicil.

Sphæria convexa minora hirsuta. Hall. helv. n. 2159.

Valsa miniata. Scop. carn.

Lichenoides tuberculosum amœne purpureum. Dill. musc. 127. t. 18. f. 6.

Habitat in Arborum ramis moribundis & emortuis,

An Agarici initium? minime. Weig. l. c. R.

8. **TREMELLA adnata**, T. rotunda imbricata livida. *Fl. succ.*

Habitat in Oceani rupibus submarinis.

1321. FUCUS.

Masc. Vesiculæ villis intertextæ.

Fem. Vesiculæ adpersæ granis immerlis apice prominulis. Sem. Solitaria.

1. **FUCUS natans** F. caule filiformi ramoso, foliis lanceolatis ferratis, fructificationibus globosis pedunculatis. *Syst. veg. 811.*

F. caule tereti ramossissimo, foliis lanceolato - ferratis, fructificationibus globosis pedunculatis subaristatis. *Sp. pl. 2. p. 1628. n. 13.*

F. folliculaceus, ferrato folio. *Bauh. pin. 365.*

Lenticula marina, ferratis foliis. Daleck. hist. 1397.

Sargazo. Pis. bras. 2. p. 266.

Habitat in Pelago libere natans, nec radicatus.

Fructificationes in aliis muticæ, in aliis filo brevi aristatæ; in quibusdam individuis ex alis foliorum ræsemi brevissimi filiculis minutis verrucosis.

Non libere natat. Gmel. l. c. R.

2. **FUCUS acinarius**. F. caule filiformi ramoso, foliis linearibus integerrimis, fructificationibus globosis pedunculatis. *Gmel. fuc. 99.*

F. caule tereti ramoso, foliis linearibus integerrimis, fructificationibus globosis pedunculatis. *Sp. pl.*

F. folliculaceus, linearæ folio. *Bauh. pin. 365.*

Lenticula marina serapionis. Lob. ic. 2. p. 256.

*Aciparia imperati. Donat marin. 35. t. 4. f. 1. **

Habitat in Italia & Oceano australiori.

Precedenti affinis. Cartilagineus, rubens, filiformis, compressus. Fructificationes ad ramorum summorum mar-

gines, globosæ, pedunculatæ, magnitudine seminis Brassicæ; sæpe ad basin foliolo subulato.

6. FUCUS *serratus* F. fronde plana dichotoma costata serrato-dentata, fructificationibus terminalibus tuberculosis. F. fronde plana dichotoma serrata ad apices tuberculata. *succ. Gmel. fuc.*

F. folio dichotomo plano serrato laciniato. *Lepp.*

F. fronde oblonga, plana, dentata, utrinque pilifera: tuberculis terminalibus. *Neck. meth. 27. R.*

Fucus s. Alga latifolia major dentata. *Morif. hist. 3. p. 648. f. 15. t. 9. f. 1. Reaum. act. gall. 1711. p. 284. a. 9. f. 1.*

Habitat in Oceano.

Frons costata.

7. FUCUS *volubilis*. F. fronde plana spirali perfoliata repando dentata. *Gmel. fuc.*

F. fronde spirali-convoluta-marginæ dentata. *Neck. meth.*

F. palica spiralis minor. *Ginan. adriat. 1. p. 26. t. 27.*

Alga spiralis maritima. *Bocc. sic. 70. t. 38. f. 2.*

Habitat in Mari Mediterraneo.

8. FUCUS *vesiculosus*. F. fronde plana dichotoma costata integerrima: vesiculis axillaribus geminis; terminalibus tuberculatis.

F. fronde dichotoma integra, caule medium folium transcurrente, vesiculis verrucosis terminalibus. *Spec. pl. succ. stamp. carn. Gmel. fuc.*

F. fronde plana integerrima vesiculis terminalibus verrucosis, primordialibus linearibus. *Neck. meth. 18.*

F. marinus s. Quercus maritima vesiculas habens. *Bauh. pin. 365. Reaum. act. gall. 1711. p. 288. f. 2.*

Quercus marina. *Lob. ic. p. 255.*

β. Fucus maritimus s. Quercus maritima foliorum extremis tumidis. *Bauh. pin. 365.*

Habitat in Mari atlantico.

9. FUCUS *divaricatus*. F. fronde plana dichotoma integerrima, axillis divaricatis: vesiculis axillaribus geminis.

F. bullatus fruticescens, caule nudo, foliis rectis compressis bifidis. *Morif. hist. 3. p. 647. f. 15. t. 8. f. 5.*

Habitat in Anglia, Lusitania.

Præcedenti tenuior. Vesiculæ axillares geminæ, passimque solitariae.

Cum E. spirali conjungit Gmelin. R.

10. FUCUS *inflatus*. F. fronde plana dichotoma integerrima punctata ovato-lanceolata inflata: apice diviso,

F. fronde bifida : laciniis ovato-lanceolatis inflatis apice divis. *Spec. pl. suec.*

Habitat in Oceano Atlantico.

Semel aut bis tantum bifidus est.

11. **FUCUS ceranoides.** **F.** fronde plana dichotoma integerrima punctata lanceolata ; fructificationibus tuberculatis bifidis terminalibus.

F. fronde dichotoma plana integra : apicibus bifidis vesiculos. *suec. carn.*

F. fronde plana laciniato-multifida , dichotoma : globulis atris lateralibus. *Neck. meth.*

F. humilis dichotomus ceranoides , latioribus , foliis ut plurimum verrucosis. *Moris. hist. 3. p. 646. f. 15. t. 8. f. 13.*

β. Fucus lacerus. *Spec. ed.*

F. membranaceus ceranoides varie dissectus. *Rai. angl. 3. p. 44.*

Habitat in Oceano.

Præcedenti affinis , sed angustior magisque ramosus , utrinque punctis impressis adpersus.

Fucus β. hujus varietas est secundum Gmelin.

12. **FUCUS spiralis.** **F.** fronde plana dichotoma integerrima punctata : inferne lineari-canaliculata , fructificationibus tuberculatis geminis. *Flor. dan. t. 286.*

F. fronde dichotoma integra , caule folium percurrente : inferne nudo , vesiculis terminalibus. *suec.*

F. spiralis maritimus major. *Rai. angl. 3. p. 41.*

Habitat in Oceano.

Frons membranacea , plana , sed inferne angustior , hinc canaliculata. Fructificationes terminales , geminæ , pedunculatæ , oblongæ , crassiusculæ. Dum crescit in mari , contortus est in spiram.

Cum F. vesiculo conjungit Gmel. fuc. 62.

13. **FUCUS canaliculatus.** **F.** fronde plana dichotoma integerrima canaliculata lineari , fructificationibus tuberculatis bipartitis obtusis.

F. fronde ramosa hinc canaliculata , vesiculifera : apicibus obtusis bifidis rugosis. *Neck. meth.*

β. Fucus (excisus) fronde lineari dichotoma canaliculata , axillis divaricatis punctata. *angl.*

F. humilis dichotomus : segmentis ex una parte gibbosis ex altera excavatis. *Rai. angl. 3. p. 43.*

F. dichotomus membranaceus ex viridi flavescens ceranoides.

702 CRYPTOGAMIA. ALGÆ.

angulos rotundiusculos efformans. *Morif. hist. 3. p. 646. f. 15. t. 8. f. 11.*

Habuat in Oceano Europæo.

Dichotomus multoties, lævis, angustus & linearis, latere altero convexo; altero canaliculato. Vesiculæ terminales, bipartitæ s. geminæ, sessiles, tuberculis adpersæ perforatis.

14 FUCUS *distichus*. F. fronde plana dichotoma integerrima lineari: fructificationibus tuberculatis mucronatis.

F. *linearis* dichotomus planus linearis acutus: vesiculis ovatis sparsis. *angl. Oed. dan. 351.*

F. filiformis. *Gmel. fuc. 72. t. 1. f. 1.*

Habitat in Oceano septentrionali.

Differt a præcedente, quod Frons minime altero latere canaliculata, quodque. Vesiculæ, siliquæ instar, longæ & acuminatæ. Kœnig. Frons nervo medio, textura herbacea.

15. FUCUS *nodosus*. F. fronde compressa dichotoma; foliis distichis integerrimis, vesiculis innatis solitariis dilatatis. *Oed. dan. 146. Gmel. fuc. 78. t. 1. B. f. 1. Baster. opusc. 6. p. 121. t. 11. f. 5.*

F. fronde compressa, vesiculis oblongis interjecta: frondulis integerrimis. *Neck. meth. 17.*

F. caule compresso dichotomo medio ramorum in vesiculam dilatato. *Juec.*

F. caule tereti compresso dichotomo, vesicula medio ramorum innata, vesiculis axillaribus laxis. *Hort. cliff. 479:*

F. caule tereti compresso dichotomo, foliis oppositis minimis, vesicula in medio singuli rami. *Fl. lapp. Guett. stamp.*

F. maritimus, vesiculis majoribus singularibus per intervalla dispositis. *Morif. hist. 3. p. 647. f. 15. t. 8. f. 2.*

F. maritimus nodosus. *Dod. pempt. 781.*

Habitat in Mari Atlantico.

Folia lanceolata, obtusa, angulis frondis distiche insidentia. Vesiculæ acutæ, medio ramorum innatæ, ramis latiores.

17. FUCUS *siliquosus*. F. fronde compressa ramosa, foliis distichis alternis integerrimis, fructificationibus peduncularis oblongis mucronatis. *Oed. dan. t. 106. Gmel. fuc. 81. t. 2. B.*

F. caule tereti ramosissimo, pedunculis alternis vesiculis oblongis acuminatis. *Juec. stamp.*

F. fronde ramosa vesiculis oblongis siliquiformibus mucronatis, primordialibus lineari-subulatis. *Neck. meth. p. 18.*

F. maritimus alter, tuberculis paucissimis. *Bauh. pin.* 365.

F. marinus quartus. *Dod. pempt.* 480.

Habitat in Oceano.

Frons filiformis, compressa, flexuosa, utrinque angulis dentata rudimentis petiolorum aut fructificationum. Fructificationes fusiformes, alternæ, rostratæ, pedunculatæ. Folia lanceolata, fructificationibus minora.

19. *FUCUS elongatus*. *F.* fronde filiformi compressa dichotoma articulata geniculis tumidiusculis. *Gmel. fuc.* 103.

F. filiformis teretiusculis subdichotomus, geniculis tumidiusculis. *Huds. angl.* 470.

F. caule lineari aphylo dichotomo. *Roy. lugdb.* 514.

Habitat inter Angliam & Hispaniam.

Frons ad dichotomias tumidiuscula & articulata, ubi per exsiccationem discedit, ut articuli sæpe solvantur.

20. *FUCUS loreus*. *F.* fronde filiformi compressa dichotoma undique utrinque tuberculata. *Fl. dan. t.* 710.

Habitat in

Frons dichotoma elongata, linearis, compressa, adspersa undique tuberculis elevatis obsoletis.

21. *FUCUS fœniculaceus*. *F.* fronde filiformi ramossissima, vesiculis ovatis terminatis foliolis multipartitis obtusis apice fructificantibus. *Gmel. fuc.* 86. t. 2. a. f. 3.

F. folliculaceus, fœniculi foliis brevioribus. *Bauh. pin.* 365.

β. *F.* (*barbatus*) vesiculis oblongis pedunculatis: terminatis foliolis linearibus. *Spec. pl.* 2.

F. maritimus. foliis tumidis barbatis. *Bauh. pin.* 365.

Quercus maritima barbata. Bauh. prodr. 154.

Habitat in Oceano.

Statura Lichenis Usneæ; Ramulis copiosissimis, capillaribus terminatis vesicula, sæpius prolifera, ultima terminata aliquot foliolis apice tuberculatis.

Fucum barbatum varietatem facit F. vesiculosi. Gmel. fuc. p. 62. R.

24. *FUCUS selaginoides*. *F.* fronde filiformi ramossissima: ramis dichotomis, foliis subulatis alternis basi vesiculosis.

F. *Abies marina. Gmel. fuc.* 83. t. 2. a. f. 2.

F. folliculaceus, foliis abrotani. *Bauh. pin.* 365.

F. crispus loricatus nigricans. *Barrel. ic.* 1290. *Bocc. musc.* 7. t. 12.

Habitat in Oceano norwegico.

Caule filiformi, flexuosus, ramossissimus. Rami imbricati. Folia subulata, brevissima, basi vesiculoso-ovata.

704 CRYPTOGAMIA. ALGÆ.

25. FUCUS *concatenatus*. F. fronde filiformi ramosissima: ramulis dichotomis, vesiculis moniliformibus distantibus innatis, foliis subulatis.

F. caule tereti ramosissimo, fructificationibus oblongis concatenatis. Roy. luglb. 514.

Habitat in Oceano.

Fucus filiformis, opposite vel alternatim ramosissimus, ultimis ramulis sæpe dichotomis. Vesiculæ ovatæ, cauli ramulisque innatæ, remotæ, cum folio subulato ad latus cujusvis vesiculæ.

26. FUCUS *aculeatus*. F. fronde filiformi compressa ramosissima: dentibus marginalibus subulatis alternis erectis. Oed. dan. 355. Pal. ii. 3. p. 34.

F. muscoides, fronde filiformi compressa: denticulis subulatis alternis. Neck. meth. p. 31.

F. angustifolius, foliis dentatis. Rai. angl. 3. p. 48.

F. tenuifolius, foliis dentatis. Moris. hist. 3. p. 648. f. 15. t. 9. f. 4.

β. Fucus (*muscoides*) fronde tereti ramosissima, ramis sparsis spinis mollibus alternis. Spec. ed. 2.

L. muscoides. Gmel. fuc. 130. t. 12. f. 1.

F. (*virgatus*) compressus, ramosissimus, foliis alternis, subulatis, adspersis spinis alternis mollibus. Gunn. norv. 93. Ad. nidros. 4. p. 83. t. 7.

F. foliis ericæ similis. Rai. hist. 73. n. 4. b.

Habitat in Oceano norvegico.

Refert caudam equinam.

27. FUCUS *lycopodioides*. F. fronde filiformi tereti subramosa, obtecta undique setis.

Conserva squarrosa, subramosa, undique obtecta setis imbricatis squarrosis. Oed. dan. t. 357.

Habitat.

Frons erecta, parum ramosa, vix pedalis undique more Lycopodii tecta foliolis setaceis, semipollicaribus.

28. FUCUS *hirsutus*. F. fronde filiformi tereti dichotoma obtecta undique hirsutie.

F. teres ramosus, ramulis imbricatis, lacinulis secundis incurvis. Hudf. angl.

Habitat in Pelago Gunnerus.

Facies fere spongiæ. Caules pedales, crassitie fili grossi, ramosissimi; ramis obtusis, teretibus, undique vestitis vellere densissimo, brevissimo, quasi cervino. Hirsuties vestiens tam

lam parva, us oculis non distinguatur. Proxime accedit ad Fucum ericoidem.

31. FUCUS *Filum*. F. fronde filiformi subfragili opaca.

F. filiformis simplex subfragilis opacus. *Amœn. acad.* 4.

259. t. 3. f. 2.

F. filiformis simplex. *Fl. suec. stamp.*

Fucus s. Filum maritimum germanicum. *Bosc. musc.* 1. p.

271. t. 9. f. 9.

Alga nigro capillaceo folio. *Bauh. pin.* 364.

Filum maritimum germanicum. *Bauh. pin.* 355.

Habitat in Oceano Atlantico.

32. FUCUS *lanosus*. F. frondibus dichotomis ramosissimis scabris.

Habitat in Oceano Islandico. *Kœnig.*

Refert lanam atram; palmariis, oculo armato scabē punctis subverticillatis.

33. FUCUS *fastigiatus*. F. fronde filiformi dichotoma ramosissima fastigiata obtusa. *Oed. dan.* 393.

F. dichotomus ramosissimus teres uniformis fastigiatus. *suec. Gmel. fuc.* 106. t. 6. f. 1.

F. fronde filiformi ramosissima dichotoma: vesiculis terminalibus, primordialibus lateralibus. *Neck. meth.*

F. caule lineari dichotomo. *Roy. lugdb.*

F. palmaris tenuis in orbem expansus, in segmenta bifida breviora teretia divisus. *Morif. hist.* 3. p. 649. f. 15. t. 9. f. 9.

F. marinus polyschides. *Læs. pruss.* 77. t. 15.

Habitat in Oceano Baltico.

34. FUCUS *furcellatus*. F. fronde filiformi dichotoma ramosissima acuminata. *Oed. dan.* 499.

F. *caprinus*, fronde dichotoma, teretiuscula, filiformi fastigiata. *Gunn. Act. nidros.* 4. p. 82. t. 5. f. 4.

F. lumbricalis. *Gmel. fuc.* 108. t. 6. f. 2. angl.

F. fronde tereti lineari-subulata, apice dichotoma inflata. *Neck. meth.*

F. parvus, segmentis prælongis teretibus acutis. *Morif. hist.* 3. p. 648. f. 15. t. 9. f. 4.

Habitat in Oceano Anglico.

F. fastigiato proxime accedit; sed altior, ramis crassioribus apice acuminatis. Color sæpius ruber.

Color constans purpureus. *Gunn. l. c. R.*

35. FUCUS *palmatus*. F. fronde palmata plana. *Knipf. cent.* 1. n. 30.

Linnaei Spec. plant. Europ. Pars. II,

Y y

706 CRYPTOGAMIA. ALGÆ.

F. (ovinus) purpureus, caule brevissimo, foliis inæqualibus profunde fissis, laciniis inferne angustatis, apice hinc fissis illine dentatis. *Ad. nidros.* 4. t. 9.

F. (dulcis) fronde palmata margine undata: primordialibus marginalibus. *Neck. meth. Gmel. fuc.* 189. t. 26. R.

F. membranaceus ceranoides. *Bauh. prodr.* 155.

F. foliaceus humilis palmam humanam referens. *Moris. hist.* 3. p. 646. f. 15. t. 8. f. 1.

Habitat in Oceano.

37. *FUCUS digitatus*. *F.* fronde palmata, foliolis ensiformibus, stirpe tereti. *angl. Oed. dan.* 392.

F. (hyperboreus) fronde simplici palmata, caule longissimo. *Gunn. flor.* 61. t. 3.

F. palmatus. *Gmel. fuc.* 202. t. 30.

F. (polyschides) fronde spiraliter convoluta apice palmata: primordialibus ovato-lanceolatis simplicissimis. *Neck. meth. musc.* p. 38.

F. scoparius. *Strom. sond.* 93. *

F. arboreus polyschides edulis. *Bauh. pin.* 64.

F. membranaceus polyphyllus major. *Rai. angl.* 46.

Habitat in Oceano Atlantico.

Caulis crassitie baculi ambulatorii.

38. *FUCUS esculentus*. *F.* fronde simplici indivisa ensiformi, stirpe tetragona pinnata folium percurrente. *Oed. dan.* 417. *Gmel. fuc.* 200. t. 29. f. 1.

Fucus fimbriatus. *Gmel. fuc.* 200. t. 29. f. 1.

F. p. ctus, caule medium folium terminale planum percurrente, inferne vesicis lineari-lanceolatis petiolatis pinnato. *Gunn. norv. n.* 313. *Ad. nidros.* 4. t. 8. f. 1.

F. fronde indivisa basi pinnulifera: rachi petioliformi quadrangulari. *Neck. meth. musc.*

F. folio plano ensiformi, caule pinnato. *Flor. lapp.*

F. longissimo latissimo tenuique folio. *Bauh. pin.* 364. *prodr.* 154.

Habitat in Mari Atlantico. Equis & hominibus esculentus. D. Gunnerus. Kænig.

Frons maxima per quam longitudinaliter ab utraque pagina transit. Petiolus crassus, quadrangularis, latus, cui infra folium utrinque insident. Pinnulæ ensiformes, planæ. Differt a F. saccharino nervo folii & pinnulis. Mant. 135.

39. *FUCUS saccharinus*. *F.* fronde subsimplici ensiformi, stirpe tereti brevissima. *Gmel. fuc.* 194. t. 27. *Oed. dan.* 416.

F. caule tereti brevissimo, folio maximo ensiformi subsimplici. suec. angl.

F. fronde indivisa simplicissima lanceolata plana: marginibus plicato-sinuatis. Neck. meth.

F. longissimo latissimo crassoque folio. Bauh. prodr. 154. Rai. hist. 74. E. N. C. 1748. p. 450. t. 9. f. 1. 2.

F. alatus f. phasnagoides. Bauh. pin. 364.

Habitat in Mare Atlantico.

40. *FUCUS sanguineus. F. frondibus membranaceis ovato-oblongis integerrimis petiolatis, caule tereti ramoso. Hudsf. angl. Oed. dan. 349. Gmel. fuc. 185. t. 24. f. 2. Giseke. ic. fasc. 1. t. 24.*

F. fronde ramosa undato-oblonga, integerrima dilatata: primordialibus racemosis. Neck. meth.

Fucus f. Alga folio membranaceo purpureo lapathi sanguineæ figura & magnitudine. Moris. hist. 3. p. 645. f. 15. t. 8. f. 6. Rai. angl. 47.

Habitat in Oceano atlantico. Kænig.

41. *FUCUS ciliatus. F. frondibus membranaceis lanceolatis proliferis ciliatis. angl. Oed. dan. t. 353.*

F. fronde plana laciniata margine ciliata, primordialibus linearibus. Neck. meth.

F. caulescens. Gmel. fuc. 173. t. 20. f. 2.

F. membranaceus rubens angustifolius, marginibus ligulis armatis. Rai. angl. 47.

F. humilis membranaceus acaules elegantissimus ruber capillis longis fimbriatus. Moris. hist. 3. p. 646.

β. *F. membranaceus purpureus latifolius pinnatus. Rai. angl. 47.*

Habitat in Oceano atlantico.

42. *FUCUS crispus. F. frondibus submembranaceis dichotomis: laciniis dilatatis crispis.*

Fucus f. Alga membranacea candida: segmentis plurimum laciniatis. Moris. hist. 3. p. 649. f. 15. t. 8. f. 6.

Habitat in Oceano atlantico. Kænig.

Laciniæ purpurascentes aut albidæ, superne latiores, apice obtuse dentatæ.

43. *FUCUS alatus. F. frondibus membranaceis subdichotomis costatis: laciniis alternis decurrentibus bifidis. angl. Oed. t. 352.*

F. fronde ramosissima nervosa: laciniis linearibus decurrentibus. Neck. meth.

F. dichotomus parvus costatus membranaceus. Rai. angl. 44.

F. purpureus tenuiter divisus non geniculatus. *Morif. hist. 3. p. 646.*

Habitat in Oceano septentrionali. Kænig.

Frondes *ramosi*, *purpurei*, *diaphani*, *lineares*, *nervo medio crassiore*, *apicibus subdentatis*.

45. *FUCUS dentatus*. *F.* frondibus membranaceis caerviis alternatim pinnatifidis : sinibus obtusis, laciniis apice erosis. *Mant. 135.*

F. pinnatifidus. angl. Oed. t. 354. sine dubio huc referendus.

F. (atomarius) fronde alternatim pinnatifida ; laciniarum denticulis erectis. *Neck. meth.*

F. atomarius. Gmel. fuc. 125. t. 10. f. 1:

F. membranaceus rubens, foliis latiusculis ad extrema dentatis. *Morif. hist. 3. p. 646. f. 15. t. 8. f. 5.*

Habitat in Oceano atlantico.

Frons *rubra*, *diaphana*, *laciniarum sinibus obtusis* : *laciniis apice dentatis*.

48. *FUCUS vittatus*. *F.* frondibus membranaceis divisus enfiformibus dentato crispatis. *Oed. dan. t. 353.*

β. *Fucus (ovatus)* fronde plana oblonga prolifera ciliata, ramentis foliaceis confertissimis. *Mant. 2. p. 312.*

Habitat β. *ad Cap. b. spei.*

Frons *var. β. lineares* s. *oblongæ* s. *lanceolatæ*, *sanguineæ* *aveniæ*, *apice proliferæ* frondibus & sic *uterius*. Margo frondium *densissime ciliatus* ciliis *concoloribus densissime congestis*, *hæc cilia ex foliolis minutissimis laceris* s. *incisis*.

49. *FUCUS ramentaceus*. *F.* frondibus filiformibus simplicibus, hinc ramentis foliaceis confertis.

Ulva sobolifera, coriacea, simplex, tubulosa, undique appendiculata. *Oed. fl. dan. t. 356.*

Habitat in Oceano septentrionali.

Striptes *palmares*, *simplices*, *ab altero tantum latere ramentis foliaceis, confertis, linearibus, coloratis*. Stirps *tubulosa*.

50. *FUCUS plumosus*. *F.* frondibus cartilagineis lanceolatis bipinnatis plumosis, caule filiformi compresso ramoso. *angl. Oed. dan. t. 350.*

F. fronde ramosa cartilaginea : laciniis pinnatifidis, superioribus terminalibus tubercularis. *Neck. meth.*

Fucoides purpureum eleganter plumosum. *Rai. angl. 37.*

Habitat in Oceano atlantico. Kænig.

Similis sequenti, sed Hypnī instar pinnatus; parvus.
Mant. 134.

56. FUCUS *confervoides*. F. frondibus membranaceis linearibus compressis ramosis, fructificationibus sparsis sessilibus subrotundis. *Kænig.*

F. (*verrucosus*) fronde tereti filiformi: ramis longis, subulatis, globuliferis. *Neck. meth. p. 28. n. 20. R.*

F. verrucosus. *Gmel. fuc. 136. t. 14. f. 1.*

F. filiformis teres ramosissimus, ramis simplicibus, vesiculis rotundis lateralibus. *Huds. angl. 470.*

F. teres rubens ramosissimus. *Rai. angl. 3. p. 51.*

Habitat in Mari anglico.

57. FUCUS *ericoides*. F. filiformis ramosissimus hirtus.

F. fronde nodosa, ramulis imbricatis globuliferis, denticulis subularis. *Neck. meth. 28.*

Erica marina. Gmel. fuc. 128. t. 11. f. 2. Gort. belg. 340. Huds. angl. 469.

Fucus foliis ericæ s. tamarisci. Rai. angl. 3. p. 49.

Tamarisco similis maritima. Bauh. pin. 365.

Erica marina quibusdam. Bauh. hist. 3. p. 799.

Habitat in Oceano Europæo.

Omissus in Syst. vegetabilium, forsan sub alia specie latens. R.

1322. ULVA.

Fructificationes in membrana diaphana.

1. ULVA *pavonia*. U. plana reniformis sessilis decussatim striata. *Syst. Veg. 817.*

Fucus (pavonius) fronde sessili reniformi decussatim striata. Spec. ed.

Fucus fronde plana reniformi decussatim striata: primordiis simplicissimis. Neck. meth. 40.

Fucus maritimus gallopavonis pennas referens. Bauh. prodr. 155. Moris. hist. 3. p. 645. f. 15. t. 8. f. 7. Ellis. cor. 88. t. 33. f. C.

Alga marina gallopavonius plumas referens. Bauh. pin. 364.

Fungus articulatus. Bauh. pin. 368.

Habitat in Mari Europæ australis.

2. ULVA *umbilicalis*. U. plana orbicularis sessilis peltata coriacea. *Syst. Veg. 817.*

710 CRYPTOGRAMIA. ALGÆ.

U. fronde dilatata subsinuata , centro radicata. *Spec. pl.*

Fucus fronde concaviuscula , peltata , coriacea. *Neck. meth.* 39.

Fucus Tremella umbilicalis. *Gmel. fuc.* 219.

Tremella marina umbilicata. *Dill. musc.* 45. t. 8. f. 3.

Fucus umbilicus marinus dictus. *Bauh. pin.* 364.

Habitat in Oceano.

3. *ULVA intestinalis.* *U. tubulosa simplex.* *Fl. suec. gallob. Weis. crypt.* 27. burg.

U. tubulis cylindricis , tuberosis. *Hall. helu. n.* 2128.

Conferva tubulosa simplex : primordialibus filamentosis. *De Neck. meth.* 12.

Tremella marina tubulosa intestinorum figura. *Dill. musc.* 47. t. 9. f. 7.

Fucus cavus. *Bauh. pin.* 364.

Habitat in Mari omni.

5. *ULVA compressa.* *U. tubulosa ramosa compressa.* *suec. carn. gallob.*

Conferva simplex ramosaque. *Neck. meth.*

Tremella marina tenuissima & compressa. *Dill. musc.* 48. t. 9. f. 8.

Habitat in Europæ mari & tectis maritimis.

7. *ULVA confervoides.* *U. filiformis articulata : articulis alternatim compressis.*

Conferva marina fistulosa. *Dill. musc.* 34. t. 6. f. 39.

Habitat in Mari.

8. *ULVA latissima.* *U. oblonga plana undulata membranacea viridis.* *suec. carn.*

Alga saccharifera. *Strom. sond.* 92.

Fucus longissimo latissimo tenuique folio. *Bauh. pin.* 364. prodr. 154.

Habitat in Mari Europæ.

Frons magnitudine folii Musæ.

9. *ULVA Lactuca.* *U. palmata prolifera membranacea : ramentis inferne angustatis* *suec.*

Fucus Tremella Lactuca. *Gmel. fuc.* 215.

Fucus (Lactuca) fronde imbricata : frondibus partialibus erectiusculis , sinuato laciniatis , obtusis , *Neck. meth.* p. 39.

Ulva marina lactucæ similis. *Raj. angl.* 3. p. 62.

Tremella marina vulgaris , lactucæ similis. *Dill. musc.* t. 8. f. 1.

Muscus marinus lactucæ similis. *Bauh. pin.* 364.

Habitat in Oceano.

Fronde*s aggregatæ , membranacæ , pallidæ , palmatæ , segmentis singulis iterum in frondes palmatas enatis , undatis obovatis , obtusis , pellucidis.*

11. ULVA *lanceolata*. U. lanceolata plana.

Tremella marina porrifolio. *Dill. musc. 46. t. 9. f. 5.*

Habitat in Oceano.

12. ULVA *labyrinthiformis*. U. cellulis labyrinthiformibus prominentiisque clavatis.

U. thermalis valvulosa erecta simplex, capitulo subrotundo.

Vandell. therm. 120. t. 2.

Habitat in Thermis Patavinis gradu coloris 49 , pulchræ ari incisa cum descriptione a Vandellio.

Membranæ binæ, horizontales, distantes superiore tenuiore, intertexta parietibus numerosis, variisque formantibus cellulas labyrinthiformes. E. superficie exsurgunt Mucrones, multiplici basi, trunci clavati, obtusi, ex 7. membranis conflati, intus cavi. In membranis & cavitatibus dispersa sunt Grana plurima, concava.

13. ULVA *Linza*. U. fronde oblonga bullata. *succ.*

Tremella marina fasciata. *Dill. musc. 46. t. 9. f. 6.*

Fucus Linza dictus, lactucæ marinæ similis. *Bauh. hist. 3. p. 81.*

Muscus lactucæ marinæ similis. *Bauh. pin. 364.*

Habitat in Oceano.

14. ULVA *pruniformis*. U. subglobosa solitaria, intus succulenta. *succ. Gunn. norv. n. 737. t. 2. f. 6. 7. & it. 9. f. 4. 6.*

Tremella pruniformis. *Weber. spicil. 279.*

Conferva pollinea globosa gelatinosa. *Hall. helv. n. 2110.*

Fucus globosus sessilis mollis. *Gleditsch. monogr.*

Habitat in lacubus Sueciæ , Borussia.

15. ULVA *granulata*. U. sphærica aggregata. *succ. gallob. herb. pal. Fl. dan. t. 705. * Burg. lith.*

Tremella (globosa) sphærica, vesiculis aggregatis. *Veis. crypt. 28.*

Tremella palustris, vesiculis sphæricis fungiformibus. *Dill. musc. 55. t. 10. f. 17.*

Habitat in Europa ad ripas fluviorum.

16. ULVA *Pisum*. V. (granulata) globosa sarcta viridis.

Conferva (Pisum) sphærica filamentis concentricis. *Flor. dan. t. 660. f. 2.*

Habitat ad ripas paludum fontiumque Europæ.

712 CRYPTOGAMIA. ALGÆ.

Rejicitur ad littora paludum natans viridis, magnitudinis seminis cannabis, copiosissima, farcta pulpa subviscosa.

Hæc Ulva, quæ granulata in Mant. p. 136. dicitur, longe diversa est ab Ulva præcedente; quare mutato nomine trivialem hanc Ulvam Pilum appellavi. R.

1323. CONFERVA.

Tubercula inæqualia, in fibris capillaribus, longissimis.

* *Filamentis simplicibus, æqualibus, geniculis destitutis.*

1. CONFERVA rivularis. C. filamentis simplicissimis æqualibus longissimis. *succ. carn. Weis. crypt. 20. pal. nass. burg. lithu.*

C. filamentis simplicissimis æqualibus, longissimis. *Hall. helu. n. 2115.*

C. capillaris simplicissima enodis. *Guett. stamp.*

C. fluviatilis sericea vulgaris & fluitans. *Dill. musc. 12. t. 2. f. 2.*

Byssus palustris, confervoides, non ramosa viridis, sericum referens, filamentis longis, tenuissimis, *Mich. gen. 210. n. 3. t. 89. f. 7. Hall. R.*

Habitat in Europæ rivulis fluviiisque pacatioribus.

Filamenta simplicissima, nitida, saturate viridia, longiora in aquis fluentibus. Weis.

2. CONFERVA fontinalis. C. filamentis simplicissimis æqualibus digito brevioribus. *succ. Fl. dan. t. 651. f. 3.*

C. cespitosa, filis rectis, undique divergentibus. *Hall. helu. n. 2114.*

C. minima, byssi facie. *Dill. musc. 14. t. 2. f. 3.*

Byssus. *Mich. gen. 211. n. 8. t. 89. f. 8. n. 14. t. 89. f. 19. & n. 15. t. 89. f. 11.* ●

Habitat in Europæ fontibus.

Filamenta uncialia, circa centrum congregata. Fasciculus in ambitu atrovirens, in medio flavesce.

* * *Filamentis ramosis, æqualibus,*

3. CONFERVA bullosa. C. filamentis æqualibus ramosis, aereas bullas includentibus. *carn. gallob. nass. pal. lithu.*

C. filis ramosis æqualibus radicatis. *Fl. succ.*

C. pilis tenerrimis, vellere gossypiaceo. *Hall. helu. n. 2120.*

C. palustris bombycina. Dill. musc. 18. t. 3. f. 11.

Alga bombycina. Bauh. pin. 363.

Muscus aquaticus bombycinus, tenuissimis filamentis. Læs. pruss. 173. t. 55.

Habitat in Europæ aquis stagnantibus.

4. *CONFERVA canalicularis. C. filamentis æqualibus basin versus ramosioribus. suec. Weis. crypt. nass.*

C. filamentis tenuibus ramosis intricatis fuscis. Hall. helu. n. 2118.

C. densissima congesta : filamentis basin versus ramosioribus. Neck. meth. 5.

C. rivulorum capillacea, densissime congestis ramulis. Dill. musc. 21. t. 4. f. 15.

Alga in tubulis aquam fontanam ducentibus. Bauh. pin. 364.

Habitat in Europæ rivulis, tubulis, canalibus molendinariis.

Saturate virens, luci objecta diaphana. Weis.

5. *CONFERVA amphibia. C. filamentis æqualibus ramosis, exsiccatione coeuntibus in aculeos. suec. Weis. crypt. pal. lithu.*

C. amphibia fibrillosa & spongiosa. Dill. musc. 22. t. 4. f. 17.

Habitat in Europæ udis, fossis, umbrosis, ubi aquæ vicissitudines servant.

Fibræ rigidulæ, intertextæ in formam spongiæ. Weis.

6. *CONFERVA littoralis. C. filamentis æqualibus ramosissimis elongatis scabriusculis.*

C. maritima filiformis longissima fusca, ramis alternis simplicibus brevioribus pallidis. Fl. suec.

C. marina capillacea longa ramosissima. Dill. musc. 23. t. 4. f. 19. Rai. syn. 59.

Habitat in Europæ maritimis rupibus.

7. *CONFERVA æruginosa. C. filamentis ramosis mollibus digito brevioribus viridissimis. carn.*

C. marina capillacea brevis viridissima mollis. Dill. musc. 23. t. 4. f. 20.

Habitat in Angliæ fucis; in mari Adriatico.

8. *CONFERVA dichotoma. C. filamentis æqualibus dichotomis.*

C. dichotoma, setis porcinis similis. Dill. musc. 17. t. 3. f. 9. Hall, helu. n. 2117.

Habitat in Angliæ fossis pratorum.

9. *CONFERVA scoparia. C. filamentis proliferis fastigiatis hirtis.*

714 CRYPTOGAMIA ALGÆ

C. filamentis fasciculato-plumosis. Neck. meth. 17.

C. marina pennata. Dill. musc. 24. t. 4. f. 23.

Habitat in mari Europæo.

10. *CONFERVA cancellata. C. filis ramosis : filamentis alternis brevibus multipartitis digitatis. suec.*

C. marina cancellata. Dill. musc. 24. t. 4. f. 22.

Habitat in mari Europæo.

Filamenta lateralibus quasi intra totidem vesiculas aerem recondunt.

* * * *Filamentis anastomosantibus.*

11. *CONFERVA reticulata. C. filamentis reticulato-coadunatis. gallob. Weis. crypt. 24. lithu.*

Conferva reticulata. Dill. musc. 20. t. 4. f. 14. suec. stamp. Hall. helu. n. 2119.

Conferva reticulata crispa. Pluk. alm. 113. t. 24. f. 2. Moris. hist. 3. p. 644. f. 15. t. 4. f. 4.

Muscus aquaticus bombycinus retiformis. Læs. pruss. 173. t. 54.

Habitat in Europæ fluviiis ad eorum littora.

Sæpe natans. Weis.

* * * * *Filamentis nodosis.*

12. *CONFERVA fluviatilis. C. filis simplicissimis setiformibus rectis ; geniculis crassioribus angulatis. lithu.*

Conferva filis simplicissimis setiformibus rectis , tuberculis sparsis. Fl. suec.

C. setiformis simpliciuscula , articulata : erecta : nodis subangulatis. Neck. meth. p. 11. n. 20.

Chara sertularioides. Weber. spicil. p. 30.

Conferva fluviatilis lubrica setosa , equiseti facie. Dill. musc. 39. t. 7. f. 47. & 48.

Habitat in Europæ fluviiis confragosis.

13. *CONFERVA gelatinosa. C. filis ramosis moniliformibus ; articulis globosis gelatinosis. Fl. suec. nass.*

C. gelatinosa globulis punctatis concatenatis. Hall. helu. n. 2124.

Chara (batrachosperma) caule ramoso filamentoso ; ramis foliaceis capillaceis , in glomeres lubricos geniculatos articulatim circa caulem dispositis. Weis. crypt. 33. c. fig.

Conferva fontana nodosa , spermatis ranarum instar lubrica , major & fusca. Dill. musc. 36. t. 7. f. 42. stamp.

C. fontalis geniculata lubrica major. Dill. E. n. C. cent. f. 6. app. 60. t. 13. f. 3.

C. fontana nodosa , spermatis ranarum instar lubrica , minor & viridis. *Dill. musc.* 37. t. 7. f. 43.

Corallina pinguis ramosa viridis. *Vaill. paris.* 40. t. 7. f. 6.

Habitat in Europæ fontibus præstantissimis limpidissimis.

Confer. *Conferva stagnalis* , globulis virescentibus mucosis. *Dill. musc.* 38. t. 7. f. 44.

Conferva alpina lubrica , filamentis nodosis cæruleis. *Dill. musc.* 38. t. 7. f. 45.

Conferva fontana nodosa lubrica , filamentis tenuissimis nigris. *Dill. musc.* 39. t. 7. f. 46.

* * * * * *Filamentis geniculatis.*

14. *CONFERVA capillaris.* *C.* filis geniculatis simplicibus : articulis alternatim compressis. *herb. pal. lithu.*

C. filamentis simplicibus invicem contortis. *Neck. meth.* p. 8.

C. filamentis longis geniculatis simplicibus. *Dill. musc.* 25. t. 5. f. 25.

C. fluitans , filamentis geniculatis. *Pluk. alm.* 113. t. 8. f. 9. *Morif. hist.* 3. p. 644. f. 15. t. 4. f. 3.

Habitat in Europæ lacubus , stagnis , vadis , fossis.

15. *CONFERVA corallina.* *C.* filis geniculatis dichotomis. *carn.*

C. (*corallinoïdes*) ramosa , articulata , apice dichotoma : filis nodosis. *Neck. meth.*

C. geniculata dichotoma , ramulis acutis , articulis pelucidis albis , geniculis purpureis lævibus. *Zæga. island.* 244.

C. marina gelatinosa , corallinæ instar geniculata crassior. *Dill. musc.* 33. t. 6. f. 36.

Habitat in Mari Europæ , colore albo & rubenti.

Confer. *Conferva marina* gelatinosa , corallinæ instar geniculata tenuior. *Dill. musc.* 33. t. 6. f. 37.

16. *CONFERVA catenata.* *C.* filis geniculatis : articulis cylindricis.

C. ramosa articulis remotiusculis subcylindricis. *Neck. meth.* p. 11.

C. ramosa , geniculis longioribus cateniformibus. *Dill. musc.* 27. t. 5. f. 27.

Habitat in Mari Europam australem & Americam meridionalem alluente.

17. *CONFERVA polymorpha.* *C.* filamentis geniculatis , ramis fasciculatis. *Fl. dan.* t. 395. *Ellis. act. angl.* 1767. vol. 57. p. 424. t. 18. f. A. B.

C. marina articulata capillacea ramosissima viridis, *Fl. suec.*

716 CRYPTOGANIA ALGÆ

C. filamentosa geniculata distincta, ramis fasciculatis apice dichotomis. Neck. meth. p. 8.

C. manna geniculata nigra palmata. Dill. musc. 32. t. 6. f. 35. t. 5. c.

Cerillæ affinis f. Muscus coralloides, multifido capillaceo folio, palmites pelagica nigra. Pink. alm. 119. t. 4. f. 10.

Muscus tridactylites capillaceus maritimus multifidus niger, radice fibrosa. Bocc. musc. 1. p. 268. t. 5. f. 11.

Muscus marinus crispus pachycaulos loricatus niger. Bocc. musc. 517. t. 7. f. 1.

Fucus crispus loricatus nigricans. Barr. rar. 1323. t. 1301. Habitat in Mari Europæo.

Est fixa, ut Conferva plumosa. Ellis. l. c. t. 18. f. C.D.

18. CONFERVA *agabunda*. *C. filamentis geniculatis flexuosis, ramis ramulisque brevioribus. Suec.*

C. filamentis capillaceis expansis subarticulatis. Neck. meth. p. 9.

C. manna trichodes, lanæ instar expansa. Dill. musc. 30. t. 5. f. 32. Ray. syn. 603

Habitat in Mari Europæo, libera vagatur in aquarum mediis.

19. CONFERVA *glomerata*, *C. filamentis geniculatis, ramulis brevioribus multifidis. Weis. crypt. 25. pal. Fl. dan. t. 651. f. 2. herb.*

C. filamentis geniculatis, ramosis: ramis fasciculatis. Neck. meth. p. 9.

C. fontinalis ramosissima glomeratim congesta. Dill. musc. 28. t. 5. f. 32. & t. 5. f. 28. 29.

C. viridis capillacea, brevioribus setis, ramosior f. Conferva minor ramosa. Moris. hist. 3. p. 644. f. 15. t. 4. f. 2.

C. fontinalis trichodes. Bauh. pin. 22.

Alga fontalis trichodes. Bauh. pin. 364.

Habitat in Europæ fontibus, fossis, rivulis.

20. CONFERVA *rupestris*. *C. filamentis geniculatis ramosissimis viridibus. Suec. norv. 2. t. 3. f. 3.*

C. filamentosa, geniculata; ramis subplumosis. Neck. meth. p. 10.

C. marina trichodes ramosior. Dill. musc. 27. t. 5. f. 29.

C. marina trichodes f. Muscus marinus virens tenuifolius. Pluk. mant. 53. t. 182. f. 6.

Habitat in Europæ marinis rupibus copiosissima.

11. CONFERVA *ægagropila*. C. filamentis geniculatis ramosissimis e centro confertissimis globum constituentibus.

C. globosa, filamentis articulatis ramosissimis e centro prodeuntibus rectis. *Fl. suec.* *

Habitat in Sueciæ lacubus, Dannemoræ, Stockholmæ &c.

4324. BYSSUS.

Lanugo aut pulvis simplicissimis.

** Filamentosa.*

1. BYSSUS *septica*. B. capillacea mollissima parallela fragilissima pallida.

B. latissima filtrum s. pannum laneum simulans, filamentis tenuissimis non ramosis. *Mich. gen.* 211. n. 10.

Habitat in Domibus sub pavimentis, ubi aer mephiticus s. suffocatus, summe septicus corrodit tanquam menstruo naturali domos ligneas, durissimosque truncos, ut fatiscentes cadant damno colonum.

Similis *Lanæ tenuissimæ gossypinæ*, longitudinaliter contexte filis araneorum tenuioribus, albido-cinerascentibus, non tenacibus, lævissimis, fumis instar accensibilibus. Mucoris septicæ pessima soror. Agarici varietas.

2. BYSSUS *flor. aquæ*. B. filamentis plumosis natantibus. *gallob. carn. pal. burg. lithu.*

B. farinacea virescens aquæ inspersa. *suec. stamp.*

Conserva crustacea tenerrima subviridis. *Hall. helu. n.* 2113.

β. Byssus membranacea aquatica. *Fl. lapp.*

B. latissima, papyri instar supra aquam expansa. *Dill. musc.* 2. t. 1. f. 1.

Habitat in Mari & omni aqua, prima æstate; nocte descendit parum, die natat fere.

Constat farina, cujus molecule, Conservæ instar, capillares, ramosæ, plumæ similes, nivæ P. Bergius.

Non est planta, sed materies collecta e particulis plantarum aquaticarum putredine destructarum. Hæc crypt. p. 19.

3. BYSSUS *cancellata*. B. filis exacte undique cancelata.

B. cancellata. *Lederm. micro.* 1. 72.

Habitat in aquis dulcibus; natans tanquam mucor virescenti luteus.

4. BYSSUS *phosphorea*. B. lanuginosa violacea lignis adnascens. *Dill. musc.* 4. t. 1. f. 6. *succ. gallob. pal. lithu.*

718 CRYPTOGAMIA. ALGÆ.

Habitat in Europæ lignis putrescentibus.

5. BYSSUS *velutina*. B. filamentosa viridis : filamentis ramosis. *suec. carn. gallob. pal. Hall. helu. n. 2101. burg. lithu.*

B. repens simplicissima ramosaque : primordialibus convolutis. *Neck. meth. p. 2.*

B. tenerrima viridis velutum referens. *Dill. musc. 7. t. 1. f. 14.*

B. terrestris viridis herbacea & mollissima, filamentis ramosis & non ramosis. *Mich. gen. 211. t. 89. f. 5.*

Habitat in Terra.

6. BYSSUS. *aurea*. B. capillacea pulverulenta, fructificationibus sparsis, filamentis simplicibus ramosisque. *carn. gallob. pal. flor. dan. t. 718. f. 1.*

B. cespitosa, pulverulenta, crocea. *Hall. helu. n. 2106.*

B. erecta, glomeratim congesta simplex ramosaque : primordialibus sparsis. *neck. meth.*

B. petræa crocea, glomerulis lanuginosis. *Dill. musc. 8. t. 1. f. 16.*

B. aurea derbensis humifusa. *Pet. gaz. 1. t. 15.*

B. minima saxatilis aurea inodora, filamentis partim simplicibus partim ramosis. *Mich. gen. 210. t. 89. f. 2.*

Habitat in rupibus Arvonix, Italiæ.

7. BYSSUS *cryptarum*. B. capillacea perennis cinerea tenax rupi innata. *suec.*

Habitat in spelunca rupestri Medelpadiæ.

* * *Pulverulenta.*

8. BYSSUS *antiquitatis*. B. pulverulenta atra. *suec. lithu.*

B. petræa integerrima fibrosa. *Dill. musc. 9. t. 1. f. 18.*

Lichen crustaceus saxatilis, omnium minimus, niger. *Hall. helu. n. 2081. R.*

Habitat in Muris antiquis.

Filamenta tenuissima tecta pulvere atro. *Weis.*

9. BYSSUS *saxatilis*. B. pulverulenta cinerea rupes operiens. *suec. carn. burg. lithu.*

Lichen (*segestria*) farinosus cinereus rupes operiens. *Neck. meth. 113.*

Habitat in Saxo omni, diutius aeri exposito, quod perenni cinereo colore obducit, ipsa vix manifesta.

10. BYSSUS *Jolithus*. B. pulverulenta sanguinea saxi innascens. *gallob.*

B. perennis pulverulenta rubra lapidibus innascens. *lapp. suec. * stamp.*

Lichen crustaceus, iridis odore, ruberrimus. *Hall. helu. n.* 2090.

Lapis violaceus sylvæ hercyniæ. *Bruckm. epist. cent. 2. p.* 110. *Hall. R.*

Byssus germanica minima saxatilis aurea violæ odorem spirans. *Mich. gen. 210. t. 89. f. 3.*

Jolithus s. lapis violaceus. *Swenck. foss. 382. It. w. got. 9. 25. * 254. 263.*

Habitat in Europæ frigida sylvis opacis, supra saxa ante unum alterumve annum eversa & supinata.

11. BYSSUS candelaris. B. pulverulenta flava lignis adnascens. *Dill. musc. 3. t. 1. f. 4. succ. carn. pal. burg. lithu.*

Lichen crusta tenuissima flava. *Hall. helu. n. 2096.*

Lichen farinaceus flavus. *Neck. meth.*

Lichen crustaceus, arboribus adnascens tenuissimus, pulverulentus, pallide luteus, s. ochroleucus. *Mich. gen. 100. n. 70.*

Habitat per omnes quatuor mundi plagas in corticibus arborum, parietibus antiquis, tectis diutius vento humido expositis.

12. BYSSUS botryoides. B. pulverulenta viridis. *fl. succ. gall. pal. lithu.*

Lichen farinosus viridis. *Neck. meth.*

Lichen crustaceus, verrucosus viridis. *Hall. helu. n. 2089.*

Byssus botryoides saturate virens. *Rai. angl. 3. p. 5. 6. Dill. musc. 3. t. 1. f. 5.*

Habitat in terra diutius humida, umbrosa, ut in ollis Hortulanorum.

13. BYSSUS incana. B. pulverulenta incana, farinæ instar strata. *Dill. musc. 3. t. 1. f. 3. succ. gallob. carn. lithu.*

Lichen farinaceus mollis albicans, glebis difformibus. *Hall. helu. n. 2094.*

Lichen pulverulentus albus, globulis rotundatis. *Schreb. spicil. 139.*

Lichen farinaceus candicans. *Neck. meth. 112.*

Habitat in solo glareoso, ad latera fossularum, juxta vias publicas.

14. BYSSUS lactea. B. pulverulento-crustacea albissima. *succ. carn. pal.*

Lichen farinaceus, mollis, albus, globulis sphaericis. *Hall. helu. n. 2093.*

Lichen crustoso-farinosus candidus. *Neck. meth.*

Lichen crustaceus, saxatilis, farinaceus, albus, globulis

720 CRYPTOGAMIA. FUNGI.
minutissimis undique refertus. *Mich. gen. 99. n. 36.*
Hall. R.

Byssus candidissima, calcis instar muscos vestiens. *Dill.
musc. 2. t. 1. f. 2.*

Habitat in Muscis & arborum corticibus.

F U N G I.

1325. A G A R I C U S.

Fungus horizontalis, subtus lamellosus.

Stipitati, pileo orbiculato.

1. AGARICUS *Cantharellus*. A. stipitatus, lamellis ramosis
decurrentibus. *succ. * paris. ged. pal. Fl. dan. t. 264.*
Schæff. fung. t. 82. burg. lithu.

A. caulescens fulvus, pileo concavo: margine reflexo,
lamellis erectis, stipite brevissimo. *Fl. lapp.*

A. luteus, pileo, plicato, lobato; lamellis venosis, ramosis,
decurrentibus. *carn.*

A. luteus, pileo turbinato, in margine anguloso, lacinioso
& varie contorto. *Gleditsch. fung. 103. n. 12.*

Merulius chantarellus. Scop. carn. ed. 2. n. 1581.

*Merulius flavus, oris contortis, & laceris. Hall. helv.
n. 2326.*

*Elvela pediculata pileata, pileo subtus reticulato. Guett.
stamp. 1. p. 12.*

Fungus minimus flavescens infundibuliformis. *Bauh. pin.
373. Vaill. paris. t. 11. f. 9. 10.*

β. Fungus pileo per maturitatem instar agaryci intybacei.
Vaill. paris. t. 11. f. 11. 12. 13.

γ. Fungus angulosus & velut in lacinias scetus. *Vaill. paris.
t. 11. f. 14. 15.*

Habitat in pratis.

Variat mire ex Agarico in Pezizam Acetabulum.

*Pilei lamellæ crassæ, ab ipsa substantia fungi non dis-
sindæ.*

2. AGARICUS *quinquepartitus*. A. stipitatus, pileo subfla-
vescente partito, lamellis albidioribus interne dentato-
connexis. *pal.*

A.

CRYPTOGAMIA: FUNGI. 521

A. caulescens, pileo cinereo quinquepartito, lamellis albis.
Fl. suec.

Habitat in pratis.

3 *AGARICUS integer*. *A.* stipitatus; lamellis omnibus magnitudine æqualibus. *suec.* * *nass. pal.*

A. caulescens, pileo plano-concavo purpureo, lamellis stipiteque albis. *Fl. lapp.*

Amanita petiolo satis, pileolo convexo, umbilicato; oris rugosis, lamellis albis. *Hall. helv. n. 2374. var. a.*

Agaricus Russula. Scop. carn. Schæff. fung. t. 58.

A. pileo orbiculari, in margine rugoso & striato; vertice tumido; petiolo brevi. *Gled. fung. 106. n. 14.*

A. roseus. Schæff. t. 75. cassipes. Ei. t. 87. 88. ruber. t. 92. cyanoxanthus. t. 93. & virescens. t. 94. hujus varietates sunt.

A. major, pediculo longo modice crasso; lamellis albis creberrimis superne ad margines apparentibus. *Rai. suppl. 18. Buxb. cent. 4. p. 12. t. 19.*

β. *A. caulescens* petiolo crassiusculo albo; pileo plano sanguineo; lamellis albis. *Fl. lapp. 487.*

Habitat in sylvis.

Amanita Hall. helv. n. 2374. var. β. est Agaricus emeticus Schæff. fung. t. 15. & 16. ab A. integro L. probe distinguendus, dum emest excitat. A. integer carnem solidiorem habet, quæ in A. emetico laxa & porosa existit. Schæff. l. c. R.

4. *AGARICUS muscarius. A.* stipitatus; lamellis dimidiatis solitariis, stipite volvato: apice dilatato, basi ovata. *suec.* * *Mich. gen. t. 78. f. 2. pal. nass. Schæff. fung. t. 27. & 28. var. absque verrucis: burg. lithu.*

Agaricus caulescens pileo sanguineo: verrucis lamellisque albis, stipite albo basigloboso. *Fl. lapp.*

Amanita petiolo anulato; sanguineus, lamellis albis. *Hall. helv. n. 2373.*

Agaricus volva exceptus, pileolo conico; in margine striato, petiolo anulato, in basin squamoso tuberosam delinente. *Gled. fung. 82. n. 1. & p. 83. var. b. absque verrucis.*

A. pileo plano miniato verrucoso; verrucis & lamellis albis; stipite albo, volvato, bulboso, annulato; longo, apice dilatato, pleno. Scop. carn. 1459.

A. olivaceus. Schæff. fung. t. 204. Hall. l. c. var. n. & A. xerampelinus. Ejusd. t. 215. Hall. l. c. var. δ.

Fungorum perniciosorum genus. 12. n. 4. *Clus. hist. 2. p. 280. Linnæ. spec. plant. Europæ, Pars II.*

22 CRYPTOGAMIA. FUNGI

Habitas in pascua.

Cinices necat & eradicat promptissime. Ic. scan. 430.

5. **AGARICUS densatus.** A. stipitatus, pileo convexo, lamellis basi mucrone dentatis. herb. pal.

A. caulescens, pileo, flavo levi margine inflexo, lamellis pallidioribus basi angularis. El. succ.

A. pistacinus Schaff. fung. 1. 301. coccineus Hall. bot. n. 2421.

Amanita flavus, viscidus, pileolo aculeo. Hall. bot. n. 2421.

Habitas in Sylvis.

Pileus florescens, levis, subviscidus. Lamellae breviter & Scipius distantes, ubi denticulo acuminato inflexae. Stipes fistulosus, apertus, cylindricus, niger, vix pilis striatus, confertim natus.

6. **AGARICUS deliciosus.** A. stipitatus, pileo velutato, succo lactescente, succ. n. pal. Schaff. fung. 1. 311. Hall. bot. n. 2419.

Amanita solvus, lacte croceo. Hall. bot. n. 2419.

Agaricus lateralis coloris, pileo concavo, croceo & lacte turgido; lamellis ramosis; stipite cylindrico, muculato, brevi. Scop. carn.

A. pileo hemisphaerico, in centro depresso & in margine fornicato; petiolo brevi & crasso. Gled. fung.

Amanita lateralis coloris, croceo succo turgens. Dill. giff.

Fungus esculentus lateralis colore immutabili, succum acrem & croceum fundens. Mich. ger. 141.

Habitas in Montibus, sterilibus, sylvest.

7. **AGARICUS lachryans.** A. stipitatus, pileo plano carne lactescente, lamellis natis, stipite longo carneo. pal. Schaff. fung. 1. 3.

Amanita solvus, lacte croceo. Hall. bot. n. 2419. cum praecedente conjungit.

Agaricus (testaceus) ferrugineus; pileo concavo; albo & dulci lacte turgido; lamellis ramosis; stipite cylindrico, brevi. Scop. carn.

Habitas in sylvis.

8. **AGARICUS piperatus.** A. stipitatus, pileo, plantisculo lactescente; margine deflexo, lamellis iscarneo-pallidis; succ. pal. burg. litha.

Amanita umbilicatus, albus, lacte acri natus. Hall. bot. n. 2328.

Agaricus pileo concavo, sæpius lactescente, lamellis ramosis, stipite nudo, cylindraceo. *Scop. carn.*

A. pileo pulvinato, integerrimo, & in centro profundius depresso, lamellis connexis angustis & compactis petiolo continuo, cylindrico & crasso. *Gled. fung.*

Fungus piperatus albus crassus, lacteo succo turgens. *Mich. gen. 141.*

Fungus albus acris. *Bauh. pin. 371.*

Habitat in Pascuis, sylvis.

9. AGARICUS campestris. A. stipitatus, pileo convexo squamato albido, lamellis ruffis. *succ. carn. norv. pal. nass. dan. t. 714. Schæff. fung. t. 33. & t. 310. 311. burg. lithu.*

A. caulescens, pileolo sordido lacero, pileo albo membranaceo, lamellis rufescentibus. *Fl. lapp.*

Amanita petiolo anulato, pileo campaniformi, albido, laminae roseis. *Hall. helu. n. 2443.*

Agaricus pileo amplo, hemisphærico, fornicato; petiolo brevi, annulo permanente cincto. *Gled. fung.*

A. pediculo annulato, pileo superne albus, inferne rubens. *Guett. stamp.*

Fungus campestris albus superne, inferne rubens. *Bauh. hist. 3. p. 824. Mich. gen. 174. n. 8.*

Habitat in pratis.

10. AGARICUS Georgii. A. stipitatus, pileo flavo convexo, lamellis albis. *succ. lithu.*

A. caulescens, pileo plano sulphureo, lamellis albis, stipite crassiusculo. *Fl. lapp.*

Amanita pileo flavo, oris striatis & lanuginosis, lamellis albis. *Hall. helu. n. 2367.*

Agaricus pileolo orbiculari, pulvinato, pallido, vel candido, margine hirsuto, aureo, striato, lamellis albis; petiolo brevi crasso & villoso. *Gled. fung.*

Fungus orbicularis exalbida pratensis. *Bauh. pin. 370.*

Fungus divi georgii, coloris exalbidi cum pauca flavitie; esculenti pratensis i. *Bauh. hist. 3. p. 824.*

Fungus croceus magnus. *Sterb. fung. 63. t. II. f. 3.*

Habitat in sylvis.

11. AGARICUS violaceus. A. stipitatus, pileo ramoso; margine violaceo tomentoso, stipite cærulescente lana ferruginea. *succ. * paris. lithu.*

Amanita violaceus, petiolo bulboso, annulato, pileo plano. *Hall. helu. n. 2435.*

724 CRYPTOGAMIA. FUNGI.

Agaricus violaceus. *Schæff. fung. t. 3. cærulescens. t. 34. amethystinus. t. 56.*

Fungus esculentus bulbosus dilute purpureus. Mich. gen. 149. t. 74. f. 1.

Fungus cæruleus major. Buxb. cent. 4. p. 7. t. 11.

Fungus lividus, pediculo bulboso. Ei. cent. 4. t. 22. Hall. R.

Habitat ad margines Sylvarum:

12. *AGARICUS cinnamomeus*. *A. stipitatus, pileo sordide flavo lamellis luteo-rufis. suec. pal.*

A. caulescens flavus, pileo plano. Fl. lapp.

Amanita sicca, leoninus, anulo araneoso. Hall. helu. n. 2432.

Amanita dura, ex fusco rubens quercina. Raj. syn. 3. p. 3. n. 23. sec. Hall. R.

Habitat in sylvis.

13. *AGARICUS viscidus. A. stipitatus, pileo purpurascente fusco viscido, lamellis fusco purpurascens. pal. burg.*

A. caulescens, pileo convexo purpurascens, lamellis rufis, stipite brevi crassiusculo albo. suec.

Habitat in Sylvis.

14. *AGARICUS equestris. A. stipitatus, pileo pallido: disco stellatim luteo, lamellis sulphureis. suec. * pal.*

A. caulescens, pileo convexo sordido: margine inflexo, lamellis pallidis: basi remotis, stipite cylindraco. Fl. lapp. 502.

Habitat in pascuis, sylvis.

15. *AGARICUS mammosus. A. stipitatus, pileo convexo acuminato griseo, lamellis convexis griseis crenatis, stipite nudo. pal. burg.*

A. caulescens, pileo convexo sordido: umbilico prominente lamellis pallidis, stipite cylindraco. Fl. lapp. 501. Fl. suec.

Fungus minor, capitulo pileum nauticum referente, pediculo longiore. Buxb. cent. 4. p. 13, t. 21. f. 1.

Habitat in sylvis.

16. *AGARICUS clypeatus. A. stipitatus, pileo hemisphaerico viscido acuminato lamellis albis, stipite longo cylindrico albo. suec. * paris. lithu.*

Amanita petiolo procero, squamoso, anulato, pileo conico, squamoso, subspadiceo, laminis albis. Hall. helu. n. 2366.

Amanita petiolo procero, anulato, in acetabulum pilei immisso, pileo squamoso & maculato, lamellis albis. *Hall. helu. n. 2371.*

Agaricus (*procerus*) pileo plano papillari, stipite convergente, procero, anulato, fistuloso, anulo peculiari inserto. *Scop. carn. Schæff. fung. t. 22. 23. Flor. dan. t. 772.*

Fungus viscidus papillaris striatus terreus, subtus albus, *Hall. helu. 41.*

Habitat in pratis sylvaticis.

Agaricum suum, procerum cum synonymis Halleri & Scopoli huc trahit ven. Schæfferus, an recte. R.

17. *AGARICUS exstindorius*. A. stipitatus, pileo campaniformi albido lacero, lamellis niveis, stipite subbulboso subulato nudo. *herb.*

A. caulescens albus, pileo conico, stipite cylindraceo. *suec. **

A. caulescens albus, petiolo, longissimo, pileo conico. *Fl. lapp.*

Amanita albus, petiolo villosus, pileolo umbonato striato, *Hall. helu. n. 2342. &*

Amanita albus, hemisphaericus, ambitu striato. *Et. n. 2343.*

Fungus albus viscidus, pediculo longo. *Buxb. cent. 4. t. 30. f. 2. sec. Haller. R.*

Habitat in fimetis & ad pagos.

19. *AGARICUS fimetarius*. A. stipitatus, pileo campanulato lacero, lamellis nigris lateraliter flexuosis, stipite fistuloso. *ged. pal. nass. Schæff. fung. t. 7. 8. 46. 47. burg. lithu.*

A. caulescens, pileo conico cinereo, lamellis nigris: margine lacero, stipite longissimo albo. *Fl. lapp. 503. Fl. suec.*

Amanita albus, campanulatus, squamosus, nigrescens. *Hall. helu. n. 2357.*

Agaricus solitarius; pileo campanulato lanuginoso striato griseo; lamellis demum atris, diffluentibus; stipite procero fistuloso, anulo fugaci cincto. *Scop. carn.*

A. pileo campanulato, sulcato, sericeo & griseo, in margine obtuse denticulato, lamellis cinereis; petiolo ex una basi multiplici, cylindrico fistuloso & albo. *Gled. fung.*

Fungus sterquilineus. *Mich. gen. 181. t. 80. f. 3. & t. 74. f. 6. Scop. R.*

725 CRYPTOGAMIA. FUNGI.

Fungus conatus. *Sterb. fung.* 218. t. 22. f. J. K. L.

Fungus albus ovum referens. *Buxb. cent.* 4. p. 16. t. 27. f. 1.

Fungorum perniciosorum genus, 14. spec. 3. *Clus. hist.* 2. p. 282.

Habitat in fœmetis.

20. AGARICUS *campanulatus*. A. stipitatus, pileo campanulato striato pellucido, lamellis adscendentibus, stipite nudo. *Scholl. barb. pal. burg. lithu.*

A. caulescens pileo conico cinereo, lamellis albis, stipite longo. *Fl. lapp.* 507. *Fl. suec.* 1054. 1213. *

Fungus multiplex obtuse conicus colore griseo murino. *Vaill. paris.* 71. t. 12. f. 1. 2.

Habitat in pratis.

21. AGARICUS *separatus*. A. stipitatus, pileo lævi livido, lamellis nigricantibus separatis, stipite pulbofo volvato. *suec.*

Habitat in stercoretis.

22. AGARICUS *fragilis*. A. stipitatus, pileo convexo viscido pellucido lamellisq̃ luteis, stipite nudo. *herb. pal. Schæff. fung.* t. 230.

Amanita flavocinnamæus, viscidus, striatus papillatus. *Hall. helu. n.* 2425.

Amanita flavus, petiolo gracillissimo & procerissimo pileo conico viscido. *Hall. l. c.* 2426.

Fungus pileo crocei splendoris. *Vaill. paris.* t. 11. f. 16. 17. 18.

Habitat ad ambulacra.

23. AGARICUS *umbelliferus*. A. stipitatus, pileo plicato membranaceo, lamellis basi latioribus. *suec. pal. Schæff. fung. tab.* 309. *lithu.*

Amanita albus, tener, pileolo campanulato striato. *Hall. helu. n.* 2346.

Agaricus pileo convexo plicato membranaceo tenero; lamellis remotis; stipite procero, capillaceo, nudo, lamellis concolore. *Scop. carn.*

A. exiguus, niveus, pileo hemisphærico, a vertice versus marginem striato, lamellis rarioribus, petiolo longiore & tenui. *Gled. fung.*

A. caulescens albus parvus, stipite longo, pileo plano pellucido: margine multifido. *Fl. lapp.*

A. minimus, capitulo turbinato plano albo, lamellis margine fuscis. *It. al.*

CRYPTOGAMIA FUNGI 729

Fungus minimus totus albus, pileo hemisphaerico utrinque striato, lamellis rarioribus. *Mich. gen.* 166. t. 80. f. 11.
Habitat inter Folia congesta, semiputrida.

24. AGARICUS *Androsaceus*. A. stipitatus albus, pileo plicato membranaceo, stipite nigro. *suec. parif. pal. Schæff. fung.* t. 239. *nass. lithu.*

A. caulescens albus minimus, petiolo angusto, pileo plano verrucoso: margine octofido. *Fl. lapp.*

Amanita petiolo nigro, pileolo albo striato, excarni. *Hall. helu. n.* 2351.

Agaricus pileo convexo plicato membranaceo; lamellis remotis; stipite nigro procero, capillaceo nudo. *Scop. carn.*

A. pileolo striato hemisphaerico parvo, lamellis angustissimis paucis petiolo capillaceo altissimo. *Gled. fung.*

Fungus caule nigro capillari, androsaces capitulo. *Bocc. mus.* 143 t. 104.

Fungus pileo candicante, lamellis paucis, pediculo fusco splendente. *Vaill. parif.* 69. t. 11. f. 21. 22. 23.

Habitat in foliis dejectis Pini.

Stipes capillaris. Lamellæ tenuissimæ.

25. AGARICUS *clavus*. A. stipitatus, pileo luteo convexo striato, lamellis stipiteque albis. *suec. * pal. Schæff. fung.* t. 59. *burg.*

Amanita minimus, oris adtractis, flavus, lamellis albis. *Hall. helu. n.* 2370.

Fungus minimus aurantius mammillaris. *Vaill. parif.* 76. t. 11. f. 19. 20.

Habitat in Nemoribus, inter folia decidua.

* * *Parasitici, acaules, dimidiati.*

26. AGARICUS *quercinus*. A. acaulis, lamellis labyrinthiformibus. *suec. carn. pal. nass. Schæff. fung.* t. 57. *burg. lithu.*

Amanita sessilis durissimis, lamellis cartilagineis intricatis. *Hall. helu. n.* 2330. R.

Agaricus coriaceus durus sessilis squamosus & multiplex, lamellis ramosis, sinuosis & sibi invicem implexis. *Gled. fund.*

Agaricus villosus, lamellis sinuosis & invicem implexis. *Buxb. cent.* 5. p. 1, 4. f. 1.

Habitat in Quercubus.

27. AGARICUS *betulinus*. A. acaulis coriaceus villosus:

728 **GYRTOGANYA FUNGUS**
margine obtuso, lamellis ramosis antistipulantibus. *Fr.*
Flor. dan. t. 776. f. 1.

Merulius sessilis, albus, hirsutus, subtus croceus, venis
serpentinis intricatis. *Hall. helv. n. 2329. R.*

Habitat in Betulis.

28. **AGARICUS alneus**. A. acaplis lamellis bifidis pulver-
ulentis. *succ. burg.*

Amanita sessilis, lamina pulverulenta, bifidis. *Hall. helv.*
n. 2333.

Agaricus sessilis, coriaceus, villosus; lamellis pulverulentis
ramosis, bifidis. *carn.*

A. sessilis squamosus lobatus & villosus; lamellis dissectis.
Gled. fung.

A. alneus. *Schæff. fung. t. 246. An. R.*

Habitat in Alno.

*Paucas species Fungorum recenseo, ut evitavi varietates,
quæ plures sunt, quam vulgo creditur, & quidem sæpe aliena
genera constituentes, adeoque in his etiamnum valde deficiat
Res herbaria.*

1326. **BOLETUS**

Fungus horizontalis, subtus prorolus.

* *Parasitici, acaules.*

2. **BOLETUS suberosus**. B. acaulis pulvinatus albus levis,
poris acutis difformibus. *herb.*

B. acaulis coriaceo-convexus villosus albus, poris diffor-
mibus, rotundis flexuosisque. *Fl. succ. 1093. 1253.*

Habitat in Betulis.

3. **BOLETUS fomentarius**. B. acaulis pulvinatus inæqualis
obtus, poris teretibus æqualibus glaucis. *succ. t.*

Habitat in Betulis.

4. **BOLETUS igniarius**. B. acaulis pulvinatus levis, poris
tenuissimis. *succ. carn. nass. burg. lithu.*

Polyporus sessilis, convexo-planus, durissimus, cinereus,
inferne albidus. *Hall. helv. n. 2288.*

Bolus subrotundus, petiolo horizontali brevissimo, tubulis
subtilissimis. *Gled. fung.*

B. unguatus. *Schæff. fung. t. 137. 138.*

Agaricus pedis equini facie. *Tournef. inst. 562.*

Fungus in caudicibus, nascent, unguis equini figura. *Bauh.*
pin. 372.

Polyporus sessilis, convexo planus, inferne albidus, superne fulvus, zonis discoloribus. *Hall. helu. n. 2289. Schæff. fung. t. 138. β. ex Schæff. R.*

Habitat in Betulis aliisque arboribus.

6. *BOLETUS versicolor*. B. acaulis : fasciis dicoloribus poris albis. *suec. ged. gallob. pal. nass. Schæff. fung. t. 136.*

Agarico polyporus versicoloribus lineis variis, poris albis tenuissimis. Hall, helu. ed. pr. 26.

Polyporus sessilis, cespitosus, planus, hirsutus, anulis versicoloribus. *Hall. helu. n. 2282.*

Boletus acaulis ; pileo coriaceo, villosa, lineis concentricis dicoloribus, fasciato tubulis; tenuissimis. *Scop. carn.*

B. coriaceus, squamosus, sessilis & multiplex, lobis varie divisis & expansis, tubulis tenuissimis, connexis angulosis. *Gled. fung.*

B. variegatus. *Schæff. fung. t. 263. atrofuscus t. 268. multicolor. t. 269. varietates hujus esse videntur.*

Habitat ad truncos arborum antiquarum.

7. *BOLETUS suaveolens*. B. acaulis, superne lævis, salicinus. *Fl. suec. gallob. lithu.*

Habitat in Salice.

* * *Stipitati.*

8. *BOLETUS perennis*. B. stipitatus perennis, pileo utrinque planiusculo. *suec. paris. pal. lithu.*

Fungus lignosus fasciatus. *Vaill. paris. t. 12. f. 7.*

Habitat in sylvis super terram, subiectis caudicibus arborum putridis.

9. *BOLETUS viscidus*. B. stipitatus, pileo pulvinato viscido, poris teretibus convexis immersis distinctis, stipite lacero.

B. caulescens, pileo luteo, poris teretibus distinctis lividis, stipite lacero. *Fl. suec. **

Habitat in Sylvis.

10. *BOLETUS luteus*. B. stipitatus, pileo pulvinato subviscido, poris rotundatis convexis flavissimis, stipite albido. *pal. Schæff. fung. t. 114.*

B. caulescens, pileo livido subviscido, poris flavis convexis, stipite albido. *Fl. suec. * paris.*

Polyporus carne secedente, petiolatus; annulatus. Hall. helu. n. 2301.

Boletus pileo plano carnosio glabro; tubulis difformibus; stipite anulato, longo. Scop. carn.

Boletus luteus. Dill. giff. 18. t. 10. n. 1.

730 CRYPTOGRAMIA FUNGI.

Pilsen. Müller oratio om suampe, Hafn. 1763. 4to.

Fungus porosus autumnalis viscidus, Buxb. cent. 5. p. 7. t. 14.

Habitat in Sylvis.

11. BOLETUS *bovinus*. B. stipitatus, pileo glabro pulvinato marginato, poris compositis acutis: porulis angulatis brevioribus. gallob. lithu.

B. caulescens, pileo testaceo, poris compositis angulatis griseis. Fl. suec. * paris.

Polyporus carne secedente, petiolatus, fulvus, inferne virescens. Hall. helu. n. 2302.

Boletus crassipes. Schæff. fung. t. 112.

B. stipitatus, pileo pulvinato, glabro; tubulis connexis, angulatis, carne pilei brevioribus. Scop. carn.

Suillus esculentus crassus, superne fulvus, inferne initio albidus. Mich. gen. 127.

Boletus julii mensis. Dill. giff. 188.

Suillus fulvus inferne ex flavo virescens. Hall. helu. ed. pr. 29.

b. Polyporus carne secedente, petiolatus, petiolo terreo, inferne aurantio. Hall. helu. n. 2307. R.

Boletus luridus. Schæff. fung. t. 107.

c. Polyporus carne secedente, purpureus, inferne flavus, Hall. helu. n. 2305. var β.

Boletus reticulatus. Schæff. t. 108.

d. Polyporus. Hall. helu. n. 2308.

Boletus bulbosus. Schæff. t. 134.

B. bovinus. Ejusd. t. 104.

e. B. olivaceus. Ejusd. t. 105.

f. Polyporus. Hall. helu. n. 2310.

Boletus rufus. Schæff. t. 103.

Habitat in Pratis.

Polyporos. Hall. helu. n. 2302-2313. recte huc retulit Scopoli. Ab his tamen Polyporus n. 2311. qui cum Boletis subtomentoso Lin. idem. videtur, omnino recedit. Wigg. primi. p. 103.

12. BOLETUS *granulatus*. B. stipitatus pileo viscido pulvinato, poris teretibus subangulatisque truncatis angula granulato. suec.

Habitat in Sylvis.

Pileus convexus, viscidus, carnosus, lividus margine acuto cinctus. Pori flavi, diffformes, inæquales, teretes & subangulati, truncati: margine laterali, ad angulos, quasi farinæ particula foliaria auti. Stipes flavus, filica brevior.

CRYPTOGAMIA. FUNGI. 731

13. **BOLETUS** *subtomentosus*. B. stipitatus, pileo flavo subtomentoso, poris subangulatis difformibus fulvis planis, stipite flavo. *succ.*

Habitat in Sylvis.

14. **BOLETUS** *subsquamosus*. B. stipitatus, pileo albido, poris difformibus oblongo-flexuosis niveis. *succ.* *

Habitat in Sylvis.

1327. HYDNUM.

Fungus horizontalis, subtus echinatus.

1. **HYDNUM** *imbricatum*. H. stipitatum, pileo convexo imbricato. *succ. pal. nass. Fl. dan. t. 176. Schæff. fung. t. 140.*

Echinus petiolatus albicans, superne squamosus. *Hall. helu. n. 2324. b.*

Erinaceus esculentus albus crassus. *Mich. gen. 132. t. 72. f. 2.*

Fungus pene candidus, prona parte erinaceus. *Bauh. hist. 3. p. 828.*

6. **Hydnum** *squamosum*. *Schæff. fung. t. 273. Hall. l. c. a. Habitat in Sylvis acerosis.*

2. **HYDNUM** *repandum*. H. stipitatum, pileo convexo lævi flexuoso. *succ. paris. carn. pal. Schæff. fung. t. 141. Flor. dan. t. 310. burg.*

H. pediculatum, pileo inæqualiter laciniato scabro flavo. *Guett. stamp.*

Echinus petiolatus, lamellatus subrufus, petiolo lævi. *Hall. helu. n. 2325.*

Boletus pileolo pulvinato, integerrimo, tubulis connexis, echinato denticulatis. *Gled. fung.*

Erinaceus coloris pallide citrei. *Dill. giss. t. 1.*

Hydnum *flavidum*. *Schæff. fung. t. 318. varietas est.*

Erinaceus esculentus pallide luteus. *Mich. gen. 132. t. 72. f. 3.*

Habitat in Sylvarum desertis.

3. **HYDNUM** *tomentosum*. H. stipitatum, pileo plano infundibuliformi. *Fl. succ. ped. herb. Fl. dan. t. 534. f. 3. Habitat in Sylvis acerosis.*

4. **HYDNUM** *auriscalpium*. H. stipitatum, pileo dimidiato. *succ. pal. nass. Schæff. fung. t. 143.*

732 CRYPTOGAMIA. FUNGI.

Echinus petiolo gracili, laterali, pileolo plano, obscuro; Hall. helu. n. 2321.

Boletus pileo semiorbiculari, leviter fastigiato, tubulis connexis, echinato-denticulatis; petiolo præalto & tenui. Gled. fung.

Erinaceus parvus hirsutus ex fusco fulvus, pileo semiorbiculari, pediculo tenuiore. Mich. gen. 132. t. 72. f. 8.

Fungus erinaceus parvus in conis abietis nascens. Buxb. cent. I. t. 57. f. 1.

Fungus erinaceus parvus, pediculo longiore auriscalpium referens buxei coloris. Buxb. hall. 129. t. 129.

Habitat in Sylvis acerosis supra terram, subjacente rama aut strobilo; an varietas sola H. imbricati?

5. *HYDNUM parasiticum. H. acaule arcuato-rugosum tomentosum. Fl. dan. t. 465.*

Habitat in Europæ arboribus.

1328. PHALLUS.

Fungus supra reticulatus, subtus lavis.

1. *PHALLUS esculentus. P. pileo ovato celluloso, stipite nudo rugoso. succ. paris. ged. pal. nass. Schæff. fung. 199. 298. 299. 300. burg. lithu.*

P. petiolo nudo, pileo subtus laxo. Hort. cliff. 479.

P. petiolo nudo, pileo subtus undique annexo. Hort. cliff.

Boletus capite terete reticulato. Hall. n. helu. 2247.

Phallus avolvatus, pileo plicato. Scop. carn.

Ph. capitulo fastigiato, subtus operculato, petiolo nudo. Gled. fung.

Boletus esculentus rugosus albicans, quasi fuligine inferus. Tournef. inst. 561. t. 329. f. A.

Boletus esculentus rugosus 1, 2, 3. Mich. gen. 230. t. 85. f. 1. 2.

Fungi esculenti species tres. Clus. hist. 2. p. 164.

Habitat in Sylvis antiquis.

Varietas pileo bicolli picta est in fl. dan. t. 53.

2. *PHALLUS impudicus. Ph. volvatus stipitatus, pileo celluloso. ged. Fl. dan. t. 175. Schæff. fung. t. 196. 198. burg. lithu.*

Ph. volvatus pileo apice clauso. succ.

Ph. volva exceptus, pilei apice pervio. Guett.

Ph. volva exceptus, capituli apice patulo. Gled. fung. Hall. helu. n. 2248.

CRYPTOGAMIA FUNGI. 733

Ph. volvatus ; pileo celluloso , utrinque pervio. *Scop. carn.*

Ph. vulgaris totus albus , volva rotunda , pileo cellutato ac summa parte umbilico pervio ornato. *Mich. gen.*

201. t. 83.

Ph. hollandicus s. batavicus. *Daleck. hist.* 1398.

Fungus foetidus penis imaginem referens. *Bauh. pin.* 374.

β. Phallus volvo exceptus capituli apice clauso. *Gled. fung.*

54. n. 1;

Habitat in Sylvis.

1329. CLATHRUS.

Fungus subrotundus , cancellatus

* *Acaules.*

1. **CLATHRUS cancellatus.** C. acaulis subrotundus. *Col. ecphr.* 1. p. 337. t. 336. *Barr. ic.* 1265. *carn. Sciaff. fung. rom.* IV. tab. in titulo expressa.

C. sessilis ovato-turbinatus. *Gl. fung.*

C. ruber. *Mich. gen.* 214. t. 93.

Boletus cancellatus purpureus. *Tournef. inst.* 561. t. 329. f. 6.

Fungus coralloides cancellatus. *Clus. app. als.*

Fungus rotundus cancellatus. *Bauh. pin.* 375.

Habitat in Europa australiori.

* * *Stipitati.*

2. **CLATHRUS denudatus.** C. stipitatus , capitulo oblongo volvato. *Jacq. miscell. austr.* 1. p. 136. tab. 6.

C. pediculatus purpureus , capite oblongo. *Guett. stamp.*

Trichia gregaria , purpurea , petiolata , capitulo sphaerico. *Hall. helv. n.* 2164. t. 48. f. 6. *Jacq. R.*

Clathroides purpureum pediculo donatum. *Mich. gen.* 142. t. 94. f. 1.

Habitat in Europa australiore.

3. **CLATHRUS nudus.** C. stipitatus , capitulo oblongo axi longitudinali adnato. *Fl. succ. Fl. dan.* t. 216.

Clathroidastrum. *Mich. gen.* 214. t. 94.

Habitat in Italiae lignis putridis.

4. **CLATHRUS recutitus.** C. stipitatus , capitulo globoso glande ovali. *Fl. succ.*

Habitat in Sueciae truncis arborum.

1330. HELVELLA.

*Fungus turbinatus.** *Stipitata.*

1. **HELVELLA** *Mitra*. H. pileo deflexo adnato lobato difformi. *suec. ged. pal. Fl. dan. t. 116. Schæff. fung. t. 154. 159. 162.*

Boletus capitulo explanato laciniato. *Hall. helv. n. 2246.*

Elvela ferme sessilis & parvo, lamina in formam capinli deorsum plicato-laciniata & crispa. *Gled. fung. C.*

Fungoides fungiforme crispum laciniatum & varie complicatum. *Mich. gen. 204. t. 86. f. 7.*

Fungus autumnalis velut apex staminis. *Mentz. pug. t. 6.*
Habitat in Truncis putridis.

* * *Acaulis.*

2. **HELVELLA** *pineti*. H. acaulis.

Agaricus acaulis utrinque planiusculus. *Fl. lapp. 517. Fl. suec. 1084.*

Habitat in Pinu, Abiete.

1331. PEZIZA.

Fungus campanulatus, sessilis.

1. **PEZIZA** *lentifera*. P. campanulata lentifera. *pal. Fl. dan. t. 105. burg.*

Cyathus sericeus intus lævis. *Hall. helv. n. 2215.*

Peziza cyathiformis. *Scop. carn. 2. n. 1629.*

P. sericea. *Schæff. fung. t. 180.*

P. caliciformis lentifera lævis. *Dill. giss. 195. Fl. suec. 1109. 1271.*

Cyathoides cyathiforme cinereum & veluti sericeum. *Mich. gen. 222. t. 102. f. 1.*

Fungoides infundibuli forma, semine fœtum. *Vaill. paris. 56. t. 11. f. 6. 7.*

Fungus campaniformis niger parvus, multa semine in se continens. *Pluk. phyt. 184. f. 9.*

Fungus pyxoides seminifer. *Næf. pruss. 98. t. 16.*

Fungus spermelias. *Bocc. musc. 1. t. 301. f. 1.*

Fungus calyciformis seminiferus. *Mentz. pug. t. 6.*

CRYPTOGAMIA: FUNGI. 735

P. *Peziza calyce campanulato, interne striato, externæ squamoso. Guett. stamp. 1. p. 15. Dalib. paris.*

Cyathus hirsutus intus striatus. *Hall. helu. n. 2214. Fl. dan. t. 469. f. 1.*

Peziza hirsuta. Schæff. t. 178.

Habitat in Agris, Lignis, Sepimentis.

2. PEZIZA punctata. *P. turbinata truncata, disco punctato succ. gallob. Fl. dan. t. 288.*

Elvela petiolo brevissimo pileo, turbinato, disco punctato. *Hort. cliff.*

Sphæria nivea, plana, punctis nigricantibus. *Hall. helu. n. 2184.*

Sphæria stercoracea. *Wig. primit. p. 85.*

Elvela turbinata, patula, disco foraminulis pertuso, basi brevissima. *Gled. fung. 44. n. 7. R.*

Fungus minimus infundibuliformis, superne nigris punctis notatus. *Rai. suppl. 21.*

Musculus minimus lignosus, disco punctato. *Bosc. musc. p. 25. t. 107.*

Habitat in Stercore equino, nec alibi, anne varietas Boleti?

Puncta atra e seminibus immersis.

3. PEZIZA cornucopioides. *P. infundibuliformis, disco patiente sinuato punctato. pal. Fl. dan. t. 384. Schæff. fung. t. 165.*

P. membranacea, tubæ similis, marginibus replicatis. *Hall. helu. n. 2216.*

Elvela cornucopiæ. *Schæff. fung. t. 165.*

Elvela punctata. *Ei. t. 166.*

Elvela stipitata; pileo infundibuliformi, punctato margine reflexo. *Strop. carn.*

Elvela corpore infundibuliformi, intus fusco, extrus cinereo, margine tumido, extrorsum reflexo. *Gled. fung. 41. b.*

Elvela infundibuliformis, ore expanso laciniato punctato. *Dalib. paris.*

Fungoides nigricans majus cornucopiæ forma. *Vaill. paris. t. 13. f. 2. 3.*

Habitat in Gallia.

Caulescens hæc est cum sequenti.

4. PEZIZA acetabulum. *P. cyathiformis: extus angulata venis ramosis. Schæff. fung. t. 150. 155. & 156.*

B. bracteata, fordide lutea, subvillosa. *Hall. helu. n. 2223.*

736 CRYPTOGAMIA. FUNGI.

Elvela corpore hemisphaerico, excavato, exterius colla
ramosis & sibi invicem implexis circumdata; basi brevissi-
ma. *Gled. fung.* 42.

Fungoides fuscum acetabuliforme: externe ramificatum.
Vaill. paris. 57. t. 13. f. 1.

Fungoides fuscum majus. *Burb. n. 4. p. 283. s. 29. f. 1. 2.*

Fungoides maximum pyxidatum. *Vaill. paris.* 57. t. 13. f. 1.

Fungoides maximum vasis fusorii forma. *Mich. gen.* 206.
t. 86. f. 1.

Habitat in Europa australi.

Varietas an hæc Agarici Chantarelli.

5. PEZIZA cyathoides. P. cyathiformis: margine obtuso
erecto. *succ. Gunn. norv. n. 328. pal.*

P. lutea parva: marginibus lævibus. *Rai. angl.* 3. p. 479.
t. 24. f. 4.

Habitat in Terra.

6. PEZIZA cupularis. P. globoso campanulato: margini
crenato. *succ. pal. Fl. dan. t. 469. f. 3.*

P. petiolata, hemisphaerica, ora dentata. *Hall. helv. n.*
2228.

P. calyce campanulato globoso: ore crenato. *Dalib. paris.*

Fungoides glandis cupulam referens, margine dentato. *Vaill.*
paris. 57. t. 11. f. 1. 2. 3.

Habitat in Gallia, Helvetia, Germania, &c.

7. PEZIZA scutellata. P. plana margine convexo piloso.
succ. pal.

P. lutea parva, marginibus pilosis. *Rai angl.* 3. p. 18.
t. 24. f. 3.

P. plana sessilis aurantia anullata. *Vaill. paris.* t. 13. j. 14.
Hall. helv. n. 2242.

Elvela (ciliata) sessilis, disco coccineo, plano; oris ci-
liatis. *Scop. carn. Schæff. fung. t. 284.*

Elvela orbicularis minima sessilis & coccinea, plana margine
ciliato vel nudo. *Gled. fung. b.*

Elvela sessilis scutellata. *Guett. stamp.*

Fungoides, qui Fungus minimus scutellatus coloris aurantii.
Vaill. paris. t. 13. f. 13. 14.

Habitat in Parietibus putridis.

8. PEZIZA cochleata. P. turbinata cochleata. *Leers. herb.*

P. bracteata, petiolo rugoso, fronde sanguinea, sericea
Hall. helv. n. 2218. *An.*

Elvela sessilis multis anfractibus cochleata. *Dalib. paris.*

Fungoides

CRYPTOGAMIA FUNGUS 737

Fungoides auriculam judæ referens, intus rufescens, extus candicans & qui farinosum. *Vaill. paris. 57. t. 11. f. 8.*

Habitat in umbrosis.

9. PEZIZA *Auricula*. P. concava rugosa auriformis. *burg. Tremella (Auricula) sessilis membranacea auriformis cinerea suec. ged. nass. Blackw. t. 334.*

Peziza bracteata cespitosa plana sericea. Hall. helu. n. 2220.

Elvela (sambucina) sessilis auriformis gelatinosa cinerea Scop. carn.

Elvela sessilis membranacea rugosa & crispa, forma concava Gled. fung.

Agaricum auriculæ forma. Mich. gen. 124. t. 66. f. 1.

Fungus membranaceus, auriculam judæ referens f. sambucinus. Bauh. pin. 372.

Fungorum perniciosum genus 1. Clus. hist. 2. p. 276.

Habitat ad Arbores putridas.

Autoritate Cl. Gleditschii ad fungos arboreos hanc speciem referre cogor, qui antea cum Dillenio pro Lichene habui, Vaill. paris. t. 11. f. 3.

1332. CLAVARIA.

Fungus lævis, oblongus.

* *Indivisa.*

1. CLAVARIA *pistillaris*. C. clavæformis simplicissima *suec. paris. carn. pal. Schæff. fung. t. 169. burg. lithu.*

C. indivisa flavescens plicata. Hall. helu. n. 2204.

C. erecta, corpore simplici, obtuso, integro. Gled. fung. 26.

C. clavata integerrima obtusa erecta. Hort. cliff.

C. alba pistilli forma. Vaill. paris. 39. t. 8. f. 5.

C. major lutea. Mich. gen. 208. t. 87. f. 1.

Fungus clavatus albidus pistillaris. Becc. musc. 1. 307.

Habitat in Sylvis umbrosis.

2. CLAVARIA *militaris*. C. clavata integerrima, capite squamoso. *paris. Fl. dan. t. 657. f. 1.*

C. gemmata. Schæff. fung. t. 290.

C. militaris crocea. Vaill. paris. 39. t. 7. f. 4.

Habitat in sylvis australibus.

3. CLAVARIA *ophioglossoides*. C. clavata integerrima compressa obtusa. *stamp. paris. pal. Schæff. fung. t. 327.*

Linnaei spec. plant. Europ. Pars II.

A a a

738 CRYPTOGAMIA. FUNGI.

C. indivisa nigra compressa. Hall. helu. n. 2206.

C. ophioglossoides nigra. Mich. gen. 208. t. 87. f. 4. Vall. paris. 39. t. 7. f. 3.

Muscus clavatus ceranoides, nigredine splendens. Pluk. t. 47. f. 3.

Habitat in sylvis australibus. ¶

* * *Ramosæ.*

4. *CLAVARIA digitata. C. ramosa lignea nigra stamp. paris. burg. lithu.*

Sphæria nigerrima, aspera, petiolata, conica. Hall. helu. n. 2193.

Sphæria (carcharia) composita caulescens atra aspera, medulla fibrosa. Weig. of. 49. t. 3. f. 6. Oed. dan. t. 405. & 540. sec. Weigel.

Sphæria clavata. Weber. spicil. 287.

Valsa clavata. Scop. carn. ed. 2. n. 1410.

Lichen-Agaricus terrestris, &c. Mich. gen. 104. t. 54. f. 4.

Agaricus digitatus niger. Tournef. inst. 562.

Habitat in sylvis australibus.

5. *CLAVARIA Hypoxylon. C. ramoso-cornuta compressa. succ. gallob. nass. pal. Schæff. fung. t. 328. burg.*

Sphæria nigerrima aspera palmata, cornubus planis, carnosus, pulverulentis. Hall. helu. n. 2194.

Sphæria digitata. Web. spicil. 287.

Valsa digitata. Scop. carn.

Lichenagaricus nigricans ligno innascens, &c. Mich. gen. 104. t. 55. f. 1.

Fungus ramosus niger compressus parvus: apicibus albidis. Rai. angl. 3. p. 15.

6. *Agaricus ramosus cornu rangiferi referens. Mich. Ehret. tab.*

Hypoxylon digitatum. Bruck. monogr.

Fungus cornu dorcadis facie. E. N. C. dec. 1. ann. 4. p. 195.

Habitat in cellis, navibus, aliisque nunquam sole illustratis.

6. *CLAVARIA coralloides. C. ramis confertis ramosissimis inæqualibus. succ. paris. pal. nass. Schæff. fung. t. 175. 177. Barrel. ic. 1260. 1262. 1266. burg. lithu.*

C. caule crassissimo, ramis innumeris, cespitosus, breviter dentatis. Hall. helu. n. 2201.

C. corpore multifido inæquali & ramosissimo, in apices graciles terminato. Gled. fung.

C. ramosissima. Roy. lugd. 518.

Corallo-fungus flavus. Vaill. paris. 41. t. 8. f. 4.

Coralloides flava albida. Tournef. inst. 564. t. 332. f. B.

Barba caprina. Sterb. fung. 96. 97. t. 11. f. A. B. C. D.

Habitat in Sylvis opacis.

7. *CLAVARIA fastigiata. C. ramis confertis ramosissimis fastigiatis obtusis luteis. Fl. suec.*

Fungoides fungiforme luteum foetidum & minus ramosum.

Rai. angl. 3. p. 479. t. 24. f. 5.

Habitat in Sylvis.

8. *CLAVARIA muscoides. C. ramis ramosis acuminatis inæqualibus luteis. suec. Flor. dan. t. 775. f. 3.*

C. caule ramoso, ramis flexuosis acutis. Hall. helv. n. 2199.

C. ramis paucioribus laxioribus longioribus. Reyg. ged. 1. p. 180. sec. Hall. R.

C. corniculata. Schæff. fung. t. 173.

Fungus parvus ramosus luteus. Rai. angl. 3. p. 16. t. 24. f. 7.

Habitat in Muscos.

1333. LYCOPERDON.

Fungus subrotundus, seminibus farinaceis repletus;

** Solida, subterranea absque radice.*

1. *LYCOPERDON Tuber. L. globosum solidum muricatum, radice destitutum. Mat. med. 231. burg.*

L. solidum. Fl. suec. Dalib. paris.

L. subterraneum, conglomeratum, cortice nigro. Hall. helv. n. 2177.

L. globosum, subterraneum, solidum & scabrum, basi & radice carens; capsulis seminalibus magnis. Gled. fung. p. 157. n. 6.

L. gulosorum. Scop. carn. 2. n. 1635.

Tuber brumale, pulpa obscura odorata. Mich. gen. 221. t. 164.

Tuber. Matth. diosc. 544. Mich. gen. 221. t. 102.

Habitat sub Terra.

2. *LYCOPERDON cervinum. L. globosum solidiusculum lacerum centro farinifero, radice destitutum.*

Sphæria subrotunda, tuberculis minimis, medulla atra pulverulenta. Hall. helv. 2191.

Sphæria cervina, subglobosa solitaria terrestris, sphæruis atris medullæ nigræ immersis. Wigg. prim. p. 85.

Lycoperdastum tuberosum arrhizon fulvum, cortice duriora

740 CRYPTOGAMIA. FUNGI.

crasso & granulato, medulla ex albo purpurascens; semine nigro crassiore. *Mich. gen.* 220. t. 99. f. 4.

Tubera cervina. *Bauh. pin.* 376. *Lob. ic.* 276.

Tuberum genus quibusdam Cervi Boletus. *Bauh. hist.* 3. p. 851.

Habitat in Bohemia, Silesia, Vermelandia.

* * *Pulverulenta, radicata supra terram.*

3. LYCOPERDON *Bovista*. L. subrotundum: laceratodeliscens. *Mat. med.* 232. *Fl. suec. paris. pal. nass. carn. Schaff.* t. 292. 295. & *alac. burg. lithu.*

L. leccatum, cinereum, polline obscure viridi. *Hall. helu.* n. 2172.

L. vulgare. *Tournef. inst.* 563.

Fungus maximus rotundus pulverulentus, dictus germanis Bösst. *Bauh. hist.* 3. p. 848.

Habitat in Campis sterilibus.

Permultæ hujus dantur varietates.

4. LYCOPERDON *aurantium*. L. sphaeroidale basi rugosum stipitatum: laciniis obtuse emarginatis dehiscens. *Dalib. paris. burg.*

L. Aurantii coloris, ad basin rugosum. *Vaill. paris.* 123. t. 16. f. 9. 10.

Habitat in Gallia.

Cum præcedente conjungit Hallerus. R.

5. LYCOPERDON *stellatum*. L. volva multifida patente, capitulo glabro: ore acuminato dentato. *suec. paris. pal. nass. Schaff fung.* t. 182. *Flor. dan.* t. 360. *burg. lithu.*

L. cortice exteriori revoluta, stellato. *Hall. helu.* n. 2174.

L. volva multifida, patente coriacea, inflexa; capitulo glabro globoso, sessili; ore acuminato dentato. *Scop. carn.*

L. globosum & sessile, volva radiata, patente; ore acuminato dentato. *Gled. fung.* p. 151. n. 5.

L. stellatum, calyce inverso. *Buxb. cent.* 2. p. 45. t. 49. f. 3.

Geaster major, umbilico fimbriato. *Mich. gen.* 220. t. 100. f. 1. 2. 3.

Fungus pulverulentus, Crepitus lupi dictus, coronatus & interne stellatus. *Rai. angl.* 3. p. 27. t. 1. f. 1.

Fungus stellatus. *Bocc. mus.* 1. t. 305. f. 4.

Habitat in Collibus.

6. LYCOPERDON *Carpobolus*. L. volva multifida, fructu globoso ex seminibus combinatis.

L. corticibus revolutis stellatis, globo projectili. *Hall. helu.* n. 2175.

Carpobolus. Mich. gen. 221. t. 101. n. 2.

Habitat in Italia, Suecia, Helvetia.

Fungus albidus, magnitudine capitis aciculæ, dehiscens in limbum patentem, 5--7 fidum; e disco exsilit vesicula ovalis, altitudine disci totius, & explodit contentum, tamquam vermis casei saltu. Forskâhl.

7. LYCOPERDON radiatum. L. disco hemisphærico, radio colorato.

Habitat in Sueciæ lignis Abietinis putridis ad prædium nostrum Saffa, ubi lectus 1760. a Filio.

Magnitudo dimidii fructus Coriandri. Discus subglobosus, fuscus, elastice dehiscens expellensque cum avolante polline lanam bombycinam, tumescentem, fuscam, rariorem, adhaerentem. Limbus campaniformis, niveus, sectus fere ad basin in partes 12. æquales, disco paulo longiores, basi integra ovali cingente discum.

8. LYCOPERDON pedunculatum. L. stipite longo, capitulo globoso glabro: ore cylindrico integerrimo. Suec. paris. carn. nass. burg. lithu.

L. petiolo gracili, capite sphærico, levi papillato, perforato. Hall. helu. n. 2170.

L. sphærico papillatum, petiolo longissimo. Gled. fung.

L. parisiense minimum pediculo donatum. Tournef. inst. 563. t. 331. f. E. F.

Fungus pulverulentus minimus, pediculo longo insidens. Rai. angl. 3. p. 27.

Habitat in Campestribus.

* * * Parasitica, in farinam satiscens.

10. LYCOPERDON cancellatum. L. parasiticum. apice pustula lateribus fissa. Jacq. austr. t. 12. Flor dan. t. 704.

Habitat in foliis Pyri.

Verruca crocea apice protuberans mammita alba lateraliter dehiscens filis apice connexis. Pulvis fuscus.

11. LYCOPERDON variolosum. L. parasiticum sessile subrotundum, cortice exteriori fulvo deciduo; farina atra compacta. Weig. 824.

L. (sepium) sessile subglobosum, cortice duplici, farina atra compacta. Weig. fl. n. 822.

Sphæria (lycoperdoides) composita convexa subsolitaria, medulla farinosa atra. Weig. obs. 47. n. 10.

Sphæria atra, convexa, carne nigra. Hall. helu. n. 2187.

Lichenoides tuberculosum compressum nigrum lignis putridis adnascens. Dill. musc. 127. t. 18. f. 7.

742 CRYPTOGAMIA. FUNGI.

Lycogala globosum grani pisi magnitudine, aeris recocti colore. *Mich. gen.* 216. t. 15. f. 2.

Habitat passim in Arborum ramis emortuis aut moribundis.

Verrucæ sparsæ; primo fulvæ, molles, succulentæ, ætate consistentiores, exuunt corticem & fuscæ evadunt, induratiæ persistunt, ac dehiscunt, licet polline nigro maxime fardæ.

Synonymon *Michelii* non huc pertinet, sed ad *Mucorem Lycogalam Scopoli* *Weber. l. c. R.*

12. *LYCOPERDON truncatum*. L. paranticum subrotundum truncatum.

Habitat in Fagetis.

Corpus subglobosum, magnitudine a Pisi ad pugni, cortice coriaceo, subgiobosum, supra truncatum & fere marginatum. Pulpa firior spongioso suberosa.

Lycoperdon truncatum est initium *Tremellæ agaricoidis*. *Retz. Ad. Holm.* 1769. p. 250. Huc pertinet *Periza polymorpha* *Oed. dan.* t. 464. fig non bona. & *Hall. helu.* t. 48. f. 8. ic. bona nomen non invenitur. *Retz. obs. fasc.* I. p. 33.

13. *LYCOPERDON pisiiformis*. L. globosum scabrum: ore perforato. *Jacq. miscell. austr. vol.* 1. tab. 7.

Habitat in truncis putridæ fagi. *Jacq. R.*

14. *LYCOPERDON Epidendrum*. L. cortice farinaque purpurea. *suec. pal. Fl. dan.* t. 720.

L. leve, miniatum, sphaericum. *Hall. helu. n.* 2173.

L. sphaericum sessile ore in apice vel integro, vel inæquali & radiato. *Gled. fung.*

L. epidendron, miniatum pulverem fundens. *Buxb. hall.* 203.

L. sanguineum sphaericum. I. *Buxb. cent.* 5. p. 15. t. 29. f. 2.

Habitat in Lignis, parietibus antiquis.

15. *LYCOPERDON epiphyllum*. L. aggregatum parasiticum: ore multifido lacero: pulvere fulvo. *Fl. suec.*

Habitat in dorso foliorum Tussilaginis.

1334. MUCOR.

Fungus vesiculis stipitatis.

* *Perennes,*

1. *MUCOR sphaerocephalus*. M. perennis, stipite filiformi nigro, capitulo globoso cinereo. *suec. pal. cent.*

Trichia petiolata, nigra, capitulo sphaerico, villo ochroleuco. *Hall. hist. helu. n.* 2161. t. 48. f. 3.

Sphærocephalos niger, villo ochroleuco. *Hall. helu.* 3. t. 1. f. 3.

Habitat in Parietibus, lapidibus, Lignis.

2. *MUCOR lichenoides*. M. perennis, stipite subulato nigro capitulo lenticulari cinereo. *succ. * pal.*

Coralloides fungiforme arborum nigrum vix crustosum. Dill. musc. 78. t. 14. f. 3.

Habitat in corticibus Pini.

Cum præcedente conjungit *Weberus spicil.* 198. R.

3. *MUCOR Embolus*. M. seta nigra, villo fusco.

Embolus seta nigra villo fusco. Fl. succ.

Embolus nigerrimus villo albo adpersus. Hall. helu. 8. t. 1. f. 1. *hist. helu. n.* 2137.

Habitat in lignis putridis.

4. *MUCOR fulvus*. M. perennis pallidus, pileo fulvo. *Fl. succ.* 2. n. 1289.

Habitat Upsaliæ *D. Solander.*

5. *MUCOR furfuraceus*. M. perennis viridis, foliis furfuraceis, stipite filiformi, capitulo globoso. *succ.* 1290. *herb.*

Trichia flava, petiolata, capitulo sphærico. *Hall. helu.* n. 2162.

Lichen capitatus. Schreb. spicil. 134.

Lichen sulphureus. Retz. Ad. Holm. 1769. p. 249. *fasc.* 1. *obs. bot. p.* 33.

Habitat in Terra nuda Suecia. *Dan. Rolande.*

* * *Fugaces.*

6. *MUCOR Mucedo*. M. stipitatus, capsula globosa. *gallob. pal. nass. Fl. dan.* t. 467. f. 4. *lithu.*

M. filamentis granulatis fugacibus putrescentibus innatis *succ.*

M. petiolatus cinereus. *Hall. helu. n.* 2147.

M. capitulo fugaci globoso; stipite setaceo, longo. *Scop. carn.* 1. p. 67. n. 5. *ed.* 2. 1644.

M. cespitosus, capitulo sphærico, petiolo setaceo longissimo. *Gled. fung.* 161. n. 4.

M. vulgaris, capitulo lucido per maturitatem nigro, pediculo griseo. *Mich. gen.* 215. t. 95. f. 1.

Mucedo capitulo diaphano, deinde viridi & postremo nigro Malp. vegetat. t. 25. f. 108. p. 2.

Habitat in variis putridis: pane, plantis &c.

7. *MUCOR leprofus*. M. setaceus, seminibus radicalibus. *Aspergillus cespitosus densissimus, initio niveus, deinde aureus, seminibus ovatis. Mich. gen.* 213. t. 91. f. 5.

744 CRYPTOGAMIA. FUNGI.

Habitat in cavernulis autumnis.

8. MUCOR *glaucus*. M. stipitatus, capitulo subrotundo aggregato. Neck. gallob. Flor. dan. t. 777. f. 2.

Byssus ipetiolata, fructificationum lineolis in capitulum radiatum dense congestis. Gled. fung. 24. n. 6.

Aspergillus capitatus, capitulo glauco, seminibus rotundis. Mich. gen. 212 t. 91. f. 1.

Habitat in Citris, Melonibus, Pomis aliisque corruptis.

9. MUCOR *crustaceus*. M. stipitatus, spicis digitatis. Neck. gallob. lithu.

M. crustaceus albus, capitulis setisque aqueis. Fl. suec.

Byssus filamentosa fugax alba, putrescentibus innata Fl. lapp. stamp.

Botrytis non ramosa alba, seminibus rotundis. Mich. gen. 212. t. 91. f. 3.

Habitat in Cibis corruptis.

10. MUCOR *cespitosus*. M. stipite ramoso, spicis ternatis.

Aspergillus albus tenuissimus, graminis dactyloidis facie, seminibus rotundis. Mich. gen. 213. t. 91. f. 3.

6. Aspergillus terrestris capitosus & ramosus albus. Mich. gen. 213. t. 91. f. 4.

Habitat in Putrescentibus.

11. MUCOR *viridescens*. M. viridis granulofus.

Habitat in Europæ; Sueciæ lignis putridis. 24

Crusta viris e Granis subrotundis, inæqualibus, viridibus, humido-gelatinosis.

12. MUCOR *Erisiphe*. M. albus capitulis fuscis sessilibus. suec. lithu.

Habitat in foliis Humuli, Aceris, Lamii, Galeopsidis, Lithospermi.

13. MUCOR *septicus*. M. unctuosus flavus. Fl. suec. Flor. dan. t. 778.

Fuligo butyracea, crocea, cauliculis ramosis, laciniatis. Hall. helu. n. 2133.

Mucor crustaceus, ramosissimus, mollis & fugax, crocei coloris. Gled. fung. 160. var. d.

Spongia fugax, mollis, flava & amœna, in pulvere coriario nascens. Marchand. act. gall. 1727. R.

Spongia fugax mollis flava amœna. March. act. 1727. p. 272.

Habitat in Vaporariis deservescantibus visibili incremento; maturus semina explodens.

*PALMÆ SPATHACEÆ,
TRIPE TALÆ.*

1335. CHAMÆROPS. dioica, *Drupæ* tres.

* * *Pennatifoliæ.*

1339. PHŒNIX, dioica, *Drupa* 1. sperma.

P A L M Æ.

1335. C H A M Æ R O P S.

Hermaphrod. *Cal.* 3. partitus. *Cor.* 3. petala. *Stam.* 6. *Pist.* 3. *Drupæ* 3. monospermæ.

Mascul. diœici, ut in hermaphrodito.

1. CHAMÆROPS *humilis*. C. frondibus palmatis plicatis; stipitibus spinosis.

Palma minor. *Bauh. pin.* 506.

Palma humilis s. Chamæriphe. *Bauh. hist.* 1. p. 368.

Chamæriphe. *Dod. pempt.* 820.

Habitas in Europa australi, præsertim Hispania.

1339. P H Œ N I X.

Masc. *Cal.* 3. partitus. *Cor.* 3. petala. *Stam.* 3.

Fem. *Cal.* 3. partitus. *Cor.* 3. petala. *Pist.* 1.
Drupa ovata.

1. PHŒNIX *daçtylifera*. P. frondibus pinnatis: foliolis complicatis ensiformibus. *med. Kniph. cent.* 2. n. 55. *fem.*

Phoenix frondibus pinnatis; foliolis alternis ensiformibus basi complicatis, stipitibus compressis dorso rotundatis. *Hort.*

P. frondibus pinnatis, foliolis angustioribus, aculeis terminalibus. *Mill. dict.*

Daçtylis palma. *Blackw. t.* 202.

Palma major. *Bauh. pin.* 506.

Palma daçtylifera major vulgaris. *Sloan. jam.* 174.

Palma. *Bauh. hist.* 1. p. 351. *Dod. pem.* 819. *Rai. hist.* 1352.

Palma hortensis mas. *Kæmpf. amæn.* 688. t. 1. 2. f. 1. 2.

Palma hortensis semina. *Kæmpf. exot.* 668, 686. t. 1. f. 2.

16. 11.

Gemeine Dattelpalme. *Lin. pfl. syst.* 1. p. 116.

Habitat in India.

F I N I S.

ADDENDA.

P A R S I.

V Eronica maritima Lin. variat foliis ternis & oppositis, cordato-ovatis, oblongis, lanceolatis. Hinc *V. longifolia* Lin: non discernendam putat. *Wigg. primit. p. 3.*

Ad pæderotam Ageriam Linn. pertinent: *Pæd. Bonarota Jacq. hort. vol. 2 t. 121.* Bonarota *chamædrifolia Scop. carn. ed. 2. n. 26.* Veronica *Scop. carn. ed. 1. p. 303. n. 4.* Bonarota montana italica, spica lutea. *Mich. gen. 19.*

Folia opposita, interdum terna; superiora sæpe alterna. Cor. lutea. bilabiata: lab. sup. integerrimo; inferiore semitrifido. *Jacq.*

Ad Pæderotam Bonarotam Linn. spectent: *Pæd. Bonarota Jacq. austr. vol 5. appr. t. 39.* P. cœrulea. *Scop. ann. 2. p. 41.*

Folia tantum opposita. Cor. cœrulea, bilabiata: lab. sup. æqualiter tridentato; lab. inf. ovato-acuto, plerumque integro. *Jacq.* Ob flores luteos *Chamædrys Montis Sumani J. Bauh.* ad præcedentem rectius refertur.

Gratiola offic. L. Regnault bot. ic.

Valeriana Locusta, olitoria. Flor. dan. t. 738.

Crocus vernus. Jacq. austr. append. t. 36.

Iris spuria. Flor. dan. t. 734.

Cyperus compressus. Jacq. hort. vol. 3. t. 12.

Scirpus romanus L. Jacq. austr. t. 448.

Phleum schoenoides Lin. Jacq. austr. vol. 5. app. t. 7.

Poa palustris. Flor. dan. t. 750.

— compressa. *Flor. dan. t. 742.*

— nemoralis. *Flor. dan. t. 749.*

Dactylis glomerata. Flor. dan. t. 743.

Lolium perenne. Flor. dan. t. 747.

Triticum repens. Flor. dan. t. 748.

Dipsacus fullonum: var. α: addantur Dips. pilosus Regnault bot. Dips. sylvestris. Jacq. austr. t. 402.

Dipsacus laciniatus. Mill. *illustrat. Jacq. austr.* t. 403. *Hail. helu.* n. 198. var. β .

Scabiosa ficula L. est *Sc. divaricata*. *Jacq. hort.* 1. t. 15. vol. 3. p. 1. Fructus *Scab. ochroleucæ similis*.

—— *ochroleuca*. *Jacq. austr.* t. 439.

Aretia alpina. *Jacq. austr. app.* t. 18.

Primula Auricula. *Jacq. austr.* t. 415.

Menyanthes nymphoides Lin. est *Waldschmidia nymphoides*, corollis ciliatis. *Wigg. premis.* p. 20, qui & novi Generis characterem proponit.

Campanula thyrsoidea. *Jacq. austr.* t. 411.

Phiteuma comosum. *Jacq. austr. app.* t. 50.

Atropa Belladonna. *Flor. dan.* t. 758.

Lycium europæum Lin. Falla est omnino descriptio, huic stirpi a *Linnaeo* adjecta. Hucjvero pertinet descriptio, quam *Linnaeus* in *Mant. alt.* p. 341. ad præcedentem plantam addidit, & quæ ita se habet: Flores e singulis gemmis 2-5, pedunculis propriis. Cor. limbus patens, fauce pallida nigro-striata, intus purpurascens. Stam. longitudine totius corollæ: quinto brevior. Porro huc spectat *Jasminoides*: N. 3. *Duham.* p. 305. 306. & *Lycium fol. oblongo lanceolatis*, &c. *Trew. Ehret.* t. 68. (non vero florum descriptio). Neque *Micheli* stirpes in tab. 105. f. 1. & 2. pictas ad hanc, neque ad præcedentem plantam referas, quæ potius alias indicare videntur species. -- Differt vero *Lycium europæum* a L. *barbaro*. Cal. 5. fido; floribus e singulis gemmis 2. 5. ramis inermibus. Tam raro quidem *Spina* observatur, ut potius pro inermi sit stirps habenda. Hæc ill. *Medicus l. c.* p. 58. seqq.

Rhamnus cubensis L. *Jacq. hort.* 3. t. 49.

Thesium alpinum. *Jacq. austr.* t. 416.

Asclepias Vincetoxicum. *Regn. bot. Floris structuram* delineat. *Jacq. in Miscel. austr.* vol. 1. p. 17. t. 1. f. 5. *Stapeliz variegatæ & hirsutæ flores describit & icone illustrat Jacquin in Miscell. austr.* 1. p. 27. t. 3. & 4.

Selsola falsa. *Jacq. hort.* vol. 3. t. 83.

Gentiana punctata. *Jacq. austr. app.* t. 28.

—— campestris. Var. β . est etiam *Gentiana tenella*.

Fris ad. Soc. Sc. Hassn. vol X. p. 436. t. 2. f. 6.

Vix est varietas *G. campestris* L. hæc enim gerit *caulem* non tetragonum, sed quadri-alatum & *folia* magis oblonga, minime basi soluta. Ceterum figura *Frisii*. aptior est, quam *Oederi fl. dan. t. 318. Retz. fasc. obs. 1. p. 13.*

Astrantia minor. Separat Jacquinus A. minorem Scop. eamque Astr. carniolicam appellat. vid. Jacq. austr. app. t. 10.

Bupieum Odontites. Jacq. hort. vol. 3. t. 91.

Tordylium officinale. Huc pertinet T. apulum Jacq. hort. 1. t. 53. Conf. Jacq. hort. 3. p. 2.

Tordylium nodosum. Jacq. austr. app. t. 24.

Daucus carota. Flor. dan. t. 723. Sabb. hort. IV. t. 74.

Daucus Visnaga. Jacq. hort. vol 3. t. 26.

Athamanta Libanotis. Flor. dan. t. 754.

Ferula nodiflora. Jacq. austr. append. t. 5.

Ligusticum peloponense. Jacq. austr. app. t. 13.

——— peregrinum. *Jacq. hort. vol. 3. t. 18.*

Angelica lucida. Jacq. hort. 3. t. 24.

Sium majus angustifolium. Gerard. emac. p. 256. fig. mala. Retz. l. c.

Sison Amomum. Regn. bot. Jacq. hort. vol. 3. t. 17.

Bubon Galbanum. Regn. bot. Jacq. hort. vol. 3. t. 36.

Oenanthe crocata. Jacq. hort. vol. 3. t. 55.

——— prolifera. *ibid. t. 62.*

Scandix odorata. Regn. bot. Jacq. austr. app. t. 37.

Est *Myrrhis* apud *Wigg. primit. 23.*

Scandix nodosa. Jac. hort. vol. 3. t. 25.

Seseli montanum L. est S. multicaule. Jacq. hort. vol. 2. t. 129 vid. vol. 3. p. 2.

Linum austriacum. Jacq. austr. t. 418.

Allium angulosum. Jacq. austr. 423. A. senescens Jacq. vind. 57.

Lilium chalcedonium. Jacq. austr. app. t. 20.

Fritillaria Meleagris. Jacq. austr. vol. 5. app. t. 32.

Erithronium dens canis. Jacq. l. c. t. 9.

Ornithogalum narbonense Lin. Removeatur O. narbonense Floræ Mæno-Francfurt. & addatur ad O. umbellatum Lin. quod revera est.

Anthericum serotinum. Jacq. austr. app. t. 38 In. Carintia crescit.

Daphne Cneorum. Jacq. austr. t. 424

Moeringia muscosa. Jacq. austr. t. 449.

Saxifraga aspera. Jacq. austr. vol. 5. app. t. 31.

D. etamnus albus. Jacq. austr. 428.

Saxifraga petræa Lin. *Sax. petræa* Gunn. noru. t. 9. f. 1. 3. *Ced. dan.* t. 68 distinctissima planta a *Sax. petræa*. Cf. *Retz.* fasc. 1. obs. p. 18.

Gypsophila repens L. Jacq. austr. t. 407.

———— *paniculata*. Jacq. austr. vol. 5. app. t. 1. In *Hungaria* sponte crescit.

Saponaria ocymoides. Jacq. austr. vol. 5. app. t. 23.

Dianthus virgineus. Jacq. austr. vol. 5. app. t. 15.

Cucubulus catholicus. Jacq. hort. i. t. 159.

———— *pumilio*. Jacq. austr. vol. 5. app. t. 2. sub *Silene*.

Silene paradoxa. Jacq. hort. 3. t. 84.

Stellaria ceratoides Lin. vix differt figura Gunn. t. 6.

f. 2. ab ista, quam depingi curavit Nob. Friis in *Actis Soc. Reg. Scient. Hassn.* vol. 10. t. 3. f. 12. cf. *Retz.* fasc. 1. obs. p. 18.

Arenaria verna. Jacq. austr. 404.

Sedum hispanicum L. in alpidibus Carnioliae & Carinthiae etiam crescit. Jacq. austr. vol. 5. app. t. 47.

Spergula saginoides. L. Deleatur icon *flor. dan.* t. 12, quæ ad *Stellariam bifloram* L. pertinet. Cum *Stellaria biflora* L. b. ab *Haller* hanc confundere videretur, aqua tamen stylis quinque abunde differt. *Retz.* fasc. 1. obs. p. 19.

Euphorbia fegetalis. Jacq. austr. t. 450.

———— *Cyparissias*. Jacq. austr. vol. 5. t. 435.

Sempervivum globiferum. Huc citat b. *Linneus* Jacq. austr. t. 12. in qua vero *Sempervivum hirtum* depingitur. Me quidem juvice huc potius pertius pertinet *S. globiferum* Jacq. austr. app. t. 40. Differt vero *S. globiferum* a *S. hirtum*: Calyce 12. partito; Petalis Germinibusque 12. staminibus 24 Jacq. R.

Sempervivum arachnoideum Jacq. austr. app. t. 42.

Sempervivum hirtum L. Huc referatur *S. hirtum* Jacq. austr. t. 12. Calyx sex foliolis constat; Petala sex sunt, Germinaque sex, rarius septem. Stam. duodecim. Hinc solo harum partium numero a *S. globifero* differt. Delendum quoque synonymon *Halleri*. Ita *Jacquin*. R.

Sempervivum montanum. Jacq. austr. append. t. 41.

Sedum foliis semipatulis hirsutis, petalis novenis sub-

hirsutis. Hall. *helu* n. 951. Caulis villosus, ex viridi tandem totus una cum Fol. ruber. Fol. semper villosa, alias viridia apicibus rubris, alias tota rubra. Petala rosea, hirsuta, 8, 9 & 10, Jacq. Hall. R.

Phyteuma orbicularis. Jacq. austr. t. 437.

Cratægus torminalis, Jacq. austr. t. 443.

Sorbus domestica. Jacq. austr. t. 447.

Potentilla supina. Jacq. austr. t. 406.

— nitida. Jacq. austr. vol. 5. app. t. 25. In *Carinthia* quoque habitat.

Geum reptans. Jacq. austr. vol. 5. app. t. 22.

Aconitum Cammarum Jacq. austr. t. 424.

Thalictrum minus. Jacq. austr. t. 419 Non huc pertinet icon flor. dan. t. 244. monente Jacquino.

— angustifolium. Jacq. hort. vol. 3. t. 43.

— flavum. Thal. nigricans Scop, a me additum distincta species est, cuius iconem dedit Ill. Jacquini austr. 421 R.

Ranunculus Thora. Jacq. austr. t. 442.

— — — hederaceus L. Stamina semper quinque. wigg. primit p. 42.

Ranunculus aquatilis α. est *Ran. heterophyllus*, fol. submersis capillaceis, emerfis reniformi-palmatis. Variatio β & γ est *Ran. aquatilis*, fol. omnibus capillaceis, laciniis divergentibus. Et variatio δ est *Ran. fluviatilis*, fol. omnibus capillaceis, laciniis parallelis longissimis. Wigg. primit. p. 42.

P A R S. I I.

Cardamine hirsuta L. apud nos semper tetradynama est. wigg. primit. p. 48.

Cardamine amara L. Antheræ nigræ, in *C. pratensis* flavæ. Wigg. primit. p. 48.

Arabis pendula. Jacq. hort. 3. t. 34.

Barbarea hirsuta L. *Arabicos* omnino species est, etiam ex caractere Linn. a glandulis desumpto. Wigg. pr. p. 50.

Synapis pyrenaica L. Jacq. hort. vol. 3. t. 97.

Malva Sherardiana. Jacq. hort. 2. t. 142.

Polygala amara. Jacq. hort. t. 405.

— — vulgaris. L. *Polygalam majorem* C. B. separat. ill. Jacquini, eamque *Polygalam majorem* salut. Vid. Jacq. austr. 413.

Genista lusitanica Lin. Gen. caule lignoso aphylo, caule juniore folioso, toto spinoso, spinis decussatis. Medic. in Obs. Soc. æcon Lutr. 177. p. 43.

Ononis arvensis. Zorn. ic. t. 132. Var. spinosa cultura

non mutatur in mitem. *Retz. fasc. 1. obs. bot. p. 23*
 Quare recte utraque varietas, ut distinctæ species,
 separatur. *Wigg. primit. p. 47.*

Ononis rotundifolia. Jacq. austr. app. t. 49.

Hippocrepis comosa. Jacq. austr. t. 431.

Trifolium alpestre. Jacq. austr. t. 433.

Scorzonera angustifolia L. Tragopogon pinifolium his-
panicum Barrel. non huc pertinet, quoniam Caulis in
dense foliosus. Wigg. primit. t. 55.

Carduus leucographus. Jacq. hort. vol. 3. t. 23.

Artemisiæ glacialis Lin. Jacq. austr. vol. 5. app. t.

35. Huc referas *Absinthium Hall. helv. n. 125. & 126.*

— — — rupestris *Lin. Adde ad notulam meam,*
quod Artemisia Genipi sit Art. spicata Jacq. austr.
vol. 5. app. t. 34. ab Art. rupestri Lin longe diversa. R.

Erigeron Gouani. Jacq. hort 3. t. 179.

Lemna gibba. Hujus fructum describit. Wigg. primit.
p. 66. n. 705. Flores describit el. Ehrhart. in Hannov.
Mag. 1779 St. 167.

— — — polyrhiza. Flores hujus etiam descripti sunt
Cf. Wigg. primit. p. 66.

— — — arrhiza *Lin. est primum initium L. polyrhizæ.*
Wigg. l. c. p. 67.

Poterium Sanguisorba L. est Sanguisorba Poterium, spiciis sub-
rotundis, floribus polyandris polygamis. Wigg. primit. p. 14.

Valantia Aparine L. est Galium Valantia, foliis sub-
sens, pedunculis trifidis medio hermaphrodito, fructu
rugoso. Wigg. primit. p. 12.

Valantia Cruciata L. est in Gallium Cruciata, caule hir-
suto, fol. quaternis ovatis. Wigg. primit. p. 12.

Polypodium cristatum L. Cf. Retzius (fasc. 1. obs.
bot. p. 33, alia, quam ego adduxi, synonyma, nempe
Polyp. Fil fem. spinosam Veis. crypt. 316. & Polyp.
Muller fridr. n. 841. t. 2. f. 2. huc trahit.

Phascon acaulon L. var. a. Icon. Oed. dan. t. 249. f.
3. non ad hunc Muscum sed ad Veberam Ehrhartii (vid.
Hannov. Mag. 1779.) seu buzbaumiam foliosam Vel-
beri pertinet Wigg. primit. p. 80.

Marchantia conica L. Ob inflorescentiam a Marchantia
diversissimam proprium genus constituit, Conocephalum
dictum apud Wiggers primit. p. 82.

Ulna pruniformis L. Quam p. o tali pinxit Gunnaro
(norv. 2. f. 7.) est initium Fusi laevi. Cf. Wigg. primit. p. 94.

Peziza cochleata L. Hujus varietas est P. coronaria Jacq.
miscell. austr. p. 140. t. 10.

I N D E X

GENERICA NOMINA Romano Charactere
traduntur : Cursivis Litteris S Y N O N Y M A.

<i>A.</i>		<i>Aizoon.</i>	685
<i>Abies.</i>	1175	<i>Alcea.</i>	905
<i>Abrotanum.</i>	1025	<i>Alcea.</i>	906
<i>Abfinthium.</i>	1025	<i>Alchemilla.</i>	177
<i>Abutilon.</i>	902. 906	<i>Alchimilla.</i>	177
<i>Acacia.</i>	1271	<i>Alisma.</i>	495
<i>Acanthus.</i>	857	<i>Alkekengi.</i>	267
<i>Acer.</i>	1266	<i>Allium.</i>	442
<i>Acetosa.</i>	485	<i>Alnus.</i>	1147
<i>Achillea.</i>	1053	<i>Aloë.</i>	464
<i>Achyranthes.</i>	311	<i>Alopecurus.</i>	84
<i>Aconitum.</i>	737	<i>Alsinastrium.</i>	544
<i>Acorus.</i>	468	<i>Alfina.</i>	411
<i>Acrostichum.</i>	1290	<i>Alfina.</i>	617
<i>Actæa.</i>	700	<i>Alfinella.</i>	188
<i>Adhatoda.</i>	28	<i>Althæa.</i>	904
<i>Adiantum.</i>	1297	<i>Alyssum.</i>	869
<i>Adonis.</i>	756	<i>Amanita.</i>	1325
<i>Adoxa.</i>	543	<i>Amaranthus.</i>	1157
<i>Ægilops.</i>	1256	<i>Amaryllis.</i>	439
<i>Ægilops.</i>	95	<i>Ambrosia.</i>	1153
<i>Ægopodium.</i>	398	<i>Amethystea.</i>	34
<i>Æsculus.</i>	498	<i>Amethystina.</i>	37
<i>Æthusa.</i>	385	<i>Ammi.</i>	365
<i>Agaricus.</i>	1325	<i>Amorpha.</i>	933
<i>Agar.</i>	465	<i>Amygdalus.</i>	674
<i>Agrimonia.</i>	663	<i>Anagallidastrium.</i>	151
<i>Agrimonoides.</i>	663	<i>Anagallis.</i>	220
<i>Agrostema.</i>	635	<i>Anagyris.</i>	552
<i>Agrostis.</i>	86	<i>Ananas.</i>	427
<i>Aira.</i>	87	<i>Anandria.</i>	1032
<i>Ajuga.</i>	763	<i>Anastatica.</i>	862

A

INDEX.

<i>Anchusa.</i>	194	<i>Asarina.</i>	642
<i>Anemone.</i>	393	<i>Afarum.</i>	333
<i>Andropogon.</i>	1251	<i>Asclepias.</i>	457
<i>Androsace.</i>	209	<i>Asparagus.</i>	201
<i>Androsace.</i>	981	<i>Asperugo.</i>	128
<i>Androsace.</i>	994	<i>Asperula.</i>	454
<i>Andryala.</i>	752	<i>Asphodelus.</i>	1195
<i>Anemone.</i>	752	<i>Asplenium.</i>	1014
<i>Anemone.</i>	752	<i>Aster.</i>	964
<i>Anemone-Ranunculus.</i>	394	<i>Astragaloides.</i>	965
<i>Anethum.</i>	377	<i>Astragalus.</i>	357
<i>Angelica.</i>	935	<i>Astrantia.</i>	369
<i>Anoris.</i>	806	<i>Arhamanta.</i>	1023
<i>Anonymos.</i>	1052	<i>Athanasia.</i>	1009
<i>Anthemis.</i>	1052	<i>Attractylis.</i>	713
<i>Anthemis.</i>	455	<i>Atragene.</i>	1260
<i>Anthericum.</i>	808	<i>Atriplex.</i>	266
<i>Antirrhinum.</i>	936	<i>Atropa.</i>	97
<i>Anthyllis.</i>	341	<i>Avena.</i>	974
<i>Anchyilis.</i>	132	<i>Aurantium.</i>	210
<i>Aparine.</i>	946	<i>Auricula Urfi.</i>	1108
<i>Aphaca.</i>	178	<i>Ayenia.</i>	226
<i>Aphanes.</i>	441	<i>Azulea.</i>	576
<i>Aphyllanthus.</i>	397	<i>Azederach.</i>	
<i>Apium.</i>	332		B.
<i>Apocynum.</i>	184	<i>Baccharis.</i>	1029
<i>Aquifolium.</i>	741	<i>Baccharis.</i>	1023
<i>Aquilegia.</i>	842	<i>Ballota.</i>	778
<i>Arabis.</i>	417	<i>Ballote.</i>	778
<i>Aralia.</i>	1280	<i>Balsamina.</i>	1092
<i>Araliastrum.</i>	596	<i>Balsamita.</i>	1024
<i>Arbutus.</i>	1002	<i>Barba Capræ.</i>	630
<i>Arctium.</i>	618	<i>Bartia.</i>	797
<i>Arenaria.</i>	208	<i>Basella.</i>	413
<i>Aretia.</i>	705	<i>Belladonna.</i>	266
<i>Argemone.</i>	101	<i>Bellidiastrum.</i>	1039-1063
<i>Ariarum.</i>	1111	<i>Bellis.</i>	1042
<i>Aristolochia.</i>	675	<i>Bellis-leucanthemum.</i>	476
<i>Armeniacæ.</i>	1038	<i>Berberis.</i>	338
<i>Arnica.</i>	1025	<i>Beta.</i>	776
<i>Artemisia.</i>	1119	<i>Betonica.</i>	1147
<i>Arum.</i>	686	<i>Betula.</i>	1012
<i>Aruncus.</i>	99	<i>Bidens.</i>	
<i>Arundo.</i>			

INDEX.

3

<i>Widens.</i>	1011	Calla.	1121
<i>Bignonia.</i>	817	Callitriche.	13
<i>Biscutella.</i>	872	Caltha.	761
<i>Biserrula.</i>	966	<i>Caltha.</i>	1073
<i>Bistorta.</i>	537	Campanula.	234
<i>Blitum.</i>	14	<i>Camphorata.</i>	176
<i>Boletus.</i>	1326	Camphorosma.	176
<i>Boletus.</i>	1328	Canna.	1
<i>Borago.</i>	200	Cannabis.	1220
<i>Brassica.</i>	884	<i>Cannabina.</i>	1237
<i>Briza.</i>	90	<i>Cannacorus.</i>	1
<i>Bromelia.</i>	427	<i>Capnoides.</i>	920
<i>Bromus.</i>	95	Capparis.	699
<i>Browallia.</i>	834	Capraria.	827
<i>Bryonia.</i>	1194	<i>Caprifolium.</i>	250
<i>Bryum.</i>	1311	Capsicum.	269
<i>Bubon.</i>	380	<i>Caraguana.</i>	428
<i>Bufonia.</i>	180	Cardamine.	876
<i>Buglossum.</i>	194	<i>Cardiaca.</i>	780
<i>Bugula.</i>	763	Cardiospermum.	540
<i>Bulbocastanum.</i>	366	<i>Carduus.</i>	1006
<i>Bulbocodium.</i>	440	Carduus.	1004
<i>Bunias.</i>	887	Carex.	1137
<i>Bunium.</i>	366	<i>Carex.</i>	1137
<i>Buphthalmum.</i>	1052	Carica.	1232
<i>Buphthalmum.</i>	1059	Carlina.	1008
<i>Bupleurum.</i>	358	Carpesium.	1028
<i>Bursa pastoris.</i>	866	Carpinus.	1171
<i>Butomus.</i>	550	Carthamus.	1010
<i>Buxbaumia.</i>	1307	<i>Carvi.</i>	395
<i>Buxus.</i>	1148	Carum.	395
<i>Byssus.</i>	1324	<i>Caryophyllata.</i>	692
		Caryophyllus.	727
		Cassia.	557
		<i>Cassida.</i>	792
		Catananche.	999
		<i>Cataria.</i>	768
		Caucalis.	362
		<i>Cedrus.</i>	1240
		Celosia.	312
		Celsia.	815
		Celtis.	1267
		Cenchrus.	1255

C.

A 2

Centaurea.	1066	Cinara.	1007
<i>Centaurium majus.</i>	1066	Cineraria.	1036
<i>Centaurium minus.</i>	352	Circæa.	25
Centunculus.	151	Cirsium.	1004
Cena.	442	Cistus.	728
Cerastium.	637	Citreum.	974
Cerasus.	675	Citrus.	974
Ceratocarpus.	1125	Clandestina.	801
Ceratonia.	1282	Clathrus.	1329
Ceratophyllum.	1162	Clavaria.	1332
Cercis.	553	Clematis.	754
Cereus.	668	Clematitis.	754
Cerithe.	198	Cleome.	890
<i>Cerinthoides.</i>	196	Clinopodium.	782
<i>Cervi spina.</i>	284	Clypeola.	870
Chærophylum.	388	Cneorum.	52
<i>Chamæbuxus.</i>	921	Cnicus.	1005
<i>Chamæcerasus.</i>	250	Cochlearia.	867
<i>Chamædaphne.</i>	140. 593	Coffea.	247
<i>Chamædryas.</i>	764	Coix.	1135
<i>Chamælea.</i>	52	Colchicum.	492
<i>Chamælinum.</i>	419	Colocasia.	1119
<i>Chamænerion.</i>	507	Colocynthis.	1193
<i>Chamæpithys.</i>	764	Colutea.	954
<i>Chamærhododendros.</i>	592	<i>Coma aurea.</i>	1019
Chara.	1127	Comarum.	694
Cheiranthus.	879	Conferva.	1323
Chelidonium.	703	Conium.	367
Chelone.	806	Convollaria.	459
<i>Chenopodio-morus.</i>	14	Convolvulus.	231
Chenopodium.	337	Conyza.	1030
Chlora.	59	Conyzella.	1031
Chondrilla.	989	Conyzoides.	1028. 1031
<i>Cristophoriana.</i>	700	<i>Corallo-fungus.</i>	1332
Chrysanthemum.	1048	Corchorus.	730
Chrysocoma.	109	Coreopsis.	1062
Chrysosplenium.	607	Coriandrum.	386
Cicer.	949	Coriaria.	1235
Cichorium.	1000	Corindum.	540
Cicuta.	384	Coris.	260
<i>Cicuta.</i>	367	Corispermum.	12
<i>Cicutaria.</i>	376	Cornus.	155
Cimicifaga.	740	<i>Corona imperialis.</i>	444

INDEX:

<i>Corona solis.</i>	106c	Cytinus.	1113
Cotonilla.	956	Cytisus.	951
Corrigiola.	409		
<i>Corrigiola.</i>	313		
Cortusa.	211		
Corylus.	1172	Dactylis.	92
Cotinus.	399	Daphne.	526
Cotula.	105c	Datura.	263
Cotyledon.	628	Daucus.	364
Cracca.	963	Delphinium.	736
Crambe.	889	<i>Dens canis.</i>	447
Crassula.	423	<i>Dens leonis.</i>	991
Cratægus.	678	Dentaria.	875
Crepis.	993	Dianthus.	614
<i>Crepis.</i>	987	Diapensia.	27
Cressa.	341	Dictamnus.	564
Crinum.	438	Digitalis.	86
Crithmum.	371	Dipsacus.	120
<i>Crocodilium.</i>	1c66	Dodecatheon.	213
Crocus.	6	Dolichos.	941
Croton.	1181	<i>Doria.</i>	1c35
Crucianella.	133	Doronicum.	1039
<i>Cruciata.</i>	1258	Draba.	864
Cucubalus.	65	Dracocephalum.	787
Cucumis.	1193	<i>racunculus.</i>	1119
Cucurbita.	1192	Drosera.	421
<i>Cuminoides.</i>	36	Dryas.	693
Cuminum.	381	Drypis.	412
Cunila.	35		
Cupressus.	1177		
Cuscuta.	182		
Cyanus.	1c66	Echinophora.	359
Cyclamen.	214	Echinops.	1084
<i>Cydonia.</i>	682	<i>Echinopus.</i>	1084
Cynanchum.	331	Echium.	203
Cynara.	1007	Elæagnus.	168
<i>Cynocrambe.</i>	1166	Elaterium.	1128
<i>Cynoglossoides.</i>	200	Elatine.	544
Cynoglossum.	195	<i>latine.</i>	808
Cynosurus.	93	Elephantopus.	1081
<i>Cyperoides.</i>	1137	Elymus.	102
Cyperus.	72	Empetrum.	1202
Cypripedium.	1100	<i>Enula.</i>	1037

D.

E.

A 3

INDEX.

7

<i>Gossypium.</i>	90	<i>Hyacinthus.</i>	461
<i>Granadilla.</i>	1110	<i>Hyacinthus stellaris.</i>	452
<i>Gratiola.</i>	30	<i>Hydrocharis.</i>	1231
<i>Grossularia.</i>	301	<i>Hydrocotyle.</i>	355
<i>Guajacum.</i>	36	<i>Hyoscyamus.</i>	264
<i>Gypsophyla.</i>	612	<i>Hyoseris.</i>	995
		<i>Hypecoum.</i>	183
<i>H.</i>		<i>Hypericum.</i>	981
		<i>Hypnum.</i>	1312
<i>Hedera.</i>	304	<i>Hypochoeris.</i>	997
<i>Hedypnois.</i>	995	<i>Hypocistis.</i>	1115
<i>Hedysarum.</i>	961	<i>Hypopytis.</i>	583
<i>Helenium.</i>	1037	<i>Hysopus.</i>	767
<i>Helianthemum.</i>	728		
<i>Helianthus.</i>	1060	<i>I.</i>	
<i>Helichrysum.</i>	1026		
<i>Heliotropium.</i>	191	<i>Jacobaea.</i>	1033
<i>Helleborine.</i>	1097	<i>Jalapa.</i>	239
<i>Helleborus.</i>	760	<i>Jasione.</i>	1090
<i>Helxine.</i>	537	<i>Jasminum.</i>	17
<i>Hemerocallis.</i>	467	<i>Iberis.</i>	868
<i>Hepatica.</i>	1315. 752	<i>Illecebrum.</i>	313
<i>Heracleum.</i>	375	<i>Ilex.</i>	184
<i>Herba paris.</i>	542	<i>Ilex.</i>	1168
<i>Hermodactylus.</i>	65	<i>Impatiens.</i>	1093
<i>Herniaria.</i>	316	<i>Imperatoria.</i>	389
<i>Hesperis.</i>	88	<i>Inula.</i>	1017
<i>Hibiscus.</i>	911	<i>Jonthlaspi.</i>	870
<i>Hieracium.</i>	992	<i>Ipomoea.</i>	232
<i>Hieracioides.</i>	993	<i>Iris.</i>	65
<i>Hippocastanum.</i>	498	<i>Isatis.</i>	888
<i>Hippocrepis.</i>	958	<i>Ischænum.</i>	1254
<i>Hippomane.</i>	1186	<i>Isnardia.</i>	164
<i>Hippophaë.</i>	1210	<i>Isoetes.</i>	1310
<i>Hippuris.</i>	11	<i>Isopyrum.</i>	757
<i>Holcus.</i>	1252	<i>Juglans.</i>	1167
<i>Holosteum.</i>	110	<i>Juncago.</i>	488
<i>Homalocenchrus.</i>	8	<i>Juncus.</i>	471
<i>Hordeum.</i>	104	<i>Jungermannia.</i>	1313
<i>Horminum.</i>	788	<i>Juniperus.</i>	1242
<i>Horminum.</i>	42	<i>Justicia.</i>	28
<i>Hottonia.</i>	216		
<i>Humulus.</i>	1211		

X.

<i>Kali.</i>	339
<i>Ketmia.</i>	911
<i>Knautia.</i>	122
<i>Knavel.</i>	611

L.

<i>Lacryma Job.</i>	1135
<i>Lactuca.</i>	988
<i>Lagoecia.</i>	306
<i>Lagurus.</i>	08
<i>Lamium.</i>	774
<i>Lampfana.</i>	948
<i>Lantana.</i>	824
<i>Lapathum.</i>	485
<i>Lappa.</i>	1002
<i>Lapsana.</i>	998
<i>Larix.</i>	1175
<i>Laferpitium.</i>	374
<i>Lathræa.</i>	801
<i>Lathyrus.</i>	946
<i>Lavatera.</i>	927
<i>Lavandula.</i>	763
<i>Laurocerasus.</i>	675
<i>Laurus.</i>	545
<i>Ledum.</i>	591
<i>Ledum.</i>	593
<i>Lemna.</i>	1130
<i>Lens.</i>	949
<i>Lentibularia.</i>	34
<i>Lenticula.</i>	1130
<i>Leontice.</i>	456
<i>Leontondon.</i>	901
<i>Leontopetalon.</i>	456
<i>Leonurus.</i>	780
<i>Lepidium.</i>	865
<i>Leucanthemum.</i>	1048
<i>Leucoinm.</i>	434
<i>Leucosium.</i>	879

<i>Lichen.</i>	1319
<i>Lichen.</i>	1315
<i>Lichenastrum.</i>	1313
<i>Lichenoides.</i>	1319
<i>Ligusticum.</i>	376
<i>Ligustrum.</i>	18
<i>Lilac.</i>	22
<i>Lilio-asphodelus.</i>	467
<i>Lilio-hyacinthus.</i>	452
<i>Lilio-narcissus.</i>	439
<i>Lilium.</i>	443
<i>Lilium convallium.</i>	459
<i>Limnopenice.</i>	11
<i>Limodorum.</i>	1094
<i>Limodorum.</i>	1098
<i>Limon.</i>	974
<i>Limofella.</i>	837
<i>Linagrostis.</i>	74
<i>Lingua Cervina.</i>	1295
<i>Linaria.</i>	808
<i>Lindernia.</i>	828
<i>Linnæa.</i>	835
<i>Linocarpon.</i>	419
<i>Linum.</i>	419
<i>Lirium.</i>	443
<i>Lithospermum.</i>	193
<i>Littorella.</i>	1145
<i>Lobelia.</i>	1091
<i>Lobelia.</i>	244
<i>Lœflingia.</i>	58
<i>Lolium.</i>	101
<i>Lonchitis.</i>	1294
<i>Lonchitis.</i>	1296
<i>Lonicera.</i>	250
<i>Loranthus.</i>	478
<i>Lotus.</i>	969
<i>Lunaria.</i>	871
<i>Lupinaster.</i>	968
<i>Lupinus.</i>	939
<i>Luppulus.</i>	1221
<i>Luteola.</i>	664
<i>Lychnidea.</i>	229
<i>Lychnis.</i>	636

Lychni Scabiosa.
Lycium.
Lycoperdon.
Lycopersicon.
Lycopodium.
Lycopsis.
Lycopus.
Lyfimachia.
Lythrum.

M.

Majorana.
Malope.
Malphigia.
Malva.
Malva.
Malvaviscus.
Malus.
Mandragora.
Marrubium.
Marfilea.
Martynia.
Matricaria.
Mays.
Medica.
Medicago.
Medium.
Melampyrum.
Melia.
Melianthus.
Melica.
Melilotus.
Melilotus.
Melissa.
Melittis.
Melo.
Melocactus.
Melochia.
Melongena.
Melopepo.
Melotria.
Menispermum.

I N D E X.

122	<i>Mentha.</i>	771
273	<i>Menyanthes.</i>	215
1333	<i>Mercurialis.</i>	1230
268	<i>Mesembryanthemum.</i>	684
1302	<i>Mespilus.</i>	681
202	<i>Meum.</i>	385
36	<i>Micropus.</i>	1080
219	<i>Millefolium.</i>	1053
660	<i>Mimosa.</i>	1271
	<i>Minuartia.</i>	114
	<i>Mirabilis.</i>	259
	<i>Mnium.</i>	1310
784	<i>Mœuringia.</i>	536
908	<i>Moldavica.</i>	787
621	<i>Mollugo.</i>	113
906	<i>Mollucca.</i>	782
905	<i>Molucella.</i>	782
911	<i>Momordica.</i>	1191
682	<i>Monarda.</i>	37
268	<i>Monotropa.</i>	583
779	<i>Montia.</i>	107
1299	<i>Morocarpus.</i>	14
811	<i>Morsus ranæ.</i>	1231
1049	<i>Morus.</i>	1150
1133	<i>Moschatellina.</i>	543
971	<i>Mucilago.</i>	1334
971	<i>Mucor.</i>	1334
234	<i>Muscari.</i>	461
800	<i>Myagrum.</i>	860
576	<i>Myolotis.</i>	192
850	<i>Myosotis.</i>	637
88	<i>Myofuros.</i>	426
1272	<i>Myofurus.</i>	426
968	<i>Myrica.</i>	1213
786	<i>Myriophyllum.</i>	1163
787	<i>Myrtus.</i>	672
1193		
668		
	<i>N.</i>	
804	<i>Naias.</i>	1198
268	<i>Narcisso-leucoium.</i>	434
1192	<i>Narcissus.</i>	436
55	<i>Nardus.</i>	75
1238	<i>Nasturtium.</i>	865

Nepeta.	768	Oxalis.	634
Nerium.	323	Orycoccus.	523
Nicotiana.	265	Orys.	634
Nigella.	742		
Nissolia.	923		
Nolana.	206		
Nymphæa.	709		
Nymphoides.	215		
		P.	
		Padus.	675
		Pæderota.	27
		Pæonia.	732
		Paliurus.	284
		Panax.	1280
Ocimum.	790	Pancratium.	437
Odontitis.	799	Panicum.	82
Oenanthe.	381	Papaver.	704
Oenothera.	505	Patietaria.	1259
Olea.	20	Paris.	542
Omphalodes.	195	Parnassia.	415
Onagra.	505	Paronychia.	313
Onobrychis.	961	Parthenium.	1154
Ononis.	935	Passerina.	530
Onopordum.	1006	Passiflora.	1110
Onosma.	199	Pastinaca.	392
Ophiorrhiza.	222	Pavia.	458
Ophioglossum.	1288	Pedicularis.	804
Ophrys.	1096	Pedicularis species.	798
Opulus.	402	Peganum.	656
Opuntia.	668	Pelecinus.	566
Orchis.	1094	Peltaria.	871
Oreoselinum.	369	Peplis.	482
Origanum.	784	Pepo.	1192
Ornithogalum.	451	Percepier.	178
Ornithopodium.	957	Periploca.	330
Ornithopus.	957	Persica.	674
Ornus.	1273	Persicaria.	537
Orobanche.	84	Pervinca.	322
Orobus.	945	Petasites.	1032
Ortegia.	57	Potiveria.	494
Orhala.	774	Peucedanum.	370
Oryza.	48	Peziza.	1331
Osmunda.	128	Phaca.	564
Ostrya.	1271	Phalangium.	455
Oxyris.	1203	Phalaris.	80
Othonna.	1076	Phalloboletus.	1328

II

Phallus.	1328	Porrum.	442
Pharnaceum.	410	Portula.	442
Pharus.	4160	Portulaca.	659
Phascum.	1305	Potamogeton.	186
Phaseolus.	940	Potamopithys.	544
Phellandrium.	383	Potentilla.	690
Philadelphus.	669	Poterium.	1167
Phillyrea.	19	Praſium.	795
Phleum.	83	Prenanthes.	990
Phlomis.	781	Primula.	210
Phoenix.	1393	Primula veris.	210
Phyllis.	353	Provençalia.	1121
Phyſalis.	267	Prunella.	733
Phyteuma.	236	Prunus.	675
Phytolacca.	641	Pſeudoacacia.	953
Picris.	986	Pſeudocyperus.	71
Piloſella.	992	Pſeudodictamnus.	779
Pilularia.	1300	Pſoralea.	967
Pimpinella.	396	Pſyllium.	148
Pimpinella.	1167	Ptarmica.	1053
Pinaſtella.	11	Pteris.	1291
Pinguicula.	33	Pulmonaria.	196
Pinus.	1175	Pulſatilla.	752
Piſtacia.	1212	Punica.	673
Piſum.	944	Pyrethrum.	1011
Plantaginella.	837	Pyrola.	598
Plantago.	148	Pyrus.	682
Platanus.	1173		
Plumbago.	227		
Poa.	89		
Polemonium.	233		
Polianthes.	460	Quaſſia.	578
Poliifolia.	593	Quamoclit.	232
Polium.	764	Quercus.	1168
Polycarpon.	112	Queria.	115
Polycnemum.	59	Quinquefolium.	690
Polygala.	921		
Polygonatum.	459		
Polygonum.	537		
Polypodium.	1296	Radicula.	877
Polytrichum.	1309	Radiola.	419
Populago.	761	Ranunculoides.	757
Populus.	1228	Ranunculus.	757

<i>Rapa.</i>	884	<i>Sagina.</i>	188
<i>Raphanus.</i>	886	<i>Sagitta.</i>	1164
<i>Raphanistrum.</i>	886	<i>Sagittaria.</i>	1164
<i>Rapistrum.</i>	889	<i>Salicaria.</i>	669
<i>Rapunculus.</i>	236. 234	<i>Salicornia.</i>	10
<i>Rapuntium.</i>	1091	<i>Salix.</i>	1201
<i>Refeda.</i>	664	<i>Salfola.</i>	339
<i>Rhabarbarum.</i>	549	<i>Salvadora.</i>	171
<i>Rhagadiolus.</i>	998	<i>Salvia.</i>	42
<i>Rhamnoides.</i>	1210	<i>Salvinia.</i>	1299
<i>Rhamnus.</i>	284	<i>Sambucus.</i>	402
<i>Rhaponticum.</i>	1066	<i>Samolus.</i>	238
<i>Rheum.</i>	549	<i>Sanguisorba.</i>	152
<i>Rhinanthus.</i>	798	<i>Sanicula.</i>	356
<i>Rhodiola.</i>	1229	<i>Saponaria.</i>	613
<i>Rhododendrum.</i>	592	<i>Satureia.</i>	765
<i>Rhus.</i>	399	<i>Satyrium.</i>	1095
<i>Ribes.</i>	301	<i>Saururus.</i>	500
<i>Riccia.</i>	1317	<i>Saxifraga.</i>	608
<i>Ricinus.</i>	1184	<i>Scabiosa.</i>	121
<i>Ricotia.</i>	874	<i>Scandix.</i>	387
<i>Rivina.</i>	174	<i>Scheuchzeria.</i>	487
<i>Roridula.</i>	307	<i>Schoenus.</i>	71
<i>Rosa.</i>	687	<i>Scilla.</i>	452
<i>Rosmarinus.</i>	41	<i>Scirpus.</i>	73
<i>Ros solis.</i>	421	<i>Sclarea.</i>	42
<i>Rubeola.</i>	133	<i>Scleranthus.</i>	611
<i>Rubia.</i>	134	<i>Scolymus.</i>	1001
<i>Rubus.</i>	688	<i>Scoparia.</i>	149
<i>Rudbeckia.</i>	1061	<i>Scorodoprasum.</i>	442
<i>Rudbeckia.</i>	253	<i>Scorpioides.</i>	959
<i>Ruellia.</i>	847	<i>Scorpiurus.</i>	959
<i>Rumex.</i>	485	<i>Scorzonera.</i>	985
<i>Ruppia.</i>	187	<i>Scorzoneroides.</i>	985
<i>Ruppia.</i>	1123	<i>Scrophularia.</i>	814
<i>Ruscus.</i>	1246	<i>Scutellaria.</i>	792
<i>Ruta.</i>	565	<i>Secale.</i>	103
<i>Ruta muraria.</i>	1250	<i>Securidaca.</i>	922
		<i>Securidaca.</i>	956
		<i>Sedum.</i>	629
		<i>Sedi species.</i>	667
		<i>Selinum.</i>	368
		<i>Sempervivum.</i>	667
<i>Sabina.</i>	1249		
<i>Saccharum.</i>	79		

S.

INDEX.

<i>Senecio.</i>	1033	<i>Stapelia.</i>	13
<i>Senna.</i>	557	<i>Staphylæa.</i>	334
<i>Serapias.</i>	1097	<i>Statice.</i>	404
<i>Seriola.</i>	996	<i>Stellaria.</i>	418
<i>Serratula.</i>	1003	<i>Stellaria.</i>	617
<i>Sesamoides.</i>	664	<i>Stellera.</i>	13
<i>Sesamum.</i>	845	<i>Stipa.</i>	529
<i>Seseli.</i>	390	<i>Stæchas.</i>	96
<i>Sherardia.</i>	127	<i>Stramonium.</i>	769
<i>Sicyos.</i>	1195	<i>Stratiotes.</i>	263
<i>Sida.</i>	902	<i>Stratiotes.</i>	744
<i>Sideritis.</i>	770	<i>Styrax.</i>	216. 1231
<i>Sigesbeckia.</i>	1057	<i>Suber.</i>	599
<i>Silene.</i>	616	<i>Subularia.</i>	1168
<i>Siliqua.</i>	1282	<i>Succisa.</i>	863
<i>Siliquastrum.</i>	553	<i>Suillus.</i>	121
<i>Sinapi.</i>	885	<i>Symphytum.</i>	1326
<i>Sinapis.</i>	885	<i>Syringa.</i>	197
<i>Sinapistrum.</i>	890	<i>Syringa.</i>	22
<i>Sisarum.</i>	378	<i>Swertia.</i>	669
<i>Sison.</i>	379		351
<i>Sisymbrium.</i>	877		
<i>Sisyrrinchium.</i>	1103		
<i>Sisyrrinchium.</i>	65		
<i>Sium.</i>	375		
<i>Smilax.</i>	1225	<i>Tamarindus.</i>	50
<i>Solanum.</i>	268	<i>Tamariscus.</i>	405
<i>Soldanella.</i>	212	<i>Tamarix.</i>	405
<i>Solidago.</i>	1035	<i>Tamus.</i>	1224
<i>Sonchus.</i>	967	<i>Tamnus.</i>	1224
<i>Sorbus.</i>	67	<i>Tanacetum.</i>	1024
<i>Sorgum.</i>	1252	<i>Taxus.</i>	1241
<i>Sparganium.</i>	1132	<i>Telephium.</i>	48
<i>Spartium.</i>	929	<i>Terebinthus.</i>	1212
<i>Speculum veneris.</i>	234	<i>Tetrahit.</i>	775
<i>Spirgula.</i>	638	<i>Teucrium.</i>	764
<i>Spermacoce.</i>	570	<i>Thalictrum.</i>	75
<i>Sphondylium.</i>	375	<i>Thapsia.</i>	391
<i>Spigelia.</i>	22	<i>Thesium.</i>	415
<i>Spinacia.</i>	1218	<i>Thlapsi.</i>	866
<i>Spiræa.</i>	686	<i>Thlaspidium.</i>	872
<i>Stachys.</i>	77	<i>Thuya.</i>	1176
<i>Stachelina.</i>	1010	<i>Thymbra.</i>	766

T.

<i>Thymelea.</i>	526		
<i>Thymus.</i>	785		
<i>Thysselinum.</i>	368		V.
<i>Tilia.</i>	717		
<i>Tillæa.</i>	189		
<i>Tinus.</i>	547	<i>Vaccinium.</i>	523
<i>Tinus.</i>	400	<i>Valantia.</i>	1258
<i>Tithymaloides.</i>	665 1013	<i>Valeriana.</i>	48
<i>Tithymalus.</i>	665	<i>Valerianella.</i>	48
<i>Tordylium.</i>	361	<i>Vallisneria.</i>	1199
<i>Tormentilla.</i>	691	<i>Velezia.</i>	350
<i>Tozzia.</i>	803	<i>Vella.</i>	861
<i>Trachelium.</i>	237	<i>Veratrum.</i>	1249
<i>Trachelium.</i>	234	<i>Verbascum.</i>	262
<i>Tradescantia.</i>	430	<i>Verbena.</i>	35
<i>Tragacantha.</i>	965	<i>Verbesina.</i>	1058
<i>Tragopogon.</i>	984	<i>Veronica.</i>	26
<i>Tragoselinum.</i>	396	<i>Vesicaria.</i>	540 869
<i>Trapa.</i>	165	<i>Viburnum.</i>	400
<i>Tremella.</i>	1320	<i>Vicia.</i>	947
<i>Trewia.</i>	712	<i>Vinca.</i>	322
<i>Tribuloides.</i>	165	<i>Viola.</i>	1092
<i>Tribulus.</i>	580	<i>Virga aurea.</i>	1035
<i>Trichomanes.</i>	1298	<i>Virga sanguinea.</i>	155
<i>Trichomanes.</i>	1295	<i>Viscago.</i>	616
<i>Trientalis.</i>	496	<i>Viscum.</i>	1209
<i>Trifolium.</i>	968	<i>Vitex.</i>	853
<i>Trifolium.</i>	968	<i>Viticella.</i>	296.754
<i>Triglochin.</i>	488	<i>Vitis.</i>	305
<i>Trigonella.</i>	970	<i>Vitis idæa.</i>	523
<i>Triticum.</i>	105	<i>Ulex.</i>	932
<i>Trollius.</i>	758	<i>Ulmaria.</i>	686
<i>Tropæolum.</i>	502	<i>Ulmus.</i>	345
<i>Tulipa.</i>	448	<i>Ulva.</i>	1322
<i>Tuna.</i>	668	<i>Unifolium.</i>	459
<i>Tunica.</i>	614	<i>Volubilis.</i>	232
<i>Turritis.</i>	883	<i>Urtica.</i>	1149
<i>Tussilago.</i>	1032	<i>Ufnea.</i>	1319
<i>Typha.</i>	1131	<i>Utricularia.</i>	34
		<i>Uvaria.</i>	750
		<i>Urva urfi.</i>	596
		<i>Uvularia.</i>	445
		<i>Vulneraria.</i>	936

INDEX

19

X.

Z.

Xanthium.

1152

Zanichellia.

1124

Xeranthemum.

1027

Zea.

1133

Xylosteum.

250

Zinnia.

1046

Ziziphora.

36

Y.

Ziziphus.

284

Zygophyllum.

577

Yucca.


463

FINIS

SUPPLEMENTUM
PLANTARUM
EUROPÆARUM,

S E U

SPECIES QUAS NOVIS CHARAC-
TERIBUS SPECIFICIS STABI-
LIVIT. CAROLUS LINNÆUS,
Filius.



MONANDRIA.

MONOGYNIA.

Tetra- HIPPURIS foliis quaternis oblongis obtusis. —
phylla. *Habitas* in Suecia, Finlandia circa Aboam.

Distinctissima a H. vulgari foliis semper quaternis
spathulat o-ovatis, obtusis.

vulgaris. HIPPURIS foliis octonis subulatis.

Hippuris vulgaris. *Syst. Veg. ed.* 13. p. 51.

DIANDRIA.

MONOGYNIA.

saxatilis. VERONICA corymbo terminali, foliis oppositis
glabriusculis, pedunculis folia floralia supe-
rantibus.

Veronica saxatilis. *Bauh. hist. v. 3. 2. p. 284. Jacq.*
obs. v. 1. p. 200. Scop. carn. ed. 2. n. 9.

Veronica fruticans. *Clus. hist. v. 1. p. 347.*

Habitas in Alpibus. 24

Caules ramosi, diffusi.

Folia ovata, plana, integerrima.

Calyx obtusus.

Corollæ cærulæ, maximæ.

urticæfo- VERONICA racemis lateralibus, foliis ovato-lan-
lia. ceolatis hirtis argute ferratis, caule erecto.

Veronica urticæfolia. *Jacq. austr. v. 1. p. 37. t. 59.*

Veronica maxima latifolia f. Quercus folio. *Morif.*
hist. v. 2. p. 322. f. 3. t. 23. f. 18.

Chamædrys falsa maxima. *Bauh. hist. v. 3. p. 286.*

Chamædrys spuria major latifolia. *Bauh. pin. p.*
248. Buxb. cent. 1. p. 23. t. 34.

Habitas in Austria. *Jacquin.* 24

Caulis filiformis, erectus, lævis, pedalis & ultra;

Supl. spec.

A 2

DIANDRIA. *Monogynia*:

Folia opposita ; subsessile , ovato-lanceolata , ut
Urticæ folia tenuissime hirta , argute serrata.
Racemi ex summis foliis filiformes , tenues , sub
 florescentia longitudine foliorum & ultra laxi:
 pedicellis capillaceis.
Calyx tetraphyllus.
Corollæ carneæ.

montana. **VERONICA** racemis lateralibus elongatis filiformibus , foliis ovatis petiolatis obtusis grosse
 serratis , caule petiolisque hirtis.

*Veronica montana. Syst. veg. ed. 13. p. 57. Jacq
 austr. v. 2. p. 6. t. 109.*

Habitat in Germania. J. C. D. Schreberg. 24

cærulea. **PÆDEROTA** corollarum labio superiore indiviso
Habitat in Austria.

lutea. **PÆDEROTA** corollarum labio superiore bifido
Habitat in Austria.
Hæc foliis longioribus.

exaltatus. **LYCOPUS** foliis basi pinnatifido-serratis.
*Lycopus foliis in profundas lacinias incisus. Tourne
 inst. p. 191.*

Habitat in Italia. 24.

Caulis humanæ altitudinis , quo a *L. europæo* dif-
 fert , ramosus , angulatus , scaber.

Folia profunde serrata : serraturis infimis profun-
 dioribus , ut fere inferne pinnatifida : lacinis
 subserratis.

Corollæ quadrifidæ , albæ : laciniis rubro-puncta-
 tis : suprema emarginata ; infima latiori.

Stamina duo , distantia.

Antheræ didymæ , rubræ.

DYGINIA.

aculea- **ANTHOXANTHUM** spicis sub globosis involucre
tum. foliaceo brevi mucronato.

Schœnus aculeatus. Syst. veg. ed. 13. p. 81.

TRIANDRIA. Monogynia.

5

Habitat in Hispania , sibiria. H. U
Culmi plures , diffusi , palmares , ramosi , vaginis
 foliorum vestiti.
Folia graminea , lævia , margine scabra : superiora
 sensim latiora.
Spica subglobosa , fere fastigiata , terminalis , flo-
 ribus fere sessilibus in sinu involucris.
Involucrum triphyllum , foliaceum , spica paullo
 longius.
Foliola sessilia , ovata , oblonga , plano ho-
 rizontalia , acuminata in apicem mucro-
 natum , rigidula , arcata.
tertium in sinu flores paucos continet.
reliqua glumæ instar flores in capitulum
 spicatos gerunt.
Gluma flosculi bivalvis , oblonga , convoluta.
Stamina bina , calyce longiora.
Styli duo.

T R I A N D R I A.

. M O N O G Y N I A.

tenuifolia. IRIS corolla imberbi , spatha biflora , folio lon-
 gissimo filiformi , tubo filiformi elongato.
Iris tenuifolia. *Pallas. itin. v. 3. p. 714. t. C. f. 2.*
Habitat in arenosis Dauriæ & ad Volgam. *Pallas. 24*
Corolla pallide cærulea , odore caryophylleo fra-
 grans.
Tubus longus , extra spathas valde tenuis.

dichotoma. IRIS corolla tenuissime barbata , caule tereti
 elongato paniculato , ramis alternis divari-
 catis bi- f. quadrifloris.
Iris dichotoma. *Pallas. itin. v. 3. p. 714. a. A. f. 2.*
Habitat in rupestribus apricis totius Dauriæ. *Pal-
 las. . .*
 Habitu refert Ixiam.
Caulis inferne tectus foliis sex f. septem , alter-
 nantibus , subradicatis , ensiformi-amplex an-

TRIANDRIA. *Monogynia.*

tibus , distichis , brevibus ; superne teres ;
glaberrimus , ramosus , folio brevissimo ad
quamque ramificationem.

Rami patentissimi , elongati , nudi , simplicis-
si , terminati spatha parva , diphylla , &
& floribus binis , tribus s. quatuor.

Flores longe pedunculati , omnium congenerum
minimi , dilute purpurascentes.

Monti. CYPERUS culmo tereti , umbella supradecom-
posita , foliis carina lævibus.

Cyperus ferotinus odoratus , radice longa , pa-
nicula e locustis latis ferrugineis compactis.

Mont. gram. t. 1. f. 2.

Habitat in India , nunc in Italia. 24

Culmus teres , lævis , aphyllus.

Folia longissima , margine aculeolis deorsum se-
cantia , cæterum glabra , etiam carina.

Umbella supradecomposita.

Involucrum hexaphyllum , longissimum , simile
ei in reliquis Cyperis triquetris.

Spiculæ ovatæ , ferrugineæ.

Pannoni- CYPERUS culmo triquetro , spicis sessilibus ;
cus. glumis lateralibus sphacelatis.

Habitat in Austria. *D. Jacquin.*

Altitudo digitalis.

Bæo- SCIRPUS culmo tereti striato nudo basi vagi-
thryon. nato , vagina truncata , spica terminali
pauciflora bivalvi : valvulis ovatis spica bre-
vioribus margine scariosis : exteriori bre-
viori.

Bæotryon. *Ehrh. phytoph. n 31.*

Habitat in Suecia , Germania. *Ehrhart.* 24

DIGYNIA.

asper. BROMUS panicula nutante scabra , spiculis vil-
losis aristatis , foliis scabris.

TRIANDRIA. Digynia.

Bromus ramosus. *Syst. veg. ed. 13. p. 1024*
(non p. 103.

Habitat in Anglia, Germania, Helvetia.

Loflingiana. AVENA panicula contracta, flosculis binis hirsutis: altero pedunculato apice biaristato, arista intermedia longiori.

Avena loslingiana. *Syst. veg. ed. 13. p. 104.*

Habitat in Hispania. Varietas major in Capibonæ spei.

incurvata. ROTTBOLLA spica tereti subulata, gluma calycina subulata appressa bipartita.

Ægilops incurvata. *Syst. veg. ed. 13. p. 762.*

Habitat in Europæ maritimis. ☉

prostratum, TRITICUM calycibus multifloris, spica disticha ovata, flosculis imbricatis aristato-mucronatis.

Secale prostratum. *Pallas. it. v. 1. p. 485. Jacq. hort. v. 3. p. 25. t. 44.*

Habitat in desertis Caspicis, solo aridissimo. *Pallas.*

Matura Spica integra cadit & germinat, tot caulibus quot spiculæ.

Ipsa spica vicem radicis agit.

Spiculæ multifloræ.

TRIGYNIA.

tetraphylum. POLYCARPON caule ramoso prostrato, foliis quaternis.

Polycarpon. Tetraphylon. *Syst. veg. ed. 13. p. III.*

TETRANDRIA.

MONOGYNIA.

aristata. ASPERULA foliis linearibus subcarnosis: inferioribus quaternis, floribus subternis aristatis.



TETRANDRIA, Monogynia?

Habitat in Europa australi.

Caulis erectus.

Flores pallidi, flavescentes, parallele positi & laciniis obtuse aristatis.

Pyrenai-cum. **GALIUM** foliis senis, floribus lateralibus oppositis subsessilibus solitariis.

Galium caulibus sulcatis flaccidis, foliis senis basi tumidis, floribus axillaribus solitariis subsessilibus, fructibus glabris. *Gouan. bot.* 5. t. 1. f. 4.

Habitat in Pyrenæis.

Caules debiles, digitales, ramosi: internodiis fere folio brevioribus.

Folia glabra, convexiuscula, mucronata.

pumila. **PLANTAGO** caule ramoso herbaceo, foliis integerrimis carnosiss, ramis lævibus.

Habitat in Europa australi. ☉

maritima. **HYDROPHYLAX.**

Habitat in arena volatili ad littora *Maris* prope Guduluhr. *Konig.* 24

Facies Arenariæ rubræ maritimæ, at major.

Radix simplex, filiformis, sanguinea, longa & carnosâ, dulcis.

Caulis repens, filiformis, glaber, coloratus, articulatus, vaginatus: vaginis obtusis, membranaceis, persistentibus; longissimus.

Folia opposita patentia, petiolata, ovata, acuta integerrima, approximata, carnosâ, nitida, exasperata: callis minimis, albicantibus, pellucidis.

Petoli breves, membranacei, marginati, caulem vaginantes, persistentes; hi evadunt foliis decidentibus vaginæ caulium permanentes.

Flores axillares, subsessiles, plerumque binati, (non oppositi) erecti.

Corolla pallide-cærulea.

Antheræ cæruleæ.

TETRANDRIA. *Monogynia.*

natans. TRAPA nucibus quadricornibus : spinis patentibus.

Trapa natans. *Syst. veg. ed. 13. p. 136.*

PENTANDRIA.

MONOGYNIA.

glutinosa. PRIMULA involucro longitudine florum sessilium , foliis lanceolatis glutinosis.

Auricula ursi montana , angustis & leviter crenatis foliis , floribus purpureis. *Pluk. phyt. 149. f. 6.*

Habitat in summis Alpibus Carinthiæ superioris , Volkartenfibus , Fladnizensibus , Lißtalensibus. *P. Xav. Wulfen.*

Statura P. farinosæ.

Folia ferrata , glabra , viscosa , odorata.

Umbella floribus quinque ad septem , sessilibus.

Involucrum penta ad heptaphyllum , longitudine tubi corollæ.

Calyx obovatus , rugosus.

Corollæ lacinia obcordatæ , violaceæ.

Odor Orchidis conopseæ.

Carpatica. CAMPANULA foliis glabris cordatis ferratis petiolatis , pedunculis elongatis , calyce reflexo , glutinoso , corolla patula.

Campanula carpatica. *Jacq. hort. v. 1. p. 22. t. 57.*

Habitat in montibus Carpaticis. *Jacquin.*

grandiflora. CAMPANULA foliis ternis oblongis ferratis , caule unifloro , flore patulo.

Campanula grandiflora. *Jacq. hort. v. 3. p. 4. t. 2.*

Habitat in Sibiria , Tartaria.

Flos figura C. carpaticæ ; an ejusdem varietas ?

Variat foliis ternis & sparsis , sed nunquam cordatis.

verticil- lata. **CAMPANULA** foliis floribusque verticillatis.
Campanula verticillata. *Pall. it. v. 3. p.*

719. *t. G. * f. 1.*

Habitat in Dauria. 24

Folia quina s. fena, verticillata, lanceolata, ferrata.

Flores verticillati, plures, pedunculati, nutantes.

cæruleum. **TRACHELIUM** ramosum, erectum, foliis ovatis ferratis planis.

Trachelium cæruleum. Syst. veg. ed. 13. p. 177.

Europæus. **EVONYMUS** floribus plerisque quadrifidis, foliis sessilibus.

Evonymus europæus. Syst. veg. ed. 13. p. 198.

leprofus. γ. cortice leproso.

Evonymus verrucosus. Scopol. carn. ed. n.

268. *Jacq. austr. v. 1. p. 30. t. 47.*

Evonymus flore pheniceo. Bauh. hist. v. 1. 2. p. 203.

Evonymus secundus. Clus. hist. 1. p. 55.

Habitat in Carniolia rupibus asperis. *D. Jacquin. H*

Facies & habitus ita *E. europæi*, ut non dubitare liceat, hunc ab illo olim natum, adeoque varietatem constantem potiusquam speciem constituere permittunt limites.

diacan **RIBES** foliis incis, aculeis geminis ad gemmas.

tha. *Ribes diacantha. Pall. it. v. 3. p. 722. t. I. f. 2.*

Grossularia daurica montana, uvæ crispæ folio. Amm. ruth. p. 198.

Habitat in Dauria. H

H. U.

Aculei tantum gemini ad gemmas, ceterum inermis.

Media inter *Grossularia* & *Ribes*ia.

Quotannis in Horto Upsaliensi floruit, sed nondum baccas habuit.

lappacea. **ACHYRANTHES** caule fruticoso diffuso prostrato, spicis interruptis, flosculis lateralibus utrinque fasciculo setarum uncinato.

Achyranthes lappacea. Syst. veg. ed. 13. p. 204.

Differt ab *A. patula* caule prostrato, fere nudo,

PENTANDRIA. *Dygina.* 11

foliis nudis , exquisitè acuminatis ; lappis
f. fasciculis rubris , nec flavis.

DYGINIA.

decumbens. GENTIANA corollis quinquefidis campanulatis , foliis radicalibus lanceolatis longissimis , caule decumbente.

Gentiana corollis quinquefidis campanulatis.
Gmel. sib. v. 4. p. 103. t. 51. f. A. quoad figuram.

Habitat in Sibiria. *H. U.*

Folia radicalia lanceolata , trinervia , tota planta longiora , *caulina* duo f. tres , parce linearia.

Caulis spithamæus , procumbens , filiformis.

Flores subverticillati , sed inferiores ramulis elevati.

Calyx latere superiori dehiscens , apice quadridentato.

Corollæ cæruleæ , campanulatæ , quinquefidæ : dente inter lacinias obtuso.

Primo intuitu similis *G. Pneumonanthæ* , sed caules decumbentes.

Epipactis. ASTRANTIA foliis quinquepartitis obtusis ferratis.

Astrantia Epipactis. *Scopol. carn. ed. 2. n. 303.*

t. 6. Jacq. austr. v. 5. p. 37. t. 11.

Helleborine Saniculæ facie. *Bauh. pin. p. 186.*

Elleborine alpina , *Saniculæ* & *Ellebori nigri* facie. *Lob. ic. p. 664.*

Habitat circa Idriam. *24*

Radix prœnat folium unum & scapum unum.

Folium tripartitum : lobo medio trifido ; lateralibus bifidis ; omnibus obtusis , ferratis.

Scapus uniflorus , nudus.

Involutum pentaphyllum , superne ferratum.

Floræ in capitulo , lutei. *Scopol.*

Spinofum. BUPLEURUM frutescens, ramis paniculæ senilibus nudis spinescens, foliis linearibus.

Bupleurum spinofum. *Gouan. bot. 9. t. 2. f. 3.*

Bupleurum hispanicum fruticosum aculeatum, gramineo folio. *Tourn. inst. p. 310.*

Habitat in Hispania. 24

Caulis humidis, fruticosus, inæqualis : ramis divaricatis.

Folia linearia, acuta, trinervia.

Panicula aphylla, lignosa, dichotoma, spinescens, deflorata in spinas transiens, more Abyssi spinosi.

Umbella trifida.

Umbellulæ quinquefidæ.

Involucra numero radiorum, minuta.

Seguieri. SELINUM involucrio universalis nullo, pistillis divaricatissimis.

Selinum Seguieri. *Syst. veg. ed. 13. p. 230.*

Ligusticum perenne, ferulæ folio, floribus albis.

Segu. ver. 2. p. 41. t. 13.

Habitat in Italia.

Caulis multiplex, quadripedalis, lævis non striatus, nisi in summitate, ubi obsoletissime.

Rami superiores, uti folia, oppositi.

Folia tripinnata, patentia.

partialia divaricata s. patentissima.

Pinnæ lineares, acutæ, sessiles, subdecurrentes, acumine inermi.

Umbella composita : radiis ad quadraginta.

Umbellulæ radiis multis.

Pedunculus communis obsolete striatus.

Involucrum nullum.

Involucella fetacea, multa, flosculis breviora.

Corollæ explicatæ.

Petala alba.

Styli fructus horizontaliter patentes.

Sylvestre. SELINUM radice fusiformi multiplici, stylis erectis, corollulis explicatis.

Selinum sylvestre, *Syst. veg. ed. 13. p. 229.*

Corollula explicatae.

Styli erecti.

Carvifolia. SELINUM caule fulcato acutangulo, involucro universaliter evanido, stylis erectis, petalis conniventibus.

Selinum Carvifolia. *Syst. veg. ed. 13. p. 230.*

Corollula. petalis conniventibus, receptaculum suum occultantibus.

Styli erecti.

odontalgica. CACHRYS foliis radicalibus supradecompositis tomentoso-canis, caule nudo umbellifero, feminibus cortice suberoso laevissimo.

Cachrys odontalgica. *Pall. it. v. 3. p. 720. t. G. f. 1. 2. 3.*

Habitat copiosissime in desertis limosis aridissimis inter Wolgam & Jaicum. *Pallar.*

Radix longissima, saporis acerrimi & aromatici, unde vi sialagoga praestat usum in Odontalgia catarrhali.

repens. SIUM caule repente, foliolis subrotundis dentato-angulatis.

Sium repens. *Jacq. austr. v. 3. p. 34. t. 260.*

Habitat in Europa.

salsum. SISON foliis radicalibus compositis : foliolis subverticillato-fasciculatis, lanceolatis, caule aphylo, ramis umbelliferis dichotomis.

Habitat in Salsis limosis aridissimis circa Wolgam. *Pallas.*

Folia tantum radicalia, caulina nulla, fasciculata, breviora, pinnata.

Pinnæ oppositæ, oblongæ : par primum decompositum ; reliqua vero non.

Foliola pinnis lanceolatis, viridia, margine membranaceo-alba fasciculata.

Caulis constans ex ramulis umbelliferis cum bracteis subulatis ad ramificationes.

ramuli divaricati.

Umbellæ parvæ , erectæ.

Involucra parva , subulata.

Folia vere vigent , & postquam evanuerunt ex-
crescit caulis , & mense augusto florescit ,
quasi aphyllus.

H E X A N D R I A.

M O N O G Y N I A.

biflora. TULIPA floribus erectis planiusculis , caule
diphyлло bi-f. trifloro , foliis lineari - su-
bulatis.

Tulipa biflora. *Pall. it. v. 3. p. 727. t. D. f. 3.*

Habitat ad Wolgam locis desertis , maxime ar-
gillofis , imo saluginosis , cum Tulipa syl-
vestri , frequens. *Pallas.*

Planta magnitudine Ornithogali lutei.

Caulis palmaris , glaberrimus.

Folia bina , alterna , linearia , utrinque atte-
nuata , recurva , canaliculata : inferius
caulem vaginans.

Pedunculi foliis breviores , bini s. terni , quasi
in umbella,

Flores Ornith. lutei aliquoties majores , patentes.

Petala tria exteriora lanceolata , dilute cya-
nea s. virescentia.

interiora alba , basi macula magna , fulva.

Filamenta basi barbata.

Bulbus quotannis post deflorescentiam perpe-
diculariter demittit bulbillum , qui succe-
dit post florescentiam exsucco.

bulbi-ferum. ORNITHOGALUM bulbis axillaribus , caule
poliphyлло unifloro.

Ornithogalum bulbiferum. *Pall. it. v. 2. p.*
736. t. Q.

Habitat in aridis limosis ad Wolgam & Jai-
cum. *Pallas.* 24

Distinctissima planta caule simplici unifloro ,
foliis alternis filiformibus , bulbulo in
axillis omnibus , floribus O. minimi lu-
teis : petalis subulatis.

Circina- ORNITHOGALUM hirsutiae canescens , foliis
tum, linearibus recurvatis canaliculatis : radi-
cali solitario ; caulinis tribus , caule tri-f.
quadrifloro.

Ornithogalum reticulatum. *Pall. it. v. 3. p.*
727. t. D. f. 2.

Habitat in siccis versus Astracaniam. *Pallas.*

Simillimum O. minimo , at distinctissimum.

Vagina laxa , fibrosa , reticulata , ferruginea ,
folium & scapum quasi cucullo circumdans.

Folium radicale unicum , gracile , recurvatum ,
lineare , canaliculatum , scapo & cyma
longius.

Folia caulina tria f. quatuor , tot ac flores qui-
bus adfident , figura radicalis.

Cyma divisione & numero flor. ut in O. minimo.

Flores majores & speciosiores quam in O. luteo.

Petala exteriora valde subulata.

Tota planta , flore excepto , hirsutiae minima
canescente tecta.

Cl. Pallas descripsit nomine O. reticulati , at
misit nomine O: circinati , alias non mu-
tassem. Perinde quale sit nomen , modo
omnibus retineatur , ne iterum confusio
scientiam laedat.

Latifolium. ORNITHOGALUM racemo longissimo foliis
lanceolato-ensiformibus.

Ornithogalum latifolium. *Syst. veg. ed. 13. p.*
271.

Folia radicalia , pedalia , pollicis transversa la-
titudine.

Scapus sesquipedalis , teres , viridis.

Racemus longissimus,

Pedunculi pollicis transversa longitudine , seu

bracteis vix longiores.

Flos utrinque albissimus.

Petala lanceolata, patentia.

Filamenta erecta, lanceolata, petalis dimidio breviora, alba.

Antheræ flavæ.

Germen virescens.

Stylus simplex, albus.

Stigma simplex.

Hyacinthoides. SCILLA racemo longissimo, floribus subverticillatis, pedunculis patentibus.

Scylla Hyacinthoides. *Syst. veg. ed.* 13. p. 272. *Gouan. bot.* 26.

Ornithogalum eriophorum orientale. *Tourn. inst. p.* 381.

Bulbus eriophorus orientalis. *Bauh. pin.* 47.

Bulbus eriophorus. *Clus. hist.* 1. p. 172.

Habitat . . . *Ex Byzantio missa.*

Bulbi valde multiplicandi.

Scapus tripedalis, a medio ad apicem florens, cærulescens.

Pedicelli cærulei, pedicellis tribus s. quatuor proximis subverticillatis.

Stipulæ minimæ, a pedicello parum remotæ.

Corollæ minimæ, violaceo-cærulescentes.

Germen album.

Nisi sæpius bulbili avellantur, vix florebit.
Gouan.

horridus. ASPARAGUS aphyllus, fruticosus, pentagonus, aculeis tetragonis compressis striatis.

Asparagus hispanicus, aculeis crassioribus horridus. *Tourn. inst. p.* 300.

Habitat in Hispania. *D. Jacquin.* 7

Caules lignosi, pentagoni, aphylli.

Aculei alterni, solitarii, remoti, lignosi, robusti, interstitiis longiores s. longitudine digiti, tetragoni, compressi, striati.

Folia vix ulla.

Tenageia

Tenageia. JUNCUS annuus, culmo ramofo paniculato, floribus folitariis feffilibus, petalis ovato-oblongis, capsula fubglobofa.

Tenageia. *Ehrh. phytoph. n.* 63.

Habitat in Germania. *Ehrhart.* ☉

TRYGINIA.

Nemola- RUMEX floribus hermaphroditis: valvuli;
pathum. linearibus obtufis integerrimis graniferis
verticillis remotis, ramis patentibus,
foliis inferioribus cordato-lanceolatis,
superioribus lanceolatis.

Nemolapathum. *Ehrh. phytophyl.*

Habitat in Germania. *Ehrhart.* 24

OCTANDRIA.

MONOGYNIA.

imperf- CHLORA corollis fexfidis.

liata. *Habitat* in extrema Italia. ☉

Caulis herbaceus, fimpliciffimus, erectus, tetragonus, lævis, palmaris: articulis foliis longioribus.

Folia oppofita, feffilia, fubamplexicaulia, ovata, lævia, acuta.

Flos pedunculatus, terminalis, flavus, folio major.

Calyx monophyllus, campanulatus, longitudine corollæ, ultra medium bifidus, patulus, perfiftens: laciniis lanceolatis.

Corolla monopetala hypocrateriformis.

Tubus brevis, patulus.

Limbus longior, fexfidus: laciniis ovalibus.

Filamenta fex, fubulata, tubo impofita, eoque paulo longiora.

Suppl. fpec.

B

Antheræ subrotundæ.

Germen oblōngum.

Styli duo, conglutinati.

Stigmata obtusa.

Genus dubium ignoto etiamnum fructu.

Facies Chloræ, sed differt calyce ad basin
diviso, nec laciniis linearibus; corolla
sexfida; stylis duobus conglutinatis.

dioica. DAPHNE floribus axillaribus geminis, foliis li-
neari-lanceolatis.

Daphne dioica. *Gouan. bot.* 27. t. 17. f. 1.

Thymelæa foliis *chamelææ* minoribus subhir-
sutis. *Bauh. pin. p.* 463.

Sanamunda prima. *Clus. hist.* 88.

Habitat in Pyrenæis locis apricis. $\bar{\eta}$

Frutex pedalis, ramosus, rigens: cortice sube-
roso, cicatricibus exasperato.

Folia conferta, imbricata, glabra, lineari-
lanceolata.

Flores axillares, bini, albo-flavescentes, qua-
drifidi, acuti, tubo ventricoso, dioici.
Gouan.

OBSERVATIO.

Daphnarum cortices vim causticam corroderendi
præbent omnes.

TRIGYNIA.

Sibiricum. POLYGONUM floribus octandris trigynis ra-
cemosis, foliis ovato-lanceolatis undulatis
scabris, stipulis hirtis.

Habitat in Sibiria. 24

H. U.

Simillimum P. divaricato,

D E C A N D R I A.

M O N O G Y N I A.

albus. **DICTAMNUS** foliis pinnatis, caule simplici.
Dictamnus albus. Syst. veg. ed. 13. p. 330.

Chrysan- **RHODODENDRON** foliis oblongis impunc-
thum. tatis supra scabris venosis, corolla
 rotata irregulari, gemma florifera ferru-
 gineo-tomentosa.

Rhododendron Chrysanthum. Pall. it. v. 3. p. 729. t. N. f. 1. 2. Kolpin. monogr. cum figura Pallasii.

Andromeda foliis ovatis utrinque venosis, corollis campanulatis obliquis. *Gmel. sibir. v. 4. p. 121. t. 54.*

Habitat in jugis alpinis frigidissimis Dauriæ, totiusque Sibiricæ. *Pallas. H*

Frutex parvus, diffusus: cortice fusco.

Folia oblonga, utrinque attenuata, venosoreticulata, margine reflexo, subtus pallidiora.

Rami inter folia squamis stipulæformibus ferrugineis tecti, & terminantur pedunculis pluribus, ex axillis superiorum squamarum unifloris.

Flos magnus, luteus, rotatus, nutans, figura *Rh. daurici.*

Adeo similis *R. daurico*, ut verear illius solam varietatem esse, & pro distincta specie vix recepissem, nisi in Medicina innotuisset ut antarthriticum experimentis cum illa institutis a *Cl. D. Kolpin.*

Unedo. **ARBUTUS** caule arboreo, foliis glabris obtuse ferratis, panicula terminali, baccis polyspermis.

Arbutus Unedo. Syst. veg. ed. 13. p. 339.

DYGINIA.

rupestris. DIANTHUS floribus solitariis , squamis calycinis obtusissimis binis , petalis crenatis.
Caryophyllus primus. *Clus. hist.* 1. p. 282. figura tenuis.

Habitat in alpibus Europæ australis. 24

Folia linearia , angustissima , triquetra , canaliculata , rigidula , in cespites connata , margine carinaque scabra.

Caulis filiformis , prostratus , uni-f. biflorus.

Flores remoti , odori.

Calycis squamæ tantum duæ , orbiculatæ , appressæ.

Petala sanguinea , crenata , nec plumosa aut villosa.

Styli duo , exserti , recurvati.

liniflora. ARENARIA caulibus erectis inferne ramosis suffruticosis , foliis subulatis , floribus geminis.

Arenaria liniflora. *Syst. veg. ed.* 13. p. 355.

Habitat in Europa australi. 5

Caules perennantes , breves , plures , ramosi , distorti.

Rami erecti , simplicissimi , teretes , spithamæi.

Folia subulata f. linearia , teretia , acuta , lævia.

Pedunculi terminales , uniflori , subpubescentes , bini , unus post alterum.

Calyx foliolis lanceolatis , bistriatis , subpubescentibus.

Corolla magnitudine Lini , alba.

Petala ovata , obtusa , striata , calyce duplo longiora.

Stamina Pistillaque alba , longitudine calycis.

Differt ab A. grandiflora foliis non plano-gramineis , calycè foliolis non multistriatis , caule multo longiore , basique suffruticoso.

PENTAGYNIA.

alpestris. LYCHNIS petalis quadrifidis coronatis, foliis recurvis.

Lychnis quadrifida. Scop. carn. ed. 2. n. 519.

Lychnis alpina, foliis angustis reflexis, petalis quadripartitis. Segu. ver. 3. p. 186. t. 5. f. 1.

Viscago foliis teneris recurvis, petalis quadrifidis. Hall. hist. n. 918.

Alfina alpina glabra, folio reflexo. Pluk. alm. 23. t. 75. f. 5.

Habitat in Austria. Jacquin. 24

Radix perennis.

Caules spithamæi, erecti, læves.

Folia lineari-lanceolata, lævia.

Flores e panicula dichotoma.

Calyx lævis: dentibus obtusis.

Petala quadrifida, alba, longitudine calycis.

Styli quatuor s. quinque.

Capsula calyce longior.

ICOSANDRIA.

PENTAGYNIA.

nivalis. PYRUS foliis integerrimis, floribus corymbosis.

Pyrus nivalis. Jacq. austr. v. 2. p. 4. t. 107.

Habitat in Austria. 5

Arbor ramulis crassiusculis.

Folia elliptica s. ovalia, petiolata, integerrima, subtus sericea albidaque.

Flores terminales, corymbosi.

Fructus globosus, acerbissimus, sed maturus putrescens dulcissimus. Jacquin.

Salicifolia. PYRUS foliis lineari-lanceolatis canis subtus

albotomentosis, floribus axillaribus solitariis subsessilibus.

Pyrus salicifolia. Pall. it. v. 3. p. 734. t. N. f. 3.

Pyrus sylvestris orientalis, folio oblongo incano. Tourn. coroll. p. 43.

Habitat in arenosis desertorum Sibiriae. H

Arbor primo vere florens.

Fructus habet pyriformes, turbinatos, parvos.

Amelanchier. PALLUS incanis, foliis ovalibus obtusis ferratis subtus tomentosis calvescentibus, stipulis subulatis lateralibus deciduis, racemo simplici paucifloro, petalis sublinearibus.

Mespilus Amelanchier. Syst. veg. ed. 13. p. 388.

Habitat in Helvetiae, Austriae, Galloprovinciae, Germaniae rupibus. Ehrhart. H

Fructus orpyalis, inermis.

Ramuli praeteriti anni epidermide alba soluta recti.

Folia ovalia, obtusa, ferrata, juniora subrus valde tomentosa, adulta calvescentia, vix pollicaria.

Stipulae laterales, petiolo adnatæ, subulatae, deciduae.

Racemus simplex, quadri- s. quinqueflorus, tomentosus.

Perianthii bracteati.

Eracae subulato - filiformes, tomentosae, caducae.

Flores P. Botryopii : interiores exteriores paulo superantes.

Petala linearis - lanceolata, obtusa.

Styli quinque.

Fructus caeruleo - nigrescens, magnitudine baccæ Ribis rubri, quinquelocularis, succulentus, dulcis, sapidus.

Semina decem, in quovis loculamento duo, fusca.

Gemmae ovatae, pubescentes, rufae : squamis interne villosis.

Ulmaria. SPIRÆA foliis interrupte pinnatis : foliolis ovatis biferratis subtus canis, floribus cymosis.

Spiræa Ulmaria. *Syst. veg. ed. 13. p. 393.*

Filipendula. SPIRÆA foliis interrupte pinnatis : foliolis lineari-lanceolatis interrupte ferratis glaberrimis, floribus cymosis.

Spiræa Filipendula. *Syst. veg. ed. 13. p. 393.*

POLYGYNIA.

pumila. ROSA germinibus ovatis, petiolis pedunculisque hispidis, caule superne aculeatissimo.

Rosa pumila. *Clus. hist. 117. Jacqu. austr. v. 2. p. 59. t. 198.*

Rosa sylvestris pumila rubens. *Bauh. pin. 483.*

Habitat in Austria. h

Ramuli aculeis copiosissimis, qui in caulibus decidunt.

Fructus magni, pyriformes.

DIDYNAMIA.

ANGIOSPERMIA.

Gymnan dra. BARTSIA diandra, foliis radicalibus binis petiolatis, caule subdiphylio uniplicato : spica lineari obtusa, verticillis bracteatis coarctatis.

Gymnan dra borealis. *Pall. it. v. 3. p. 710. t. A. f. 1.*

Lagotis glauca. *Gartner. Ad. petrop. v. 14. p. 533. t. 18.*

Veronica foliis inferioribus ovatis crenatis, superioribus rotundis mucronatis, caule spica terminato. *Gmel. fib. v. 3. p. 219. n. 13.*

Gerberia. Steller. manuscr.

Habitat in terris arcticis circa Obum, in alpibus Dauriæ summis, in Kamtschatcæ rupibus calvis frigidissimis, earumque latus boreale, ubi nulla alia planta crescit, amans. 24

Radix subhorizontalis, crassiuscula, sapore dulcifaruo.

Folia radicalia bina, petiolata, crassiuscula, succulenta, ovalia, integra s. ferrata, glaberrima.

Caulis palmaris, teres, erectus, glaber, simplicissimus, diphyllus.

Folia in medio caulis, alterna, ovata, sessilia.

Spica constat ex verticillis coarctatis.

Bractæ ovatæ, cærulescentes.

Calyx coloratus, compressus, tridentatus: laciniis lateralibus emarginatis.

Corolla dilute cærulea.

Labium inferius bi-s. trifidum.

Capsula compressa, apice quadridentata.

Variat magnitudine & figura foliorum a climate & solo; etiam foliis connatis duplicato-ferratis.

Ob solum numerum staminum hanc a Bartsiæ genere, quocum in ceteris convenit, separare nolui, cum tamen facilitas methodi inde non læderetur.

O B S E R V A T I O.

Difficile separantur Genera Bartsiæ & Rhi-
nanthi.

ambigua. DIGITALIS calycinis foliolis lanceolatis, corollis galea emarginata, foliis subtus pubescentibus.

Digitalis ambigua. Syst. veg. ed. 13. p. 470.

Digitalis ochroleuca. Jacq. austr. v. 1. p. 36. t. 57.

TETRADYNAMIA. *Siliquosa.* 25

Digitalis foliis calycinis lanceolatis, galea incisa, faucibus maculosis. *Hall. hist.* n. 331.

Digitalis lutea magno flore. *Bauh. pin. p.* 244. *Riu. mon.* 104. *Bess. cyst. æst. f.* 3.

Habitat in Europæ australis montibus. 24

Similis *D. luteæ*, sed differt corollis majoribus.

TETRADYNAMIA.

SILICULOSA.

alliacea. PELTARIA foliis amplexicaulibus oblongis indivisis.

Peltaria alliacea. *Syst. veg. ed.* 13. p. 494.

SILIQUOSA.

hispida. ARABIS foliis hispidis: radicalibus runcinatis, caulinis lanceolatis.

Arabis hispida. *Syst. veg. ed.* 13. p. 501.

Habitat in Austriæ rupibus. *D. Myging.* 24

Statura *A. thalianæ*, simillima *A. lyratæ*, sed diversa.

Radix perennis.

Caulis palmaris, erectus.

Folia radicalia copiosissima, sublyrata, runcinata, hispida.

Caulina sub ramis solitaria, parva, lanceolata.

Calyx subclausus.

Corolla alba: petalis erectis.

lucida. ARABIS foliis amplexicaulibus lucidis.

Habitat in Pannonia. 24

Caulis palmaris, simplicissimus, teres, lævis.

Folia lucida, integerrima, consistentia.

radicalia obovata, petiolata, obtusa:

26 TETRADYNAMIA. *Siliquosa.*

caulina alterna, sessilia, amplexicaulia;
cordata f. cordato-oblonga.

Corymbus terminalis, racemosus evadens.

Calyx glaber, hians.

Petala erectiuscula, alba, linearia, integra;
calyce duplo longiora, basi angustiora.

Stamina longitudine petalorum.

Stigma obtusum.

Anthera flavæ.

Erucoi- RAPHANUS filiquis ovatis gibbis, rostro
des. longitudine filiquæ.

Rapica sylvestris lutea, bursæ pastoris folio,
italica. *Barr. rar.* 42. t. 1016. mala.

Habitat in Italia. ♂

Radix simplex, vix caule crassior.

Caulis sesquipedalis, scaber, basi purpurascens,
superne ramis simplicibus.

Folia petiolata, runcinata, lævia.

Lacinia acuminatæ, margine anteriore
denticulatæ: terminalis angustior.

Flores racemosi.

Calyx patulus f. dehiscens, nec clausus.

Petala lutea, emarginata.

Stamina longitudine petalorum.

Siliqua ovata, gibba, carnosæ, lævis, non
articulata, bivalvis.

Rostrum conicum, ancipiti-tetragonum,
longitudine filiquæ.

Semina quatuor ad septem utrinque. depressa:
dissepimento membranaceo, albo, inter
semina se infinuante.

DIADELPHIA.

DECANDRIA.

Hispanica. ONONIS pedunculis aristatis subunifloris;
foliis omnibus ternatis canaliculatis recur-
vatis totaliter serratis.

Anonis non spinosa, villosa & viscosa, hispanica. *Barrell. rar. 74. t. 775.*

Habitat in Hispania. \bar{h}

Planta suffruticosa, stricta, viscida, inodora.

Folia omnia ternata, secundum totam longitudinem serrata, crassiuscula, recurva, O. natricis minora, sed crassiora.

Stipulae patentēs, recurvarae.

Pedunculi laterales, arista longitudine petioli, uni-f. biflori.

Flores similes O. natricis, lutei, rubro-striati.

An O. crispae varietas?

canescens. OROBUS caule ramoso, foliis bijugis linearibus, stipulis semisagittatis subulatis.

Orobis angustifolius \mathcal{E} . *Syst. veg. ed. 13. p. 550.*

Orobis caule ramoso, foliis quaterno-pinnatis linearibus, stipulis semisagittatis subulatis.

Ger. prou. 493.

Araci f. Apios leguminosae species Tossano Carolo. *Bauh. hist. v. 2. p. 326.*

Habitat in Gallia. $2\frac{1}{2}$

Similis O. angustifolio, sed caulis ramosus, flores albo-caerulei.

albus. OROBUS foliis bijugis ensiformibus petiolatis, stipulis simplicibus, caule simplici.

Orobis pannonicus. *Jacqu. vind. p. 128. Jacqu. austr. v. 1. p. 25, t. 39.*

Habitat in Austria. *Mygind. 2\frac{1}{2}*

Radices tuberosae, sessiles.

Simillimus O. angustifolio, sed folia petiolata, nec subsessilia; flores albi, nec lutei; racemi absque filo terminali, quod in O. angustifolio.

glaber. CYTISUS foliis ternatis, floribus lateralibus solitariis subsessilibus pendulis.

Cytisus glaber, filiqua angusta. *Bauh. hist.* v.

I. 2. p. 373.

Habitat in Austria. $\bar{\eta}$

Frutex lævis, teres.

Folia petiolata, ternata.

Foliola æqualia, lanceolato - elliptica;
margine albicantia, parva, acuta, subtus
oculis vix manifeste pilosiuscula.

Stipulæ nullæ.

Pedunculi laterales, solitarii, flore breviores.

Calyx oblongus, subventricosus, trifidus.

Corolla lutea, majuscula.

pendu **CYTISUS** floribus subbinatis pedunculatis
linus. lateralibus nutantibus secundis.

Cytisus lusitanicus, medicæ folio, floribus in
foliorum alis. *Tourn. inst.* p. 648.

Genista striata. *Hill. bot.* 13. *app. t.* 13.

Habitat in Lusitania. *Vandelli.* $\bar{\eta}$

Frutex ramis virgatis, patentissimis, divari-
catis, striatis, apice aridis.

Gemmæ canescentes.

Folia ternata, bina sæpius.

Foliola ovata, lævia, subtus pubes-
centia.

Flores axillares, laterales, secundum longi-
tudinem ramorum, duo s. tres simul,
nutantes.

Pedunculi uniflori, gemini, cernui, læves,
floribus breviores.

Calyx lævis.

Corollæ flavæ.

Legumina pilosa, compressa.

Salsula. **PHACA** caulescens, erecta, canescens, foliis
pinnatis, leguminibus pedunculatis glo-
bosis cernuis.

Phaca Salsula. *Pall. it.* v. 3. p. 747. t. B. b.
f. 1, 2.

Habitat in campis falsis circa lacum siccum
Tarei Dauriæ.
Simillima herba & figura leguminum Astragalo
chinenfi.

Messanense. **TRIFOLIUM** legumin. racemosis nudis monospermis arcuato-striatis semiovatis acutis, caule erecto.

Trifolium messanense. Mant. plant. p. 275.

Melilotus messanensis, procumbens, folliculis rugosis, Rai. hist. p. 952.

Habitat in Italia. ☉

H. U.

Caulis erectus, teres.

Folia ternata, obovata, ferrulata.

Racemi axillares, breves, erecti, lutei.

Legumina T. Meliloti officinalis majora, striis arcuatis, parallelis; semiovata, gibba.

tetraphyllus. **LOTUS** leguminibus solitariis, foliolis obcordatis cuneatis, stipulis solitariis.

Habitat in Majorca. 24

Caules plures, decumbentes, palmares, filiformes, herbacei, demum adscendentes.

Folia ternata, petiolata: foliolis obcordatis, cuneatis, sessilibus, lævibus.

Stipulæ solitariae, nec pares, (altera deficiente,) foliaceæ, longitudine foliorum, patentes, lanceolato-oblongæ, obtusæ, quartum foliolum mentientes.

Pedunculus e summitate, uniflorus, internodiis longior.

Corolla lutea: vexillo venis purpureis, extus rufo.

prostrata. **MEDICAGO** diffusa, foliis ternatis cuneiformibus apice dentatis, stipulis setaceis integerrimis, leguminibus cochleatis inermibus.

Medicago prostrata. Jacq. hort. v. 1. p. 39. t. 89.

Medica maritima, saxatilis, minima, perennis,

SYNGENESIA. *Polygamia æqualis*:

foliis angustioribus in apice incis, fructu parvo non spinoso. *Till. pis.* 110.

Habitat . . . *Communicavit Cl. Jacquin.* 24
Nimis magna affinitas cum *M. polymorpha*, sed hæc perennis.

SYNGENESIA.

POLYGAMIA ÆQUALIS,

Kolpinia. LAPSANA calycibus fructus undique patentibus radiis subulatis incurvis, echinatis, foliis caulinis lanceolatis indivisis.

Kolpinia linearis. *Pall. it. v. 3. p. 755. t. L. l. f. 2.*

Habitat in Dauria. *Pallas.* ☉

Planta tota *L. stellatæ*, sed semina incurva, hinc secundum longitudinem aculeis fetaceis echinata; at *L. stellatæ* semina recta & lævia sunt.

australis. CARDUUS foliis decurrentibus runcinatis spinosis, calycibus subsessilibus terminalibus.

Habitat in Europa australi. H. U.

Caulis pedalis, teres, striatus, lanato-villosus.

Folia alterna, remota, sessilia, decurrentia vix ad proximum, oblonga, lateribus subsinuata, subarachnoideo-villosa, laterum venis in spinas albas excurrentibus.

Calyces terminales, sessiles, (laterales solitarii), plures, ovati, parvi, imbricati: squamis lanceolatis, acuminatis spina, erectis, subpatulis.

Corolla flosculis circiter octo, purpureis.

Similis *C. argentato*, sed flores non pedunculati.

Pannoni- CARDUUS foliis semidecurrentibus nudis indivisis ciliatis, flore subsolitario.

SYNGENESIA. *Polygamia superflua.* 21

Cirsium angustifolium laciniatum. *Bauh. pin. p.*
377.

Cirsium pannonicum primum. *Clus. hist. 2. p.*
148. *Bauh. hist. v. 3. 1. p. 46.*

Habitat in Austria.

Planta inermis.

Folia semidecurrentia, præsertim altero latere lævia, ciliata setis mollibus.

Calyx terminalis, magnitudine Serratulæ arvensis, inermis: squamis linea alba notatis.

Corolla pallide purpurascens.

albifrons. CACALIA herbacea, foliis cordatis biserratis acutis subtus tomentosis, stipulis oblongis rotundatis.

Cacalia tomentosa. *Jacq. austr. v. 3. p. 20. t.*
235.

Habitat in Austria. Jacquin.

Singularis & distinctissima foliis binis ad basin petioli, quæ referunt stipulas sessiles, oblongas, subdenticulatas, subtus tomentosas.

SYNGENESIA.

POLYGAMIA SUPERFLUA.

pectinata. ARTEMISIA foliis pinnatis pectinatis glabris sessilibus, floribus axillaribus solitariis sessilibus quadrifloris.

Artemisia pectinata. *Pall. it. v. 3. p. 755. t. H.*
h. f. 2.

Habitat in siccis Dauriæ cum A. palustri Pal. ☉

Planta elegans & fragrantissima.

Caulis erectus, herbaceus, dodrantalis.

Facile cognoscitur foliis simpliciter pinnatis: pinnis, setaceis, parallelis, eoque pec-

tinnatis , rarius pinna bifurca ; floribus in axillis per caulem solitariis.

alpinus. **SENECIO** corollis radiantibus , foliis cordatis lyratis grosse ferratis , petiolis auritis.

Cineraria alpina. *Syst. veg. ed. 13. p. 636.*

Senecio foliis crassis , ferratis , imis petiolatis , subrotundis , caulinis lanceolatis. *Hall. hist. hist. v. 1. p. 23. n. 67.*

Senecio foliis imis petiolatis , cordatis , obiter dentatis , caulinis lingulatis , sessilibus. *Hall. hist. v. 1. p. 23. n. 68.*

Habitat in alpibus Helveticis , Suecia & Germania. *Dick. Osbeck. 45.*

Caulis herbaceus , striato-angulatus f. teres , erectus , simplex , bipedalis & ultra , superne ramosus in paniculam corymbosam.

Folia alterna , petiolata , remota , cordata , suprema ovata , lyrata , grosse ferrata , obtusiuscula subtus incana , venosa.

Petioles longitudine foliorum , in medio dentibus foliaceis binis f. quaternis , ut in *Erysimo Barbarea*.

Corymbus terminalis & axillares solitarii , longius pedunculati.

Pedunculi proprii conspersi squamis minutis , subulatis.

Calyx calyculatus , æqualis , apice sphacelatus.

Calyculus minutus , appressus , apice sphacelatus.

Corolla lutea , S. Jacobææ.

Variat in terra fertili hortorum caule multo ramoso , foliis laciniatis , ac si alia esset planta.

cordifolia. **CINERARIA** panicula pauciflora , foliis caulinis petiolatis cordatis acute ferratis glabris , caule angulato.

Cineraria cordifolia. *Jacqu. austr. v. 2. p. 47. t. 176.*

Habitat in Austria. *Jacquin.*

Satis

Satis differt a *C. Sibirica* floribus paniculatis, nec ramosis; foliis caulinis petiolatis; calyce patulo, & ceteris.

erispa. CINERARIA floribus paniculatis, foliis caulinis spathulato-oblongis ferratis obtusis amplexicaulibus basi dilatato-dentatis.

Habitat in Austria. *Jacquin.*

Folia radicalia magna, cordata, ovato-rotundata, dentata, in petiolum decurrentia.

Petiolus longus, alatus, dentatus.

Caulis erectus, fulcatus.

Folia caulina spathulata, amplexicaulia, dilatationibus ad basin foliorum.

Pedunculi & Calyces villosi.

SYNGENESIA.

MONOGAMIA.

montana. JASIONE.

Jasione montana *Syst. veg. ed. 13. p. 666.*

perennis. *B.* radice perenni.

Habitat . . . *Semina communicavit Thouin. 24*

Radix perennis.

Caulis I. montanæ annuæ altior.

Capitula florum majora.

Variat foliis glaberrimis & hirsutissimis in eadem radice, etiam radice, etiam integris & ferratis.

GYNANDRIA.

DIANDRIA.

Xipho- SERAPIAS bulbis fibrosis, foliis ensiformibus
phyllum. distichis, bracteis minutissimis, floribus erectis: nectarii labio obruto petalis brevior.

CRYPTOGAMIA. *Algæ.*

TETRANDRIA.

- alba.* BETULA foliis acuminatis ferratis.
Betula alba. *Syst. veg. ed. 13. p. 708.*
Dalecar- β . foliis subpalmatis : laciniis dentatis.
lica. Habitat in Dalecarlia. \bar{h}
Varietas valde singularis.
- incana.* BETULA foliis incanis ovatis acutis biser-
ratis, stipulis lanceolatis, amentis spicatis.
Betula Alnus incana. *Spec. plant. ed. 2. p. 1394.*
Alnus. *Flor. lapp. p. 260.*
Arre. Alder. Leipe. Ulra. *Suec.*
Habitat in Sueciæ provinciis elevatioribus are-
nosis. \bar{h}
Distinctissima a B. Alno, tam structura partium
ac usu œconomico, sed de his plura in
Flora Suecica nova.

CRYPTOGAMIA.

ALGÆ.

- Sertula-* JUNGERMANNIA furculis repentibus, rē-
rioides. mulis undique floriferis, foliis capillaceis
æqualibus articulatis.
Habitat circa Upsaliam locis humidis inter Brya
& Hypna, quibus arcte incumbit. *Invenit.*
Ol. Swartz, Botanices Studiosus optimæ
spei.
Surculi flexuosi, longi, arcte Muscis sæpe
rependo inferati, conferti.
Folia capillacea, erecta, pulchre articulata, sed
adeo tenuia, ut absque lente articulo
non videri possit, quæ refert Sertularia
f. pilos flaminum Tradescantiæ.
Ramuli in omnibus axillis floriferi.
Calyces oblongi, sensim dilatati.

Flores simplicissimi, (quod singulate), pedicellati, in umbellam simplicissimam enati.

Chordor-CAREX spica composita : spiculis androgynis
rhiza. approximatis superne masculis, capsulis compressis, radice repente filiformi.

Chordorrhiza. Ehrh. phytoph. n. 77.

Habitat in Suecia. Ehrhart. 24

Heleonaf-CAREX spica composita : spiculis androgynis
tes. approximatis superne femininis, capsulis imbricatis lateribus integerrimis.

Heleonastes. Ehrh. phytoph. n. 28.

Habitat in Suecia. Ehrhart. 24

Leptochys-CAREX spica sexa distincta : masculis unica,
chys. fructibus pedunculatis remota perianthea uniformibus, capsulis remotis apice truncatis.

Leptochys. Ehrh. phytoph. n. 43.

Habitat in Germania. Ehrhart. 24

Dryas-CAREX spica sexa distincta : masculis unica,
fructibus pedunculatis remota perianthea uniformibus, capsulis remotis apice truncatis

Dryas. Ehrh. phytoph. n. 54.

Habitat in Germania. Ehrhart. 24

Agrostis-CAREX spica sexa distincta : masculis pluribus,
agrostis. fructibus pedunculatis remota perianthea uniformibus, capsulis remotis apice truncatis

Agrostis. Ehrh. phytoph. n. 59.

Habitat in Germania. Ehrhart. 24

Lichen Burgeffii. *Syst. veg. ed.*
Lightf. scot. p. 82. t. 26.

Lichen foliaceus, erectiusculus,
 crispus, scutellis planis aut
 crispo. *Burgeff. manusc.*

Habitat in Scotia. D. Burgeff.

ornatus. LICHEN foliaceus, erectiusculus
 crispus, scutellis marginal
 depressis rubris margine cris
Habitat in Europa.

Pulcherrimus, totus viridi-niten
Frondes parvæ & copia scutello
 elevatorium videtur frondibu

Scutella nitida, plana, rubra :
 vato, magno, crispo-ciliato
 peltæ quali in L. faccato appa

nigres- LICHEN foliaceus, gelatinosus,
 cens. lobatus, rugosus, atro-vire
 confertis rufis.

Lichen nigrescens. *Huds. angl. ed.*
 Lichenoides gelatinosum, men
 tenue, nigricans. *Dill. musc*
 19. f. 20.

Habitat in arborum truncis, rupib

F U N G I.

cancel- LYCOPERDON parasiticum fol
 lium. cellatum filis parallelis.

Lycoperdon cancellatum. *Syst. v*
 824, *Jacq. austr. v. 1. p. 13.*

Habitat supra folia Pyri. Jacquin.

Verruca crocea, folio innata, c

Mammilla verrucæ apici insidens
 matura dehiscit in fila par
 connexa.

Pulvis ferrugineus, fuscus.

crivatus. MUCOR perennis stipite filiformi
 tubo obtuso nigro glabro.

Habitat in foliis arborum.

refert puncta in frondibus Filici

POST-SCRIPTUM.

TYPOGRAPHI BENEVOLO LECTORI.

*N*U N c tandem peractō systemate plantarum Europæ , quod tibi omnium specierum vulgariū , utilium aut curiosarum , characteres genericos & específicos exhibet , desiderabis certe fundamenta Botanice immortalis Linnæi , nec-non innumeras observationes æconomicas & medicas quas in variis operibus fecunda manu sparjit ; sed spem nobis dedit indefessus editor quod proximis nundinis hæc omnia nobis collecta tradere poterit.

In secunda hac collectione invenies. 1°. Linnæi Fundamenta Botanica. 2°. Linnæi Bibliothecam botanicam exquisitis censuris Halleri ditatam. 3°. Linnæi systemata botanica , additis novis omnibus quæ ab anno 1740 , fuerant evulgata. 4°. Linnæi Criticam botanicam cum sententiis oppositis Ludwigi , Halleri , Oederi , Adansonii. 5°. Cæteras Linnæi Philosophiæ botanicæ sectiones , perpetuis commentariis ex Amœnitatibus academicis ejusdem Authoris desumptis explicatas.

*In tertia tandem collectione egregius editor conser-
vavit. 1°. Omnes observationes botanicas & descriptivas
quæ sparsæ leguntur in Flora Suecica , & Laponica , et
horto Cliffortiano , Upsaliensi , nec-non in Amœnitatibus*

academicis. 2°. Omnes annotationes practicas aut æconomicas quæ inveniuntur in Materia Medica Linnæi , in ejusdem Flora Suecica , in amœnitatibus academicis , &c.

Sic hac sola collectione cum systemate plantarum Europæ coadunata , minimo pretio obtinere poteris , benevole Lector , quidquid utile aut curiosum proposuit Linnæus in numerosis vix quæ hodie reperiundis operibus.



